Published in final edited form as:

Vis Neurosci. 2014 May ; 31(3): 275–283. doi:10.1017/S0952523814000078.

Visual-somatosensory integration in aging: Does stimulus location really matter?

JEANNETTE R. MAHONEY¹, CUILING WANG^{1,2}, KRISTINA DUMAS³, and ROEE HOLTZER^{1,3}

¹The Department of Neurology, Division of Cognitive & Motor Aging, Albert Einstein College of Medicine, Bronx, New York

²Department of Epidemiology & Population Health, Albert Einstein College of Medicine, Bronx, New York

³Ferkauf Graduate School of Psychology, Albert Einstein College of Medicine, Bronx, New York

Abstract

Individuals are constantly bombarded by sensory stimuli across multiple modalities that must be integrated efficiently. Multisensory integration (MSI) is said to be governed by stimulus properties including space, time, and magnitude. While there is a paucity of research detailing MSI in aging, we have demonstrated that older adults reveal the greatest reaction time (RT) benefit when presented with simultaneous visual-somatosensory (VS) stimuli. To our knowledge, the differential RT benefit of visual and somatosensory stimuli presented within and across spatial hemifields has not been investigated in aging. Eighteen older adults (Mean = 74 years; 11 female), who were determined to be non-demented and without medical or psychiatric conditions that may affect their performance, participated in this study. Participants received eight randomly presented stimulus conditions (four unisensory and four multisensory) and were instructed to make speeded foot-pedal responses as soon as they detected any stimulation, regardless of stimulus type and location of unisensory inputs. Results from a linear mixed effect model, adjusted for speed of processing and other covariates, revealed that RTs to all multisensory pairings were significantly faster than those elicited to averaged constituent unisensory conditions ($p < 0.01$). Similarly, race model violation did not differ based on unisensory spatial location ($p = 0.41$). In summary, older adults demonstrate significant VS multisensory RT effects to stimuli both within and across spatial hemifields.

Keywords

Multisensory integration; Sensory processing; Aging; Spatial rule

Introduction

Multisensory integration (MSI) research investigates how concurrent sensory information from the external world is processed simultaneously in the brain. Primarily, MSI has been explored in pairings of the three “major” senses (i.e., visual, auditory, and somatosensory systems), which are typically referred to as auditory-somatosensory (**AS**), auditory-visual (**AV**), and visual-somatosensory (**VS**) pairs, but tri-modal auditory-visual-somatosensory interactions have also been assessed (Diederich & Colonius, 2004). Using psychophysical, electrophysiological, and neuroimaging procedures, researchers are able to investigate these integrative effects in humans across all ages. Behavioral research focusing on reaction times (RTs) has demonstrated multisensory effects with shorter RTs for multisensory conditions compared to RTs to constituent unisensory conditions (**AV**: Harrington & Peck, 1998; Molholm et al., 2002; **AS**: Murray et al., 2005; **VS**: Pavani et al., 2000). Additionally, electrophysiological results have revealed multisensory effects of short latency that likely reflect early sensory processing across multisensory pairings (Giard & Peronnet, 1999; AV: Fort et al., 2002; AS: Foxe et al., 2002; Molholm et al., 2002; **VS**: Schurmann et al., 2002; Murray et al., 2005; Teder-Sälejärvi et al., 2005).

Much of the current work in the field of MSI has been driven by the seminal neurophysiological work of Stein, Meredith, and colleagues who studied the integration of auditory, visual, and somatosensory evoked responses using single-cell recordings in the superior colliculus (SC) of cats (e.g., Stein et al., 1975; Meredith & Stein, 1986; Meredith et al., 1987; Stein & Meredith, 1993; Stein & Wallace, 1996; Meredith & Stein, 1996; Wallace et al., 1996). The SC located superior to the brainstem and inferior to the thalamus contains seven layers of alternating white and gray matter and a high proportion of multisensory neurons (Stein et al., 1988). The deeper layers of SC contain overlapping spatial maps of the visual, auditory, and somatosensory modalities (Stein & Meredith, 1993). Additionally, the SC plays a direct role in the motor control of orientation behaviors of the eyes, ears, and head toward various sensory stimuli and is therefore considered a specialized structure for stimulus detection and subsequent gaze-orienting.

Results from Stein & Meredith’s work revealed that the neural response elicited from two or more concurrent sensory inputs causes a change in a cell’s responsiveness (i.e., excitation) that is either less than or greater than the sum of the responses to the constituent unisensory stimulus inputs. The researchers revealed that such integration appeared to be highly affected by various stimulus properties and detailed three simple governing principles critical for MSIs to occur in SC (cf., Stein & Meredith, 1993). They posited that integration of multisensory inputs in SC was greatest for inputs presented simultaneously or in close temporal proximity (the “temporal rule”). The second rule, the “inverse-effectiveness rule,” stated that the strength of multisensory responses was inversely related to the magnitude of the constituent unisensory inputs. That is, as the detectability of the constituent unisensory inputs decreased, such that multisensory SC neurons responded poorly to either sensory input in isolation, MSI effects became relatively greater. Finally, and of most relevance to the current experiment, Stein and Meredith concluded that MSI was greatest for stimuli that were presented to the same location in space (the “spatial rule”). The authors claimed that spatially aligned bimodal stimuli would fall within the excitatory receptive fields of a given

multisensory neuron, resulting in an enhancement of the bimodal neuron's response. However, as the spatial disparity of the bimodal stimuli increased, inhibitory mechanisms that could potentially suppress the neural response of either constituent unisensory modality were subsequently activated.

In the ongoing effort by researchers to detail the significance of the aforementioned cortical MSI effects, these principles have provided a framework from which to derive testable hypotheses; however, these principles are not without their limitations. That is, recent studies have revealed that cortical integrations simply do not always adhere to the "rules" set forth by Stein and colleagues (e.g., Murray et al., 2005; Holmes, 2007; Ross et al., 2007). Specifically, in an effort to determine whether multisensory interactions in early cortical regions were truly governed by the "spatial-rule," Murray et al. (2005) presented spatially aligned and misaligned **AS** stimuli to left and right while simultaneously recording evoked-related potentials. Participants were told to respond to all stimuli *via* a foot pedal. Behavioral results revealed statistically significant speeding of RTs to multisensory **AS** conditions regardless of spatial alignment. ERP findings revealed similar **AS** multi-sensory interactions in auditory association areas for spatially aligned and misaligned **AS** pairs at just 50 ms. Similarly and as part of a larger study, Teder-Sälejärvi et al. (2005) reported statistically significant RT facilitation to multisensory **AV** conditions regardless of spatial alignment. Collectively, these results revealed that **AV** and **AS** interactions in humans are not constrained by space using a simple RT task; however, other researchers argue that the spatial rule fails in some cases because behavioral analyses assessing redundant signals are not sensitive to spatial alignment (see Otto et al., 2013).

Investigations concerning MSI have predominantly been limited to young adults, and consequently very little information is known about this phenomenon in aging. Nevertheless, a handful of studies have revealed significantly greater MSI effects for old compared to young adults (Laurienti et al., 2006; Peiffer et al., 2007; Hugenschmidt et al., 2009; Stephen et al., 2010; Mahoney et al., 2011). In fact, results from our recent investigation examining the differential effects of **AV**, **AS**, and **VS** multisensory processing revealed that older adults exhibited the greatest multisensory RT benefit when presented with concurrent **VS** information (Mahoney et al., 2011). Thus, there is evidence to suggest successful MSI in older adults; however, whether such integrative RT effects vary in non-demented older adults using visual and somatosensory inputs presented within and across hemifields has yet to be reported. Attainment of this knowledge could prove useful in the development of specific rehabilitative tools designed to optimize MSI processes in older adults.

Materials and methods

Participants

Eighteen (11 female) old adults recruited from the Central Control of Mobility in Aging (CCMA) study at the Albert Einstein College of Medicine campus in Bronx, NY, participated in the current study. Sixteen participants were right-handed as assessed by the Edinburgh handedness inventory (Oldfield, 1971). Potential participants were identified from a population list of lower Westchester county, NY, and were first contacted with a

letter and then by telephone inviting them to participate. A structured telephone screening interview was administered to potential participants to assess for eligibility. Briefly, eligibility criteria required that participants be 65 years of age and older, reside in lower Westchester county, and speak English. Participants were required to see, hear, and feel all sensory stimulation at appropriate levels (see Sensory screening procedures section). Exclusion criteria included inability to independently ambulate, dementia, significant loss of vision and/or hearing, current or history of neurological or psychiatric disorders, recent or anticipated medical procedures that may affect mobility, and/or receiving hemodialysis. All participants provided written informed consent to the experimental procedures (see also Holtzer et al., 2014), which were approved by the Committee on Clinical Investigation (CCI; the institutional review board of the Albert Einstein College of Medicine).

Cognitive and disease status

All study participants took part in an initial telephone screening session where medical and psychological history was acquired by a research assistant to ensure appropriateness for the CCMA study. Participant's cognitive status was first screened using reliable cut scores from the AD8 Dementia Screening Interview (cutoff score = 2; Galvin et al., 2005; Galvin et al., 2006) and the Memory Impairment Screen (MIS; cutoff score <5; Buschke et al., 1999) and later confirmed using standardized clinical case conference procedures (see Holtzer et al., 2008 for specifics). Tests included in our in-house clinical neuropsychology battery have been validated in previous studies of this aged population (Masur et al., 1994; Sliwinski et al., 1997; Holtzer et al., 2006; Holtzer et al., 2007). Additionally, participants were screened for depression (Geriatric Depression Scale (GDS) – 30 item; Yesavage et al., 1982) and anxiety (Beck Anxiety Inventory (BAI); Beck et al., 1988).

Global disease status summary scores (range 0–10) were obtained from dichotomous rating (presence or absence) of diabetes, chronic heart failure, arthritis, hypertension, depression, stroke, Parkinson's disease, chronic obstructive pulmonary disease, angina, and myocardial infarction (see also Holtzer et al., 2006; Verghese et al., 2007; Holtzer et al., 2008; Mahoney et al., 2010; Mahoney et al., 2011).

Sensory screening procedures

All participants were required to successfully complete a sensory screening exam, where visual, auditory, and somatosensory acuity were formally tested to ensure appropriateness for the study. All participants' visual acuity was measured using a Snellen eye chart (better or equal to 20/70). A computerized tone-emitting otoscope that delivered lateral and bilateral 20, 25, and 40 dB tones at 500, 1000, 2000, and 4000 Hz using E-prime 2.0 software (Psychology Software Tools, Inc., (PST), Pittsburgh, PA) was used to assess hearing loss. In terms of the somatosensory screening, each individual received a pulse of 30 V, which was gradually increased in voltage by 5 V to determine each individual's minimal threshold of detection; this threshold was operationally defined as the minimum level of voltage necessary for each participant to feel equal stimulation in their left and right fingers at a level that was not painful (see Mahoney et al., 2011; Mahoney et al., 2012 for specifics).

Stimuli and task procedures

Participants were seated comfortably in a well-lit room and asked to fixate a fixation point (a black cross measuring 0.5 cm × 0.5 cm) visible on the center of the display while avoiding any unnecessary head movements during the experimental blocks. The viewing distance was set at 57 cm and the stimulus field was 38 × 30.5 deg of visual angle. The background field luminance was 105 cd/m². Participants' arms rested comfortably on a table and were roughly half the distance from the visual display. Their hands were about 57 cm apart, symmetrical about the vertical meridian. RT and accuracy were collected as participants performed a simple RT task, depressing a foot pedal located under their right foot as quickly as possible, in response to all stimuli, regardless of the unisensory spatial location. In total, participants responded to eight different stimulus conditions (four unisensory and four multisensory; see Fig. 1). The unisensory conditions included right- or left-sided visual (**V**) or somatosensory (**S**) stimuli. The **V** stimuli were black asterisks presented for 100 ms on a 17" computer monitor, which were 0.64 cm in diameter, 8.14 cm from the central fixation point. Lateralized unisensory **S** pulses were produced from a constant voltage linear isolated programmable stimulator (Biopac Systems, Inc., www.biopac.com) that delivered 100 ms pulses at a custom voltage (individually adjusted to a comfortable level for each participant) to electrodes placed symmetrically on the left and right index or middle fingers (depending upon pre-existing nerve or skin damage at the time of testing).

The multisensory conditions included simultaneous **VS** stimulation presented within (left or right) or across hemifields (e.g., left somatosensory and right visual stimulation – see Fig. 1). In terms of naming the **VS** conditions, the location of the somatosensory input (left or right) determines the condition type, whereby simultaneous visual inputs are presented to either the same (i.e., within hemifield) or opposite (i.e., across hemifield) side of the somatosensory inputs (see Fig. 1c and section Linear mixed effects model). The spatial separation between the visual and somatosensory inputs was equal to 49 deg within and 65 deg across hemifields. Using E-prime 2.0 software, the eight stimulus conditions were presented in random order with equal frequency over 3 blocks of 96 trials, yielding a total of 36 trials per condition. The inter-stimulus interval (ISI) varied randomly from 1.0 to 3.0 s to avoid anticipatory effects.

Behavioral data analysis

Linear mixed effects model—The multisensory effect of condition and spatial location, and their interactions on RT were assessed using a linear mixed effects model (LMEM), adjusted for gender, education, global disease status, and speed of processing (Salthouse, 1985). All statistical analyses were run using SAS 9.2 (SAS Institute Inc., Cary, NC) statistical software. A random intercept was included in the model to allow the entry point to vary across individuals.

The model consisted of a two level condition (multisensory or unisensory average), a two level spatial location (within or across hemifield), and a two level hemispace (left (L) or right (R) side). Specifically, the four multisensory conditions were: within hemi left (VL_SL); within hemi right (VR_SR); across hemi left (VR_SL); across hemi right (VL_SR); and the four unisensory averaged conditions were: within hemi left [(VL +

SL)/2]; within hemi right [(VR + SR)/2]; across hemi left [(VR + SL)/2]; and across hemi right [(VL + SR)/2] (see also Fig. 1). Inverse transformation for RT ($1/RT \times 1000$) was applied to achieve normality and variance stabilization of differences in speed of processing in old participants. The advantage of the linear mixed effects model (LME) is that the heterogeneity and correlation of repeated measures under different conditions are taken into account (Laird & Ware, 1982).

Test of the race model—Multisensory integrative processing can easily be derived through psychophysical data (e.g., RTs) because when two sources of sensory information (e.g., a visual cue and somatosensory pulse) are presented concurrently, they offer redundant signals that give rise to faster detection responses. This phenomenon is referred to as a redundant signal effect (RSE; Kinchla, 1974). Two very distinct models can be implemented to explain a RSE: race models and co-activation models (Miller, 1982). In race models, when two information sources are presented concurrently (e.g., a multisensory stimulus), the signal from the information source that is processed the fastest is the signal that produces the response (i.e., the “winner” of the race). However, co-activation models are supported when RTs to multisensory stimuli are faster than would be predicted by race models. In the latter case, the RT facilitation is accounted for by interactions that allow signals from redundant information sources to integrate or combine non-linearly. Nonetheless, tests developed to assess race model violations are inherently controversial (Eriksen et al., 1989; Mordkoff & Yantis, 1991) since these models are conservative (Miller, 1986; Gondan et al., 2004) and have inherent limitations like assumptions about independence between unisensory processes (Colonius & Diederich, 2006; see also Mahoney et al., 2011). Similarly, work from Otto and Mamassian (2012) and Otto et al. (2013) also reveals that RT facilitation can be explained by probability summation, providing evidence for multisensory integrative effects even when distinct parallel processes are not integrated.

To circumvent some of these various limitations, a more conservative test of the race model as outlined by Colonius and Diederich (2006) was used in the current study: $R_{XY} = P(RT_{XY} = t) - \min [P(RT_X = t) + P(RT_Y = t), 1]$. For any latency, the race model holds when the cumulative probability (CP) value of the multisensory condition (actual CP) is less than or equal to the sum of the CP values from each of the unisensory stimuli or the predicted CP. The model places an upper limit of 1 on the predicted CP of RT for the constituent unisensory stimuli. When the actual CP value is greater than the predicted CP value, the result of this inequality is a positive value, which is indicative of a violation of the race model.

Specifically, individual RTs were recorded for each trial. RTs were sorted in ascending order by stimulus condition and then averaged. Trials with RT responses that exceeded ± 2 standard deviations from the individual mean of each participant were considered outliers and were excluded. Trials with RTs < 100 ms were also excluded from the analysis as they were not considered to be physiologically plausible responses. For each participant, the RT range within the valid RTs was calculated across the eight stimulus conditions and quantized into twenty bins from the fastest RT (or zero percentile) to the slowest RT (hundredth percentile) in 5% increments (0%, 5%, ..., 95%, 100%). Differences between actual CP distributions $[P(RT_{XY} = t)]$ and predicted CP distributions $[\min [P(RT_X = t) + P(RT_Y = t)]]$

were calculated across each time bin for each multisensory pairing (see also Colonius & Diederich, 2006; Mahoney et al., 2011), where values greater than zero are indicative of race model violation, providing support for multisensory integrative processes.

Results

Demographics

Eighteen older individuals (mean age 74 ± 5.13 years; 11 female) participated in the current experiment. None of the participants met criteria for dementia or mild cognitive impairment using established clinical consensus case-conference procedures (Holtzer et al., 2008). All participants were deemed relatively healthy as determined by their global health status (Holtzer et al., 2006; Verghese et al., 2007; Holtzer et al., 2008; Mahoney et al., 2010). Table 1 delineates other demographic information including mean education level (in years), GDS (Yesavage et al., 1982) score, BAI (Beck et al., 1988) score, and mean value and range of the unisensory probes, and overall RT in milliseconds (ms).

RT and linear mixed effects model results

The current task required participants to quickly press a foot-pedal in response to both unisensory and multisensory stimulation (refer to Stimuli and task procedures section and Fig. 1 for details). In total, participants received 288 randomly presented trials and their RT per trial was recorded in milliseconds (ms) using E-prime 2.0 software. Individual mean values were calculated. Trials that exceeded ± 2 standard deviations from the mean and that were equal to 100 ms were excluded from the analysis to avoid the influence of outliers. The mean percentage of excluded trials ranged from 1.54 to 2.62% across the eight stimulus conditions. The mean RT (with SEM bars) to each of the multisensory conditions is displayed next to the constituent unisensory conditions for convenience in Fig. 2.

Results from the linear mixed effects model (LMEM; see Table 2), which adjusted for speed of processing, gender, education, and global disease status, revealed significant multisensory RT effects (Table 2, #1; $p < 0.001$), with RTs to **VS** multisensory stimuli faster than RTs to unisensory **V** and **S** averages. Further, the effect of spatial location (Table 2, #2; $p = .41$) was not significant; suggesting no reliable difference in RTs to unisensory stimulation presented within or across spatial hemifields. RTs to all four multisensory conditions were significantly faster than RTs to unisensory averaged conditions regardless of spatial location (Table 2, #3; $p < 0.001$). Taken together, participants demonstrated significant **VS** multisensory effects to stimuli presented within and across hemifield, indicating that spatial location of the unisensory stimulus is not a critical stimulus parameter for multisensory RT effects to occur in aging.

Race model results

Results from planned comparisons (paired *t*-tests) confirm that RTs to all **VS** pairs, within and across hemifields, were significantly shorter than any unisensory stimulus condition (see Table 3). This provides evidence for a RSE across all sensory combinations. In regards to testing the race model, CP at each time bin was group-averaged separately for each multisensory condition (actual) to form a distribution that maintained the shape of the

individuals' data and was then compared to the model (predicted) for the multisensory pairings. Differences between actual and predicted CP models of the grouped RT distribution for all four multisensory conditions are depicted in Fig. 3; where positive values represent a violation in the race model.

Discussion

To date, multisensory research in young adults has shown clearly that sensory information is simultaneously gathered in the brain and integrated in a parallel, not serial, manner (Harrington & Peck, 1998; Pavani et al., 2000; Molholm et al., 2002; Murray et al., 2005). While such integration is said to be governed by basic stimulus principles including time, space, and magnitude (Stein et al., 1975; Meredith & Stein, 1986; Meredith et al., 1987; Stein et al., 1988; Stein & Meredith, 1990; Stein & Meredith, 1993; Meredith & Stein, 1996; Wallace et al., 1996), these principles do not always hold true for young human psychophysical and electrophysiological studies (e.g., Murray et al., 2005; Holmes, 2007; Ross et al., 2007). The current available body of research on MSI in aging is quite limited and to our knowledge this is the first study to report significant MSI RT effects in older adults under conditions with varying degrees of spatial separation between unisensory visual and somatosensory constituents.

Results from this experiment demonstrate that older participants were significantly faster at responding to the **VS** multisensory conditions compared to the combined unisensory stimuli. Even after controlling for speed of processing, gender, education, and global disease status, MSI RT effects persisted across all four multisensory conditions, regardless of whether the unisensory inputs were presented within or across hemifields. These results provide evidence for a RSE (Kinchla, 1974) and are consistent with other behavioral multisensory findings in young (Harrington & Peck, 1998; Pavani et al., 2000; Molholm et al., 2002; Murray et al., 2005) and old adults (Laurienti et al., 2006; Peiffer et al., 2007; Mahoney et al., 2011). Further, this significant RSE across all **VS** conditions subsequently violated the race model; a result that is congruent with previous studies (see Molholm et al., 2002; Murray et al., 2005; Mahoney et al., 2011). However, this methodology was not without its limitations, as it did not take into account measures of noise that form as a result of repeated exposure to sensory stimulation (Otto & Mamassian, 2012). Nevertheless, these results do reveal a significant multisensory RT effect for simultaneously presented visual and somatosensory stimulation to different spatial locations, thus indicating that spatial location was not a critical stimulus parameter for successful **VS** multisensory RT effects in aging.

MSI in aging is not fully understood

Older adults demonstrate greater multisensory enhancement compared to younger adults when provided with redundant information across multiple sensory channels (Laurienti et al., 2006; Peiffer et al., 2007; Hugenschmidt et al., 2009; Stephen et al., 2010; Mahoney et al., 2011). Laurienti et al. (2006; Peiffer et al. 2007) have examined **AV** integration in young and old adults and reported shorter RTs to **AV** compared to unisensory conditions across both age groups. Results from our recent investigation (Mahoney et al., 2011) examining the differential effect of **AV**, **AS**, and **VS** multi-sensory processing revealed that older adults

exhibit the greatest multisensory RT benefit when presented with concurrent **VS** information. However, very little is known about the behavioral benefits of MSI in aging (see also Freiherr et al., 2013), and the notion as to whether increased levels of MSI are actually beneficial has yet to be determined, as it could be the case that older adults simply require greater MSI.

In the limited number of multisensory aging experiments, the effect of MSI has been attributed to basic degenerative changes in neuronal architecture during the aging process. However, this speculative interpretation has not been systematically assessed in aging. Given that compensatory models of aging suggest that alternate brain networks are recruited to help older adults compensate for age-related differences (Stern et al., 2005), it could be argued that larger multisensory RT benefits in older adults might be a compensatory process used to overcome age-related physiological declines in unisensory processing. However, clearly further investigations are required to better understand the biological basis of the aging process, as it relates to MSI, including specific contrasts between young and old adults as well as individual variations in MSI processing across the adult lifespan.

Future directions and implications for daily functioning for elders

VS integration is not well understood in aging. We believe that MSI is an integral aspect of functioning and mobility in the real world, given known overlaps in direct cortico-cortico and cortico-thalamic connections involved in successful MSI and motor functions. The finding that older adults demonstrate MSI effects regardless of whether stimuli are presented within or across hemifields has major public health implications. Given the current finding of multisensory RT speeding to **VS** cues regardless of spatial location in older adults, perhaps implementation of multisensory cross-walk stimulators (either **VS** or even **AV**) would be beneficial in senior-living communities. These stimulators will increase public safety, as older adults will undoubtedly demonstrate faster responses (i.e., start crossing the street earlier) to these external cues, regardless of the stimulator's physical location. Although admittedly speculative, these findings could also be implemented as assistive or rehabilitative tools designed to enhance functional independence in older adults.

Conclusions

The present study expands upon previous findings delineating the occurrence of successful **VS** integration in aging (Mahoney et al., 2011) by demonstrating that older adults experience such integrative RT effects regardless of whether constituent stimuli are presented within or across hemifields. The speeded multisensory **VS** responses were indicative of a RSE that subsequently violated the race model, regardless of spatial location. Taken together, our results suggest that spatial location of the unisensory constituents is not critical for the occurrence of **VS** integration in older adults and could prove useful in the development of future MSI rehabilitative tools.

Acknowledgments

Research was supported by funding from the Albert Einstein College of Medicine's Resnick Gerontology Center Pilot Grant awarded to Dr. Mahoney. Additional funding was given to Dr. Holtzer, who is supported by the National Institute on Aging (R01AG036921).

References

- Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: Psychometric properties. *Journal of Consulting & Clinical Psychology*. 1988; 56:893–897. [PubMed: 3204199]
- Buschke H, Kuslansky G, Katz M, Stewart WF, Sliwinski MJ, Eckholdt HM, Lipton RB. Screening for dementia with the memory impairment screen. *Neurology*. 1999; 52:231–238. [PubMed: 9932936]
- Colonius H, Diederich A. The race model inequality: Interpreting a geometric measure of the amount of violation. *Psychological Review*. 2006; 113:148–154. [PubMed: 16478305]
- Diederich A, Colonius H. Bimodal and trimodal multisensory enhancement: Effects of stimulus onset and intensity on reaction time. *Perception and Psychophysics*. 2004; 66:1388–1404. [PubMed: 15813202]
- Eriksen CW, Goettl B, St James JD, Fournier LR. Processing redundant signals: Coactivation, divided attention, or what? *Perception and Psychophysics*. 1989; 45:356–370. [PubMed: 2710636]
- Fort A, Delpuech C, Pernier J, Giard MH. Early auditory-visual interactions in human cortex during nonredundant target identification. *Brain Research Cognitive Brain Research*. 2002; 14:20–30. [PubMed: 12063127]
- Foxe JJ, Wylie GR, Martinez A, Schroeder CE, Javitt DC, Guilfoyle D, Ritter W, Murray MM. Auditory-somatosensory multisensory processing in auditory association cortex: An fMRI study. *Journal of Neurophysiology*. 2002; 88:540–543. [PubMed: 12091578]
- Freiherr J, Lundström JN, Habel U, Reetz K. Multisensory integration mechanisms during aging. *Frontiers in Human Neuroscience*. 2013; 7:863. [PubMed: 24379773]
- Galvin JE, Roe CM, Powlishta KK, Coats MA, Muich SJ, Grant E, Miller JP, Storandt M, Morris JC. The AD8: A brief informant interview to detect dementia. *Neurology*. 2005; 65:559–564. [PubMed: 16116116]
- Galvin JE, Roe CM, Xiong C, Morris JC. Validity and reliability of the AD8 informant interview in dementia. *Neurology*. 2006; 67:1942–1948. [PubMed: 17159098]
- Giard MH, Peronnet F. Auditory-visual integration during multimodal object recognition in humans: A behavioral and electrophysiological study. *Journal of Cognitive Neuroscience*. 1999; 11:473–490. [PubMed: 10511637]
- Gondan M, Lange K, Rosler F, Roder B. The redundant target effect is affected by modality switch costs. *Psychonomic Bulletin and Review*. 2004; 11:307–313. [PubMed: 15260198]
- Harrington LK, Peck CK. Spatial disparity affects visual-auditory interactions in human sensorimotor processing. *Experimental Brain Research*. 1998; 122:247–252. [PubMed: 9776523]
- Holmes NP. The law of inverse effectiveness in neurons and behaviour: Multisensory integration versus normal variability. *Neuropsychologia*. 2007; 45:3340–3345. [PubMed: 17663007]
- Holtzer R, Friedman R, Lipton RB, Katz M, Xue X, Verghese J. The relationship between specific cognitive functions and falls in aging. *Neuropsychology*. 2007; 21:540–548. [PubMed: 17784802]
- Holtzer R, Verghese J, Wang C, Hall CB, Lipton RB. Within-person across-neuropsychological test variability and incident dementia. *The Journal of the American Medical Association*. 2008; 300:823–830.
- Holtzer R, Verghese J, Xue X, Lipton RB. Cognitive processes related to gait velocity: Results from the Einstein Aging study. *Neuropsychology*. 2006; 20:215–223. [PubMed: 16594782]
- Holtzer R, Wang C, Verghese J. Performance variance on walking while talking tasks: Theory, findings, and clinical implications. *Age (Dordrecht)*. 2014; 36:373–81.
- Hugenschmidt CE, Mozolic JL, Laurienti PJ. Suppression of multisensory integration by modality-specific attention in aging. *Neuroreport*. 2009; 20:349–353. [PubMed: 19218871]
- Kinchla R. Detecting target elements in multielement arrays: A confusability model. *Perception and Psychophysics*. 1974; 15:149–158.
- Laird NM, Ware JH. Random-effects models for longitudinal data. *Biometrics*. 1982; 38:963–974. [PubMed: 7168798]
- Laurienti PJ, Burdette JH, Maldjian JA, Wallace MT. Enhanced multisensory integration in older adults. *Neurobiology of Aging*. 2006; 27:1155–1163. [PubMed: 16039016]

- Mahoney JR, Li PC, Oh-Park M, Verghese J, Holtzer R. Multisensory integration across the senses in young and old adults. *Brain Research*. 2011; 1426:43–53. [PubMed: 22024545]
- Mahoney JR, Verghese J, Dumas K, Wang C, Holtzer R. The effect of multisensory cues on attention in aging. *Brain Research*. 2012; 1472:63–73. [PubMed: 22820295]
- Mahoney JR, Verghese J, Goldin Y, Lipton R, Holtzer R. Alerting, orienting, and executive attention in older adults. *Journal of the International Neuropsychological Society*. 2010; 16:877–889. [PubMed: 20663241]
- Masur DM, Sliwinski M, Lipton RB, Blau AD, Crystal HA. Neuropsychological prediction of dementia and the absence of dementia in healthy elderly persons. *Neurology*. 1994; 44:1427–1432. [PubMed: 8058143]
- Meredith MA, Nemitz JW, Stein BE. Determinants of multisensory integration in superior colliculus neurons. I. Temporal factors. *The Journal of Neuroscience*. 1987; 7:3215–3229. [PubMed: 3668625]
- Meredith MA, Stein BE. Visual, auditory, and somatosensory convergence on cells in superior colliculus results in multisensory integration. *Journal of Neurophysiology*. 1986; 56:640–662. [PubMed: 3537225]
- Meredith MA, Stein BE. Spatial determinants of multisensory integration in cat superior colliculus neurons. *Journal of Neurophysiology*. 1996; 75:1843–1857. [PubMed: 8734584]
- Miller J. Divided attention: Evidence for coactivation with redundant signals. *Cognitive Psychology*. 1982; 14:247–279. [PubMed: 7083803]
- Miller J. Timecourse of coactivation in bimodal divided attention. *Perception and Psychophysics*. 1986; 40:331–343. [PubMed: 3786102]
- Molholm S, Ritter W, Murray MM, Javitt DC, Schroeder CE, Foxe JJ. Multisensory auditory-visual interactions during early sensory processing in humans: A high-density electrical mapping study. *Brain Research Cognitive Brain Research*. 2002; 14:115–128. [PubMed: 12063135]
- Mordkoff JT, Yantis S. An interactive race model of divided attention. *Journal of Experimental Psychology Human Perception and Performance*. 1991; 17:520–538. [PubMed: 1830090]
- Murray MM, Molholm S, Michel CM, Heslenfeld DJ, Ritter W, Javitt DC, Schroeder CE, Foxe JJ. Grabbing your ear: rapid auditory-somatosensory multisensory interactions in low-level sensory cortices are not constrained by stimulus alignment. *Cerebral Cortex*. 2005; 15:963–974. [PubMed: 15537674]
- Oldfield RC. The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*. 1971; 9:97–113. [PubMed: 5146491]
- Otto TU, Dassy B, Mamassian P. Principles of multisensory behavior. *Journal of Neuroscience*. 2013; 33:7463–74. [PubMed: 23616552]
- Otto TU, Mamassian P. Noise and correlations in parallel perceptual decision making. *Current Biology*. 2012; 22:1391–1396. [PubMed: 22771043]
- Pavani F, Spence C, Driver J. Visual capture of touch: Out-of-the-body experiences with rubber gloves. *Psychological Science*. 2000; 11:353–359. [PubMed: 11228904]
- Peiffer AM, Mozolic JL, Hugenschmidt CE, Laurienti PJ. Age-related multisensory enhancement in a simple audiovisual detection task. *Neuroreport*. 2007; 18:1077–1081. [PubMed: 17558300]
- Ross LA, Saint-Amour D, Leavitt VM, Javitt DC, Foxe JJ. Do you see what I am saying? exploring visual enhancement of speech comprehension in noisy environments. *Cerebral Cortex*. 2007; 17:1147–1153. [PubMed: 16785256]
- Salthouse, TA. Speed of Behavior and its Implication for Cognition in *Handbook of the Psychology of Aging*. Birren, JE.; Schaie, KW., editors. New York: Van Nostrand Reinhold Co; 1985. p. 400–426.
- Schurmann M, Kolev V, Menzel K, Yordanova J. Spatial coincidence modulates interaction between visual and somatosensory evoked potentials. *Neuroreport*. 2002; 13:779–783. [PubMed: 11997686]
- Sliwinski M, Buschke H, Stewart WF, Masur D, Lipton RB. The effect of dementia risk factors on comparative and diagnostic selective reminding norms. *Journal of the International Neuropsychological Society*. 1997; 3:317–326. [PubMed: 9260441]

- Stein BE, Huneycutt WS, Meredith MA. Neurons and behavior: The same rules of multisensory integration apply. *Brain Research*. 1988; 448:355–358. [PubMed: 3378157]
- Stein BE, Magalhaes-Castro B, Kruger L. Superior colliculus: Visuotopic-somatotopic overlap. *Science*. 1975; 189:224–226. [PubMed: 1094540]
- Stein BE, Meredith MA. Multisensory integration. Neural and behavioral solutions for dealing with stimuli from different sensory modalities. *Annals of the New York Academy of Sciences*. 1990; 608:51–65. discussion, 65–70. [PubMed: 2075959]
- Stein, BE.; Meredith, MA. *The Merging of the Senses*. Cambridge, Massachusetts: MIT Press; 1993.
- Stein BE, Wallace MT. Comparisons of cross-modality integration in midbrain and cortex. *Progress in Brain Research*. 1996; 112:289–99. [PubMed: 8979836]
- Stephen JM, Knoefel JE, Adair J, Hart B, Aine CJ. Aging-related changes in auditory and visual integration measured with MEG. *Neuroscience Letters*. 2010; 484:76–80. [PubMed: 20713130]
- Stern Y, Habeck C, Moeller J, Scarmeas N, Anderson KE, Hilton HJ, Flynn J, Sackeim H, van Heertum R. Brain networks associated with cognitive reserve in healthy young and old adults. *Cerebral Cortex*. 2005; 15:394–402. [PubMed: 15749983]
- Teder-Sälejärvi WA, Di Russo F, McDonald JJ, Hillyard SA. Effects of spatial congruity on audio-visual multimodal integration. *Journal of Cognitive Neuroscience*. 2005; 17:1396–1409. [PubMed: 16197693]
- Verghese J, Wang C, Lipton RB, Holtzer R, Xue X. Quantitative gait dysfunction and risk of cognitive decline and dementia. *Journal of Neurology Neurosurgery and Psychiatry*. 2007; 78:929–935.
- Wallace MT, Wilkinson LK, Stein BE. Representation and integration of multiple sensory inputs in primate superior colliculus. *Journal of Neurophysiology*. 1996; 76:1246–1266. [PubMed: 8871234]
- Yesavage JA, Brink TL, Rose TL, Lum O, Huang V, Adey M, Leirer VO. Development and validation of a geriatric depression screening scale: A preliminary report. *Journal of Psychiatric Research*. 1982; 17:37–49. [PubMed: 7183759]

a. Apparatus b. Unisensory Conditions c. Multisensory Conditions **Fig. 1.**

Experimental Procedures. (a) Apparatus: Participants were seated in a comfortable chair and were required to make speeded foot-pedal responses to all stimuli, regardless of sensory modality and spatial location. They responded to all stimuli by pressing the pedal under their right foot. (b) Unisensory conditions: Participants received visual (V) and somatosensory (S) alone stimuli that were presented to either the left or right side of space. (c) Multisensory conditions: Participants received a total of four multisensory VS stimulus conditions, where both V and S information were presented within or across hemifields.

Fig. 2. Averaged RT data by modality. Mean values (with SEM bars) of each **VS** multisensory pairing, placed directly next to constituent unisensory **V** and **S** mean values. * $p < 0.01$.

Fig. 3.

Results of Miller's Test of the Race Model. The cumulative probability difference waves (actual minus predicted probability), over the trajectory of averaged responses, are depicted for each of the four multisensory VS pairings. Note that violations of the race model were evident across all four multisensory conditions over the fastest quartile (0–25%) of RT responses (see shaded box).

Table 1

Participant Demographics (n=18)

Variable	M±SD	Range
Age (years)	73.94 (5.12)	66 - 87
Education (years)	13.56 (1.98)	12 - 18
Global Disease Status Score (0-10)	0.83 (0.86)	0 - 2
Beck Anxiety Score	5.17 (7.97)	0 - 34
Geriatric Depression Score	4.67 (4.78)	0 - 18
Somatosensory Pulse (volts)	86.94 (20.73)	50 - 135
Overall RT (ms)	313.81 (39.97)	217 – 373

Table 2
Linear mixed effects model results for: 1) Multisensory RT Effect, 2) Spatial Location, 3) Multisensory RT Effect by spatial location, and 4) Adjustments.*

#	Group/Adjustment	Condition	Estimate	95% Confidence Interval		SE	P value
				lower	upper		
1)	Multisensory RT Effect	Multisensory vs. Unisensory Average	0.77	0.67	0.86	0.05	<.001
2)	Spatial Location	Within vs. Across Hemifield	0.04	-0.05	0.12	0.04	NS
3)	Multisensory RT Effect and Spatial Location	Within Hemi (Left) vs. Uni Average [VL, SL]	0.77	0.69	0.84	0.04	<.001
		Within Hemi (Right) vs. Uni Average [VR, SR]	0.74	0.61	0.87	0.07	<.001
		Across Hemi (Left) vs. Uni Average [VR, SL]	0.81	0.67	0.95	0.07	<.001
		Across Hemi (Right) vs. Uni Average [VL, SR]	0.76	0.62	0.90	0.07	<.001
4)	Adjustments	Gender	-0.18	-0.63	0.27	0.23	0.44
		Education	0.11	-0.09	0.32	0.10	0.28
		Global Disease Status	-0.31	-0.65	0.03	0.17	0.08

* This Linear Mixed Effects Model adjusted for processing speed by using the inverse transformation of reaction time (RT: 1/RT*1000).

Table 3
Multisensory RT effect by Spatial Location^a

Spatial Location	RTs to Simultaneous conditions	RTs to constituent VS unisensory conditions		T Value _(df) ; p-Value
Within Hemi Right [VR_SR]	275 ms	SR:	308 ms	$t_{(17)} = 3.36$; $p < 0.001$
		VR:	399 ms	$t_{(17)} = 11.53$; $p < 0.001$
Within Hemi Left [VL_SL]	273 ms	SL:	307 ms	$t_{(17)} = 3.99$; $p < 0.001$
		VL:	403 ms	$t_{(17)} = 14.18$; $p < 0.001$
Across Hemi Left [VL_SR]	270 ms	SL:	307 ms	$t_{(17)} = 3.87$; $p < 0.001$
		VR	399 ms	$t_{(17)} = 13.23$; $p < 0.001$
Across Hemi Right [VR_SL]	276 ms	SR	308 ms	$t_{(17)} = 3.36$; $p < 0.001$
		VL	403 ms	$t_{(17)} = 11.51$; $p < 0.001$

^aResults from planned comparisons (paired t-tests) confirm that RTs to all VS pairs, within and across hemi fields, were significantly shorter than any unisensory stimulus condition; thus, providing evidence for a RSE across all sensory combinations.