Published in final edited form as:

Anim Behav. 2012 July 1; 84(1): 21–27. doi:10.1016/j.anbehav.2012.03.010.

Evidence for intra-sexual selection in wild female baboons

Dorothy L. Cheney^{1,*}, Joan B. Silk², and Robert M. Seyfarth³

¹Department of Biology, University of Pennsylvania, Philadelphia, PA 19104

²Department of Anthropology, University of California, Los Angeles, CA 90095

³Department of Psychology, University of Pennsylvania, Philadelphia, PA 19104

Abstract

Because female reproductive success in social mammals is determined largely by parental rather than mating effort, intra-sexual competition among females has typically been assumed to occur primarily over food. Recently, however, renewed attention has been paid to the importance of other sources of variation in female fitness, with a concomitant focus on a broader definition of intra-sexual selection that encompasses both competition for resources and competition for mates and social partners. We present behavioural and demographic data gathered over 15 years on a group of wild chacma baboons (*Papio hamadryas ursinus*) which show that females incur several costs when living in groups with many females and comparatively few males: increased intra-sexual aggression, less stable bonds with female partners, and increased mortality. Female-female aggression was higher in years when the adult sex ratio was more highly skewed toward females, with low-ranking female exhibiting particularly high rates of aggression toward other females. Females' social bonds with other females also became less stable in years when the group contained many females. Finally, female mortality rates were highest in years when the group contained comparatively more females and fewer males. The negative correlation between the number of males and female mortality suggests that the increase in mortality was due to increased predation rather than food competition. Results indicate that intra-sexual competition for social partners, mates, paternal investment, and perhaps also male protection occurs even in species where females exhibit low reproductive skew.

Keywords

female; baboon; intra-sexual selection; mate competition; social bonds; paternal investment; mortality; predation

Discussions about the costs and benefits of female sociality in mammals have tended to emphasize selective pressures related to food competition. Because female reproductive success is determined largely by parental effort and nutritional condition (Trivers 1972),

© 2012 The Association for the Study of Animal Behaviour. Published by Elsevier Ltd. All rights reserved.

*corresponding author cheney@sas.upenn.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

female sociality is thought to reflect a trade-off between benefits associated with enhanced between-group competition and reduced predation risk and costs associated with increased within-group food competition (Emlen & Oring 1977; Wrangham 1980; van Schaik 1983; Janson & van Schaik 1988; Sterck et al. 1997; Koenig 2002; Majolo et al. 2008). As the number of females in a group increases, within-group competition is expected to intensify, leading groups to become less cohesive and even to fission.

Recently, however, renewed attention has been paid to the importance of other sources of variation in female fitness, with a concomitant focus on a broader definition of intra-sexual selection that encompasses variation in access not just to food but also to mates and same-sexed social partners (Carranza 2009; Clutton-Brock 2009; Clutton-Brock & McAuliffe 2009; Stockley & Bro-Jorgensen 2011; Rosvall 2011). For example, female chacma baboons (*Papio hamadryas ursinus*) who are able to maintain strong bonds with other females experience higher longevity and offspring survival than females with weaker bonds, and these benefits are largely independent of dominance rank (Silk et al. 2003, 2009; 2010b; see also Silk et al. 2003). Similar correlations between strong same-sex bonds and female reproductive success have been documented in horses (*Equus equus*: Cameron et al. 2009); bottlenosed dolphins (*Tursiops truncatus*: Frère et al. 2010), house mice (*Mus musculus*: Weidt et al. 2008), rats (*Rattus norvegicus*: Yee et al. 2008), and humans (e.g. Holt-Lunstad et al. 2010). Just as increasing numbers of females may intensify resource competition, however, so may they constrain females' ability to maintain all of their possible social relationships. Literature surveys suggest that, as the number of females in a group size increases, the relative proportion of grooming that females are able to devote to each other group member decreases, thereby decreasing group cohesion (Dunbar 1991; Henzi et al. 1997; Lehmann et al. 2007).

In many species, females also compete actively for access to mates and paternal investment (reviewed by Carranza 2009; Clutton-Brock 2009; Clutton-Brock & McAuliffe 2009; Stockley & Bro-Jorgensen 2011; Rosvall 2011). Such competition is particularly intense in socially monogamous species with high degrees of paternal investment (e.g. many birds; West-Eberhard 1983; Kraaijeveld et al. 2007; Rosvall 2011) and cooperatively breeding species with high reproductive skew (e.g. meerkats, *Suricata suricatta*, Clutton-Brock et al. 2006; Alpine marmots, *Marmota marmota*: Hacklander et al. 2003). However, female mate competition also occurs in species with low reproductive skew, and strongly biased sex ratios heighten such competition (e.g. antelope: Milner-Gulland et al. 2003; ponies, *Equus caballus*: Rutberg & Greenberg 1990; Bro-Jorgensen 2007; house mice: Rusu & Krackow 2004; gorillas, *Gorilla gorilla*: Watts 1990; chimpanzees, *Pan troglodytes*: Niemeyer & Anderson 1983; bonobos, *Pan paniscus*: Hohmann & Fruth 2003; baboons, *Papio hamadryas* spp: Seyfarth 1978; Dunbar & Sharman 1983; reviewed by Stockley & Bro-Jorgensen 2011; Rosvall 2011).

In one study of chacma baboons, rates of female-female aggression increased as the proportion of sexually receptive females in the group increased, with pregnant females exhibiting the highest rates of aggression (Huchard & Cowlshaw 2011). One interpretation of this behaviour is that food competition favors females who attempt to suppress the reproduction of rivals (see also Wasser & Starling 1988). However, such aggression may

also reflect competition for paternal investment in offspring. Lactating female chacma baboons compete to form ‘friendships’ with resident adult males (Palombit et al. 2001) that may function to deter infanticide (Palombit et al. 1997; Weingrill 2000; Clarke et al. 2009). Competition for male friends may exacerbate mate competition because lactating females form friendships preferentially with their primary consort partner, who is often their infants’ father (Moscovice et al. 2010). Increased competition for mates and the paternal investment they provide may represent another cost of increasing female numbers, and this will be exacerbated when the adult sex ratio is heavily skewed toward females.

While most groups of mammals exhibit adult female-biased sex ratios, it remains unclear whether this skew benefits or harms females. Most species of polygynous mammals are sexually dimorphic, with the result that males are dominant to females and can exclude them from desirable resources. Females living in groups with many males and a more evenly balanced sex ratio may therefore incur costs associated with decreased access to food. Conversely, if males contribute to parental care, or if predation rates are high, large numbers of males may enhance female and offspring survival. There is some indication that, as predation risk increases, primate groups experience a disproportionate increase in the number of males over what would be predicted by the number of females (Hill & Lee 1998). If the presence of many males deters predation, females may experience lower mortality rates when there are more males present and the group’s sex ratio is more evenly balanced. The degree to which variation in sex ratios influence female fitness, however, has seldom been considered.

In this paper, we use behavioural and demographic data collected over a 15 year-period on a group of wild chacma baboons to examine how variation in the relative numbers of males and females affects female intra-sexual competition, the strength of female bonds, and female mortality.

Methods

Study group and demographic analysis

Research was conducted between 1992–2007 on a group of wild chacma baboons in the Moremi Game Reserve in the Okavango Delta of Botswana (Cheney et al. 2004; Cheney & Seyfarth 2007). All research was conducted using noninvasive methods approved by the Animal Care and Use Committee of the University of Pennsylvania (Protocol no. 19001). The group had been observed since 1978 and was fully habituated to humans on foot. Information about maternal kinship was available for all individuals, and paternal relatedness was known for some. From 1992–2007 group size (i.e. individuals > 1 year) ranged from 55 to 76. The primary causes of mortality in this population are infanticide and predation by leopards, lions, and other predators. Predation is estimated to account for more than 96% of juvenile and adult deaths (Busse 1982; Cheney et al. 2004; Cheney & Seyfarth 2007).

As in other species of cercopithecine primates, female baboons remain in their natal groups throughout their lives and assume dominance ranks similar to their mothers’ (Melnick & Pearl 1987; Cheney & Seyfarth 2007). Adult female dominance ranks were calculated

monthly based on the direction of approach-retreat interactions. For most of the study, the female dominance hierarchy remained stable.

Females in this population reach puberty between 4–5 years of age and usually give birth for the first time between the ages of 6 to 8. All females who reached 4 years of age ($N = 64$) were included in the demographic analyses. By the age of four years, males begin to dominate adult females (Engh et al. 2009). However, males do not enter the adult male dominance hierarchy or reach adult size until around 9 years of age, when they usually emigrate from their natal group.

Because they do not emigrate, females and juveniles who disappeared were assumed to be dead. In many cases, mortality could be ascribed to confirmed or strongly suspected predation (Cheney et al. 2004; Cheney & Seyfarth 2007). We did not attempt to calculate mortality rates for males because males who disappeared may have emigrated.

The annual number of individuals in each age/sex class was calculated as of January 1 each year. All analyses were restricted to individuals aged more than 4 years. The annual number of females aged 4 or more varied from 24 to 36. The annual number of males aged 4 or more varied from 16 to 25. The annual number of resident immigrant and natal males aged more than 9 varied from 6 to 12. When calculating the adult sex ratio, we included females aged more than 4 years, fully grown immigrant males, and natal males aged at least 9 years. Thus, subadult males (aged 4 through 8) were included in the “number of males”, but not in the “adult sex ratio”, category. The annual adult male:female ratio ranged from 1:2.5 to 1:4.5. Because group size was positively correlated with the number of females (two-tailed Spearman rank correlation: $r_s = 0.74$, $N = 15$ years, $P < 0.01$), the number of adult males ($r_s = 0.65$, $P < 0.02$), and the number of males ($r_s = 0.38$, $P < 0.20$), and in particular because we wanted to examine separately the effects of female and male numbers on behaviour and mortality, group size was not included as a predictor variable in statistical tests.

Behavioural and statistical analyses

Data on affiliative and aggressive behaviour were gathered using focal animal sampling (Altmann 1974) during 1992–93 and 2001–2007 (7 years). Behavioural analyses were restricted to females aged at least 5 years because younger individuals were not consistently sampled in all years. Aggressive behaviour included head bobs, lunges, chases, and bites. Because high-ranking females have more available targets for aggression than do low-ranking females, each female’s annual rate of aggression was corrected for the number of females who ranked below her.

To examine whether cycling and pregnant females experienced more aggression than lactating females (Wasser & Starling 1988, Huchard & Cowlshaw 2011), we compared the frequency of each female’s actual aggression given and received in each of three reproductive states (cycling, lactating, and pregnant) with the amount that would have been expected given the proportion of time that she spent in that state. This analysis was restricted to data gathered between 2003–2005, the three years for which we had the most accurate and continuous information about females’ reproductive states. Females who were observed in only one reproductive state were not included in this analysis.

We were also interested in determining whether the consistency of females' preferences for their top partners varied in relation to the number of females in the group. To assess changes in the consistency of females' preferences from one year to the next, we modified the procedures used in Silk et al. (2010b, submitted). First, we identified each female's top three partners (as measured by rate of approaches, groom presents, grooming initiations, and grooming duration; see Silk et al. (2010b) in each year. Then, for females who were present during at least two years, we assessed the consistency of females' preferences from one year to the next using the formula:

$$\frac{NS - U}{NS - S - X}$$

where N is the number of years in which partner choice was evaluated, S is the number of top partners being evaluated, U is the number of different top partners that the female had in those two years, and X is the number of top partners in the first year who died in second year. (No female ever lost all three partners from one year to the next.) Here, $N = 2$ and $S = 3$, so the formula simplifies to $(6 - U) / (3 - X)$. Importantly, this index controls for changes in female' top partners that were due to deaths. Thus, females with low values were ones who switched partners from one year to the next even though the previous year's partners were still present in the group. The value of the Partner Preference Index (PPI) varied from 1, for females who had the same partners from one year to the next, to 0, for females who had a completely different set of partners in each year. PPI values were scored across 6 years (2001–2007).

In statistical tests, females were classified according to their dominance rank (calculated as the proportion of other females dominated) and the presence or absence of close kin (mothers, adult daughters, or adult maternal sisters) that were present in each year.

Statistical analyses were conducted with R statistical software (version 2.9.0, R Foundation for Statistical Computing, R Development Core Team, 2009). We used linear mixed models (LMM) for analyses in which individuals were sampled repeatedly and did not contribute equally to the data set. LMMs were calculated using the function 'lmer' of the R package 'lme4' (R package version 0.999375–31). We used a linear model (LM) for the analysis of annual mortality rates across the 15 complete years of observation (1993–2007). To begin analyses, we first examined a full model that used a single dependent measure and all possible predictors and their interactions. If the overall result was significant, we eliminated non-significant predictors and ran a second model that included all significant predictors and any possible interactions. Where we found a significant interaction between two effects, we split the data set and carried out post hoc comparisons to explore the nature of the interaction in more detail. In some simple analyses where individuals were not sampled repeatedly and contributed equally to the data set, we used a binomial test, chi-square test, or a Spearman rank correlation coefficient. All correlations were corrected for ties, and all reported probability values are two-tailed.

Results

Female-female aggression

To examine which factors best predicted annual rates of female-female aggression, we first ran a LMM with each female's annual corrected rate of aggression toward other females as the dependent variable and number of females, number of males, sex ratio, female dominance rank, and their interactions as predictor variables. Female identity was entered as a random factor. Only sex ratio and dominance rank were significant predictors. There were no significant interactions (Table 1a).

To examine more closely the role of these significant predictors, we next ran a LMM using only sex ratio, rank, and their interaction as predictor variables. There was a significant interaction between sex ratio and rank, indicating that rates of female-female aggression were higher in years when the sex ratio was more skewed toward females, and that, as the sex ratio became more skewed, low-ranking females showed a relatively greater increase in rates of aggression toward other females than did high-ranking females (Table 1b; Fig. 1).

Although baboons are not seasonal breeders, two or more females may be in estrus simultaneously (Bulger 1993), and when that happens they cannot all form consortships with the alpha male. Information on the identity and rank of each female's primary consort partner was available for 22 conceptive cycles from 2004 to 2007 (Moscovice et al. 2010). There was a significant positive correlation between the dominance ranks of male and female consort partners ($r_s = 0.77$, $P < 0.01$), suggesting either that high-ranking females were able to exclude lower-ranking females from access to dominant partners and/or that high-ranking males preferred to mate with high-ranking females.

Mate competition also appears to have been associated with rare mob attacks on other females. As in other female cercopithecines, most aggressive interactions among female baboons are dyadic, and coalitions seldom involve more than two females (Wittig et al. 2007). The years from 2000 to 2005, however, comprised a period when the sex ratio was more skewed than average (1:4 vs. 1:3 in other years) and female aggression rates were also higher than average (0.344/female/hr vs. 0.168/female/hr). This period was also characterized by unusual mob attacks by females on other females and rare challenges to the female dominance hierarchy (described by Engh et al. 2006; Cheney & Seyfarth 2007; Engh et al. 2009). Most unusually for cercopithecine primates, the instigators often ranked lower than their opponents, and they typically targeted estrous females who were in consort with an adult male (81% of 31 mob attacks). The attacks eventually led several adult and adolescent females to fall below their matriline's rank, while eight adolescent females rose above theirs (Engh et al. 2006, 2009).

With the exception of rare mob attacks, however, cycling females did not generally receive more aggression than would have been expected given the length of time that they spent in that reproductive state (binomial test: $N = 22$, 63.6% more, 36.4% less, $P = 0.286$; cf. Huchard & Cowlishaw 2011). Also in contrast to Huchard & Cowlishaw's study (2011), we did not find that lactating females as a group received less aggression than expected ($N = 24$,

more = 33.3%, less = 66.7%, $P = 0.152$), or that pregnant females gave more aggression ($N = 24$, 50% more, 50% less, $P > 0.999$).

Although there was little variation from year to year in the proportion of multiparous females that were cycling, pregnant, or lactating, there was considerable annual variation in the proportion of females that were nulliparous (*i.e.*, either cycling before their first conception or experiencing their first pregnancy: range: 21–43%). A LMM revealed that female aggression rates were significantly higher in years when the group contained a higher proportion of nulliparous females ($r = 0.24$, $t = 3.49$, $N = 15$, $P < 0.001$).

Consistency of social bonds among females

To examine the relationship between group composition and the consistency of females' partner preferences, we first ran a LMM with each female's partner preference index (PPI) as the dependent variable and number of females, number of males, adult sex ratio, female dominance rank, presence or absence of kin (a categorical variable), and their interactions as predictors. Female identity was entered as a random factor (Table 2a).

Because this model suggested that dominance rank was unimportant, we deleted this predictor and ran a second model with number of females, number of males, adult sex ratio, and the presence of kin as predictors. All were significant; there were no significant interactions (Table 2b). Additional post hoc LMMs in which PPI was the dependent variable and each of the significant predictor variables was tested alone revealed a significant negative correlation only between PPI and the number of females ($t = -2.505$, $N = 148$, $P < 0.05$; number of males: $t = .374$, *NS*; sex ratio: $t = -0.821$, *NS*; presence of kin: $t = 1.474$, *NS*). Thus, as the number of females in the group increased, females were less likely to retain their top social partners from one year to the next (Fig. 2).

Mortality

To examine the factors associated with female mortality, we ran a LM with annual female mortality rate as the dependent variable and the number of females, number of males, adult sex ratio, and their interactions as predictors. This analysis revealed that annual mortality rates were significantly negatively correlated with the number of males ($t = -2.947$, $N = 15$, $P < 0.05$) and significantly positively correlated with the number of females ($t = 2.509$, $P < 0.05$) (Fig. 3).

There was some indication that, with the exception of very old females (>20 years), females in the top third of the dominance hierarchy experienced lower mortality rates than females in the bottom two thirds of the hierarchy. Only 18% of the 33 females (aged < 20) that died between 1993 and 2007 ranked in the top third of the dominance hierarchy (one-sample chi square test: $X^2 = 4.55$, $P = 0.066$). By contrast, 48% of females in the bottom third of the dominance hierarchy died during this time. Similarly, of the eight lactating females with young infants that died during this time, only one (13%) ranked in the top third of the dominance hierarchy.

Finally, there was no relationship between annual infant mortality and the number of females ($t = -0.015$, *NS*) or the number of males ($t = -0.383$, *NS*) in the group.

Discussion

Taken together, results suggest that female baboons incur several, probably independent, costs when living in groups with many females and few males: increased intra-sexual aggression, less stable bonds with female partners, and increased mortality rates. These costs appear to be unrelated to food competition.

Our results are consistent with previous research (Huchard & Cowlshaw 2011) suggesting that female baboons actively compete for access to males, and that they may do so because of the paternal investment they can potentially provide. Rates of female aggression, and in particular rates of aggression by low-ranking females, increased as the sex ratio became more skewed toward females, suggesting that when males were in relatively short supply there was more competition among females, and that this cost was borne primarily by low-ranking females.

Female-female competition for males may be particularly intense among chacma baboons due to the threat of infanticide by immigrant males (Palombit et al. 1997; Weingrill 2000; Clarke et al. 2009). Lactating females compete to establish friendships with males (Palombit et al. 2001), and individuals who are unable to do so show elevated glucocorticoid levels when a new male enters the group (Beehner et al. 2005; Engh et al. 2006). Females form friendships preferentially with former consort partners, who are often their infants' fathers (Moscovice et al. 2010). The fact that dominant females can displace subordinate females from male friends, and in some cases prevent them from forming friendships entirely, suggests that paternal investment is a limited commodity that cannot be shared across a large number of females. Thus, any females who become pregnant within the same year may pose a competitive threat that potentially affects infant survival. As a result, it may be advantageous for females in all reproductive states to attempt to prevent other females from mating and forming friendships with males.

Rates of female aggression were higher in years when the group included a larger proportion of nulliparous females. Access to mates and, subsequently, male friends may be especially important for young females, who often experience higher infant mortality rates (Nicolson 1987; Cheney et al. 2004). Older mothers tend to form weaker friendships than younger mothers and in some cases even prefer associating with male and female matrilineal kin, especially when they have subadult sons (Palombit et al. 2001). For all of these reasons, nulliparous females may be more likely than older females to mount challenges to the dominance hierarchy and attack other cycling females (see also Samuels & Hendrickson 1983; Samuels et al. 1987 for similar observations on rhesus macaques, *Macaca mulatta*, and yellow baboons, *P.h. cynocephalus*).

An additional cost of living with large numbers of females may be a reduction in the consistency of social bonds. The partner preference index (PPI) can be regarded as a psychological measure of females' ability or motivation to retain preferred female partners from one year to the next when those partners are still present in the group. Results indicate that females' partner consistency decreased as the number of females in the group increased. Whether this decrease resulted from increased competition for partners, the need to interact

with more partners (Lehmann et al. 2007), or some combination of these and other factors remains to be determined. Previous analyses on the same social group have indicated that females with strong, consistent bonds experience lower mortality and higher offspring survival than those weaker bonds (Silk et al. 2010a,b). Thus, to the extent that the presence of many females causes females' partner preferences to become less stable, large numbers of females may represent a fitness cost independent of those associated with increased competition for food, mates, and male friends.

Finally, female mortality rates were significantly higher in years when the group contained high numbers of females and low numbers of males, suggesting that reduced survival may constitute a third cost of living in such groups. It seems unlikely that the increase in female mortality was due primarily to heightened food competition, because female mortality rates were *negatively* correlated with the number of males. If food competition had played an important role, this correlation would have been positive. Instead, it seems possible that females living in groups with relatively few males and relatively large numbers of females are at increased risk of predation, the primary cause of adult and juvenile mortality in this population (Cheney et al. 2004). Primate groups that confront high rates of predation typically include more males than would be expected given the number of females (Hill & Lee 1998). The presence of many males may reduce predation risk not only through early detection but also through active defense, because male baboons frequently attack and even injure predators such as leopards (*Panthera pardus*) and hyenas (*Crocuta crocuta*) (Cowlshaw 1994; Cheney et al. 2004). As a result, any decrease in the number of males relative to females may put females at increased predation risk; such groups may even attract predators. More data on dispersion patterns and group cohesion (e.g. King & Cowlshaw 2009) will be needed to investigate these questions.

High-ranking females may be buffered to some degree from these risks. High-ranking females tend to live longer than low-ranking females (Silk et al. 2010b) and also appear to experience lower mortality rates. High-ranking females may be able to minimize the threat of predation by foraging and resting in proximity to males, thereby forcing more subordinate individuals to the group's periphery, where they may be more vulnerable to attack (Ron et al. 1996). Such competitive exclusion may have contributed to low-ranking females' heightened aggression in years when the sex ratio was most skewed.

Infant mortality rates were unaffected by variation in the numbers of males and females, probably because the primary cause of infant mortality is infanticide by immigrant males. In any given year, as many as eight males may immigrate into the group. Though these males may aid in defense against predator attacks, they also represent an infanticidal threat. Whether a male is able to commit infanticide depends on whether he can attain high dominance status (Palombit et al. 2000), which seems to be unaffected by the number of males in the group.

Because female reproductive success in social mammals is determined largely by parental rather than mating effort, intra-sexual competition among females has typically been assumed to occur primarily over food. Recently, however, renewed attention has been paid to the importance of mate competition among females, with a concomitant focus on a

broader definition of intra-sexual selection that encompasses both competition for resources and competition for mates and social partners (Carranza 2009; Clutton-Brock 2009; Clutton-Brock & McAuliffe 2009; Stockely & Bro-Jorgensen 2011; Huchard & Cowlshaw 2011; Rosvall 2011). Such competition is predicted to be particularly intense in monogamous species and in species that exhibit high reproductive skew, where females compete for a single breeding position. The results described here indicate that intra-sexual competition for social partners, mates, paternal investment, and perhaps also male protection occurs even in species where females exhibit low skew, and that females living in groups with many other females and an imbalanced sex ratio suffer the costs of increased aggression, less stable social relationships, and increased mortality.

Acknowledgments

We thank the Office of the President of the Republic of Botswana and the Botswana Department of Wildlife and National Parks for permission to conduct research in the Moremi Reserve. We are grateful to J. Beehner, T. Bergman, R. Palombit, C. Crookford, A. Engh, J. Fischer, M. Heesen, S. Johnson, D. Kitchen, L. Moscovice, J. Nicolson, D. Rendall, R. Wittig, K. Seyfarth, C. Shaw, M. Mokopi, and A. Mokopi and for their assistance with data collection. We also thank J. Beehner, T. Bergman, N. Snyder-Mackler, and two anonymous reviewers for comments. Research was supported by the National Science Foundation, the National Institutes of Health, the Leakey Foundation, the National Geographic Society, and the University of Pennsylvania. Research was approved by the Animal Care and Use Committee of the University of Pennsylvania (Protocol no. 19001).

References

- Altmann J. Observational study of behavior: Sampling methods. *Behaviour*. 1974; 49:229–267.
- Beehner JC, Bergman TJ, Cheney DL, Seyfarth RM, Whitten PL. The effect of new alpha males on female stress in free-ranging baboons. *Animal Behaviour*. 2005; 69:1211–1221.
- Bro-Jorgensen J. Reversed sexual conflict in a promiscuous antelope. *Current Biology*. 2007; 17:2157–2161. [PubMed: 18060785]
- Bulger JB. Dominance rank and access to estrous females in male savanna baboons. *Behaviour*. 1993; 127:67–103.
- Busse C. Leopard and lion predation upon chacma baboons living in the Moremi Wildlife Reserve. *Botswana Notes and Records*. 1982; 12:15–21.
- Cameron EZ, Setsaas TH, Linklater WL. Social bonds between unrelated females increase reproductive success in feral horses. *Proceedings of the National Academy of Sciences USA*. 2009; 106:13850–13853.
- Carranza J. Defining sexual selection as sex-dependent selection. *Animal Behaviour*. 2009; 77:749–751.
- Cheney, DL.; Seyfarth, RM. *Baboon Metaphysics: The Evolution of a Social Mind*. Chicago: Chicago University Press; 2007.
- Cheney DL, Seyfarth RM, Fischer J, Beehner J, Bergman T, Johnson SE, Kitchen DM, Palombit RA, Rendall D, Silk JB. Factors affecting reproduction and mortality among baboons in the Okavango Delta, Botswana. *International Journal of Primatology*. 2004; 25:401–428.
- Clarke PMR, Henzi SP, Barrett L. Sexual conflict in chacma baboons, *Papio hamadryas ursinus*: absent males select for proactive females. *Animal Behaviour*. 2009; 77:1217–1225.
- Clutton-Brock TH. Sexual selection in females. *Animal Behaviour*. 2009; 77:3–11.
- Clutton-Brock TH, McAuliffe K. Female mate choice in mammals. *The Quarterly Review of Biology*. 2009; 84:3–27. [PubMed: 19326786]
- Clutton-Brock TH, Hodge SJ, Spong G, Russell AF, Jordan NR, Bennett NC, Sharpe LL, Manser MB. Intrasexual competition and sexual selection in cooperative mammals. *Nature*. 2006; 444:1065–1068. [PubMed: 17183322]
- Cowlshaw G. Vulnerability to predation in baboon populations. *Behaviour*. 1994; 131:293–304.

- Dunbar RIM. Functional significance of social grooming in primates. *Folia primatologica*. 1991; 57:121–131.
- Dunbar RIM, Sharman M. Female competition for access to males affects birth rate in baboons. *Behavioral Ecology and Sociobiology*. 1983; 13:157–159.
- Emlen ST, Oring LW. Ecology, sexual selection and the evolution of mating systems. *Science*. 1977; 197:215–223. [PubMed: 327542]
- Engh AE, Beehner JC, Bergman TJ, Whitten PL, Hoffmeier RR, Seyfarth RM, Cheney DL. Female hierarchy instability, male immigration, and infanticide increase glucocorticoid levels in female chacma baboons. *Animal Behaviour*. 2006; 71:1227–1237.
- Engh AL, Hoffmeier RR, Seyfarth RM, Cheney DL. O brother, where art thou? The varying influence of older siblings in rank acquisition by female baboons. *Behavioral Ecology and Sociobiology*. 2009; 64:97–104.
- Frere CH, Krutzen M, Mann J, Connor RC, Bejder L, Sherwin WB. Social and genetic interactions drive fitness variation in a free-living dolphin population. *Proceedings of the National Academy of Sciences USA*. 2010; 107:19949–19954.
- Hacklander K, Mostl E, Walter A. Reproductive suppression in female Alpine marmots, *Marmota marmota*. *Animal Behaviour*. 2003; 65:1133–1140.
- Henzi SP, Lycett JE, Piper SE. Fission and troop size in a mountain baboon population. *Animal Behaviour*. 1997; 53:525–535.
- Hill RA, Lee PC. Predation risk as an influence on group size in cercopithecoid primates: implications for social structure. *Journal of Zoology, London*. 1998; 245:447–456.
- Hohmann G, Fruth B. Intra- and inter-sexual aggression by bonobos in the context of mating. *Behaviour*. 2003; 140:1389–1413.
- Holt-Lunstad J, Smith TB, Layton JB. Social relationships and mortality risk: a meta-analytic review. *PLoS Medicine*. 2010; 7:e1000316.
- Huchard H, Cowlshaw G. Female-female aggression around mating: an extra cost of sociality in a multimale primate society. *Behavioral Ecology*. 2011; 22:1003–1011.
- Janson C, van Schaik C. Recognizing the many faces of food competition: methods. *Behaviour*. 1988; 105:165–186.
- King AJ, Cowlshaw G. All together now: behavioural synchrony in baboons. *Animal Behaviour*. 2009; 78:1381–1387.
- Koenig A. Competition for resources and its behavioral consequences among female primates. *International Journal of Primatology*. 2002; 23:759–783.
- Lehmann J, Korstjens A, Dunbar R. Group size, grooming and social cohesion in primates. *Animal Behaviour*. 2007; 74:1617–1629.
- Majolo B, De Bortol Vizioli A, Schino G. Costs and benefits of group living in primates: group size effects on behaviour and demography. *Animal Behaviour*. 2008; 76:1235–1247.
- Majolo B, Koyama N. Seasonal effects on reconciliation in *Macaca fuscata yakui*. *International Journal of Primatology*. 2006; 27:1383–1397.
- Melnick, DJ.; Pearl, MC. Cercopithecines in multi-male groups: genetic diversity and population structure. In: Smuts, BB.; Cheney, DL.; Seyfarth, RM.; Wrangham, RW.; Struhsaker, TT., editors. *Primate Societies*. Chicago: University of Chicago Press; 1987. p. 121–134.
- Milner-Gulland EJ, Bukreeva OM, Coulson T, Lushchekina AA, Kholodova MV, Bekenov AB, Grachev IA. Reproductive collapse in saiga antelope harems. *Nature*. 2003; 422:135. [PubMed: 12634775]
- Moscovice LR, DiFiore A, Crockford C, Kitchen DW, Wittig R, Seyfarth RM, Cheney DL. Hedging their bets? Male and female chacma baboons form friendships based on likelihood of paternity. *Animal Behaviour*. 2010; 79:1007–1015.
- Nicolson, NA. Infants, mothers, and other females. In: Smuts, BB.; Cheney, DL.; Seyfarth, RM.; Wrangham, RW.; Struhsaker, TT., editors. *Primate Societies*. Chicago: University of Chicago Press; 1987. p. 330–342.
- Niemeyer CL, Anderson JR. Primate harassment of matings. *Ethology and Sociobiology*. 1983; 4:205–220.

- Palombit RA, Seyfarth RM, Cheney DL. The adaptive value of “friendships” to female baboons: experimental and observational evidence. *Animal Behaviour*. 1997; 54:599–614. [PubMed: 9299045]
- Palombit R.; Cheney, D.; Seyfarth, R.; Rendall, D.; Silk, J.; Johnson, S.; Fischer, J. Male infanticide and defense of infants in chacma baboons. In: van Schaik, C.; Janson, C., editors. *Infanticide by Males and Its Implications*. Cambridge: Cambridge University Press; 2000. p. 123–152.
- Palombit RA, Cheney DL, Seyfarth RM. Female-female competition for male “friends” in wild chacma baboons (*Papio cynocephalus ursinus*). *Animal Behaviour*. 2001; 61:1159–1171.
- Ron T, Henzi PS, Motro U. Do female chacma baboons compete for a safe spatial position in a southern woodland habitat? *Behaviour*. 1996; 133:475–490.
- Rosvall KA. Intrasexual competition in females: evidence for sexual selection? *Behavioral Ecology*. 2011; 22:1131–1140. [PubMed: 22479137]
- Rusu A, Krackow S. Kin-preferential cooperation, dominance-dependent reproductive skew, and competition for mates in communally nesting female house mice. *Behavioral Ecology and Sociobiology*. 2004; 56:298–305.
- Rutberg AT, Greenberg SA. Dominance, aggression frequencies and modes of aggressive competition in feral pony mares. *Animal Behaviour*. 1990; 40:322–331.
- Samuels A, Hendrickson RV. Outbreak of severe aggression in captive *Macaca mulatta*. *American Journal of Primatology*. 1983; 5:277–281.
- Samuels A, Silk JB, Altmann J. Continuity and change in dominance relations among female baboons. *Animal Behaviour*. 1987; 35:785–793.
- Seyfarth RM. Social relationships among adult male and female baboons. I. Behaviour during sexual consortship. *Behaviour*. 1978; 64:204–226.
- Silk JB, Alberts SC, Altmann J. Social bonds of female baboons enhance infant survival. *Science*. 2003; 302:1231–1234. [PubMed: 14615543]
- Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, Wittig RM, Seyfarth RM, Cheney DL. The benefits of social capital: close social bonds among female baboons enhance offspring survival. *Proceedings of the Royal Society of London, B*. 2009; 276:3099–3104.
- Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, Wittig RM, Seyfarth RM, Cheney DL. Female chacma baboons form strong, equitable, and enduring social bonds. *Behavioral Ecology and Sociobiology*. 2010a; 64:1733–1747. [PubMed: 20976293]
- Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, Wittig RM, Seyfarth RM, Cheney DL. Strong and consistent social bonds enhance the longevity of female baboons. *Current Biology*. 2010b; 20:1359–1361. [PubMed: 20598541]
- Sterck EHM, Watts DP, van Schaik CP. The evolution of female social relationships in nonhuman primates. *Behavioral Ecology and Sociobiology*. 1997; 41:291–309.
- Stockley P, Bro-Jorgensen J. Female competition and its evolutionary consequences in mammals. *Biological Reviews of the Cambridge Philosophical Society*. 2011; 86:341–366. [PubMed: 20636474]
- Trivers, RL. Parental investment and sexual selection. In: Campbell, B., editor. *Sexual Selection and the Descent of Man, 1871–1971*. Chicago: Aldine; 1972. p. 136–179.
- van Schaik CP. Why are diurnal primates living in groups? *Behaviour*. 1983; 87:120–144.
- Wasser SK, Starling AK. Proximate and ultimate causes of reproductive suppression among female yellow baboons at Mikumi National Park, Tanzania. *American Journal of Primatology*. 1988; 16:97–121.
- Watts DP. Mountain gorilla life histories, reproductive competition, and some implications for captive husbandry. *Zoo Biology*. 1990; 9:185–200.
- Weidt A, Hofmann SE, Koenig B. Not only mate choice matters: fitness consequences of social partner choice in female house mice. *Animal Behaviour*. 2008; 75:801–808.
- Weingrill T. Infanticide and the value of male-female relationships in mountain chacma baboons. *Behaviour*. 2000; 137:337–359.
- Wittig RM, Crockford C, Seyfarth RM, Cheney DL. Vocal alliances in chacma baboons, *Papio hamadryas ursinus*. *Behavioral Ecology and Sociobiology*. 2007; 61:899–909.

- Wrangham RW. An ecological model of female-bonded primate groups. *Behaviour*. 1980; 75:262–300.
- Yee JR, Cavigelli SA, Delgado B, McClintock M. Reciprocal affiliation among adolescent rats during a mild group stressor predicts mammary tumors and lifespan. *Psychosomatic Medicine*. 2008; 70:1050–1059. [PubMed: 18842748]

Highlights

- Female–female competition in baboons increases with increasing sex ratio skew.
- Female baboons' bonds weaken with increasing female numbers.
- Female mortality increases as female numbers increase and male numbers decrease.
- Females incur costs when living in groups with many females and skewed sex ratios.

Figure 1.

The mean annual rate of females' corrected rates of aggression toward other females, compared with the annual female:male ratio. Black circles depict mean rates of aggression by females in the bottom half of the female dominance hierarchy; open circles depict mean rates of aggression by females in the top half of the dominance hierarchy. Mean values are presented for illustrative purposes only. All statistical analyses were based on continuous measures. Lines represent the linear model that minimises the sum of the squares of the residuals.

Figure 2.

Females' mean annual partner preference indices (PPI), compared with the annual number of females in the group. Legend as in Fig. 1.

Figure 3.

The mean annual female mortality rate, compared with annual number of males (open circles) and females (black circles) in the group. Legend as in Fig. 1.

Table 1

a. Results of a linear mixed model (LMM) in which each female's annual corrected rate of aggression toward other females was the dependent variable (N = 195) and number of females, number of males, sex ratio, and female dominance rank were predictor variables (all continuous measures). Female identity (N = 57) was entered as a random factor.

Predictors	Estimate	Standard Error	t value
# females aged >4	-0.009	0.019	-0.460
# males aged >4	0.056	0.068	0.822
Sex ratio	0.821	0.270	3.044**
Rank	-0.634	0.260	-2.440*

b. Results of a LMM using only the significant predictors (sex ratio, rank, and their interaction).

Predictors	Estimate	Standard Error	t value
Sex ratio	1.118	0.243	4.596**
Dominance rank	3.010	1.570	1.917
Sex ratio × rank	-0.998	0.425	-2.349*

* = $P < 0.05$;

** = $P < 0.01$

Legend as in Table 1a.

Table 2

a. Results of a LMM in which each female's PPI was the dependent variable (N = 148) and the number of females, number of males, adult sex ratio, female dominance rank (all continuous variables), and presence or absence of kin (a categorical variable) were predictors. Female identity (N = 44) was entered as a random factor.			
Predictors	Estimate	Standard Error	t value
# females aged >4	−0.018	0.006	−3.011**
# males aged >4	−0.148	0.056	−2.663*
Adult sex ratio	−0.451	0.169	−2.676*
Rank	0.037	0.089	0.412
Presence of kin	0.122	0.062	1.988

b. Results of a LMM in which female PPI was the dependent variable and the number of females, the number of males, and sex ratio were predictors.			
Predictors	Estimate	Standard Error	t value
# females aged >4	−0.019	0.006	−3.065**
# males aged >4	−0.149	0.056	−2.684*
Sex ratio	−0.454	0.168	−2.695*
Presence of kin	0.133	0.056	2.391*

Legend as in Table 1a.