Published in final edited form as:

Microbiol Spectr. 2014 ; 2(2): . doi:10.1128/microbiolspec.PLAS-0006-2013.

Plasmid-mediated quinolone resistance

George A. Jacoby¹, Jacob Strahilevitz², and David C. Hooper³

¹Lahey Hospital and Medical Center, Burlington, Massachusetts 01805

²Hadassah-Hebrew University, Jerusalem 91120, Israel

³Massachusetts General Hospital, Boston, Massachusetts 02114

Summary

Three mechanisms for plasmid-mediated quinolone resistance (PMQR) have been discovered since 1998. Plasmid genes *qnrA*, *qnrB*, *qnrC*, *qnrD*, *qnrS*, and *qnrVC* code for proteins of the pentapeptide repeat family that protects DNA gyrase and topoisomerase IV from quinolone inhibition. The *qnr* genes appear to have been acquired from chromosomal genes in aquatic bacteria, are usually associated with mobilizing or transposable elements on plasmids, and are often incorporated into *sulI*-type integrons. The second plasmid-mediated mechanism involves acetylation of quinolones with an appropriate amino nitrogen target by a variant of the common aminoglycoside acetyltransferase AAC(6')-Ib. The third mechanism is enhanced efflux produced by plasmid genes for pumps QepAB and OqxAB. PMQR has been found in clinical and environmental isolates around the world and appears to be spreading. The plasmid-mediated mechanisms provide only low-level resistance that by itself does not exceed the clinical breakpoint for susceptibility but nonetheless facilitates selection of higher-level resistance and makes infection by pathogens containing PMQR harder to treat.

Introduction

Plasmid-mediated quinolone resistance (PMQR) was late in being discovered. Nalidixic acid, the first quinolone to be used clinically, was introduced in 1967 for urinary tract infections. Resistance was soon observed and could also be readily selected in the laboratory. It was produced by amino acid substitutions in the cellular targets of quinolone action: DNA gyrase and topoisomerase IV (1–3). Later, decreased quinolone accumulation due to pump activation and porin loss were added as additional resistance mechanisms. Search for transferable nalidixic acid resistance in over 500 gram-negative strains in the 1970s was unrevealing (4). In the 1980's fluoroquinolones became available that were more potent and broader in spectrum. Quinolone usage increased with subsequent parallel increases in quinolone resistance (5, 6). In 1987 PMQR was reported to be present in a nalidixic acid resistant isolate of *Shigella dysenteriae* from Bangladesh (7), but this claim was later withdrawn (8). True PMQR was reported in 1998 in a multiresistant urinary *Klebsiella pneumoniae* isolate at the University of Alabama that could transfer low level resistance to nalidixic acid, ciprofloxacin, and other quinolones to a variety of gram-negative recipients (9). The responsible gene was termed *qnr*, later amended to *qnrA*, as additional *qnr* alleles were discovered. Investigation of a *qnrA* plasmid from Shanghai that provided more than the expected level of ciprofloxacin resistance led to the discovery in

2006 of a second mechanism for PMQR: modification of certain quinolones by a particular aminoglycoside acetyltransferase, AAC(6')-Ib-cr (10). A third mechanism for PMQR was added in 2007 with the discovery of plasmid-mediated quinolone efflux pumps QepA (11, 12) and OqxAB (13). A multiplex PCR assay for eight PMQR genes (lacking only *qnrVC*) has recently been perfected (14). In the past decade these genes have been found in bacterial isolates from around the world. They reduce the susceptibility of bacteria to quinolones, usually not to the level of non-susceptibility, but facilitating the selection of more quinolone resistant mutants and treatment failure. PMQR has been frequently reviewed (15–20).

Qnr Structure and Function

Cloning and sequencing *qnrA* revealed that it encoded a 218 residue protein with a tandemly repeating unit of five amino acids that indicated membership in the large (more than 1000 members) pentapeptide repeat family of proteins (21). Knowledge of the sequence allowed search for *qnrA* by PCR, and it was soon discovered in a growing number of organisms, including other *K. pneumoniae* strains in the United States (22, 23), *Escherichia coli* isolates in Shanghai (24), and *Salmonella enterica* strains in Hong Kong (25). *qnrA* was subsequently followed by discovery of plasmid-mediated *qnrS* (26), *qnrB* (27), *qnrC* (28), and *qnrD* (29). The *qnrVC* gene from *Vibrio cholerae* can also be located in a plasmid (30–33) or in transmissible form as part of an integrating conjugative element (34). These *qnr* genes generally differ in sequence by 35 % or more from *qnrA* and each other. Allelic variants have also been described in each family differing by 10% or less: 5 alleles for *qnrVC*, 7 alleles for *qnrA*, 9 for *qnrS*, and 71 for *qnrB* (35)(<http://www.lahey.org/qnrstudies/>, accessed 12/09/13). *qnr* genes are also found on the chromosome of both gram negative and gram positive bacteria from both clinical and environmental sources (36–38).

The sequence of pentapeptide repeat proteins can be represented as [S,T,A,V][D,N][L,F][S,T,R][G] (39). The first such protein to have its structure determined by x-ray crystallography was MfpA, encoded on the chromosome of mycobacterial species including *M. smegmatis* where its deletion increased fluoroquinolone susceptibility (40). MfpA is a dimer linked C-terminus to C-terminus and folded into a right-handed quadrilateral β helix with size, shape, and charge mimicking the β form of DNA (41). The middle, usually hydrophobic, amino acid (i) of the pentapeptide repeat and the first polar or hydrophobic residue (i–2) point inward while the remaining (i–1, i+1, i+2) amino acids are oriented outward presenting a generally anionic surface. Extensive hydrogen bonding between backbone atoms of neighboring coils stabilizes the helix. The structures of three Qnr proteins are known: EfsQnr from *Enterococcus faecalis* (42), AhQnr from *Aeromonas hydrophila* (43), and plasmid-mediated QnrB1 (44). All are rod-like dimers (Fig. 1). The monomers of QnrB1 and AhQnr have projecting loops of 8 and 12 amino acids that are important for their activity (Fig. 1). Deletion of the smaller A loop reduces quinolone protection while deletion of the larger B loop or both loops destroys protective activity (43, 44). Deletion of even a single amino acid in the larger loop compromises protective activity (45). MfpA and EfsQnr lack loops, but EfsQnr differs from MfpA in having an additional β -helical rung, a capping peptide, and a 25-amino acid flexible extension required for full protective activity.

Topoisomerases twist and untwist the DNA helix by binding to it and introducing a pair of staggered, single-strand breaks in one segment through which a second DNA segment is passed (46). Quinolones bind to the complex of enzyme and DNA stabilizing the cleavage or cleaved complex, blocking religation, and leading ultimately to lethal double stranded breaks (47). In cell-free systems QnrA (21, 48), QnrB (27, 45, 49), QnrS (50), AhQnr (43), and EfsQnr (42) have been shown to protect *E. coli* DNA gyrase from quinolone inhibition (Fig. 2). Protection of topoisomerase IV by QnrA has been demonstrated as well (51). Protection occurs at low concentrations of Qnr relative to quinolone. Fig. 2 shows that for DNA gyrase inhibited by 6 μM (2 $\mu\text{g/ml}$) ciprofloxacin, half protection required only 0.5 nM QnrB1 and some protective effect was seen with as little as 5 pM (27). At high QnrB concentrations (25–30 μM in the same system) gyrase inhibition occurs (27, 49). In contrast MfpA inhibits *M. tuberculosis* or *E. coli* gyrase with an IC_{50} of 1.75 to 3 μM and lacks any protective effect against ciprofloxacin. EfsQnr is intermediate. It partially protect *E. coli* gyrase against ciprofloxacin inhibition but also inhibits ATP-dependent supercoiling activity of gyrase with an IC_{50} of 1.2 μM (42). In a gel displacement assay QnrA binds to DNA gyrase and its GyrA and GyrB subunits and also to topoisomerase IV and its ParC and ParE subunits (48, 51). Competition between MfpA and substrate DNA for binding to gyrase has been proposed as the mechanism for its inhibitory effect (41). Qnr proteins with their additional structural features (loops, N-terminal extension) are proposed to bind to gyrase and topoisomerase IV targets in such a way as to destabilize the cleavage complex between enzyme, DNA, and quinolone causing its release, religation of DNA, and regeneration of active topoisomerase (43, 44).

Qnr Origin

Qnr homologs can be found on the chromosome of many γ -Proteobacteria, Firmicutes, and Actinomycetales, including species of *Bacillus*, *Enterococcus*, *Listeria*, and *Mycobacteria*, as well as anaerobes such as *Clostridium difficile* and *C. perfringens* (36, 38, 52, 53). Nearly 50 allelic variants have been found on the chromosome of *Stenotrophomonas maltophilia* (36, 54–57). Aquatic bacteria are especially well represented including species of *Aeromonas*, *Photobacterium*, *Shewanella*, and *Vibrio* (17, 58, 59). QnrA1 is 98% identical to the chromosomally determined Qnr of *Shewanella algae* (58), QnrS1 is 83% identical to Qnr from *Vibrio splendidus* (60), and QnrC is 72% identical to chromosomal Qnr in *Vibrio orientalis* or *V. cholerae*. (28). QnrB homologs, on the other hand, are found on the chromosome of members of the *Citrobacter freundii* complex, including *Citrobacter braakii*, *Citrobacter werkmanii* and *Citrobacter youngae* of both clinical (61) and environmental origin. The small, non-conjugative plasmids that carry *qnrD* can be found in other Enterobacteriaceae but are especially likely to be found in Proteaceae, such as *Proteus mirabilis*, *P. vulgaris*, and *Providencia rettgeri* (62) and may have originated there (63, 64).

The wide distribution of *qnr* suggests an origin well before quinolones were discovered. Indeed, *qnrB* genes and pseudogenes have been discovered on the chromosome of *Citrobacter freundii* strains collected in the 1930s (65)

Qnr Plasmids

Genes for PMQR have been found on plasmids varying in size and incompatibility specificity (Table 1), indicating that the spread of multiple plasmids has been responsible for the dissemination of this resistance around the world. Such plasmid heterogeneity may also have contributed to the variety of bacterial hosts for PMQR and indicates that plasmid acquisition of *qnr* and other quinolone resistance determinants occurred independently multiple times.

A mobile or transposable element is almost invariably associated with *qnr* genes, (Table 2 and Fig. 3). *qnrA1* has usually been associated with *ISCR1* (66), although 63% of *qnrA1*-positive *K. pneumoniae* strains in a study from South Korea were negative for *ISCR1* by PCR (67). The *ISCR1* element is not only involved in gene mobilization. It also provides an active promoter for resistance gene expression (68). Often a single copy of *ISCR1* is found upstream from *qnrA1*, but in pMG252 and related plasmids, the *qnrA1* gene is bracketed by two copies of *ISCR1* (69, 70). The *qnrA1* *ISCR1* complex is inserted in turn into a *sul1*-type integron containing several other resistance gene cassettes (24). In pSZ50 from Mexico the integron containing *ISCR1* and *qnrA1* is duplicated in tandem (71). The *qnrA3* and *qnrA6* alleles are also linked to *ISCR1*. The gene for *qnrB1*, however, is often associated not with *ISCR1* but with *orf1005*, encoding a putative transposase (27). *qnrB1* has also been found linked to an upstream truncated *orf1005* and a downstream IS26 (72, 73), while the *qnrB20* allele is sandwiched between an upstream IS26 and a downstream *orf1005* (72). Alleles *qnrB2*, *qnrB4*, *qnrB6*, and *qnrB10* are associated with *ISCR1*, usually as a single copy (73–75), but in some plasmids two copies of *ISCR1* surround *qnrB2* (76, 77). *qnrB19* has been found in three genetic environments: within large plasmids associated with *ISEcp1C*-based transposons, in large plasmids bracketed by IS26, and in small ColE1-type plasmids (~3-kb) lacking insertion sequences in which a flanking *oriT* locus and Xer recombination site have been proposed to be involved in site-specific recombination (73, 78–82). In all three settings *qnrB19* is linked to a fragment of *pspF* implying that the putative mobilization pathways may be related. In plasmid pLRM24 *qnrB19* linked to *ISEcp1* has inserted into a Tn3-like element also containing a mobile element encoding KPC-3 carbapenemase (83). The *qnrS1* gene is not linked to *ISCR1* but is associated with an upstream Tn3-like transposon, in several plasmids containing an active TEM-1 gene (26, 73, 84–86). In other plasmids *qnrS1* is associated with IS26 (87), IS2 (88), or *ISEcl2*, a novel insertion element belonging to the IS3 family (89). On the other hand *qnrS2* has been found as part of a mobile insertion cassette, an element with bracketing inverted repeats but lacking a transposase (90). The *qnrC* gene is found downstream from *ISPmi1*, an insertion sequence also belonging to the IS3 family (28). *qnrD* has typically been found on small, nonconjugative plasmids and is also located inside a mobile insertion cassette (29, 62–64, 91). *qnrVC* is so far the only *qnr* gene located in a cassette with a linked *attC* site (92). *qnrVC* genes have been found on plasmids in *A. punctata* (30), and *V. fluvialis* (32), within integrons in *A. baumannii* (93) and *P. aeruginosa* (94), and within the transmissible SXT integrating element of *V. cholerae* (34, 95).

qnr genes are usually found in multiresistance plasmids linked to other resistance determinants. β -lactamase genes have been conspicuously common, including genes for

AmpC β -lactamases (22, 96–101), CTX-M enzymes (96, 97, 99, 100, 102–114), IMP enzymes (74, 115), KPC enzymes (116–118), LAP-1 or LAP-2 (88, 89, 119–121), SHV-12 (96, 99, 104, 105, 107, 109, 115, 119, 121–127), VEB-1 (126, 128–130), and VIM-1 (111, 131). *qnrB4* and *bla*_{DHA-1} have been found near each other on similar plasmids from around the world (96, 108, 127, 132–135). *qnrB* alleles are also frequently found in plasmids linked to variable portions of the operons for *psp* (phage shock protein) and *sap* (peptide ABC transporter, ATP-binding protein) genes. These genes flank *qnrB* on the chromosome of several *Citrobacter* spp. and their co-acquisition with *qnrB* is one of the arguments for *Citrobacter* as the source of *qnrB* alleles (61). Molecular studies with I-CeuI and S1 nuclease followed by double hybridization for *qnr* and 23s rRNA genes have identified alleles for *qnrB6* (136), *qnrB12* (137), and *qnrB16* (136) on the chromosome of several *Citrobacter* spp. Note that these same alleles have also been reported on plasmids in species other than *Citrobacter* (Table 2).

Spread of *qnr* plasmids

PMQR genes have been found in a variety of Enterobacteriaceae, especially *Escherichia coli* and species of *Enterobacter*, *Klebsiella*, and *Salmonella* (Table 2). They have been conspicuously rare in non-fermenters but have occasionally been reported in *Pseudomonas aeruginosa*, other *Pseudomonas* spp. (138, 139), *Acinetobacter baumannii* (139, 140), and *Stenotrophomonas maltophilia* (139). Genes for *qnrA1* and *qnrB6* have also been found in *Haemophilus parasuis* from pigs in South China (141). *qnr* genes are found in a variety of gram-positive organisms but are chromosomal and not plasmid-mediated (38, 53). Of the various *qnr* varieties, *qnrB* seems somewhat more common than *qnrA* or *qnrS*, which are more common than *qnrD*. Only a single isolate of *qnrC* is known (28). The relative frequency of various alleles can be judged by the number of references in Table 2: i.e. for *qnrB* the most frequently detected alleles are *qnrB1*, *qnrB2*, *qnrB4*, *qnrB6*, *qnrB10*, and *qnrB19*. The earliest known *qnr* outside of *Citrobacter* spp, dates from 1988 (142). Studies in the last decade suggest that *qnr* detection is increasing, but is still less than 10% in unselected clinical isolates, with usually greatest prevalence in *E. cloacae*, less in *K. pneumoniae*, and least in *E. coli* (127, 143–146). Higher frequencies result if samples are preselected for ESBL or other resistance phenotypes (127, 147), but the prevalence of *qnr* genes has reached as high as 39% in an unselected sample of *E. cloacae* isolates at one hospital in China (127).

Although most prevalence studies have surveyed hospital isolates, animals have not been neglected. PMQR genes have been found in samples from domestic or wild birds (75, 86, 148, 149), cats (75, 81, 112, 150), cattle (151, 152), chickens (75, 86, 153–160), dogs (75, 81, 112, 150, 156, 161), ducks (101, 156, 160, 162), fish (163–165), geese (101, 156, 160, 162), horses (81, 86, 166), pigs (75, 101, 155, 156, 160, 162, 167), rabbits (168), reptiles (155, 169), sheep (155), turkeys (155), and zoo animals (170–172).

Regulation of *qnr*

Not surprisingly, *qnrA* expression is influenced by the strength of its promoter. A *qnrA1* plasmid from Shanghai was found to give 8-fold higher ciprofloxacin MICs than other

qnrA1 plasmids and had a 12-fold stronger promoter attributed to a 7-bp deletion between the +1 transcription initiation site and the start of the *qnrA1* gene (144). Environmental conditions have also been found to affect expression of *qnr* genes and may offer clues concerning the native function of these genes. Expression of the *qnrA* gene of *S. algae*, an organism adapted to growth at low temperature, is stimulated up to 8-fold by cold shock but not by other conditions such as DNA damage, oxidative or osmotic stress, starvation, or heat shock (173). Expression of *qnrB* alleles, on the other hand, is augmented up to 9-fold by exposure to DNA damaging agents such as ciprofloxacin or mitomycin C via an upstream LexA binding site and the classical SOS system (174, 175). *qnrD* and the chromosomal *qnr* of *S. marcescens* are similarly regulated (176). Expression of plasmid-mediated *qnrS1* or the related chromosomal *qnrVS1* of *V. splendidus* is also stimulated by ciprofloxacin up to 30-fold, but by a mechanism independent of the SOS system. No LexA binding site is found upstream from these *qnr* genes, but upstream sequence is required for quinolone stimulation to occur (177). Some naturally occurring quinolone-like compounds such as quinine, 2-hydroxyquinoline, 4-hydroxyquinoline, or the *Pseudomonas* quinolone signal for quorum sensing also induce *qnrS1*, but not *qnrVS1* (178).

AAC(6')-Ib-cr

AAC(6')-Ib-cr is a bifunctional variant of a common acetyltransferase active on such aminoglycosides as amikacin, kanamycin, and tobramycin but also able to acetylate those fluoroquinolones with an amino nitrogen on the piperazinyl ring, such as ciprofloxacin and norfloxacin (10). Compared to other AAC(6')-Ib enzymes, the -cr variant has two unique amino acid substitutions: Trp102Arg and Asp179Tyr, both of which are required for quinolone acetylating activity. Models of enzyme action suggest that the Asp179Tyr replacement is particularly important in permitting π -stacking interactions with the quinolone ring to facilitate quinolone binding. The role of Trp102Arg is to position the Tyr face for optimal interaction (179) or to hydrogen bond to keto or carboxyl groups of the quinolone to fix it in place (180). Both AGG and CGG have been found as the Arg codon at 102 allowing variants of the *aac(6')-Ib-cr* gene to be distinguished (73). A lower level of *aac(6')-Ib-cr* expression has been found in a strain with an upstream 12-bp deletion displacing the promoter -10 box (181). A 26 amino acid larger AAC(6')-Ib-cr4 enzyme with consequent Trp128Arg and Asp205Tyr substitutions (182) and non-functional truncated *aac(6')-Ib-cr* genes (183) have also been reported.

The *aac(6')-Ib-cr* gene is usually found in a cassette as part of an integron in a multiresistance plasmid, which may contain other PMQR genes. Association with ESBL CTX-M-15 is particularly common (110, 184–190). A mobile genetic element, especially IS26, is often associated (191). *aac(6')-Ib-cr* may also be chromosomal (192, 193). The gene has been found world-wide (Table 2) in a variety of Enterobacteriaceae and even in *P. aeruginosa* (138). It is more prevalent in *E. coli* than other Enterobacteriaceae (145, 184, 194, 195), and has often been more common than *qnr* alleles (14, 75, 184, 192, 194, 196).

QepA and OqxAB

QepA is a plasmid-mediated efflux pump in the major facilitator (MFS) family that decreases susceptibility to hydrophilic fluoroquinolones, especially ciprofloxacin and norfloxacin (11, 197). *qepA* has often been found on plasmids also encoding aminoglycoside ribosomal methylase *rmtB* (12, 167, 198). Substantial differences in quinolone resistance produced by different *qepA* transconjugants suggest variability in the level of *qepA* expression, by mechanisms as yet to be defined (167). IS26 elements and ISCR3C have been implicated in mobilizing the *qepA* gene to plasmids (199). A variant differing in two amino acids (QepA2) has also been described (199).

OqxAB is an efflux pump in the resistance-nodulation-division (RND) family that was initially recognized on transmissible plasmids responsible for resistance to olaquinox used for growth enhancement in pigs (200, 201). It has a wide substrate specificity including chloramphenicol, trimethoprim, and quinolones such as ciprofloxacin, flumequin, norfloxacin, and nalidixic acid (13). *oqxAB* has been found on plasmids in clinical isolates of *E. coli* and *K. pneumoniae* and in the chromosome and on plasmids of *S. enteritis* flanked in both locations by IS26-like elements (202–207). In *E. coli* isolates from farms in China where olaquinox was in use, *oqxAB* was found on transmissible plasmids in 39% of isolates from animals and 30% of isolates from farm workers (204). Linkage of *oqxAB* with genes for CTX-M-14 and other plasmid-mediated CTX-M alleles has been noted (160). It is common (usually 75% or more) on the chromosome of *K. pneumoniae* isolates, where up to 20-fold variation in expression implies the presence of regulatory control (73, 203, 206, 208, 209). In *K. pneumoniae* overexpression of the nearby *rarA* gene is associated with increased *oqxAB* expression, while increased expression of adjacent *oqxR* gene downregulates OqxAB production (210, 211). Sequence variants *oqxA2*, *oqxB2* and *oqxB3* have been described (208).

Other plasmid-mediated efflux pumps active on quinolones have been reported but as yet little studied. Plasmid pRSB101 isolated from an uncultivated organism in activated sludge at a wastewater treatment plant contained a MDR transport system with an RND-type membrane fusion protein conferring resistance to nalidixic acid and norfloxacin (212). It differs in sequence from QepA and OqxAB. Plasmids in *S. aureus* (especially MRSA) encoding the QacBIII variant belonging to the MFS family confer decreased susceptibility to norfloxacin and ciprofloxacin (213).

Resistance produced by PMQR determinants

Table 3 shows the minimum inhibitory concentration (MIC) produced in a common *E. coli* strain by PMQR genes. *qnr* genes produce about the same resistance to ciprofloxacin and levofloxacin as single mutations in *gyrA*, but have less effect on susceptibility to nalidixic acid. *aac(6')-Ib-cr* and *qepA* give lower levels, which is confined to ciprofloxacin in the case of *aac(6')-Ib-cr* because of its substrate specificity. All provide a decrease in susceptibility that does not reach the CLSI breakpoint for even intermediate resistance. How then can PMQR genes be clinically important?

The answer is that PMQR genes facilitate the selection of higher levels of quinolone resistance. Fig. 4 shows the effect of plasmid pMG252 encoding QnrA on survival of *E. coli* J53 at increasing concentrations of ciprofloxacin. Survivors occur until a concentration of more than 1 µg/ml ciprofloxacin is reached. This limiting concentration has been termed the mutant prevention concentration (MPC), and the concentration between the MIC and MPC at which mutants are selected is the mutant selection window (214). PMQR genes exert their influence by widening the mutant selection window and elevating the MPC, as shown for both *qnr* (215, 216) and *aac(6')-Ib-cr* (10). The same augmentation of resistance selection has been found in clinical isolates of *E. cloacae* (147). Surprisingly, in *qnr*-harboring *E. coli* *gyrA* resistance mutants are rarely selected (217), although resistance produced by *qnr* and *gyrA* is additive (218–220), clinical isolates with *qnr* alleles or *aac(6')-Ib-cr* and *gyrA*, *parC*, and other resistance mutations are common (221), and *gyrA* mutants have developed in *qnr* positive isolates from patients treated with quinolone for *E. coli* (222) or *S. enterica* (223) infections.

It should be noted that higher levels of quinolone resistance are seen if a plasmid or strain carries two or more genes for quinolone resistance, such as both *qnr* and *aac(6')-Ib-cr*, and that ciprofloxacin MICs of 2 µg/ml can be reached with *qnrA* in *E. coli* overexpressing the AcrAB multi-drug efflux pump (224). A fully resistant *E. coli* with a ciprofloxacin MIC of 4 µg/ml has been reported with plasmid-mediated *qnrS1* and *oqxAB* as well as overexpression of AcrAB and other efflux pumps (225).

While the frequency of quinolone resistance in clinical isolates has paralleled quinolone usage, the appearance of PMQR has also played a role. At Hadassah hospitals in Israel ciprofloxacin resistance was uncommon in *E. coli*, *K. pneumoniae*, and *Enterobacter* spp. until the mid-1990s, just when *qnr* and *aac(6')-Ib-cr* genes became prevalent in these strains (143, 226). Similarly, in Korea the increasing frequency of ciprofloxacin resistance in Enterobacteriaceae since 2000 has been associated with an increasing prevalence of PMQR genes (145). In Spain also the prevalence of PMQR in clinical isolates of *E. coli* and *K. pneumoniae* increased between 2000 and 2006 (227). In Canada as well the prevalence of *aac(6')-Ib-cr* increased in the Calgary Health Region between 2004 and 2007 (184).

Clinical importance

In animal model infections the presence of a *qnr* gene makes an infecting agent harder to treat with quinolones. This detrimental effect has been shown in mice with pneumonia produced by *K. pneumoniae* or *E. coli* (228, 229) and in *E. coli* UTI models (230, 231).

Patients treated with levofloxacin for bloodstream infections caused by gram-negative organisms with elevated quinolone MICs that were still within the susceptible category had worse outcomes than similar patients infected with more susceptible organisms (232), but a specific effect of PMQR carriage on outcome has been harder to document. Chong et al. evaluated 351 blood culture isolates of *Enterobacter* or *Klebsiella* at a health center in Korea and found 26 positive for *qnrA*, *qnrB*, or *qnrS* genes. The *qnr* positive patients were hospitalized longer, but there was no difference in in-hospital or 30-day mortality between the *qnr* positive or negative patient groups (233). Liao et al. studied 227 blood culture

isolates of *Klebsiella* from a hospital in Taiwan and found nine positive for *qnrB* or *qnrS*. 14-day mortality was similar in patients infected with or without *qnr*-containing isolates, but there was a trend for increased in-hospital mortality (234). Further studies are needed to distinguish a specific effect of *qnr* from the effect of other resistance genes so often linked to it.

Coda

The varieties of PMQR that eventually emerged exemplify the three general mechanism of bacterial resistance to any antimicrobial agents: target alteration (Qnr), drug modification (AAC(6′)-Ib-cr), and efflux pump activation (QepA and OqxAB). AAC(6′)-Ib-cr arose by mutations altering two amino acid encoded by a common aminoglycoside resistance gene. Acquisition of the other PMQR genes illustrates the variety of genetic elements now available on plasmids for resistance gene mobilization and the frequent obscurity of the ultimate gene donors. Who would have guessed that aquatic bacteria harbored *qnr* genes without evident selective advantage thus encoding a protein that blocked the action of a synthetic antimicrobial agent that they would never be expected to encounter? Unfortunately, knowledge of resistance mechanisms has not led to new therapeutic strategies. Acetylation of ciprofloxacin is inhibited competitively by aminoglycoside substrates of AAC(6′)-Ib-cr (179), but other inhibitors of the enzyme are not yet known and no efflux pump inhibitors are commercially available. Overcoming Qnr blockage of quinolone inhibition will require deeper knowledge of how this DNA mimic interacts with topoisomerase as well as the development of other bacterial topoisomerase poisons that escape the action of Qnr.

Acknowledgments

This work was supported by grant R01 AI057576 from the US Public Health Service, National Institutes of Health to D.C.H. and G.A.J.

References

1. Gellert M, Mizuuchi K, O’Dea MH, Itoh T, Tomizawa JI. Nalidixic acid resistance: a second genetic character involved in DNA gyrase activity. *Proc Natl Acad Sci USA*. 1977; 74:4772–4776. [PubMed: 337300]
2. Yoshida H, Kojima T, Yamagishi J, Nakamura S. Quinolone-resistant mutations of the *gyrA* gene of *Escherichia coli*. *Mol Gen Genet*. 1988; 211:1–7. [PubMed: 2830458]
3. Hooper, DC. Mechanisms of quinolone resistance. In: Hooper, DC.; Rubinstein, E., editors. *Quinolone antimicrobial agents*. 3. ASM Press; Washington, D.C.: 2003. p. 41-67.
4. Burman LG. Apparent absence of transferable resistance to nalidixic acid in pathogenic Gram-negative bacteria. *J Antimicrob Chemother*. 1977; 3:509–516. [PubMed: 332684]
5. Neuhauser MM, Weinstein RA, Rydman R, Danziger LH, Karam G, Quinn JP. Antibiotic resistance among gram-negative bacilli in US intensive care units: implications for fluoroquinolone use. *JAMA*. 2003; 289:885–888. [PubMed: 12588273]
6. Linder JA, Huang ES, Steinman MA, Gonzales R, Stafford RS. Fluoroquinolone prescribing in the United States: 1995 to 2002. *Am J Med*. 2005; 118:259–268. [PubMed: 15745724]
7. Munshi MH, Sack DA, Haider K, Ahmed ZU, Rahaman MM, Morshed MG. Plasmid-mediated resistance to nalidixic acid in *Shigella dysenteriae* type 1. *Lancet*. 1987; 2:419–421. [PubMed: 2887725]

8. Ashraf MM, Ahmed ZU, Sack DA. Unusual association of a plasmid with nalidixic acid resistance in an epidemic strain of *Shigella dysenteriae* type 1 from Asia. *Can J Microbiol.* 1991; 37:59–63. [PubMed: 2021898]
9. Martínez-Martínez L, Pascual A, Jacoby GA. Quinolone resistance from a transferable plasmid. *Lancet.* 1998; 351:797–799. [PubMed: 9519952]
10. Robicsek A, Strahilevitz J, Jacoby GA, Macielag M, Abbanat D, Park CH, Bush K, Hooper DC. Fluoroquinolone-modifying enzyme: a new adaptation of a common aminoglycoside acetyltransferase. *Nat Med.* 2006; 12:83–88. [PubMed: 16369542]
11. Yamane K, Wachino J, Suzuki S, Kimura K, Shibata N, Kato H, Shibayama K, Konda T, Arakawa Y. New plasmid-mediated fluoroquinolone efflux pump, QepA, found in an *Escherichia coli* clinical isolate. *Antimicrob Agents Chemother.* 2007; 51:3354–3360. [PubMed: 17548499]
12. Périchon B, Courvalin P, Galimand M. Transferable resistance to aminoglycosides by methylation of G1405 in 16S rRNA and to hydrophilic fluoroquinolones by QepA-mediated efflux in *Escherichia coli*. *Antimicrob Agents Chemother.* 2007; 51:2464–2469. [PubMed: 17470656]
13. Hansen LH, Jensen LB, Sorensen HI, Sorensen SJ. Substrate specificity of the OqxAB multidrug resistance pump in *Escherichia coli* and selected enteric bacteria. *J Antimicrob Chemother.* 2007; 60:145–147. [PubMed: 17526501]
14. Ciesielczuk H, Hornsey M, Choi V, Woodford N, Wareham DW. Development and evaluation of a multiplex PCR for eight plasmid-mediated quinolone-resistance determinants. *J Med Microbiol.* 2013; 62:1823–1827. [PubMed: 24000223]
15. Guan X, Xue X, Liu Y, Wang J, Wang Y, Wang J, Wang K, Jiang H, Zhang L, Yang B, Wang N, Pan L. Plasmid-mediated quinolone resistance—current knowledge and future perspectives. *J Int Med Res.* 2013; 41:20–30. [PubMed: 23569126]
16. Ruiz J, Pons MJ, Gomes C. Transferable mechanisms of quinolone resistance. *Int J Antimicrob Agents.* 2012; 40:196–203. [PubMed: 22831841]
17. Poirel L, Cattoir V, Nordmann P. Plasmid-mediated quinolone resistance; interactions between human, animal, and environmental ecologies. *Front Microbiol.* 2012; 3:24. [PubMed: 22347217]
18. Rodríguez-Martínez JM, Cano ME, Velasco C, Martínez-Martínez L, Pascual A. Plasmid-mediated quinolone resistance: an update. *J Infect Chemother.* 2011; 17:149–182. [PubMed: 20886256]
19. Hernandez A, Sanchez MB, Martinez JL. Quinolone resistance: much more than predicted. *Front Microbiol.* 2011; 2 article 22.
20. Strahilevitz J, Jacoby GA, Hooper DC, Robicsek A. Plasmid-mediated quinolone resistance: a multifaceted threat. *Clin Microbiol Rev.* 2009; 22:664–689. [PubMed: 19822894]
21. Tran JH, Jacoby GA. Mechanism of plasmid-mediated quinolone resistance. *Proc Natl Acad Sci U S A.* 2002; 99:5638–5642. [PubMed: 11943863]
22. Rodríguez-Martínez JM, Pascual A, García I, Martínez-Martínez L. Detection of the plasmid-mediated quinolone resistance determinant *qnr* among clinical isolates of *Klebsiella pneumoniae* producing AmpC-type β -lactamase. *J Antimicrob Chemother.* 2003; 52:703–706. [PubMed: 12951342]
23. Wang M, Sahn DF, Jacoby GA, Hooper DC. Emerging plasmid-mediated quinolone resistance associated with the *qnr* gene in *Klebsiella pneumoniae* clinical isolates in the United States. *Antimicrob Agents Chemother.* 2004; 48:1295–1299. [PubMed: 15047532]
24. Wang M, Tran JH, Jacoby GA, Zhang Y, Wang F, Hooper DC. Plasmid-mediated quinolone resistance in clinical isolates of *Escherichia coli* from Shanghai, China. *Antimicrob Agents Chemother.* 2003; 47:2242–2248. [PubMed: 12821475]
25. Cheung TK, Chu YW, Chu MY, Ma CH, Yung RW, Kam KM. Plasmid-mediated resistance to ciprofloxacin and cefotaxime in clinical isolates of *Salmonella enterica* serotype Enteritidis in Hong Kong. *J Antimicrob Chemother.* 2005; 56:586–589. [PubMed: 16033804]
26. Hata M, Suzuki M, Matsumoto M, Takahashi M, Sato K, Ibe S, Sakae K. Cloning of a novel gene for quinolone resistance from a transferable plasmid in *Shigella flexneri* 2b. *Antimicrob Agents Chemother.* 2005; 49:801–803. [PubMed: 15673773]
27. Jacoby GA, Walsh KE, Mills DM, Walker VJ, Oh H, Robicsek A, Hooper DC. *qnrB*, another plasmid-mediated gene for quinolone resistance. *Antimicrob Agents Chemother.* 2006; 50:1178–1182. [PubMed: 16569827]

28. Wang M, Guo Q, Xu X, Wang X, Ye X, Wu S, Hooper DC, Wang M. New plasmid-mediated quinolone resistance gene, *qnrC*, found in a clinical isolate of *Proteus mirabilis*. *Antimicrob Agents Chemother*. 2009; 53:1892–1897. [PubMed: 19258263]
29. Cavaco LM, Hasman H, Xia S, Aarestrup FM. *qnrD*, a novel gene conferring transferable quinolone resistance in *Salmonella enterica* serovar Kentucky and Bovismorbificans strains of human origin. *Antimicrob Agents Chemother*. 2009; 53:603–608. [PubMed: 19029321]
30. Xia R, Guo X, Zhang Y, Xu H. *qnrVC*-like gene located in a novel complex class 1 integron harboring the *ISCR1* element in an *Aeromonas punctata* strain from an aquatic environment in Shandong Province, China. *Antimicrob Agents Chemother*. 2010; 54:3471–3474. [PubMed: 20516288]
31. Ming L. Mechanisms of fluoroquinolone resistance in *Vibrio parahaemolyticus*. 2013 GenBank KC540630.
32. Singh R, Rajpara N, Tak J, Patel A, Mohanty P, Vinothkumar K, Chowdhury G, Ramamurthy T, Ghosh A, Bhardwaj AK. Clinical isolates of *Vibrio fluvialis* from Kolkata, India, obtained during 2006: plasmids, the *qnr* gene and a mutation in gyrase A as mechanisms of multidrug resistance. *J Med Microbiol*. 2012; 61:369–374. [PubMed: 22034163]
33. Fonseca EL, Vicente AC. Epidemiology of *qnrVC* alleles and emergence out of the Vibrionaceae family. *J Med Microbiol*. 2013; 62:1628–1630. [PubMed: 23800600]
34. Kim HB, Wang M, Ahmed S, Park CH, LaRocque RC, Faruque AS, Salam MA, Khan WA, Qadri F, Calderwood SB, Jacoby GA, Hooper DC. Transferable quinolone resistance in *Vibrio cholerae*. *Antimicrob Agents Chemother*. 2010; 54:799–803. [PubMed: 19949057]
35. Jacoby G, Cattoir V, Hooper D, Martínez-Martínez L, Nordmann P, Pascual A, Poirel L, Wang M. *qnr* gene nomenclature. *Antimicrob Agents Chemother*. 2008; 52:2297–2299. [PubMed: 18426894]
36. Sánchez MB, Hernández A, Rodríguez-Martínez JM, Martínez-Martínez L, Martínez JL. Predictive analysis of transmissible quinolone resistance indicates *Stenotrophomonas maltophilia* as a potential source of a novel family of Qnr determinants. *BMC Microbiol*. 2008; 8:148–161. [PubMed: 18793450]
37. Boulund F, Johnning A, Pereira MB, Larsson DG, Kristiansson E. A novel method to discover fluoroquinolone antibiotic resistance (*qnr*) genes in fragmented nucleotide sequences. *BMC Genomics*. 2012; 13:695. [PubMed: 23231464]
38. Jacoby GA, Hooper DC. Phylogenetic analysis of chromosomally determined Qnr and related proteins. *Antimicrob Agents Chemother*. 2013; 57:1930–1934. [PubMed: 23318805]
39. Vetting MW, Hegde SS, Fajardo JE, Fiser A, Roderick SL, Takiff HE, Blanchard JS. Pentapeptide repeat proteins. *Biochemistry*. 2006; 45:1–10. [PubMed: 16388575]
40. Montero C, Mateu G, Rodriguez R, Takiff H. Intrinsic resistance of *Mycobacterium smegmatis* to fluoroquinolones may be influenced by new pentapeptide protein MfpA. *Antimicrob Agents Chemother*. 2001; 45:3387–3392. [PubMed: 11709313]
41. Hegde SS, Vetting MW, Roderick SL, Mitchenall LA, Maxwell A, Takiff HE, Blanchard JS. A fluoroquinolone resistance protein from *Mycobacterium tuberculosis* that mimics DNA. *Science*. 2005; 308:1480–1483. [PubMed: 15933203]
42. Hegde SS, Vetting MW, Mitchenall LA, Maxwell A, Blanchard JS. Structural and biochemical analysis of the pentapeptide repeat protein *EfsQnr*, a potent DNA gyrase inhibitor. *Antimicrob Agents Chemother*. 2011; 55:110–117. [PubMed: 20937785]
43. Xiong X, Bromley EH, Oelschlaeger P, Woolfson DN, Spencer J. Structural insights into quinolone antibiotic resistance mediated by pentapeptide repeat proteins: conserved surface loops direct the activity of a Qnr protein from a gram-negative bacterium. *Nucleic Acids Res*. 2011; 39:3917–3927. [PubMed: 21227918]
44. Vetting MW, Hegde SS, Wang M, Jacoby GA, Hooper DC, Blanchard JS. Structure of QnrB1, a plasmid-mediated fluoroquinolone resistance factor. *J Biol Chem*. 2011; 286:25265–25273. [PubMed: 21597116]
45. Jacoby GA, Corcoran MA, Mills DM, Griffin CM, Hooper DC. Mutational analysis of quinolone resistance protein QnrB1. *Antimicrob Agents Chemother*. 2013; 57:5733–5736. [PubMed: 23979738]

46. Kampranis SC, Bates AD, Maxwell A. A model for the mechanism of strand passage by DNA gyrase. *Proc Natl Acad Sci U S A*. 1999; 96:8414–8419. [PubMed: 10411889]
47. Drlica K, Malik M, Kerns RJ, Zhao X. Quinolone-mediated bacterial death. *Antimicrob Agents Chemother*. 2008; 52:385–392. [PubMed: 17724149]
48. Tran JH, Jacoby GA, Hooper DC. Interaction of the plasmid-encoded quinolone resistance protein Qnr with *Escherichia coli* DNA gyrase. *Antimicrob Agents Chemother*. 2005; 49:118–125. [PubMed: 15616284]
49. Mérens A, Matrat S, Aubry A, Lascols C, Jarlier V, Soussy CJ, Cavallo JD, Cambau E. The pentapeptide repeat proteins MfpA_{MI} and QnrB4 exhibit opposite effects on DNA gyrase catalytic reactions and on the ternary gyrase-DNA-quinolone complex. *J Bacteriol*. 2009; 191:1587–1594. [PubMed: 19060136]
50. Tavio MM, Jacoby GA, Hooper DC. Structural determinants of QnrS1 resistance. 52nd Intersci Conf Antimicrob Agents Chemother C1-683a. 2012
51. Tran JH, Jacoby GA, Hooper DC. Interaction of the plasmid-encoded quinolone resistance protein QnrA with *Escherichia coli* topoisomerase IV. *Antimicrob Agents Chemother*. 2005; 49:3050–3052. [PubMed: 15980397]
52. Arsène S, Leclercq R. Role of a *qnr*-like gene in the intrinsic resistance of *Enterococcus faecalis* to fluoroquinolones. *Antimicrob Agents Chemother*. 2007; 51:3254–3258. [PubMed: 17620379]
53. Rodríguez-Martínez JM, Velasco C, Briaes A, García I, Conejo MC, Pascual A. Qnr-like pentapeptide repeat proteins in gram-positive bacteria. *J Antimicrob Chemother*. 2008; 61:1240–1243. [PubMed: 18343805]
54. Shimizu K, Kikuchi K, Sasaki T, Takahashi N, Ohtsuka M, Ono Y, Hiramatsu K. Smqnr, a new chromosome-carried quinolone resistance gene in *Stenotrophomonas maltophilia*. *Antimicrob Agents Chemother*. 2008; 52:3823–3825. [PubMed: 18644963]
55. Sánchez MB, Martínez JL. SmQnr contributes to intrinsic resistance to quinolones in *Stenotrophomonas maltophilia*. *Antimicrob Agents Chemother*. 2010; 54:580–581. [PubMed: 19841154]
56. Gordon NC, Wareham DW. Novel variants of the Smqnr family of quinolone resistance genes in clinical isolates of *Stenotrophomonas maltophilia*. *J Antimicrob Chemother*. 2010; 65:483–489. [PubMed: 20071366]
57. Zhang R, Sun Q, Hu YJ, Yu H, Li Y, Shen Q, Li GX, Cao JM, Yang W, Wang Q, Zhou HW, Hu YY, Chen GX. Detection of the Smqnr quinolone protection gene and its prevalence in clinical isolates of *Stenotrophomonas maltophilia* in China. *J Med Microbiol*. 2012; 61:535–539. [PubMed: 22096133]
58. Poirel L, Rodriguez-Martinez JM, Mammeri H, Liard A, Nordmann P. Origin of plasmid-mediated quinolone resistance determinant QnrA. *Antimicrob Agents Chemother*. 2005; 49:3523–3525. [PubMed: 16048974]
59. Poirel L, Liard A, Rodriguez-Martinez JM, Nordmann P. Vibrionaceae as a possible source of Qnr-like quinolone resistance determinants. *J Antimicrob Chemother*. 2005; 56:1118–1121. [PubMed: 16227349]
60. Cattoir V, Poirel L, Mazel D, Soussy CJ, Nordmann P. *Vibrio splendidus* as the source of plasmid-mediated QnrS-like quinolone resistance determinants. *Antimicrob Agents Chemother*. 2007; 51:2650–2651. [PubMed: 17452482]
61. Jacoby GA, Griffin CM, Hooper DC. *Citrobacter* spp. as a source of qnrB alleles. *Antimicrob Agents Chemother*. 2011; 55:4979–4984. [PubMed: 21844311]
62. Zhang S, Sun J, Liao XP, Hu QJ, Liu BT, Fang LX, Deng H, Ma J, Xiao X, Zhu HQ, Liu YH. Prevalence and plasmid characterization of the qnrD determinant in *Enterobacteriaceae* isolated from animals, retail meat products, and humans. *Microb Drug Resist*. 2013; 19:331–335. [PubMed: 23557071]
63. Guillard T, Cambau E, Neuwirth C, Nenninger T, Mbadi A, Brasme L, Vernet-Garnier V, Bajolet O, de Champs C. Description of a 2,683-base-pair plasmid containing qnrD in two *Providencia rettgeri* isolates. *Antimicrob Agents Chemother*. 2012; 56:565–568. [PubMed: 21986831]

64. Guillard T, Grillon A, de Champs C, Cartier C, Madoux J, Lozniewski A, Berçot B, Riahi J, Vernet-Garnier V, Cambau E. Mobile insertion cassetts as a source of *qnrD* mobilization onto small non-transmissible plasmids in. *Proteaeae PLoS One*. (in press).
65. Saga T, Sabtcheva S, Mitsutake K, Ishii Y, Tateda K, Yamaguchi K, Kaku M. Characterization of *qnrB*-like genes in *Citrobacter* species of the American Type Culture Collection. *Antimicrob Agents Chemother*. 2013; 57:2863–2866. [PubMed: 23529729]
66. Toleman MA, Bennett PM, Walsh TR. *ISCR* elements: novel gene-capturing systems of the 21st century? *Microbiol Mol Biol Rev*. 2006; 70:296–316. [PubMed: 16760305]
67. Park YJ, Yu JK, Kim SY, Lee S, Jeong SH. Prevalence and characteristics of *qnr* determinants and *aac(6')-Ib-cr* among ciprofloxacin-susceptible isolates of *Klebsiella pneumoniae* in Korea. *J Antimicrob Chemother*. 2010; 65:2041–2043. [PubMed: 20615929]
68. Rodriguez-Martinez JM, Poirel L, Canton R, Nordmann P. Common region CR1 for expression of antibiotic resistance genes. *Antimicrob Agents Chemother*. 2006; 50:2544–2546. [PubMed: 16801441]
69. Robicsek A, Jacoby GA, Hooper DC. The worldwide emergence of plasmid-mediated quinolone resistance. *Lancet Infect Dis*. 2006; 6:629–640. [PubMed: 17008172]
70. Rodriguez-Martinez JM, Velasco C, Garcia I, Cano ME, Martinez-Martinez L, Pascual A. Characterisation of integrons containing the plasmid-mediated quinolone resistance gene *qnrA1* in *Klebsiella pneumoniae*. *Int J Antimicrob Agents*. 2007; 29:705–709. [PubMed: 17368003]
71. Garza-Ramos U, Barrios H, Hernandez-Vargas MJ, Rojas-Moreno T, Reyna-Flores F, Tinoco P, Othon V, Poirel L, Nordmann P, Cattoir V, Ruiz-Palacios G, Fernandez JL, Santamaria RI, Bustos P, Castro N, Silva-Sanchez J. Transfer of quinolone resistance gene *qnrA1* to *Escherichia coli* through a 50 kb conjugative plasmid resulting from the splitting of a 300 kb plasmid. *J Antimicrob Chemother*. 2012; 67:1627–1634. [PubMed: 22514263]
72. Teo JW, Ng KY, Lin RT. Detection and genetic characterisation of *qnrB* in hospital isolates of *Klebsiella pneumoniae* in Singapore. *Int J Antimicrob Agents*. 2009; 33:177–180. [PubMed: 18993034]
73. Andres P, Lucero C, Soler-Bistue A, Guerriero L, Albornoz E, Tran T, Zorreguieta A, Group P, Galas M, Corso A, Tolmasky ME, Petroni A. Differential distribution of plasmid-mediated quinolone resistance genes in clinical enterobacteria with unusual phenotypes of quinolone susceptibility from Argentina. *Antimicrob Agents Chemother*. 2013; 57:2467–2675. [PubMed: 23478955]
74. Espedido BA, Partridge SR, Iredell JR. *bla_{IMP-4}* in different genetic contexts in *Enterobacteriaceae* isolates from Australia. *Antimicrob Agents Chemother*. 2008; 52:2984–2987. [PubMed: 18490506]
75. Ma J, Zeng Z, Chen Z, Xu X, Wang X, Deng Y, Lu D, Huang L, Zhang Y, Liu J, Wang M. High prevalence of plasmid-mediated quinolone resistance determinants *qnr*, *aac(6')-Ib-cr* and *qepA* among ceftiofur-resistant *Enterobacteriaceae* isolates from companion and food-producing animals. *Antimicrob Agents Chemother*. 2009; 53:519–524. [PubMed: 18936192]
76. Chen YT, Liao TL, Liu YM, Lauderdale TL, Yan JJ, Tsai SF. Mobilization of *qnrB2* and *ISCR1* in plasmids. *Antimicrob Agents Chemother*. 2009; 53:1235–1237. [PubMed: 19075060]
77. Garnier F, Raked N, Gassama A, Denis F, Ploy MC. Genetic environment of quinolone resistance gene *qnrB2* in a complex *sul1*-type integron in the newly described *Salmonella enterica* serovar Keurmassar. *Antimicrob Agents Chemother*. 2006; 50:3200–3202. [PubMed: 16940131]
78. Cattoir V, Nordmann P, Silva-Sanchez J, Espinal P, Poirel L. *ISEcp1*-mediated transposition of *qnrB*-like gene in *Escherichia coli*. *Antimicrob Agents Chemother*. 2008; 52:2929–2932. [PubMed: 18519717]
79. Dionisi AM, Lucarelli C, Owczarek S, Luzzi I, Villa L. Characterization of the plasmid-borne quinolone resistance gene *qnrB19* in *Salmonella enterica* serovar Typhimurium. *Antimicrob Agents Chemother*. 2009; 53:4019–4021. [PubMed: 19528272]
80. Hordijk J, Bosman AB, van Essen-Zandbergen A, Veldman K, Dierikx C, Wagenaar JA, Mevius D. *qnrB19* gene bracketed by *IS26* on a 40-kilobase *IncR* plasmid from an *Escherichia coli* isolate from a veal calf. *Antimicrob Agents Chemother*. 2011; 55:453–454. [PubMed: 20956595]

81. Schink AK, Kadlec K, Schwarz S. Detection of *qnr* genes among *Escherichia coli* isolates of animal origin and complete sequence of the conjugative *qnrB19*-carrying plasmid pQNR2078. *J Antimicrob Chemother.* 2012; 67:1099–1102. [PubMed: 22334601]
82. Tran T, Andres P, Petroni A, Soler-Bistue A, Albornoz E, Zorreguieta A, Reyes-Lamoth R, Sherratt DJ, Corso A, Tolmasky ME. Small plasmids harboring *qnrB19*: a model for plasmid evolution mediated by site-specific recombination at *oriT* and Xer sites. *Antimicrob Agents Chemother.* 2012; 56:1821–1827. [PubMed: 22290975]
83. Rice LB, Carias LL, Hutton RA, Rudin SD, Endimiani A, Bonomo RA. The KQ element, a complex genetic region conferring transferable resistance to carbapenems, aminoglycosides, and fluoroquinolones in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother.* 2008; 52:3427–3429. [PubMed: 18573935]
84. Kehrenberg C, Friederichs S, de Jong A, Michael GB, Schwarz S. Identification of the plasmid-borne quinolone resistance gene *qnrS* in *Salmonella enterica* serovar Infantis. *J Antimicrob Chemother.* 2006; 58:18–22. [PubMed: 16720566]
85. Kehrenberg C, Hopkins KL, Threlfall EJ, Schwarz S. Complete nucleotide sequence of a small *qnrS1*-carrying plasmid from *Salmonella enterica* subsp. enterica Typhimurium DT193. *J Antimicrob Chemother.* 2007; 60:903–905. [PubMed: 17652106]
86. Dolejska M, Villa L, Hasman H, Hansen L, Carattoli A. Characterization of IncN plasmids carrying *bla*_{CTX-M-1} and *qnr* genes in *Escherichia coli* and *Salmonella* from animals, the environment and humans. *J Antimicrob Chemother.* 2013; 68:333–339. [PubMed: 23060365]
87. Chen YT, Shu HY, Li LH, Liao TL, Wu KM, Shiau YR, Yan JJ, Su IJ, Tsai SF, Lauderdale TL. Complete nucleotide sequence of pK245, a 98-kilobase plasmid conferring quinolone resistance and extended-spectrum-beta-lactamase activity in a clinical *Klebsiella pneumoniae* isolate. *Antimicrob Agents Chemother.* 2006; 50:3861–3866. [PubMed: 16940067]
88. Hu FP, Xu XG, Zhu DM, Wang MG. Coexistence of *qnrB4* and *qnrS1* in a clinical strain of *Klebsiella pneumoniae*. *Acta pharmacologica Sinica.* 2008; 29:320–324. [PubMed: 18298896]
89. Poirel L, Cattoir V, Soares A, Soussy CJ, Nordmann P. Novel Ambler class A β -lactamase LAP-1 and its association with the plasmid-mediated quinolone resistance determinant QnrS1. *Antimicrob Agents Chemother.* 2007; 51:631–637. [PubMed: 17116662]
90. Picão RC, Poirel L, Demarta A, Silva CS, Corvaglia AR, Petrini O, Nordmann P. Plasmid-mediated quinolone resistance in *Aeromonas allosaccharophila* recovered from a Swiss lake. *J Antimicrob Chemother.* 2008; 62:948–950. [PubMed: 18772162]
91. Mazzariol A, Kocsis B, Koncan R, Kocsis E, Lanzafame P, Cornaglia G. Description and plasmid characterization of *qnrD* determinants in *Proteus mirabilis* and *Morganella morganii*. *Clin Microbiol Infect.* 2012; 18:E46–48. [PubMed: 22192340]
92. Fonseca EL, Dos Santos Freitas F, Vieira VV, Vicente AC. New *qnr* gene cassettes associated with superintegron repeats in *Vibrio cholerae* O1. *Emerg Infect Dis.* 2008; 14:1129–1131. [PubMed: 18598639]
93. Wu K, Wang F, Sun J, Wang Q, Chen Q, Yu S, Rui Y. Class 1 integron gene cassettes in multidrug-resistant Gram-negative bacteria in southern China. *Int J Antimicrob Agents.* 2012; 40:264–267. [PubMed: 22817917]
94. Overlapping outbreaks of imipenem resistant *Pseudomonas aeruginosa* in a Tunisian burn unit – emergence of new integrons encompassing *bla*_{VIM-2} [Internet]. 2012.
95. Kumar P, Thomas S. Presence of *qnrVC3* gene cassette in SXT and class 1 integrons of *Vibrio cholerae*. *Int J Antimicrob Agents.* 2011; 37:280–281. [PubMed: 21295446]
96. Pai H, Seo MR, Choi TY. Association of QnrB determinants and production of extended-spectrum β -lactamases or plasmid-mediated AmpC β -lactamases in clinical isolates of *Klebsiella pneumoniae*. *Antimicrob Agents Chemother.* 2007; 51:366–368. [PubMed: 17074790]
97. Castanheira M, Mendes RE, Rhomberg PR, Jones RN. Rapid emergence of *bla*_{CTX-M} among Enterobacteriaceae in U.S. Medical Centers: molecular evaluation from the MYSTIC Program (2007). *Microb Drug Resist.* 2008; 14:211–216. [PubMed: 18707552]
98. Jeong HS, Bae IK, Shin JH, Jung HJ, Kim SH, Lee JY, Oh SH, Kim HR, Chang CL, Kho WG, Lee JN. Prevalence of plasmid-mediated quinolone resistance and its association with extended-

- spectrum beta-lactamase and AmpC beta-lactamase in *Enterobacteriaceae*. Korean J Lab Med. 2011; 31:257–264. [PubMed: 22016679]
99. Luo Y, Yang J, Zhang Y, Ye L, Wang L, Guo L. Prevalence of β -lactamases and 16S rRNA methylase genes amongst clinical *Klebsiella pneumoniae* isolates carrying plasmid-mediated quinolone resistance determinants. Int J Antimicrob Agents. 2011; 37:352–355. [PubMed: 21376540]
100. Yang HF, Cheng J, Hu LF, Ye Y, Li JB. Plasmid-mediated quinolone resistance in extended-spectrum- β -lactamase- and AmpC β -lactamase-producing *Serratia marcescens* in China. Antimicrob Agents Chemother. 2012; 56:4529–4531. [PubMed: 22664965]
101. Liu BT, Liao XP, Yue L, Chen XY, Li L, Yang SS, Sun J, Zhang S, Liao SD, Liu YH. Prevalence of β -lactamase and 16S rRNA methylase genes among clinical *Escherichia coli* isolates carrying plasmid-mediated quinolone resistance genes from animals. Microb Drug Resist. 2013; 19:237–245. [PubMed: 23289437]
102. Soge OO, Adeniyi BA, Roberts MC. New antibiotic resistance genes associated with CTX-M plasmids from uropathogenic Nigerian *Klebsiella pneumoniae*. J Antimicrob Chemother. 2006; 58:1048–1053. [PubMed: 16997844]
103. Cattoir V, Weill FX, Poirel L, Fabre L, Soussy CJ, Nordmann P. Prevalence of *qnr* genes in *Salmonella* in France. J Antimicrob Chemother. 2007; 59:751–754. [PubMed: 17307773]
104. Jiang Y, Zhou Z, Qian Y, Wei Z, Yu Y, Hu S, Li L. Plasmid-mediated quinolone resistance determinants *qnr* and *aac(6')-Ib-cr* in extended-spectrum β -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* in China. J Antimicrob Chemother. 2008; 61:1003–1006. [PubMed: 18299311]
105. Guessennd N, Bremont S, Gbonon V, Kacou-Ndouba A, Ekaza E, Lambert T, Dosso M, Courvalin P. Résistance aux quinolones de type *qnr* chez les entérobactéries productrices de bêta-lactamases à spectre élargi à Abidjan en Côte d'Ivoire. Pathol Biol (Paris). 2008; 56:439–446. [PubMed: 18926644]
106. Lavigne JP, Marchandin H, Delmas J, Bouziges N, Lecaillon E, Cavalie L, Jean-Pierre H, Bonnet R, Sotto A. *qnrA* in CTX-M-producing *Escherichia coli* isolates from France. Antimicrob Agents Chemother. 2006; 50:4224–4228. [PubMed: 16982788]
107. Lavilla S, Gonzalez-Lopez JJ, Sabate M, Garcia-Fernandez A, Larrosa MN, Bartolome RM, Carattoli A, Prats G. Prevalence of *qnr* genes among extended-spectrum β -lactamase-producing enterobacterial isolates in Barcelona, Spain. J Antimicrob Chemother. 2008; 61:291–295. [PubMed: 18029415]
108. Yang H, Chen H, Yang Q, Chen M, Wang H. High prevalence of plasmid-mediated quinolone resistance genes *qnr* and *aac(6')-Ib-cr* in clinical isolates of *Enterobacteriaceae* from nine teaching hospitals in China. Antimicrob Agents Chemother. 2008; 52:4268–4273. [PubMed: 18809939]
109. Iabadene H, Messai Y, Ammari H, Ramdani-Bouguessa N, Lounes S, Bakour R, Arlet G. Dissemination of ESBL and Qnr determinants in *Enterobacter cloacae* in Algeria. J Antimicrob Chemother. 2008; 62:133–136. [PubMed: 18385142]
110. Perilli M, Forcella C, Celenza G, Frascaria P, Segatore B, Pellegrini C, Amicosante G. Evidence for *qnrB1* and *aac(6')-Ib-cr* in CTX-M-15-producing uropathogenic *Enterobacteriaceae* in an Italian teaching hospital. Diagn Microbiol Infect Dis. 2009; 64:90–93. [PubMed: 19249178]
111. Miró E, Segura C, Navarro F, Sorlí L, Coll P, Horcajada JP, Álvarez-Lerma F, Salvadó M. Spread of plasmids containing the *bla_{TEM-1}* and *bla_{CTX-M}* genes and the *qnr* determinant in *Enterobacter cloacae*, *Klebsiella pneumoniae* and *Klebsiella oxytoca* isolates. J Antimicrob Chemother. 2010; 65:661–665. [PubMed: 20089541]
112. Albrechtova K, Dolejska M, Cizek A, Tausova D, Klimes J, Bebora L, Literak I. Dogs of nomadic pastoralists in northern Kenya are reservoirs of plasmid-mediated cephalosporin- and quinolone-resistant *Escherichia coli*, including pandemic clone B2-O25-ST131. Antimicrob Agents Chemother. 2012; 56:4013–4017. [PubMed: 22508313]
113. Shibl AM, Al-Agamy MH, Khubnani H, Senok AC, Tawfik AF, Livermore DM. High prevalence of acquired quinolone-resistance genes among *Enterobacteriaceae* from Saudi Arabia with CTX-M-15 β -lactamase. Diagn Microbiol Infect Dis. 2012; 73:350–353. [PubMed: 22633335]

114. Li DX, Zhang SM, Hu GZ, Wang Y, Liu HB, Wu CM, Shang YH, Chen YX, Du XD. Tn3-associated *rmtB* together with *qnrS1*, *aac(6')-Ib-cr* and *bla*_{CTX-M-15} are co-located on an F49:A-B- plasmid in an *Escherichia coli* ST10 strain in China. *J Antimicrob Chemother.* 2012; 67:236–238. [PubMed: 22010207]
115. Wu JJ, Ko WC, Tsai SH, Yan JJ. Prevalence of plasmid-mediated quinolone resistance determinants QnrA, QnrB, and QnrS among clinical isolates of *Enterobacter cloacae* in a Taiwanese hospital. *Antimicrob Agents Chemother.* 2007; 51:1223–1227. [PubMed: 17242140]
116. Chmelnitsky I, Navon-Venezia S, Strahilevitz J, Carmeli Y. Plasmid-mediated *qnrB2* and carbapenemase gene *bla*_{KPC-2} carried on the same plasmid in carbapenem-resistant ciprofloxacin-susceptible *Enterobacter cloacae* isolates. *Antimicrob Agents Chemother.* 2008; 52:2962–2965. [PubMed: 18519713]
117. Endimiani A, Carias LL, Hujer AM, Bethel CR, Hujer KM, Perez F, Hutton RA, Fox WR, Hall GS, Jacobs MR, Paterson DL, Rice LB, Jenkins SG, Tenover FC, Bonomo RA. Presence of plasmid-mediated quinolone resistance in *Klebsiella pneumoniae* isolates possessing *bla*_{KPC} in the United States. *Antimicrob Agents Chemother.* 2008; 52:2680–2682. [PubMed: 18426899]
118. Mendes RE, Bell JM, Turnidge JD, Yang Q, Yu Y, Sun Z, Jones RN. Carbapenem-resistant isolates of *Klebsiella pneumoniae* in China and detection of a conjugative plasmid (*bla*_{KPC-2} plus *qnrB4*) and a *bla*_{IMP-4} gene. *Antimicrob Agents Chemother.* 2008; 52:798–799. [PubMed: 18086850]
119. Poirel L, Leviandier C, Nordmann P. Prevalence and genetic analysis of plasmid-mediated quinolone resistance determinants QnrA and QnrS in *Enterobacteriaceae* isolates from a French university hospital. *Antimicrob Agents Chemother.* 2006; 50:3992–3997. [PubMed: 16982792]
120. García-Fernández A, Fortini D, Veldman K, Mevius D, Carattoli A. Characterization of plasmids harbouring *qnrS1*, *qnrB2* and *qnrB19* genes in *Salmonella*. *J Antimicrob Chemother.* 2009; 63:274–281. [PubMed: 19001452]
121. Potron A, Poirel L, Bernabeu S, Monnet X, Richard C, Nordmann P. Nosocomial spread of ESBL-positive *Enterobacter cloacae* co-expressing plasmid-mediated quinolone resistance Qnr determinants in one hospital in France. *J Antimicrob Chemother.* 2009; 64:653–654. [PubMed: 19549669]
122. Cambau E, Lascols C, Sougakoff W, Bebear C, Bonnet R, Cavallo JD, Gutmann L, Ploy MC, Jarlier V, Soussy CJ, Robert J. Occurrence of *qnrA*-positive clinical isolates in French teaching hospitals during 2002–2005. *Clin Microbiol Infect.* 2006; 12:1013–1020. [PubMed: 16961639]
123. Corkill JE, Anson JJ, Hart CA. High prevalence of the plasmid-mediated quinolone resistance determinant *qnrA* in multidrug-resistant *Enterobacteriaceae* from blood cultures in Liverpool, UK. *J Antimicrob Chemother.* 2005; 56:1115–1117. [PubMed: 16260446]
124. Rodriguez-Martinez JM, Poirel L, Pascual A, Nordmann P. Plasmid-mediated quinolone resistance in Australia. *Microb Drug Resist.* 2006; 12:99–102. [PubMed: 16922624]
125. Szabó D, Kocsis B, Rókusz L, Szentandrassy J, Katona K, Kristóf K, Nagy K. First detection of plasmid-mediated, quinolone resistance determinants *qnrA*, *qnrB*, *qnrS* and *aac(6')-Ib-cr* in extended-spectrum β -lactamase (ESBL)-producing *Enterobacteriaceae* in Budapest, Hungary. *J Antimicrob Chemother.* 2008; 62:630–632. [PubMed: 18477707]
126. Poirel L, Villa L, Bertini A, Pitout JD, Nordmann P, Carattoli A. Expanded-spectrum β -lactamase and plasmid-mediated quinolone resistance. *Emerg Infect Dis.* 2007; 13:803–805. [PubMed: 18044054]
127. Zhao X, Xu X, Zhu D, Ye X, Wang M. Decreased quinolone susceptibility in high percentage of *Enterobacter cloacae* clinical isolates caused only by Qnr determinants. *Diagn Microbiol Infect Dis.* 2010; 67:110–113. [PubMed: 20227223]
128. Mammeri H, Van De Loo M, Poirel L, Martinez-Martinez L, Nordmann P. Emergence of plasmid-mediated quinolone resistance in *Escherichia coli* in Europe. *Antimicrob Agents Chemother.* 2005; 49:71–76. [PubMed: 15616277]
129. Nazic H, Poirel L, Nordmann P. Further identification of plasmid-mediated quinolone resistance determinant in *Enterobacteriaceae* in Turkey. *Antimicrob Agents Chemother.* 2005; 49:2146–2147. [PubMed: 15855552]

130. Poirel L, Van De Loo M, Mammeri H, Nordmann P. Association of plasmid-mediated quinolone resistance with extended-spectrum β -lactamase VEB-1. *Antimicrob Agents Chemother.* 2005; 49:3091–3094. [PubMed: 15980408]
131. Aschbacher R, Doumith M, Livermore DM, Larcher C, Woodford N. Linkage of acquired quinolone resistance (*qnrS1*) and metallo-beta-lactamase (*bla_{VIM-1}*) genes in multiple species of Enterobacteriaceae from Bolzano, Italy. *J Antimicrob Chemother.* 2008; 61:515–523. [PubMed: 18184647]
132. Wu JJ, Ko WC, Wu HM, Yan JJ. Prevalence of Qnr determinants among bloodstream isolates of *Escherichia coli* and *Klebsiella pneumoniae* in a Taiwanese hospital, 1999–2005. *J Antimicrob Chemother.* 2008; 61:1234–1239. [PubMed: 18356152]
133. Lee CH, Liu JW, Li CC, Chien CC, Tang YF, Su LH. Spread of *ISCR1* elements containing *bla_{DHA-1}* and multiple antimicrobial resistance genes leading to increase of flomoxef resistance in extended-spectrum- β -lactamase-producing *Klebsiella pneumoniae*. *Antimicrob Agents Chemother.* 2011; 55:4058–4063. [PubMed: 21746945]
134. Verdet C, Benzerara Y, Gautier V, Adam O, Ould-Hocine Z, Arlet G. Emergence of DHA-1-producing *Klebsiella* spp. in the Parisian region: genetic organization of the *ampC* and *ampR* genes originating from *Morganella morganii*. *Antimicrob Agents Chemother.* 2006; 50:607–617. [PubMed: 16436717]
135. Hidalgo L, Gutierrez B, Ovejero CM, Carrilero L, Matrat S, Saba CK, Santos-Lopez A, Thomas-Lopez D, Hofer A, Suarez M, Santurde G, Martin-Espada C, Gonzalez-Zorn B. *Klebsiella pneumoniae* sequence type 11 from companion animals bearing ArmA methyltransferase, DHA-1 β -lactamase, and QnrB4. *Antimicrob Agents Chemother.* 2013; 57:4532–4534. [PubMed: 23752506]
136. Sanchez-Cespedes J, Marti S, Soto SM, Alba V, Melción C, Almela M, Marco F, Vila J. Two chromosomally located *qnrB* variants, *qnrB6* and the new *qnrB16*, in *Citrobacter* spp. isolates causing bacteraemia. *Clin Microbiol Infect.* 2009; 15:1132–1138. [PubMed: 19392890]
137. Kehrenberg C, Friederichs S, de Jong A, Schwarz S. Novel variant of the *qnrB* Gene, *qnrB12*, in *Citrobacter werkmanii*. *Antimicrob Agents Chemother.* 2008; 52:1206–1207. [PubMed: 18086833]
138. Ogbolu DO, Daini OA, Ogunledun A, Alli AO, Webber MA. High levels of multidrug resistance in clinical isolates of Gram-negative pathogens from Nigeria. *Int J Antimicrob Agents.* 2011; 37:62–66. [PubMed: 21074376]
139. Wang F, Wu K, Sun J, Wang Q, Chen Q, Yu S, Rui Y. Novel *ISCR1*-linked resistance genes found in multidrug-resistant Gram-negative bacteria in southern China. *Int J Antimicrob Agents.* 2012; 40:404–408. [PubMed: 22890194]
140. Touati A, Brasme L, Benallaoua S, Gharout A, Madoux J, De Champs C. First report of *qnrB*-producing *Enterobacter cloacae* and *qnrA*-producing *Acinetobacter baumannii* recovered from Algerian hospitals. *Diagn Microbiol Infect Dis.* 2008; 60:287–290. [PubMed: 18036760]
141. Guo L, Zhang J, Xu C, Zhao Y, Ren T, Zhang B, Fan H, Liao M. Molecular characterization of fluoroquinolone resistance in *Haemophilus parasuis* isolated from pigs in South China. *J Antimicrob Chemother.* 2011; 66:539–542. [PubMed: 21217127]
142. Jacoby GA, Gacharna N, Black TA, Miller GH, Hooper DC. Temporal appearance of plasmid-mediated quinolone resistance genes. *Antimicrob Agents Chemother.* 2009; 53:1665–1666. [PubMed: 19164145]
143. Strahilevitz J, Engelstein D, Adler A, Temper V, Moses AE, Block C, Robicsek A. Changes in *qnr* prevalence and fluoroquinolone resistance in clinical isolates of *Klebsiella pneumoniae* and *Enterobacter* spp. collected from 1990 to 2005. *Antimicrob Agents Chemother.* 2007; 51:3001–3003. [PubMed: 17526754]
144. Xu X, Wu S, Ye X, Liu Y, Shi W, Zhang Y, Wang M. Prevalence and expression of the plasmid-mediated quinolone resistance determinant *qnrA1*. *Antimicrob Agents Chemother.* 2007; 51:4105–4110. [PubMed: 17724159]
145. Kim HB, Park CH, Kim CJ, Kim EC, Jacoby GA, Hooper DC. Prevalence of plasmid-mediated quinolone resistance determinants over a 9-year period. *Antimicrob Agents Chemother.* 2009; 53:639–645. [PubMed: 19064896]

146. Kim NH, Choi EH, Sung JY, Oh CE, Kim HB, Kim EC, Lee HJ. Prevalence of plasmid-mediated quinolone resistance genes and ciprofloxacin resistance in pediatric bloodstream isolates of *Enterobacteriaceae* over a 9-year period. *Jpn J Infect Dis.* 2013; 66:151–154. [PubMed: 23514914]
147. Robicsek A, Sahm DF, Strahilevitz J, Jacoby GA, Hooper DC. Broader distribution of plasmid-mediated quinolone resistance in the United States. *Antimicrob Agents Chemother.* 2005; 49:3001–3003. [PubMed: 15980384]
148. Literak I, Micudova M, Tausova D, Cizek A, Dolejska M, Papousek I, Prochazka J, Vojtech J, Borleis F, Guardone L, Guenther S, Hordowski J, Lejas C, Meissner W, Marcos BF, Tucakov M. Plasmid-mediated quinolone resistance genes in fecal bacteria from rooks commonly wintering throughout Europe. *Microb Drug Resist.* 2012; 18:567–573. [PubMed: 22731858]
149. Tausova D, Dolejska M, Cizek A, Hanusova L, Hrusakova J, Svoboda O, Camlik G, Literak I. *Escherichia coli* with extended-spectrum β -lactamase and plasmid-mediated quinolone resistance genes in great cormorants and mallards in Central Europe. *J Antimicrob Chemother.* 2012; 67:1103–1107. [PubMed: 22323500]
150. Shaheen BW, Nayak R, Foley SL, Boothe DM. Chromosomal and plasmid-mediated fluoroquinolone resistance mechanisms among broad-spectrum-cephalosporin-resistant *Escherichia coli* isolates recovered from companion animals in the USA. *J Antimicrob Chemother.* 2013; 68:1019–1024. [PubMed: 23302578]
151. Kirchner M, Wearing H, Teale C. Plasmid-mediated quinolone resistance gene detected in *Escherichia coli* from cattle. *Vet Microbiol.* 2011; 148:434–435. [PubMed: 20884136]
152. Madec JY, Poirel L, Saras E, Gourguechon A, Girlich D, Nordmann P, Haenni M. Non-ST131 *Escherichia coli* from cattle harbouring human-like *bla*_{CTX-M-15}-carrying plasmids. *J Antimicrob Chemother.* 2012; 67:578–581. [PubMed: 22210752]
153. Wu CM, Wang Y, Cao XY, Lin JC, Qin SS, Mi TJ, Huang SY, Shen JZ. Emergence of plasmid-mediated quinolone resistance genes in *Enterobacteriaceae* isolated from chickens in China. *J Antimicrob Chemother.* 2009; 63:408–411. [PubMed: 19029095]
154. Cerquetti M, Garcia-Fernandez A, Giufre M, Fortini D, Accogli M, Graziani C, Luzzi I, Caprioli A, Carattoli A. First report of plasmid-mediated quinolone resistance determinant *qnrS1* in an *Escherichia coli* strain of animal origin in Italy. *Antimicrob Agents Chemother.* 2009; 53:3112–3114. [PubMed: 19451298]
155. Veldman K, Cavaco LM, Mevius D, Battisti A, Franco A, Botteldoorn N, Bruneau M, Perrin-Guyomard A, Cerny T, De Frutos Escobar C, Guerra B, Schroeter A, Gutierrez M, Hopkins K, Myllyniemi AL, Sunde M, Wasyl D, Aarestrup FM. International collaborative study on the occurrence of plasmid-mediated quinolone resistance in *Salmonella enterica* and *Escherichia coli* isolated from animals, humans, food and the environment in 13 European countries. *J Antimicrob Chemother.* 2011; 66:1278–1286. [PubMed: 21393198]
156. Chen X, Zhang W, Pan W, Yin J, Pan Z, Gao S, Jiao X. Prevalence of *qnr*, *aac(6')-Ib-cr*, *qepA*, and *oxxAB* in *Escherichia coli* isolates from humans, animals, and the environment. *Antimicrob Agents Chemother.* 2012; 56:3423–3427. [PubMed: 22391545]
157. Du XD, Li DX, Hu GZ, Wang Y, Shang YH, Wu CM, Liu HB, Li XS. Tn1548-associated *armA* is co-located with *qnrB2*, *aac(6')-Ib-cr* and *bla*_{CTX-M-3} on an IncFII plasmid in a *Salmonella enterica* subsp. *enterica* serovar Paratyphi B strain isolated from chickens in China. *J Antimicrob Chemother.* 2012; 67:246–248. [PubMed: 21965429]
158. Literak I, Reitschmied T, Bujnakova D, Dolejska M, Cizek A, Bardon J, Pokludova L, Alexa P, Halova D, Jamborova I. Broilers as a source of quinolone-resistant and extraintestinal pathogenic *Escherichia coli* in the Czech Republic. *Microb Drug Resist.* 2013; 19:57–63. [PubMed: 23020862]
159. Kim JH, Cho JK, Kim KS. Prevalence and characterization of plasmid-mediated quinolone resistance genes in *Salmonella* isolated from poultry in Korea. *Avian pathology : journal of the WVPA.* 2013; 42:221–229.
160. Liu BT, Yang QE, Li L, Sun J, Liao XP, Fang LX, Yang SS, Deng H, Liu YH. Dissemination and characterization of plasmids carrying *oxxAB-bla*_{CTX-M} genes in *Escherichia coli* isolates from food-producing animals. *PLoS One.* 2013; 8:e73947. [PubMed: 24040123]

161. Pomba C, da Fonseca JD, Baptista BC, Correia JD, Martinez-Martinez L. Detection of the pandemic O25-ST131 human virulent *Escherichia coli* CTX-M-15-producing clone harboring the *qnrB2* and *aac(6')-Ib-cr* genes in a dog. *Antimicrob Agents Chemother*. 2009; 53:327–328. [PubMed: 19001117]
162. Yue L, Jiang HX, Liao XP, Liu JH, Li SJ, Chen XY, Chen CX, Lu DH, Liu YH. Prevalence of plasmid-mediated quinolone resistance *qnr* genes in poultry and swine clinical isolates of *Escherichia coli*. *Vet Microbiol*. 2008; 132:414–420. [PubMed: 18573620]
163. Verner-Jeffreys DW, Welch TJ, Schwarz T, Pond MJ, Woodward MJ, Haig SJ, Rimmer GS, Roberts E, Morrison V, Baker-Austin C. High prevalence of multidrug-tolerant bacteria and associated antimicrobial resistance genes isolated from ornamental fish and their carriage water. *PLoS One*. 2009; 4:e8388. [PubMed: 20027306]
164. Jiang HX, Tang D, Liu YH, Zhang XH, Zeng ZL, Xu L, Hawkey PM. Prevalence and characteristics of β -lactamase and plasmid-mediated quinolone resistance genes in *Escherichia coli* isolated from farmed fish in China. *J Antimicrob Chemother*. 2012; 67:2350–2353. [PubMed: 22809702]
165. Han JE, Kim JH, Choresca CH Jr, Shin SP, Jun JW, Chai JY, Park SC. First description of ColE-type plasmid in *Aeromonas* spp. carrying quinolone resistance (*qnrS2*) gene. *Letters in applied microbiology*. 2012; 55:290–294. [PubMed: 22862417]
166. Dolejska M, Duskova E, Rybarikova J, Janoszowska D, Roubalova E, Dibdakova K, Maceckova G, Kohoutova L, Literak I, Smola J, Cizek A. Plasmids carrying *bla*_{CTX-M-1} and *qnr* genes in *Escherichia coli* isolates from an equine clinic and a horseback riding centre. *J Antimicrob Chemother*. 2011; 66:757–764. [PubMed: 21393204]
167. Liu JH, Deng YT, Zeng ZL, Gao JH, Chen L, Arakawa Y, Chen ZL. Coprevalence of plasmid-mediated quinolone resistance determinants QepA, Qnr, and AAC(6')-Ib-cr among 16S rRNA methylase RmtB-producing *Escherichia coli* isolates from pigs. *Antimicrob Agents Chemother*. 2008; 52:2992–2993. [PubMed: 18490500]
168. Dotto G, Giacomelli M, Grilli G, Ferrazzi V, Carattoli A, Fortini D, Piccirillo A. High Prevalence of *oqxAB* in *Escherichia coli* isolates from domestic and wild lagomorphs in Italy. *Microb Drug Resist*. 2013
169. Guerra B, Helmuth R, Thomas K, Beutlich J, Jahn S, Schroeter A. Plasmid-mediated quinolone resistance determinants in *Salmonella* spp. isolates from reptiles in Germany. *J Antimicrob Chemother*. 2010; 65:2043–2045. [PubMed: 20601355]
170. Ahmed AM, Motoi Y, Sato M, Maruyama A, Watanabe H, Fukumoto Y, Shimamoto T. Zoo animals as reservoirs of gram-negative bacteria harboring integrons and antimicrobial resistance genes. *Appl Environ Microbiol*. 2007; 73:6686–6690. [PubMed: 17720829]
171. Wang Y, He T, Han J, Wang J, Foley SL, Yang G, Wan S, Shen J, Wu C. Prevalence of ESBLs and PMQR genes in fecal *Escherichia coli* isolated from the non-human primates in six zoos in China. *Vet Microbiol*. 2012; 159:53–59. [PubMed: 22487457]
172. Dobiasova H, Dolejska M, Jamborova I, Brhelova E, Blazkova L, Papousek I, Kozlova M, Klimes J, Cizek A, Literak I. Extended spectrum beta-lactamase and fluoroquinolone resistance genes and plasmids among *Escherichia coli* isolates from zoo animals, Czech Republic. *FEMS microbiology ecology*. 2013; 85:604–611. [PubMed: 23679004]
173. Kim HB, Park CH, Gavin M, Jacoby GA, Hooper DC. Cold shock induces *qnrA* expression in *Shewanella algae*. *Antimicrob Agents Chemother*. 2011; 55:414–416. [PubMed: 21078945]
174. Wang M, Jacoby GA, Mills DM, Hooper DC. SOS regulation of *qnrB* expression. *Antimicrob Agents Chemother*. 2009; 53:821–823. [PubMed: 19029320]
175. Da Re S, Garnier F, Guerin E, Campoy S, Denis F, Ploy MC. The SOS response promotes *qnrB* quinolone-resistance determinant expression. *EMBO Rep*. 2009; 10:929–933. [PubMed: 19556999]
176. Briaies A, Rodriguez-Martinez JM, Velasco C, Machuca J, Diaz de Alba P, Blazquez J, Pascual A. Exposure to diverse antimicrobials induces the expression of *qnrB1*, *qnrD* and *smaqrn* genes by SOS-dependent regulation. *J Antimicrob Chemother*. 2012; 67:2854–2859. [PubMed: 22915457]

177. Okumura R, Liao CH, Gavin M, Jacoby GA, Hooper DC. Quinolone Induction of *qnrVS1* in *Vibrio splendidus* and plasmid-carried *qnrS1* in *Escherichia coli*, a mechanism independent of the SOS system. *Antimicrob Agents Chemother.* 2011; 55:5942–5945. [PubMed: 21930884]
178. Kwak YG, Jacoby GA, Hooper DC. Induction of plasmid-carried *qnrS1* in *Escherichia coli* by naturally occurring quinolones and quorum-sensing signal molecules. *Antimicrob Agents Chemother.* 2013; 57:4031–4034. [PubMed: 23689721]
179. Vetting MW, Park CH, Hegde SS, Jacoby GA, Hooper DC, Blanchard JS. Mechanistic and structural analysis of aminoglycoside *N*-acetyltransferase AAC(6′)-Ib and its bifunctional fluoroquinolone-active AAC(6′)-Ib-cr variant. *Biochemistry.* 2008; 47:9825–9835. [PubMed: 18710261]
180. Maurice F, Broutin I, Podglajen I, Benas P, Collatz E, Dardel F. Enzyme structural plasticity and the emergence of broad-spectrum antibiotic resistance. *EMBO Rep.* 2008; 9:344–349. [PubMed: 18292754]
181. Ruiz E, Ocampo-Sosa AA, Alcoba-Florez J, Roman E, Arlet G, Torres C, Martinez-Martinez L. Changes in ciprofloxacin resistance levels in *Enterobacter aerogenes* isolates associated with variable expression of the *aac(6′)-Ib-cr* gene. *Antimicrob Agents Chemother.* 2012; 56:1097–1100. [PubMed: 22106222]
182. de Toro M, Rodriguez I, Rojo-Bezares B, Helmuth R, Torres C, Guerra B, Saenz Y. pMdT1, a small ColE1-like plasmid mobilizing a new variant of the *aac(6′)-Ib-cr* gene in *Salmonella enterica* serovar Typhimurium. *J Antimicrob Chemother.* 2013; 68:1277–1280. [PubMed: 23361643]
183. Guillard T, Duval V, Moret H, Brasme L, Vernet-Garnier V, de Champs C. Rapid detection of *aac(6′)-Ib-cr* quinolone resistance gene by pyrosequencing. *J Clin Microbiol.* 2010; 48:286–289. [PubMed: 19923484]
184. Pitout JD, Wei Y, Church DL, Gregson DB. Surveillance for plasmid-mediated quinolone resistance determinants in Enterobacteriaceae within the Calgary Health Region, Canada: the emergence of *aac(6′)-Ib-cr*. *J Antimicrob Chemother.* 2008; 61:999–1002. [PubMed: 18296438]
185. Jones GL, Warren RE, Skidmore SJ, Davies VA, Gibreel T, Upton M. Prevalence and distribution of plasmid-mediated quinolone resistance genes in clinical isolates of *Escherichia coli* lacking extended-spectrum β -lactamases. *J Antimicrob Chemother.* 2008; 62:1245–1251. [PubMed: 18827034]
186. Oteo J, Cuevas O, López-Rodríguez I, Banderas-Florido A, Vindel A, Pérez-Vázquez M, Bautista V, Arroyo M, García-Caballero J, Marin-Casanova P, González-Sanz R, Fuentes-Gómez V, Oña-Compán S, García-Cobos S, Campos J. Emergence of CTX-M-15-producing *Klebsiella pneumoniae* of multilocus sequence types 1, 11, 14, 17, 20, 35 and 36 as pathogens and colonizers in newborns and adults. *J Antimicrob Chemother.* 2009; 64:524–528. [PubMed: 19525516]
187. Baudry PJ, Nichol K, DeCorby M, Lagace-Wiens P, Olivier E, Boyd D, Mulvey MR, Hoban DJ, Zhanel GG. Mechanisms of resistance and mobility among multidrug-resistant CTX-M-producing *Escherichia coli* from Canadian intensive care units: the 1st report of QepA in North America. *Diagn Microbiol Infect Dis.* 2009; 63:319–326. [PubMed: 19216943]
188. Coelho A, Mirelis B, Alonso-Tarrés C, Nieves Larrosa M, Miró E, Clivillé Abad R, Bartolomé RM, Castañer M, Prats G, Johnson JR, Navarro F, González-López JJ. Detection of three stable genetic clones of CTX-M-15-producing *Klebsiella pneumoniae* in the Barcelona metropolitan area, Spain. *J Antimicrob Chemother.* 2009; 64:862–864. [PubMed: 19628471]
189. Literacka E, Bedenic B, Baraniak A, Fiett J, Tonkic M, Jajic-Bencic I, Gniadkowski M. *bla*_{CTX-M} genes in *Escherichia coli* strains from Croatian hospitals are located in new (*bla*_{CTX-M-3a}) and widely spread (*bla*_{CTX-M-3a} and *bla*_{CTX-M-15}) genetic structures. *Antimicrob Agents Chemother.* 2009; 53:1630–1635. [PubMed: 19188377]
190. Sabtcheva S, Kaku M, Saga T, Ishii Y, Kantardjiev T. High prevalence of the *aac(6′)-Ib-cr* gene and its dissemination among *Enterobacteriaceae* isolates by CTX-M-15 plasmids in Bulgaria. *Antimicrob Agents Chemother.* 2009; 53:335–336. [PubMed: 19001110]
191. Ruiz E, Rezusta A, Saenz Y, Rocha-Gracia R, Vinue L, Vindel A, Villuendas C, Azanedo ML, Monforte ML, Revillo MJ, Torres C. New genetic environments of *aac(6′)-Ib-cr* gene in a

- multiresistant *Klebsiella oxytoca* strain causing an outbreak in a pediatric intensive care unit. *Diagn Microbiol Infect Dis*. 2011; 69:236–238. [PubMed: 21251575]
192. Ruiz E, Saenz Y, Zarazaga M, Rocha-Gracia R, Martinez-Martinez L, Arlet G, Torres C. *qnr*, *aac(6)-Ib-cr* and *qepA* genes in *Escherichia coli* and *Klebsiella* spp.: genetic environments and plasmid and chromosomal location. *J Antimicrob Chemother*. 2012; 67:886–897. [PubMed: 22223228]
 193. Musumeci R, Rausa M, Giovannoni R, Cialdella A, Bramati S, Sibra B, Giltri G, Vigano F, Cocuzza CE. Prevalence of plasmid-mediated quinolone resistance genes in uropathogenic *Escherichia coli* isolated in a teaching hospital of northern Italy. *Microb Drug Resist*. 2012; 18:33–41. [PubMed: 21711147]
 194. Park CH, Robicsek A, Jacoby GA, Sahm D, Hooper DC. Prevalence in the United States of *aac(6)-Ib-cr* encoding a ciprofloxacin-modifying enzyme. *Antimicrob Agents Chemother*. 2006; 50:3953–3955. [PubMed: 16954321]
 195. Kim ES, Jeong JY, Jun JB, Choi SH, Lee SO, Kim MN, Woo JH, Kim YS. Prevalence of *aac(6)-Ib-cr* encoding a ciprofloxacin-modifying enzyme among *Enterobacteriaceae* blood isolates in Korea. *Antimicrob Agents Chemother*. 2009; 53:2643–2645. [PubMed: 19289526]
 196. Yang J, Luo Y, Li J, Ma Y, Hu C, Jin S, Ye L, Cui S. Characterization of clinical *Escherichia coli* isolates from China containing transferable quinolone resistance determinants. *J Antimicrob Chemother*. 2010; 65:453–459. [PubMed: 20056688]
 197. Périchon B, Bogaerts P, Lambert T, Frangeul L, Courvalin P, Galimand M. Sequence of conjugative plasmid pIP1206 mediating resistance to aminoglycosides by 16S rRNA methylation and to hydrophilic fluoroquinolones by efflux. *Antimicrob Agents Chemother*. 2008; 52:2581–2592. [PubMed: 18458128]
 198. Yamane K, Wachino J, Suzuki S, Arakawa Y. Plasmid-mediated *qepA* gene among *Escherichia coli* clinical isolates from Japan. *Antimicrob Agents Chemother*. 2008; 52:1564–1566. [PubMed: 18285488]
 199. Cattoir V, Poirol L, Nordmann P. Plasmid-mediated quinolone resistance pump QepA2 in an *Escherichia coli* isolate from France. *Antimicrob Agents Chemother*. 2008; 52:3801–3804. [PubMed: 18644958]
 200. Sorensen AH, Hansen LH, Johannesen E, Sorensen SJ. Conjugative plasmid conferring resistance to olaquinox. *Antimicrob Agents Chemother*. 2003; 47:798–799. [PubMed: 12543696]
 201. Hansen LH, Johannesen E, Burmolle M, Sorensen AH, Sorensen SJ. Plasmid-encoded multidrug efflux pump conferring resistance to olaquinox in *Escherichia coli*. *Antimicrob Agents Chemother*. 2004; 48:3332–3337. [PubMed: 15328093]
 202. Norman A, Hansen LH, She Q, Sørensen SJ. Nucleotide sequence of pOLA52: a conjugative IncX1 plasmid from *Escherichia coli* which enables biofilm formation and multidrug efflux. *Plasmid*. 2008; 60:59–74. [PubMed: 18440636]
 203. Kim HB, Wang M, Park CH, Kim EC, Jacoby GA, Hooper DC. *oqxAB* encoding a multidrug efflux pump in human clinical isolates of *Enterobacteriaceae*. *Antimicrob Agents Chemother*. 2009; 53:3582–3584. [PubMed: 19528276]
 204. Zhao J, Chen Z, Chen S, Deng Y, Liu Y, Tian W, Huang X, Wu C, Sun Y, Zeng Z, Liu JH. Prevalence and dissemination of *oqxAB* in *Escherichia coli* Isolates from animals, farmworkers, and the environment. *Antimicrob Agents Chemother*. 2010; 54:4219–4224. [PubMed: 20696876]
 205. Wong MH, Chen S. First detection of *oqxAB* in *Salmonella* spp. isolated from food. *Antimicrob Agents Chemother*. 2013; 57:658–660. [PubMed: 23147728]
 206. Rodríguez-Martínez JM, Díaz de Alba P, Briales A, Machuca J, Lossa M, Fernández-Cuenca F, Rodríguez Baño J, Martínez-Martínez L, Pascual A. Contribution of OqxAB efflux pumps to quinolone resistance in extended-spectrum- β -lactamase-producing *Klebsiella pneumoniae*. *J Antimicrob Chemother*. 2013; 68:68–73. [PubMed: 23011289]
 207. Li L, Liao X, Yang Y, Sun J, Li L, Liu B, Yang S, Ma J, Li X, Zhang Q, Liu Y. Spread of *oqxAB* in *Salmonella enterica* serotype Typhimurium predominantly by IncHI2 plasmids. *J Antimicrob Chemother*. 2013; 68:2263–2268. [PubMed: 23737490]

208. Yuan J, Xu X, Guo Q, Zhao X, Ye X, Guo Y, Wang M. Prevalence of the *oqxAB* gene complex in *Klebsiella pneumoniae* and *Escherichia coli* clinical isolates. *J Antimicrob Chemother.* 2012; 67:1655–1659. [PubMed: 22438434]
209. Perez F, Rudin SD, Marshall SH, Coakley P, Chen L, Kreiswirth BN, Rather PN, Hujer AM, Toltzis P, van Duin D, Paterson DL, Bonomo RA. OqxAB, a quinolone and olaquinox efflux pump, is widely distributed among multidrug-resistant *Klebsiella pneumoniae* isolates of human origin. *Antimicrob Agents Chemother.* 2013; 57:4602–4603. [PubMed: 23817374]
210. Veleba M, Higgins PG, Gonzalez G, Seifert H, Schneiders T. Characterization of RarA, a novel AraC family multidrug resistance regulator in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother.* 2012; 56:4450–4458. [PubMed: 22644028]
211. De Majumdar S, Veleba M, Finn S, Fanning S, Schneiders T. Elucidating the regulon of multidrug resistance regulator RarA in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother.* 2013; 57:1603–1609. [PubMed: 23318802]
212. Szczepanowski R, Krahn I, Linke B, Goesmann A, Puhler A, Schluter A. Antibiotic multiresistance plasmid pRSB101 isolated from a wastewater treatment plant is related to plasmids residing in phytopathogenic bacteria and carries eight different resistance determinants including a multidrug transport system. *Microbiology.* 2004; 150:3613–3630. [PubMed: 15528650]
213. Nakaminami H, Noguchi N, Sasatsu M. Fluoroquinolone efflux by the plasmid-mediated multidrug efflux pump QacB variant QacBIII in *Staphylococcus aureus*. *Antimicrob Agents Chemother.* 2010; 54:4107–4111. [PubMed: 20660673]
214. Drlica K. The mutant selection window and antimicrobial resistance. *J Antimicrob Chemother.* 2003; 52:11–17. [PubMed: 12805267]
215. Jacoby GA. Mechanisms of resistance to quinolones. *Clin Infect Dis.* 2005; 41(Suppl 2):S120–126. [PubMed: 15942878]
216. Rodríguez-Martínez JM, Velasco C, García I, Cano ME, Martínez-Martínez L, Pascual A. Mutant prevention concentrations of fluoroquinolones for *Enterobacteriaceae* expressing the plasmid-carried quinolone resistance determinant *qnrA1*. *Antimicrob Agents Chemother.* 2007; 51:2236–2239. [PubMed: 17404007]
217. Cesaro A, Bettoni RR, Lascols C, Merens A, Soussy CJ, Cambau E. Low selection of topoisomerase mutants from strains of *Escherichia coli* harbouring plasmid-borne *qnr* genes. *J Antimicrob Chemother.* 2008; 61:1007–1015. [PubMed: 18325893]
218. Martínez-Martínez L, Pascual A, García I, Tran J, Jacoby GA. Interaction of plasmid and host quinolone resistance. *J Antimicrob Chemother.* 2003; 51:1037–1039. [PubMed: 12654766]
219. Briales A, Rodríguez-Martínez JM, Velasco C, Díaz de Alba P, Domínguez-Herrera J, Pachon J, Pascual A. *In vitro* effect of *qnrA1*, *qnrB1*, and *qnrS1* genes on fluoroquinolone activity against isogenic *Escherichia coli* isolates with mutations in *gyrA* and *parC*. *Antimicrob Agents Chemother.* 2011; 55:1266–1269. [PubMed: 21173174]
220. Luo Y, Li J, Meng Y, Ma Y, Hu C, Jin S, Zhang Q, Ding H, Cui S. Joint effects of topoisomerase alterations and plasmid-mediated quinolone-resistant determinants in *Salmonella enterica* Typhimurium. *Microb Drug Resist.* 2011; 17:1–5. [PubMed: 20818995]
221. Lascols C, Robert J, Cattoir V, Bébér C, Cavallo JD, Podglajen I, Ploy MC, Bonnet R, Soussy CJ, Cambau E. Type II topoisomerase mutations in clinical isolates of *Enterobacter cloacae* and other enterobacterial species harbouring the *qnrA* gene. *Int J Antimicrob Agents.* 2007; 29:402–409. [PubMed: 17254753]
222. Poirel L, Pitout JD, Calvo L, Rodríguez-Martínez JM, Church D, Nordmann P. *In vivo* selection of fluoroquinolone-resistant *Escherichia coli* isolates expressing plasmid-mediated quinolone resistance and expanded-spectrum β -lactamase. *Antimicrob Agents Chemother.* 2006; 50:1525–1527. [PubMed: 16569874]
223. de Toro M, Rojo-Bezares B, Vinué L, Undabeitia E, Torres C, Sáenz Y. *In vivo* selection of *aac(6)-Ib-cr* and mutations in the *gyrA* gene in a clinical *qnrS1*-positive *Salmonella enterica* serovar Typhimurium DT104B strain recovered after fluoroquinolone treatment. *J Antimicrob Chemother.* 2010; 65:1945–1949. [PubMed: 20639314]

224. Jeong JY, Kim ES, Choi SH, Kwon HH, Lee SR, Lee SO, Kim MN, Woo JH, Kim YS. Effects of a plasmid-encoded *qnrA1* determinant in *Escherichia coli* strains carrying chromosomal mutations in the *acrAB* efflux pump genes. *Diagn Microbiol Infect Dis*. 2008; 60:105–107. [PubMed: 17889488]
225. Sato T, Yokota S, Uchida I, Okubo T, Usui M, Kusumoto M, Akiba M, Fujii N, Tamura Y. Fluoroquinolone resistance mechanisms in an *Escherichia coli* isolate, HUE1, without quinolone resistance-determining region mutations. *Front Microbiol*. 2013; 4:10.3389/fmicb.2013.00125
226. Warburg G, Korem M, Robicsek A, Engelstein D, Moses AE, Block C, Strahilevitz J. Changes in *aac(6′)-Ib-cr* prevalence and fluoroquinolone resistance in nosocomial isolates of *Escherichia coli* collected from 1991 through 2005. *Antimicrob Agents Chemother*. 2009; 53:1268–1270. [PubMed: 19104026]
227. Briales A, Rodriguez-Martinez JM, Velasco C, de Alba PD, Rodriguez-Bano J, Martinez-Martinez L, Pascual A. Prevalence of plasmid-mediated quinolone resistance determinants *qnr* and *aac(6′)-Ib-cr* in *Escherichia coli* and *Klebsiella pneumoniae* producing extended-spectrum β -lactamases in Spain. *Int J Antimicrob Agents*. 2012; 39:431–434. [PubMed: 22365240]
228. Rodríguez-Martínez JM, Pichardo C, García I, Pachón-Ibañez ME, Docobo-Pérez F, Pascual A, Pachón J, Martínez-Martínez L. Activity of ciprofloxacin and levofloxacin in experimental pneumonia caused by *Klebsiella pneumoniae* deficient in porins, expressing active efflux and producing QnrA1. *Clin Microbiol Infect*. 2008; 14:691–697. [PubMed: 18558942]
229. Domínguez-Herrera J, Velasco C, Docobo-Pérez F, Rodríguez-Martínez JM, López-Rojas R, Briales A, Pichardo C, Díaz-de-Alba P, Rodríguez-Baño J, Pascual A, Pachón J. Impact of *qnrA1*, *qnrB1* and *qnrS1* on the efficacy of ciprofloxacin and levofloxacin in an experimental pneumonia model caused by *Escherichia coli* with or without the GyrA mutation Ser83Leu. *J Antimicrob Chemother*. 2013; 68:1609–1615. [PubMed: 23493313]
230. Allou N, Cambau E, Massias L, Chau F, Fantin B. Impact of low-level resistance to fluoroquinolones due to *qnrA1* and *qnrS1* genes or a *gyrA* mutation on ciprofloxacin bactericidal activity in a murine model of *Escherichia coli* urinary tract infection. *Antimicrob Agents Chemother*. 2009; 53:4292–4297. [PubMed: 19635960]
231. Jakobsen L, Cattoir V, Jensen KS, Hammerum AM, Nordmann P, Frimodt-Møller N. Impact of low-level fluoroquinolone resistance genes *qnrA1*, *qnrB19* and *qnrS1* on ciprofloxacin treatment of isogenic *Escherichia coli* strains in a murine urinary tract infection model. *J Antimicrob Chemother*. 2012; 67:2438–2444. [PubMed: 22685162]
232. Defife R, Scheetz MH, Feinglass JM, Postelnick MJ, Scarsi KK. Effect of differences in MIC values on clinical outcomes in patients with bloodstream infections caused by gram-negative organisms treated with levofloxacin. *Antimicrob Agents Chemother*. 2009; 53:1074–1079. [PubMed: 19075047]
233. Chong YP, Choi SH, Kim ES, Song EH, Lee EJ, Park KH, Cho OH, Kim SH, Lee SO, Kim MN, Jeong JY, Woo JH, Kim YS. Bloodstream infections caused by *qnr*-positive Enterobacteriaceae: clinical and microbiologic characteristics and outcomes. *Diagn Microbiol Infect Dis*. 2010; 67:70–77. [PubMed: 20227228]
234. Liao CH, Hsueh PR, Jacoby GA, Hooper DC. Risk factors and clinical characteristics of patients with *qnr*-positive *Klebsiella pneumoniae* bacteraemia. *J Antimicrob Chemother*. 2013; 68:2907–2914. [PubMed: 23956373]
235. Buchko GW, Ni S, Robinson H, Welsh EA, Pakrasi HB, Kennedy MA. Characterization of two potentially universal turn motifs that shape the repeated five-residues fold–Crystal structure of a luminal pentapeptide repeat protein from *Cyanospora* 51142. *Protein science : a publication of the Protein Society*. 2006; 15:2579–2595. [PubMed: 17075135]
236. Lascols C, Podglajen I, Verdet C, Gautier V, Gutmann L, Soussy CJ, Collatz E, Cambau E. A plasmid-borne *Shewanella algae* gene, *qnrA3*, and its possible transfer in vivo between *Kluyvera ascorbata* and *Klebsiella pneumoniae*. *J Bacteriol*. 2008; 190:5217–5223. [PubMed: 18515416]
237. Mnif B, Ktari S, Chaari A, Medhioub F, Rhimi F, Bouaziz M, Hammami A. Nosocomial dissemination of *Providencia stuartii* isolates carrying *bla*_{OXA-48}, *bla*_{PER-1}, *bla*_{CMY-4} and *qnrA6* in a Tunisian hospital. *J Antimicrob Chemother*. 2013; 68:329–332. [PubMed: 23014719]

238. Robicsek A, Strahilevitz J, Sahm DF, Jacoby GA, Hooper DC. *qnr* prevalence in ceftazidime-resistant *Enterobacteriaceae* isolates from the United States. *Antimicrob Agents Chemother*. 2006; 50:2872–2874. [PubMed: 16870791]
239. Quiroga MP, Andres P, Petroni A, Soler Bistué AJ, Guerriero L, Vargas LJ, Zorreguieta A, Tokumoto M, Quiroga C, Tolmasky ME, Galas M, Centrón D. Complex class 1 integrons with diverse variable regions, including *aac(6')-Ib-cr*, and a novel allele, *qnrB10*, associated with *ISCR1* in clinical enterobacterial isolates from Argentina. *Antimicrob Agents Chemother*. 2007; 51:4466–4470. [PubMed: 17938184]
240. Hopkins KL, Wootton L, Day MR, Threlfall EJ. Plasmid-mediated quinolone resistance determinant *qnrS1* found in *Salmonella enterica* strains isolated in the UK. *J Antimicrob Chemother*. 2007; 59:1071–1075. [PubMed: 17434878]
241. Bönemann G, Stiens M, Pühler A, Schlüter A. Mobilizable IncQ-related plasmid carrying a new quinolone resistance gene, *qnrS2*, isolated from the bacterial community of a wastewater treatment plant. *Antimicrob Agents Chemother*. 2006; 50:3075–3080. [PubMed: 16940104]
242. Cattoir V, Poirel L, Aubert C, Soussy CJ, Nordmann P. Unexpected occurrence of plasmid-mediated quinolone resistance determinants in environmental *Aeromonas* spp. *Emerg Infect Dis*. 2008; 14:231–237. [PubMed: 18258115]
243. Rajpara N, Patel A, Tiwari N, Bahuguna J, Antony A, Choudhury I, Ghosh A, Jain R, Bhardwaj AK. Mechanism of drug resistance in a clinical isolate of *Vibrio fluvialis*: involvement of multiple plasmids and integrons. *Int J Antimicrob Agents*. 2009; 34:220–225. [PubMed: 19427174]
244. Boyd DA, Tyler S, Christianson S, McGeer A, Muller MP, Willey BM, Bryce E, Gardam M, Nordmann P, Mulvey MR. Complete nucleotide sequence of a 92-kilobase plasmid harboring the CTX-M-15 extended-spectrum β -lactamase involved in an outbreak in long-term-care facilities in Toronto, Canada. *Antimicrob Agents Chemother*. 2004; 48:3758–3764. [PubMed: 15388431]
245. Jacoby G, Chow N, Waites K. Prevalence of plasmid-mediated quinolone resistance. *Antimicrob Agents Chemother*. 2003; 47:559–562. [PubMed: 12543659]
246. Jonas D, Biehler K, Hartung D, Spitzmüller B, Daschner FD. Plasmid-mediated quinolone resistance in isolates obtained in German intensive care units. *Antimicrob Agents Chemother*. 2005; 49:773–775. [PubMed: 15673764]
247. Jeong JY, Yoon HJ, Kim ES, Lee Y, Choi SH, Kim NJ, Woo JH, Kim YS. Detection of *qnr* in clinical isolates of *Escherichia coli* from Korea. *Antimicrob Agents Chemother*. 2005; 49:2522–2524. [PubMed: 15917562]
248. Honoré S, Lascols C, Malin D, Targaouchi R, Cattoir V, Legrand P, Soussy CJ, Cambau E. Émergence et diffusion chez les entérobactéries du nouveau mécanisme de résistance plasmidique aux quinolones Qnr (résultats hôpital Henri-Mondor 2002–2005). *Pathol Biol (Paris)*. 2006; 54:270–279. [PubMed: 16473479]
249. Saito R, Kumita W, Sato K, Chida T, Okamura N, Moriya K, Koike K. Detection of plasmid-mediated quinolone resistance associated with *qnrA* in an *Escherichia coli* clinical isolate producing CTX-M-9 β -lactamase in Japan. *Int J Antimicrob Agents*. 2007; 29:600–602. [PubMed: 17239568]
250. Poirel L, Nordmann P, De Champs C, Eloy C. Nosocomial spread of QnrA-mediated quinolone resistance in *Enterobacter sakazakii*. *Int J Antimicrob Agents*. 2007; 29:223–224. [PubMed: 17000086]
251. Park YJ, Yu JK, Lee S, Oh EJ, Woo GJ. Prevalence and diversity of *qnr* alleles in AmpC-producing *Enterobacter cloacae*, *Enterobacter aerogenes*, *Citrobacter freundii* and *Serratia marcescens*: a multicentre study from Korea. *J Antimicrob Chemother*. 2007; 60:868–871. [PubMed: 17660263]
252. Paauw A, Verhoef J, Fluit AC, Blok HE, Hopmans TE, Troelstra A, Leverstein-van Hall MA. Failure to control an outbreak of *qnrA1*-positive multidrug-resistant *Enterobacter cloacae* infection despite adequate implementation of recommended infection control measures. *J Clin Microbiol*. 2007; 45:1420–1425. [PubMed: 17314218]
253. Ellington MJ, Hope R, Turton JF, Warner M, Woodford N, Livermore DM. Detection of *qnrA* among *Enterobacteriaceae* from South-East England with extended-spectrum and high-level AmpC β -lactamases. *J Antimicrob Chemother*. 2007; 60:1176–1178. [PubMed: 17827140]

254. Castanheira M, Pereira AS, Nicoletti AG, Pignatari AC, Barth AL, Gales AC. First report of plasmid-mediated *qnrA1* in a ciprofloxacin-resistant *Escherichia coli* strain in Latin America. *Antimicrob Agents Chemother.* 2007; 51:1527–1529. [PubMed: 17220425]
255. Cavaco LM, Hansen DS, Friis-Møller A, Aarestrup FM, Hasman H, Frimodt-Møller N. First detection of plasmid-mediated quinolone resistance (*qnrA* and *qnrS*) in *Escherichia coli* strains isolated from humans in Scandinavia. *J Antimicrob Chemother.* 2007; 59:804–805. [PubMed: 17284539]
256. Xiong Z, Wang P, Wei Y, Wang H, Cao H, Huang H, Li J. Investigation of *qnr* and *aac(6')-Ib-cr* in *Enterobacter cloacae* isolates from Anhui Province, China. *Diagn Microbiol Infect Dis.* 2008; 62:457–459. [PubMed: 18814992]
257. Tamang MD, Seol SY, Oh JY, Kang HY, Lee JC, Lee YC, Cho DT, Kim J. Plasmid-mediated quinolone resistance determinants *qnrA*, *qnrB*, and *qnrS* among clinical isolates of *Enterobacteriaceae* in a Korean hospital. *Antimicrob Agents Chemother.* 2008; 52:4159–4162. [PubMed: 18725444]
258. Öktem IM, Gülay Z, Biçmen M, Gür D. *qnrA* prevalence in extended-spectrum β -lactamase-positive *Enterobacteriaceae* isolates from Turkey. *Jpn J Infect Dis.* 2008; 61:13–17. [PubMed: 18219128]
259. Murray A, Mather H, Coia JE, Brown DJ. Plasmid-mediated quinolone resistance in nalidixic-acid-susceptible strains of *Salmonella enterica* isolated in Scotland. *J Antimicrob Chemother.* 2008; 62:1153–1155. [PubMed: 18779257]
260. Cavaco LM, Frimodt-Møller N, Hasman H, Guardabassi L, Nielsen L, Aarestrup FM. Prevalence of quinolone resistance mechanisms and associations to minimum inhibitory concentrations in quinolone-resistant *Escherichia coli* isolated from humans and swine in Denmark. *Microb Drug Resist.* 2008; 14:163–169. [PubMed: 18500921]
261. Amin AK, Wareham DW. Plasmid-mediated quinolone resistance genes in *Enterobacteriaceae* isolates associated with community and nosocomial urinary tract infection in East London, UK. *Int J Antimicrob Agents.* 2009; 34:490–491. [PubMed: 19261447]
262. Usein CR, Palade AM, Tatu-Chitoiu D, Ciontea S, Ceciu S, Nica M, Damian M. Identification of plasmid-mediated quinolone resistance *qnr*-like genes in Romanian clinical isolates of *Escherichia coli* and *Klebsiella pneumoniae*. *Roum Arch Microbiol Immunol.* 2009; 68:55–57. [PubMed: 19507629]
263. Kim SY, Park YJ, Yu JK, Kim YS, Han K. Prevalence and characteristics of *aac(6')-Ib-cr* in AmpC-producing *Enterobacter cloacae*, *Citrobacter freundii*, and *Serratia marcescens*: a multicenter study from Korea. *Diagn Microbiol Infect Dis.* 2009; 63:314–318. [PubMed: 19216942]
264. Gunell M, Webber MA, Kotilainen P, Lilly AJ, Caddick JM, Jalava J, Huovinen P, Siitonen A, Hakanen AJ, Piddock LJ. Mechanisms of resistance in nontyphoidal *Salmonella enterica* strains exhibiting a nonclassical quinolone resistance phenotype. *Antimicrob Agents Chemother.* 2009; 53:3832–3836. [PubMed: 19596880]
265. Sjölund-Karlsson M, Folster JP, Pecic G, Joyce K, Medalla F, Rickert R, Whichard JM. Emergence of plasmid-mediated quinolone resistance among non-Typhi *Salmonella enterica* isolates from humans in the United States. *Antimicrob Agents Chemother.* 2009; 53:2142–2144. [PubMed: 19223618]
266. Cui S, Li J, Sun Z, Hu C, Jin S, Li F, Guo Y, Ran L, Ma Y. Characterization of *Salmonella enterica* isolates from infants and toddlers in Wuhan, China. *J Antimicrob Chemother.* 2009; 63:87–94. [PubMed: 18984647]
267. Ode T, Saito R, Kumita W, Sato K, Okugawa S, Moriya K, Koike K, Okamura N. Analysis of plasmid-mediated multidrug resistance in *Escherichia coli* and *Klebsiella oxytoca* isolates from clinical specimens in Japan. *Int J Antimicrob Agents.* 2009; 34:347–350. [PubMed: 19560903]
268. Fang H, Huang H, Shi Y, Hedin G, Nord CE, Ullberg M. Prevalence of *qnr* determinants among extended-spectrum β -lactamase-positive *Enterobacteriaceae* clinical isolates in southern Stockholm, Sweden. *Int J Antimicrob Agents.* 2009; 34:268–270. [PubMed: 19409763]
269. Naqvi SM, Jenkins C, McHugh TD, Balakrishnan I. Identification of the *qnr* family in *Enterobacteriaceae* in clinical practice. *J Antimicrob Chemother.* 2009; 63:830–832. [PubMed: 19218277]

270. Le TM, Baker S, Le TP, Cao TT, Tran TT, Nguyen VM, Campbell JI, Lam MY, Nguyen TH, Nguyen VV, Farrar J, Schultz C. High prevalence of plasmid-mediated quinolone resistance determinants in commensal members of the *Enterobacteriaceae* in Ho Chi Minh City, Vietnam. *J Med Microbiol*. 2009; 58:1585–1592. [PubMed: 19696153]
271. Bouchakour M, Zerouali K, Gros Claude JD, Amarouch H, El Mdaghri N, Courvalin P, Timinouni M. Plasmid-mediated quinolone resistance in expanded spectrum beta lactamase producing *enterobacteriaceae* in Morocco. *Journal of infection in developing countries*. 2010; 4:779–803. [PubMed: 21252459]
272. Xiong Z, Li J, Li T, Shen J, Hu F, Wang M. Prevalence of plasmid-mediated quinolone-resistance determinants in *Shigella flexneri* isolates from Anhui Province, China. *J Antibiot (Tokyo)*. 2010; 63:187–189. [PubMed: 20203702]
273. Kim YS, Kim ES, Jeong JY. Genetic organization of plasmid-mediated Qnr determinants in cefotaxime-resistant *Enterobacter cloacae* isolates in Korea. *Diagn Microbiol Infect Dis*. 2010; 68:318–321. [PubMed: 20846811]
274. Ferreira S, Paradela A, Velez J, Ramalheira E, Walsh TR, Mendo S. Carriage of *qnrA1* and *qnrB2*, *bla_{CTX-M15}*, and complex class 1 integron in a clinical multiresistant *Citrobacter freundii* isolate. *Diagn Microbiol Infect Dis*. 2010; 67:188–190. [PubMed: 20338709]
275. Kiiru J, Kariuki S, Goddeeris BM, Revathi G, Maina TW, Ndegwa DW, Muyodi J, Butaye P. *Escherichia coli* strains from Kenyan patients carrying conjugatively transferable broad-spectrum beta-lactamase, *qnr*, *aac(6)-Ib-cr* and 16S rRNA methyltransferase genes. *J Antimicrob Chemother*. 2011; 66:1639–1642. [PubMed: 21474478]
276. Herrera-Leon S, Gonzalez-Sanz R, Herrera-Leon L, Echeita MA. Characterization of multidrug-resistant Enterobacteriaceae carrying plasmid-mediated quinolone resistance mechanisms in Spain. *J Antimicrob Chemother*. 2011; 66:287–290. [PubMed: 21115442]
277. Garcia-Fulgueiras V, Bado I, Mota MI, Robino L, Cordeiro NF, Varela A, Algorta G, Gutkind G, Ayala JA, Vignoli R. Extended-spectrum β -lactamases and plasmid-mediated quinolone resistance in enterobacterial clinical isolates in the paediatric hospital of Uruguay. *J Antimicrob Chemother*. 2011; 66:1725–1729. [PubMed: 21685201]
278. Chowdhury G, Pazhani GP, Nair GB, Ghosh A, Ramamurthy T. Transferable plasmid-mediated quinolone resistance in association with extended-spectrum β -lactamases and fluoroquinolone-acetylating aminoglycoside-6'-N-acetyltransferase in clinical isolates of *Vibrio fluvialis*. *Int J Antimicrob Agents*. 2011; 38:169–173. [PubMed: 21683552]
279. Silva-Sanchez J, Barrios H, Reyna-Flores F, Bello-Diaz M, Sanchez-Perez A, Rojas T, Consortium BR, Garza-Ramos U. Prevalence and characterization of plasmid-mediated quinolone resistance genes in extended-spectrum β -lactamase-producing *Enterobacteriaceae* isolates in Mexico. *Microb Drug Resist*. 2011; 17:497–505. [PubMed: 21834663]
280. Muller S, Oesterlein A, Frosch M, Abele-Horn M, Valenza G. Characterization of extended-spectrum beta-lactamases and *qnr* plasmid-mediated quinolone resistance in German isolates of *Enterobacter* species. *Microb Drug Resist*. 2011; 17:99–103. [PubMed: 21190474]
281. Jeong HS, Bae IK, Shin JH, Kim SH, Chang CL, Jeong J, Kim S, Lee CH, Ryoo NH, Lee JN. Fecal colonization of *Enterobacteriaceae* carrying plasmid-mediated quinolone resistance determinants in Korea. *Microb Drug Resist*. 2011; 17:507–512. [PubMed: 21830908]
282. Frank T, Mbecko JR, Misatou P, Monchy D. Emergence of quinolone resistance among extended-spectrum β -lactamase-producing *Enterobacteriaceae* in the Central African Republic: genetic characterization. *BMC research notes*. 2011; 4:309. [PubMed: 21867486]
283. Tran QT, Nawaz MS, Deck J, Nguyen KT, Cerniglia CE. Plasmid-mediated quinolone resistance in *Pseudomonas putida* isolates from imported shrimp. *Appl Environ Microbiol*. 2011; 77:1885–1887. [PubMed: 21193671]
284. Hassan WM, Hashim A, Domany R. Plasmid mediated quinolone resistance determinants *qnr*, *aac(6)-Ib-cr*, and *qep* in ESBL-producing *Escherichia coli* clinical isolates from Egypt. *Indian J Med Microbiol*. 2012; 30:442–447. [PubMed: 23183470]
285. Paltansing S, Kraakman ME, Ras JM, Wessels E, Bernards AT. Characterization of fluoroquinolone and cephalosporin resistance mechanisms in Enterobacteriaceae isolated in a Dutch teaching hospital reveals the presence of an *Escherichia coli* ST131 clone with a specific mutation in *parE*. *J Antimicrob Chemother*. 2013; 68:40–45. [PubMed: 22991424]

286. Wang Y, Ying C, Wu W, Ye Y, Zhang H, Yu J. Investigation of quinolone resistance in *Shigella sonnei*. Chin J Infect Chemother. 2009; 9:27–31.
287. Pasom W, Chanawong A, Lulitanond A, Wilailuckana C, Kenprom S, Puang-Ngern P. Plasmid-mediated quinolone resistance genes, *aac(6′)-Ib-cr*, *qnrS*, *qnrB*, and *qnrA*, in urinary isolates of *Escherichia coli* and *Klebsiella pneumoniae* at a teaching hospital, Thailand. Jpn J Infect Dis. 2013; 66:428–432. [PubMed: 24047744]
288. Silva-Sánchez J, Cruz-Trujillo E, Barrios H, Reyna-Flores F, Sánchez-Pérez A, Consortium BR, Garza-Ramos U. Characterization of plasmid-mediated quinolone resistance (PMQR) genes in extended-spectrum β -lactamase-producing *Enterobacteriaceae* pediatric clinical isolates in Mexico. PLoS One. 2013; 8:e77968. [PubMed: 24147104]
289. Ferrari R, Galiana A, Cremades R, Rodriguez JC, Magnani M, Tognim MC, Oliveira TC, Royo G. Plasmid-mediated quinolone resistance (PMQR) and mutations in the topoisomerase genes of *Salmonella enterica* strains from Brazil. Braz J Microbiol. 2013; 44:651–656. [PubMed: 24294265]
290. Xia R, Ren Y, Xu H. Identification of plasmid-mediated quinolone resistance *qnr* genes in multidrug-resistant gram-negative bacteria from hospital wastewaters and receiving waters in the Jinan area, China. Microb Drug Resist. 2013; 19:446–456. [PubMed: 23844849]
291. Chu YW, Cheung TK, Ng TK, Tsang D, To WK, Kam KM, Lo JY. Quinolone resistance determinant *qnrA3* in clinical isolates of *Salmonella* in 2000–2005 in Hong Kong. J Antimicrob Chemother. 2006; 58:904–905. [PubMed: 16943211]
292. Dahmen S, Poirel L, Mansour W, Bouallegue O, Nordmann P. Prevalence of plasmid-mediated quinolone resistance determinants in *Enterobacteriaceae* from Tunisia. Clin Microbiol Infect. 2010; 16:1019–1023. [PubMed: 19650824]
293. Arpin C, Thabet L, Yassine H, Messadi AA, Boukadida J, Dubois V, Coulange-Mayonnove L, Andre C, Quentin C. Evolution of an incompatibility group IncA/C plasmid harboring *bla_{CMY-16}* and *qnrA6* genes and its transfer through three clones of *Providencia stuartii* during a two-year outbreak in a Tunisian burn unit. Antimicrob Agents Chemother. 2012; 56:1342–1349. [PubMed: 22155825]
294. Nazik H, Ongen B, Kuvat N. Investigation of plasmid-mediated quinolone resistance among isolates obtained in a Turkish intensive care unit. Jpn J Infect Dis. 2008; 61:310–312. [PubMed: 18653977]
295. Torpdahl M, Hammerum AM, Zachariassen C, Nielsen EM. Detection of *qnr* genes in *Salmonella* isolated from humans in Denmark. J Antimicrob Chemother. 2009; 63:406–408. [PubMed: 19056747]
296. Karah N, Poirel L, Bengtsson S, Sundqvist M, Kahlmeter G, Nordmann P, Sundsfjord A, Samuelsen O. Plasmid-mediated quinolone resistance determinants *qnr* and *aac(6′)-Ib-cr* in *Escherichia coli* and *Klebsiella* spp. from Norway and Sweden. Diagn Microbiol Infect Dis. 2010; 66:425–431. [PubMed: 20226333]
297. Meradi L, Djahoudi A, Abdi A, Bouchakour M, Perrier Gros Claude JD, Timinouni M. Résistance aux quinolones de types *qnr*, *aac(6′)-Ib-cr* chez les entérobactéries isolées à Annaba en Algérie. Pathol Biol (Paris). 2011; 59:e73–78. [PubMed: 19896287]
298. Villa L, Poirel L, Nordmann P, Carta C, Carattoli A. Complete sequencing of an IncH plasmid carrying the *bla_{NDM-1}*, *bla_{CTX-M-15}* and *qnrB1* genes. J Antimicrob Chemother. 2012; 67:1645–1650. [PubMed: 22511638]
299. Guillard T, de Champs C, Moret H, Bertrand X, Scheftel JM, Cambau E. High-resolution melting analysis for rapid characterization of *qnr* alleles in clinical isolates and detection of two novel alleles, *qnrB25* and *qnrB42*. J Antimicrob Chemother. 2012; 67:2635–2639. [PubMed: 22850691]
300. Dolejska M, Villa L, Dobiasova H, Fortini D, Feudi C, Carattoli A. Plasmid content of a clinically relevant *Klebsiella pneumoniae* clone from the Czech Republic producing CTX-M-15 and *QnrB1*. Antimicrob Agents Chemother. 2013; 57:1073–1076. [PubMed: 23229477]
301. Viana AL, Cayo R, Avelino CC, Gales AC, Franco MC, Minarini LA. Extended-spectrum β -lactamases in *Enterobacteriaceae* isolated in Brazil carry distinct types of plasmid-mediated quinolone resistance genes. J Med Microbiol. 2013; 62:1326–1331. [PubMed: 23741024]

302. Saiful Anuar AS, Mohd Yusof MY, Tay ST. Prevalence of plasmid-mediated *qnr* determinants and gyrase alteration in *Klebsiella pneumoniae* isolated from a university teaching hospital in Malaysia. *Eur Rev Med Pharmacol Sci*. 2013; 17:1744–1747. [PubMed: 23852897]
303. Gay K, Robicsek A, Strahilevitz J, Park CH, Jacoby G, Barrett TJ, Medalla F, Chiller TM, Hooper DC. Plasmid-mediated quinolone resistance in non-Typhi serotypes of *Salmonella enterica*. *Clin Infect Dis*. 2006; 43:297–304. [PubMed: 16804843]
304. Whichard J, Gay K, Stevenson JE, Joyce K, Cooper K, Omondi M, F M, Jacoby GA, Barrett TJ. Human *Salmonella* and concurrent decreased susceptibility to quinolones and extended-spectrum cephalosporins. *Emerg Infect Dis*. 2007; 13:1681–1688. [PubMed: 18217551]
305. Hopkins KL, Day M, Threlfall EJ. Plasmid-mediated quinolone resistance in *Salmonella enterica*, United Kingdom. *Emerg Infect Dis*. 2008; 14:340–342. [PubMed: 18258138]
306. Minarini LA, Poirel L, Cattoir V, Darini AL, Nordmann P. Plasmid-mediated quinolone resistance determinants among enterobacterial isolates from outpatients in Brazil. *J Antimicrob Chemother*. 2008; 62:474–478. [PubMed: 18552340]
307. Wu JJ, Ko WC, Chiou CS, Chen HM, Wang LR, Yan JJ. Emergence of *Qnr* determinants in human *Salmonella* isolates in Taiwan. *J Antimicrob Chemother*. 2008; 62:1269–1272. [PubMed: 18957397]
308. Veldman K, van Pelt W, Mevius D. First report of *qnr* genes in *Salmonella* in The Netherlands. *J Antimicrob Chemother*. 2008; 61:452–453. [PubMed: 18086694]
309. Shin JH, Jung HJ, Lee JY, Kim HR, Lee JN, Chang CL. High rates of plasmid-mediated quinolone resistance *QnrB* variants among ciprofloxacin-resistant *Escherichia coli* and *Klebsiella pneumoniae* from urinary tract infections in Korea. *Microb Drug Resist*. 2008; 14:221–226. [PubMed: 18707554]
310. Liassine N, Zulueta-Rodriguez P, Corbel C, Lascols C, Soussy CJ, Cambau E. First detection of plasmid-mediated quinolone resistance in the community setting and in hospitalized patients in Switzerland. *J Antimicrob Chemother*. 2008; 62:1151–1152. [PubMed: 18669519]
311. Gutierrez B, Herrera-Leon S, Escudero JA, Hidalgo L, Gonzalez-Sanz R, Arroyo M, San Millan A, Echeita MA, Gonzalez-Zorn B. Novel genetic environment of *qnrB2* associated with TEM-1 and SHV-12 on pB1004, an IncHI2 plasmid, in *Salmonella* Bredeney BB1047 from Spain. *J Antimicrob Chemother*. 2009; 64:1334–1336. [PubMed: 19773254]
312. Pfeifer Y, Matten J, Rabsch W. *Salmonella enterica* serovar Typhi with CTX-M β -lactamase, Germany. *Emerg Infect Dis*. 2009; 15:1533–1535. [PubMed: 19788837]
313. Cattoir V, Poirel L, Rotimi V, Soussy CJ, Nordmann P. Multiplex PCR for detection of plasmid-mediated quinolone resistance *qnr* genes in ESBL-producing enterobacterial isolates. *J Antimicrob Chemother*. 2007; 60:394–397. [PubMed: 17561500]
314. Pallecchi L, Riccobono E, Mantella A, Bartalesi F, Sennati S, Gamboa H, Gotuzzo E, Bartoloni A, Rossolini GM. High prevalence of *qnr* genes in commensal enterobacteria from healthy children in Peru and Bolivia. *Antimicrob Agents Chemother*. 2009; 53:2632–2635. [PubMed: 19364872]
315. Sjölund-Karlsson M, Howie R, Rickert R, Krueger A, Tran TT, Zhao S, Ball T, Haro J, Pecic G, Joyce K, Fedorka-Cray PJ, Whichard JM, McDermott PF. Plasmid-mediated quinolone resistance among non-Typhi *Salmonella enterica* isolates, USA. *Emerg Infect Dis*. 2010; 16:1789–1791. [PubMed: 21029547]
316. Lin CJ, Siu LK, Ma L, Chang YT, Lu PL. Molecular epidemiology of ciprofloxacin-resistant extended-spectrum β -lactamase-producing *Klebsiella pneumoniae* in Taiwan. *Microb Drug Resist*. 2012; 18:52–58. [PubMed: 22022870]
317. Nordmann P, Poirel L, Mak JK, White PA, McIver CJ, Taylor P. Multidrug-resistant *Salmonella* strains expressing emerging antibiotic resistance determinants. *Clin Infect Dis*. 2008; 46:324–325. [PubMed: 18171274]
318. Kanamori H, Yano H, Hirakata Y, Endo S, Arai K, Ogawa M, Shimojima M, Aoyagi T, Hatta M, Yamada M, Nishimaki K, Kitagawa M, Kunishima H, Kaku M. High prevalence of extended-spectrum β -lactamases and *qnr* determinants in *Citrobacter* species from Japan: dissemination of CTX-M-2. *J Antimicrob Chemother*. 2011; 66:2255–2262. [PubMed: 21733965]

319. Zhu YL, Yang HF, Liu YY, Hu LF, Cheng J, Ye Y, Li JB. Detection of plasmid-mediated quinolone resistance determinants and the emergence of multidrug resistance in clinical isolates of *Shigella* in SiXian area, China. *Diagn Microbiol Infect Dis*. 2013; 75:327–329. [PubMed: 23410262]
320. Cavaco LM, Korsgaard H, Sorensen G, Aarestrup FM. Plasmid-mediated quinolone resistance due to *qnrB5* and *qnrS1* genes in *Salmonella enterica* serovars Newport, Hadar and Saintpaul isolated from turkey meat in Denmark. *J Antimicrob Chemother*. 2008; 62:632–634. [PubMed: 18495652]
321. Lunn AD, Fabrega A, Sanchez-Cespedes J, Vila J. Prevalence of mechanisms decreasing quinolone-susceptibility among *Salmonella* spp. clinical isolates. *International microbiology : the official journal of the Spanish Society for Microbiology*. 2010; 13:15–20. [PubMed: 20890836]
322. Folster JP, Pecic G, Bowen A, Rickert R, Carattoli A, Whichard JM. Decreased susceptibility to ciprofloxacin among *Shigella* isolates in the United States, 2006 to 2009. *Antimicrob Agents Chemother*. 2011; 55:1758–1760. [PubMed: 21220535]
323. Zhao JY, Dang H. Coastal seawater bacteria harbor a large reservoir of plasmid-mediated quinolone resistance determinants in Jiaozhou Bay, China. *Microb Ecol*. 2012; 64:187–199. [PubMed: 22252223]
324. Fortini D, Fashae K, Garcia-Fernandez A, Villa L, Carattoli A. Plasmid-mediated quinolone resistance and β -lactamases in *Escherichia coli* from healthy animals from Nigeria. *J Antimicrob Chemother*. 2011; 66:1269–1272. [PubMed: 21393162]
325. Hammerl JA, Beutlich J, Hertwig S, Mevius D, Threlfall EJ, Helmuth R, Guerra B. pSGI15, a small ColE-like *qnrB19* plasmid of a *Salmonella enterica* serovar Typhimurium strain carrying *Salmonella* genomic island 1 (SGI1). *J Antimicrob Chemother*. 2010; 65:173–175. [PubMed: 19861336]
326. Pallecchi L, Riccobono E, Mantella A, Fernandez C, Bartalesi F, Rodriguez H, Gotuzzo E, Bartoloni A, Rossolini GM. Small *qnrB*-harbouring ColE-like plasmids widespread in commensal enterobacteria from a remote Amazonas population not exposed to antibiotics. *J Antimicrob Chemother*. 2011; 66:1176–1178. [PubMed: 21393218]
327. Jeong HS, Kim JA, Shin JH, Chang CL, Jeong J, Cho JH, Kim MN, Kim S, Kim YR, Lee CH, Lee K, Lee MA, Lee WG, Lee JN. Prevalence of plasmid-mediated quinolone resistance and mutations in the gyrase and topoisomerase IV genes in *Salmonella* isolated from 12 tertiary-care hospitals in Korea. *Microb Drug Resist*. 2011; 17:551–557. [PubMed: 21830947]
328. Riveros M, Riccobono E, Durand D, Mosquito S, Ruiz J, Rossolini GM, Ochoa TJ, Pallecchi L. Plasmid-mediated quinolone resistance genes in enteroaggregative *Escherichia coli* from infants in Lima, Peru. *Int J Antimicrob Agents*. 2012; 39:540–542. [PubMed: 22521526]
329. Gonzalez F, Pallecchi L, Rossolini GM, Araque M. Plasmid-mediated quinolone resistance determinant *qnrB19* in non-typhoidal *Salmonella enterica* strains isolated in Venezuela. *Journal of infection in developing countries*. 2012; 6:462–464. [PubMed: 22610715]
330. Bae IK, Park I, Lee JJ, Sun HI, Park KS, Lee JE, Ahn JH, Lee SH, Woo GJ. Novel variants of the *qnrB* gene, *qnrB22* and *qnrB23*, in *Citrobacter werkmanii* and *Citrobacter freundii*. *Antimicrob Agents Chemother*. 2010; 54:3068–3069. [PubMed: 20421404]
331. Wang D, Wang H, Qi Y, Liang Y, Zhang J, Yu L. Novel variants of the *qnrB* gene, *qnrB31* and *qnrB32*, in *Klebsiella pneumoniae*. *J Med Microbiol*. 2011; 60:1849–1852. [PubMed: 21816942]
332. Wang D, Wang H, Qi Y, Liang Y, Zhang J, Yu L. Characteristics of *Klebsiella pneumoniae* harboring *QnrB32*, *Aac(6′)-Ib-cr*, *GyrA* and *CTX-M-22* genes. *Folia Histochem Cytobiol*. 2012; 50:68–74. [PubMed: 22532138]
333. Poirel L, Nguyen TV, Weintraub A, Leviandier C, Nordmann P. Plasmid-mediated quinolone resistance determinant *qnrS* in *Enterobacter cloacae*. *Clin Microbiol Infect*. 2006; 12:1021–1023. [PubMed: 16961640]
334. Avsaroglu MD, Helmuth R, Junker E, Hertwig S, Schroeter A, Akcelik M, Bozoglu F, Guerra B. Plasmid-mediated quinolone resistance conferred by *qnrS1* in *Salmonella enterica* serovar Virchow isolated from Turkish food of avian origin. *J Antimicrob Chemother*. 2007; 60:1146–1150. [PubMed: 17881633]

335. Vasilaki O, Ntokou E, Ikonomidis A, Sofianou D, Frantzidou F, Alexiou-Daniel S, Maniatis AN, Pournaras S. Emergence of the plasmid-mediated quinolone resistance gene *qnrS1* in *Escherichia coli* isolates in Greece. *Antimicrob Agents Chemother*. 2008; 52:2996–2997. [PubMed: 18490503]
336. Pu XY, Pan JC, Wang HQ, Zhang W, Huang ZC, Gu YM. Characterization of fluoroquinolone-resistant *Shigella flexneri* in Hangzhou area of China. *J Antimicrob Chemother*. 2009; 63:917–920. [PubMed: 19297378]
337. Keddy KH, Smith AM, Sooka A, Ismail H, Oliver S. Fluoroquinolone-resistant typhoid, South Africa. *Emerg Infect Dis*. 2010; 16:879–880. [PubMed: 20409393]
338. Le TM, AbuOun M, Morrison V, Thomson N, Campbell JI, Woodward MJ, Van Vinh Chau N, Farrar J, Schultz C, Baker S. Differential phenotypic and genotypic characteristics of *qnrS1*-harboring plasmids carried by hospital and community commensal Enterobacteria. *Antimicrob Agents Chemother*. 2011; 55:1798–1802. [PubMed: 21282449]
339. Sato T, Yokota S, Uchida I, Okubo T, Ishihara K, Fujii N, Tamura Y. A fluoroquinolone-resistant *Escherichia coli* clinical isolate without quinolone resistance-determining region mutations found in Japan. *Antimicrob Agents Chemother*. 2011; 55:3964–3965. [PubMed: 21670180]
340. Mokracka J, Gruszczynska B, Kaznowski A. Integrons, β -lactamase and *qnr* genes in multidrug resistant clinical isolates of *Proteus mirabilis* and *P. vulgaris*. *APMIS*. 2012; 120:950–958. [PubMed: 23030307]
341. Kanamori H, Yano H, Hirakata Y, Hirotani A, Arai K, Endo S, Ichimura S, Ogawa M, Shimojima M, Aoyagi T, Hatta M, Yamada M, Gu Y, Tokuda K, Kunishima H, Kitagawa M, Kaku M. Molecular characteristics of extended-spectrum β -lactamases and *qnr* determinants in *Enterobacter* species from Japan. *PLoS ONE*. 2012; 7:e37967. [PubMed: 22719857]
342. Sánchez-Céspedes J, Blasco MD, Martí S, Alba V, Alcalde E, Esteve C, Vila J. Plasmid-mediated QnrS2 determinant from a clinical *Aeromonas veronii* isolate. *Antimicrob Agents Chemother*. 2008; 52:2990–2991. [PubMed: 18505860]
343. Arias A, Seral C, Navarro F, Miro E, Coll P, Castillo FJ. Plasmid-mediated QnrS2 determinant in an *Aeromonas caviae* isolate recovered from a patient with diarrhoea. *Clin Microbiol Infect*. 2010; 16:1005–1007. [PubMed: 19863591]
344. Majumdar T, Das B, Bhadra RK, Dam B, Mazumder S. Complete nucleotide sequence of a quinolone resistance gene (*qnrS2*) carrying plasmid of *Aeromonas hydrophila* isolated from fish. *Plasmid*. 2011; 66:79–84. [PubMed: 21664930]
345. Han JE, Kim JH, Cheresca CH Jr, Shin SP, Jun JW, Chai JY, Han SY, Park SC. First description of the *qnrS*-like (*qnrS5*) gene and analysis of quinolone resistance-determining regions in motile *Aeromonas* spp. from diseased fish and water. *Res Microbiol*. 2012; 163:73–79. [PubMed: 22024339]
346. Kristiansson E, Fick J, Janzon A, Grabic R, Rutgersson C, Weijdegard B, Soderstrom H, Larsson DG. Pyrosequencing of antibiotic-contaminated river sediments reveals high levels of resistance and gene transfer elements. *PLoS One*. 2011; 6:e17038. [PubMed: 21359229]
347. Li J, Wang T, Shao B, Shen J, Wang S, Wu Y. Plasmid-mediated quinolone resistance genes and antibiotic residues in wastewater and soil adjacent to Swine feedlots: potential transfer to agricultural lands. *Environmental health perspectives*. 2012; 120:1144–1149. [PubMed: 22569244]
348. Karisik E, Ellington MJ, Pike R, Warren RE, Livermore DM, Woodford N. Molecular characterization of plasmids encoding CTX-M-15 β -lactamases from *Escherichia coli* strains in the United Kingdom. *J Antimicrob Chemother*. 2006; 58:665–668. [PubMed: 16870648]
349. Machado E, Coque TM, Cantón R, Baquero F, Sousa JC, Peixe L. Dissemination in Portugal of CTX-M-15-, OXA-1-, and TEM-1-producing *Enterobacteriaceae* strains containing the *aac(6)-Ib-cr* gene, which encodes an aminoglycoside- and fluoroquinolone-modifying enzyme. *Antimicrob Agents Chemother*. 2006; 50:3220–3221. [PubMed: 16940136]
350. Pallecchi L, Bartoloni A, Fiorelli C, Mantella A, Di Maggio T, Gamboa H, Gotuzzo E, Kronvall G, Paradisi F, Rossolini GM. Rapid dissemination and diversity of CTX-M extended-spectrum β -lactamase genes in commensal *Escherichia coli* isolates from healthy children from low-resource settings in Latin America. *Antimicrob Agents Chemother*. 2007; 51:2720–2725. [PubMed: 17548490]

351. Ambroži Avguštin J, Keber R, Žerjavi K, Oražem T, Grabnar M. Emergence of the quinolone resistance-mediating gene *aac(6')-Ib-cr* in extended-spectrum-β-lactamase-producing *Klebsiella* isolates collected in Slovenia between 2000 and 2005. *Antimicrob Agents Chemother.* 2007; 51:4171–4173. [PubMed: 17846136]
352. Cordeiro NF, Robino L, Medina J, Seija V, Bado I, Garcia V, Berro M, Pontet J, Lopez L, Bazet C, Rieppi G, Gutkind G, Ayala JA, Vignoli R. Ciprofloxacin-resistant enterobacteria harboring the *aac(6')-Ib-cr* variant isolated from feces of inpatients in an intensive care unit in Uruguay. *Antimicrob Agents Chemother.* 2008; 52:806–807. [PubMed: 17999962]
353. Chmelnitsky I, Hermesh O, Navon-Venezia S, Strahilevitz J, Carmeli Y. Detection of *aac(6')-Ib-cr* in KPC-producing *Klebsiella pneumoniae* isolates from Tel Aviv, Israel. *J Antimicrob Chemother.* 2009; 64:718–722. [PubMed: 19656782]
354. Deepak RN, Koh TH, Chan KS. Plasmid-mediated quinolone resistance determinants in urinary isolates of *Escherichia coli* and *Klebsiella pneumoniae* in a large Singapore hospital. *Ann Acad Med Singapore.* 2009; 38:1070–1073. [PubMed: 20052442]
355. Morgan-Linnell SK, Becnel Boyd L, Steffen D, Zechiedrich L. Mechanisms accounting for fluoroquinolone resistance in *Escherichia coli* clinical isolates. *Antimicrob Agents Chemother.* 2009; 53:235–241. [PubMed: 18838592]
356. Bell JM, Turnidge JD, Andersson P. *aac(6')-Ib-cr* genotyping by simultaneous high-resolution melting analyses of an unlabeled probe and full-length amplicon. *Antimicrob Agents Chemother.* 2010; 54:1378–1380. [PubMed: 20008777]
357. Hidalgo-Grass C, Strahilevitz J. High-resolution melt curve analysis for identification of single nucleotide mutations in the quinolone resistance gene *aac(6')-Ib-cr*. *Antimicrob Agents Chemother.* 2010; 54:3509–3511. [PubMed: 20498320]
358. Christiansen N, Nielsen L, Jakobsen L, Stegger M, Hansen LH, Frimodt-Moller N. Fluoroquinolone resistance mechanisms in urinary tract pathogenic *Escherichia coli* isolated during rapidly increasing fluoroquinolone consumption in a low-use country. *Microb Drug Resist.* 2011; 17:395–406. [PubMed: 21668371]
359. Frasson I, Cavallaro A, Bergo C, Richter SN, Palu G. Prevalence of *aac(6')-Ib-cr* plasmid-mediated and chromosome-encoded fluoroquinolone resistance in *Enterobacteriaceae* in Italy. *Gut Pathog.* 2011; 3:12. [PubMed: 21827661]
360. Park KS, Kim MH, Park TS, Nam YS, Lee HJ, Suh JT. Prevalence of the plasmid-mediated quinolone resistance genes, *aac(6')-Ib-cr*, *qepA*, and *oqxAB* in clinical isolates of extended-spectrum β-lactamase (ESBL)-producing *Escherichia coli* and *Klebsiella pneumoniae* in Korea. *Ann Clin Lab Sci.* 2012; 42:191–197. [PubMed: 22585617]
361. Bartoloni A, Pallecchi L, Riccobono E, Mantella A, Magnelli D, Di Maggio T, Villagran AL, Lara Y, Saavedra C, Strohmeyer M, Bartalesi F, Trigo C, Rossolini GM. Relentless increase of resistance to fluoroquinolones and expanded-spectrum cephalosporins in *Escherichia coli*: 20 years of surveillance in resource-limited settings from Latin America. *Clin Microbiol Infect.* 2013; 19:356–361. [PubMed: 22414066]
362. Chen DQ, Yang L, Luo YT, Mao MJ, Lin YP, Wu AW. Prevalence and characterization of quinolone resistance in *Laribacter hongkongensis* from grass carp and Chinese tiger frog. *J Med Microbiol.* 2013; 62:1559–1564. [PubMed: 23906590]
363. Liu BT, Wang XM, Liao XP, Sun J, Zhu HQ, Chen XY, Liu YH. Plasmid-mediated quinolone resistance determinants *oqxAB* and *aac(6')-Ib-cr* and extended-spectrum β-lactamase gene *bla_{CTX-M-24}* co-located on the same plasmid in one *Escherichia coli* strain from China. *J Antimicrob Chemother.* 2011; 66:1638–1639. [PubMed: 21546384]
364. Yang H, Duan G, Zhu J, Zhang W, Xi Y, Fan Q. Prevalence and characterisation of plasmid-mediated quinolone resistance and mutations in the gyrase and topoisomerase IV genes among *Shigella* isolates from Henan, China, between 2001 and 2008. *Int J Antimicrob Agents.* 2013; 42:173–177. [PubMed: 23796894]
365. Rocha-Gracia R, Ruiz E, Romero-Romero S, Lozano-Zarain P, Somalo S, Palacios-Hernández JM, Caballero-Torres P, Torres C. Detection of the plasmid-borne quinolone resistance determinant *qepA1* in a CTX-M-15-producing *Escherichia coli* strain from Mexico. *J Antimicrob Chemother.* 2010; 65:169–171. [PubMed: 19910326]

366. Deng Y, He L, Chen S, Zheng H, Zeng Z, Liu Y, Sun Y, Ma J, Chen Z, Liu JH. F33:A-:B- and F2:A-:B- plasmids mediate dissemination of *rmtB-bla*_{CTX-M-9} group genes and *rmtB-qepA* in *Enterobacteriaceae* isolates from pets in China. *Antimicrob Agents Chemother*. 2011; 55:4926–4929. [PubMed: 21788459]
367. Tian GB, Rivera JI, Park YS, Johnson LE, Hingwe A, Adams-Haduch JM, Doi Y. Sequence type ST405 *Escherichia coli* isolate producing QepA1, CTX-M-15, and RmtB from Detroit, Michigan. *Antimicrob Agents Chemother*. 2011; 55:3966–3967. [PubMed: 21670190]

Fig. 1. The rod-like structure of the QnrB1 dimer is shown above with the sequence of the monomer below. The sequence is divided into four columns representing the four faces of the right-handed quadrilateral β -helix. Face names and color are shown at the top along with the naming convention for the five residues of the pentapeptide repeats. Loops A and B are indicated by one and two asterisks, respectively, with their sequences indicated below and the loops shown as black traces on the diagram. The N-terminal α -helix is colored salmon. The molecular 2-fold symmetry is indicated with a black diamond. Type II turn (235) containing faces are shown as spheres and type IV-containing faces as strands. Adapted from reference (44). © the American Society for Biochemistry and Molecular Biology.

Fig. 2.

QnrB1 protection of DNA gyrase from ciprofloxacin inhibition of supercoiling. Reaction mixtures of 30 μ l were analyzed by agarose gel electrophoresis. Reaction mixtures contained 0.2 μ g relaxed pBR322 DNA (lanes 1 to 14), 6.7 nM gyrase (lanes 2 to 14), 2 μ g/ml ciprofloxacin (lanes 3 to 14), and QnrB-His6 fusion protein at 25 μ M (lane 4), 5 μ M (lane 5), 2.5 μ M (lane 6), 0.5 μ M (lane 7), 50 nM (lane 8), 5 nM (lane 9), 0.5 nM (lane 10), 50 pM (lane 11), 5 pM (lane 12), or 0.5 pM (lane 13). Reprinted from (27)

Fig. 3.
Genetic environment of *qnr* alleles.

Fig. 4. Survival at increasing fluoroquinolone concentrations for *Escherichia coli* J53 and J53 pMG252. A large inoculum (10^{10} colony forming units) and appropriate dilutions were applied to Mueller-Hinton agar plates containing the indicated concentration of ciprofloxacin, and surviving colonies were counted after incubation for 72 h at 37°C [from (215)] with permission.

Table 1

Representative plasmids and transmissible PMQR genes

Plasmid	PMQR gene	Host	Year of isolation	Size (kb)	Inc group	Country	Linked <i>bla</i> genes ^a	Reference
pMG252	<i>qnrA1</i>	<i>K. pneumoniae</i>	1994	~180		USA	FOX-5	(9)
pHSH2	<i>qnrA1</i>	<i>E. coli</i>	2000–2001	85		China		(24)
pQR1	<i>qnrA1</i>	<i>E. coli</i>	2003	180	A/C ₂	France	VEB-1	(126, 128)
<i>b</i>	<i>qnrA1</i>	<i>E. cloacae</i>	2002–2005	75	H12	France	SHV-12	(119, 126)
<i>b</i>	<i>qnrA1</i>	<i>E. aerogenes</i>	2002–2005	150	FII	France	SHV-12	(119, 126)
pSZ50	<i>qnrA1</i>	<i>E. coli</i>	2004	~50	N	Mexico		(71)
pHE96	<i>qnrA3</i>	<i>K. pneumoniae</i>	2004	70	N	France		(236)
<i>b</i>	<i>qnrA6</i>	<i>P. stuartii</i>	2011	100	A/C	Tunisia	OXA-48, PER-1, CMY-4	(237)
pMG298	<i>qnrB1</i>	<i>K. pneumoniae</i>	2002–2003	340		India	CTX-M-15	(27)
pJBE401	<i>qnrB2</i>	<i>K. pneumoniae</i>	2003	>150	L/M	Australia	IMP-4	(74)
<i>b</i>	<i>qnrB2</i>	<i>S. enterica</i>	2000			Senegal	SHV-12	(77)
pMG319	<i>qnrB4</i>	<i>E. cloacae</i>	1999–2004	200		USA	DHA-1	(238)
pPMDHA	<i>qnrB4</i>	<i>K. oxytoca</i>	2002	(Tra ⁻)		France	DHA-1	(134)
<i>b</i>	<i>qnrB4</i>	<i>E. cloacae</i>	<i>b</i>	119		China	DHA-1	(108)
pHND2	<i>qnrB6</i>	<i>K. pneumoniae</i>	2006			China	CTX-M-9G ^c	(75)
pARCF702	<i>qnrB10</i>	<i>C. freundii</i>	2005			Argentina		(239)
pR4525	<i>qnrB19</i>	<i>E. coli</i>	2002	40		Columbia	SHV-12 CTX-M-12	(78)
pLRM24	<i>qnrB19</i>	<i>K. pneumoniae</i>	2007	80		USA	KPC-3	(83, 117)
pPAB19-1	<i>qnrB19</i>	<i>S. enterica</i>	2006	2.7	ColEI	Argentina		(82)
pQAR2078	<i>qnrB19</i>	<i>E. coli</i>	2005	42.4	N	Germany		(81)
pAH0376	<i>qnrS1</i>	<i>S. flexneri</i>	2003	~50		Japan		(26)
pINF5	<i>qnrS1</i>	<i>S. enterica</i>	2004	58		Europe		(84)
TP _{qnrS-2a}	<i>qnrS1</i>	<i>S. enterica</i>	2004	44	N	UK		(240)
pK245	<i>qnrS1</i>	<i>K. pneumoniae</i>	2002	98		Taiwan	SHV-2	(87)
<i>b</i>	<i>qnrS1</i>	<i>S. enterica</i>	1999–2006	>250	H12	Netherlands	LAP-2	(120)
<i>b</i>	<i>qnrS2</i>	<i>d</i>	2004	8.5	Q	Germany		(241)
p37	<i>qnrS2</i>	<i>A. punctata</i>	2006	55	U	France		(242)
pHS10	<i>qnrC</i>	<i>P. mirabilis</i>	2006	~120		China		(28)

Plasmid	PMQR gene	Host	Year of isolation	Size (kb)	Inc group	Country	Linked <i>bla</i> genes ^a	Reference
p2007057	<i>qnrD1</i>	<i>S. enterica</i>	2006–2007	4.3		China		(29)
<i>e</i>	<i>qnrVC1</i>	<i>V. cholerae</i>	2002–2008			Bangladesh		(34)
<i>b</i>	<i>qnrVC4</i>	<i>A. punctata</i>	2008			China	PER-1	(30)
pBD146	<i>qnrVC5</i>	<i>V. fluvialis</i>	1998–2002	7.5		India		(243)
pHSH10-2	<i>aac(6')-Ib-cr</i>	<i>E. coli</i>	2000–2001			China		(10)
pC15-1a	<i>aac(6')-Ib-cr</i>	<i>E. coli</i>	2000–2002	92	FII	Canada	CTX-M-15	(244)
pHPA	<i>qepA1</i>	<i>E. coli</i>	2002		FII	Japan	CTX-M-12	(11)
pPI206	<i>qepA1</i>	<i>E. coli</i>	2000–2005	168	FI	Belgium		(197)
pQep	<i>qepA2</i>	<i>E. coli</i>	2007	90	FI	France		(199)
pOLA52	<i>oqxAB</i>	<i>E. coli</i>	<i>b</i>	52	X1	Denmark		(200, 202)
pHXY0908	<i>oqxAB</i>	<i>S. typhimurium</i>	2009		HI2	China		(207)

^a Only unusual *bla* genes are shown.

^b Not specified.

^c CTX-M-9 group. See reference for details.

^d Unidentified bacteria in activated sludge

^e Transmitted as part of an integrating conjugative element

Table 2

Distribution of PMQR genes

PMQR gene	Source Country	Organism	Mobilizing element	Reference
<i>qnrA1</i>	Algeria, Australia, Belgium, Brazil, Central African Republic, China, Denmark, Egypt, France, Germany, Hungary, India, Israel, Ivory Coast, Japan, Kenya, Mexico, Morocco, Netherlands, Nigeria, Portugal, Romania, Saudi Arabia, Singapore, South Korea, Spain, Sweden, Taiwan, Thailand, Turkey, UK, Uruguay, USA, Vietnam	<i>A. baumannii</i> , <i>C. freundii</i> , <i>E. aerogenes</i> , <i>E. cloacae</i> , <i>E. sakazakii</i> , <i>E. coli</i> , <i>H. parasuis</i> , <i>K. oxytoca</i> , <i>K. pneumoniae</i> , <i>P. mirabilis</i> , <i>P. aeruginosa</i> , <i>P. oryziatransiens</i> , <i>P. putida</i> , <i>S. enterica</i> , <i>S. marcescens</i> , <i>S. sonnei</i> , <i>S. maltophilia</i> , <i>V. fluvialis</i>	ISCR1	(9, 22–24, 67, 98, 104–108, 113, 115, 119, 122–125, 127–130, 138–141, 143, 144, 147, 155, 226, 227, 238, 245–290)
<i>qnrA3</i>	China, France, Hong Kong	<i>K. pneumoniae</i> , <i>K. ascorbata</i> , <i>Kluyvera</i> spp., <i>S. enterica</i> , <i>S. algae</i>	ISCR1, IS26	(25, 153, 156, 236, 291)
<i>qnrA6</i>	France, Tunisia	<i>C. freundii</i> , <i>K. pneumoniae</i> , <i>P. mirabilis</i> , <i>P. stuartii</i>	ISCR1	(237, 292, 293)
<i>qnrB1</i>	Algeria, Argentina, Brazil, China, Czech Republic, Denmark, France, Egypt, India, Italy, Ivory Coast, Malaysia, Mexico, Morocco, Netherlands, Nigeria, Norway, Saudi Arabia, Singapore, Scotland, Singapore, South Korea, Spain, Sweden, Thailand, Tunisia, Turkey, UK	<i>C. freundii</i> , <i>C. koseri</i> , <i>E. cloacae</i> , <i>E. gergoviae</i> , <i>E. coli</i> , <i>K. ornithinolytica</i> , <i>K. pneumoniae</i> , <i>S. enterica</i> , <i>S. marcescens</i>	Orf1005, IS26	(27, 67, 72, 73, 98, 102, 105, 109, 110, 113, 162, 186, 251, 259, 263, 268, 269, 271, 279, 281, 284, 285, 287, 288, 292, 294–302)
<i>qnrB2</i>	Argentina, Australia, Bolivia, Brazil, Czech Republic, China, France, Germany, Hungary, Ireland, Israel, Kuwait, Mexico, Morocco, Netherlands, Portugal, Peru, Scotland, Senegal, South Korea, Spain, Sweden, Switzerland, Taiwan, Tunisia, UK, USA	<i>C. freundii</i> , <i>E. cloacae</i> , <i>E. coli</i> , <i>K. oxytoca</i> , <i>K. pneumoniae</i> , <i>S. enterica</i> , <i>S. typhi</i>	ISCR1	(27, 61, 67, 73, 77, 96, 98, 108, 115, 116, 120, 125, 132, 139, 145, 155, 157, 161, 227, 238, 251, 257, 259, 265, 268, 271, 273, 274, 276, 279, 281, 290, 292, 299, 301, 303–316)
<i>qnrB3</i>	USA	<i>E. coli</i>		(238)
<i>qnrB4</i>	Algeria, Australia, China, France, Germany, Ivory Coast, Japan, Morocco, Netherlands, Saudi Arabia, Singapore, South Korea, Spain, Sweden, Switzerland, Taiwan, Thailand, UK, USA	<i>C. freundii</i> , <i>C. koseri</i> , <i>E. aerogenes</i> , <i>E. cloacae</i> , <i>E. gergoviae</i> , <i>E. coli</i> , <i>K. ornithinolytica</i> , <i>K. oxytoca</i> , <i>K. pneumoniae</i> , <i>S. enterica</i> , <i>S. marcescens</i> , <i>Shigella</i> sp.	ISCR1	(61, 67, 72, 75, 96, 98, 104, 105, 108, 109, 113, 121, 127, 132, 133, 145, 155, 186, 192, 196, 220, 238, 251, 256, 257, 266, 268, 271, 276, 280, 281, 285, 287, 290, 299, 309, 310, 316–319)
<i>qnrB5</i>	Denmark, France, Mexico, South Korea, UK, USA	<i>E. coli</i> , <i>K. pneumoniae</i> , <i>S. enterica</i>		(96, 97, 145, 259, 265, 279, 288, 295, 299, 303, 305, 320)
<i>qnrB6</i>	China, France, Germany, Japan, Malaysia, Mexico, Netherlands, Poland, Singapore, South Korea, Spain, Sweden, Thailand, USA	<i>C. freundii</i> , <i>E. aerogenes</i> , <i>E. cloacae</i> , <i>E. coli</i> , <i>H. parasuis</i> , <i>K. oxytoca</i> , <i>K. pneumoniae</i> , <i>S. enterica</i> , <i>S. fonticola</i> , <i>S. marcescens</i> , <i>S. flexneri</i>	ISCR1	(67, 72, 75, 98, 100, 104, 108, 127, 139, 141, 148, 153, 155, 167, 169, 192, 196, 204, 227, 251, 257, 266–268, 273, 279, 285, 287, 288, 290, 302, 309, 321, 322)
<i>qnrB7</i>	Kuwait, Netherlands, Norway, South Korea	<i>C. freundii</i> , <i>E. cloacae</i> , <i>K. pneumoniae</i> , <i>S. enterica</i>		(98, 155, 296, 313)
<i>qnrB8</i>	Brazil, China, France, Kuwait, South Korea, UK	<i>C. freundii</i> , <i>E. aerogenes</i>		(98, 104, 269, 299, 306, 313)
<i>qnrB9</i>	China, South Korea	<i>C. freundii</i>		(98, 290, 323)

PMQR gene	Source Country	Organism	Mobilizing element	Reference
<i>qnrB10</i>	Argentina, Bolivia, China, Malaysia, Nigeria, Peru, South Korea	<i>C. braakii</i> , <i>C. freundii</i> , <i>E. amnigenus</i> , <i>E. cloacae</i> , <i>E. coli</i> , <i>K. pneumoniae</i> , <i>S. Choleraesuis</i> , <i>S. marcescens</i>	ISCR1	(73, 127, 145, 153, 281, 302, 314, 324)
<i>qnrB12</i>	Netherlands, South Korea	<i>C. freundii</i> , <i>S. enterica</i>		(98, 155)
<i>qnrB16</i>	South Korea	<i>K. pneumoniae</i>		(281)
<i>qnrB19</i>	Argentina, Bolivia, Brazil, Columbia, Czech Republic, Finland, Denmark, Germany, Italy, Mexico, Netherlands, Nigeria, Peru, Poland, South Korea, UK, USA, Venezuela	<i>E. aerogenes</i> , <i>E. fergusonii</i> , <i>E. coli</i> , <i>K. oxytoca</i> , <i>K. pneumoniae</i> , <i>K. ascorbata</i> , <i>S. enterica</i> , <i>S. sonnei</i>	ISEcp1, IS26	(73, 78–83, 86, 120, 148, 155, 158, 166, 169, 288, 289, 301, 314, 315, 322, 324–329)
<i>qnrB20</i>	Mexico, Singapore	<i>K. pneumoniae</i>	Orf1005, IS26	(72, 288)
<i>qnrB22</i>	South Korea	<i>C. werkmanii</i> ^a		(330)
<i>qnrB23</i>	South Korea	<i>C. freundii</i> ^a		(330)
<i>qnrB26</i>	China	<i>P. vulgaris</i>		(290)
<i>qnrB31</i>	China	<i>K. pneumoniae</i>		(331)
<i>qnrB32</i>	China	<i>K. pneumoniae</i>		(331, 332)
<i>qnrS1</i>	Algeria, Argentina, Belgium, Bolivia, Brazil, Canada, China, Czech Republic, Denmark, Egypt, France, Germany, Greece, Hungary, Israel, Italy, Ivory Coast, Japan, Malaysia, Mexico, Morocco, Netherlands, Nigeria, Norway, Poland, Peru, Romania, Serbia, South Africa, South Korea, Slovakia, Spain, Sweden, Switzerland, Taiwan, Thailand, Tunisia, Turkey, UK, USA, Vietnam	<i>C. freundii</i> , <i>C. koseri</i> , <i>E. aerogenes</i> , <i>E. cloacae</i> , <i>E. coli</i> , <i>K. oxytoca</i> , <i>K. pneumoniae</i> , <i>M. morganii</i> , <i>P. mirabilis</i> , <i>S. enterica</i> , <i>S. typhi</i> , <i>S. boydii</i> , <i>S. flexneri</i> , <i>S. dysenteriae</i>		(26, 67, 73, 75, 84, 86, 98, 104, 105, 107–109, 115, 119–121, 125, 127, 131, 132, 143, 145, 148, 150, 151, 153–156, 158, 166, 167, 169, 172, 184, 186, 192, 196, 204, 220, 227, 240, 251, 255–257, 259, 261, 262, 264, 265, 268–273, 276, 279, 281, 284, 285, 287, 290, 292, 294–296, 299, 301–303, 305, 307, 308, 314–316, 318–322, 324, 333–341)
<i>qnrS2</i>	China, Czech Republic, Denmark, France, Germany, India, South Korea, Spain, Switzerland, USA	<i>A. allosaccharophila</i> , <i>A. caviae</i> , <i>A. hydrophila</i> , <i>A. media</i> , <i>A. punctata</i> , <i>A. veronii</i> , <i>E. cloacae</i> , <i>Pseudoalteromonas</i> spp., <i>Pseudomonas</i> spp., <i>S. enterica</i> , <i>S. marcescens</i> , <i>Shigella</i> sp.		(90, 100, 127, 148, 156, 167, 241, 242, 266, 272, 295, 299, 303, 319, 323, 342–345)
<i>qnrS3</i>	China	<i>E. coli</i>		(162)
<i>qnrS4</i>	Denmark	<i>S. enterica</i>		(295)
<i>qnrS5</i>	South Korea	<i>Aeromonas</i> spp.		(345)
<i>qnrC</i>	China	<i>P. mirabilis</i>	ISPmi1	(28)
<i>qnrD1</i>	China, Czech Republic, France, India, Italy, Netherlands, Nigeria, Poland, Spain	<i>E. coli</i> , <i>C. freundii</i> , <i>K. pneumoniae</i> , <i>M. morganii</i> , <i>P. mirabilis</i> , <i>P. vulgaris</i> , <i>P. rettgeri</i> , <i>P. aeruginosa</i> , <i>S. enterica</i>		(29, 62, 63, 91, 138, 148, 155, 204, 323, 340, 346, 347)
<i>qnrVC1</i>	Bangladesh, Brazil, India, Tunisia	<i>P. aeruginosa</i> , <i>V. cholerae</i>		(33, 34, 92, 94, 95)
<i>qnrVC3</i>	India	<i>V. cholerae</i>		(95)
<i>qnrVC4</i>	China, Haiti, Portugal	<i>A. hydrophila</i> , <i>A. punctata</i> , <i>Aeromonas</i> sp., <i>Pseudomonas</i> sp., <i>V. cholera</i>	ISCR1	(30, 33)
<i>qnrVC5</i>	China, Haiti, India	<i>V. parahaemolyticus</i> , <i>V. cholera</i> , <i>V. fluvialis</i>		(31–33)

PMQR gene	Source Country	Organism	Mobilizing element	Reference
<i>qnrVC6</i>	China	<i>A. baumannii</i>		(93)
<i>aac(6)-Ib-cr</i>	Algeria, Argentina, Australia, Bolivia, Brazil, Bulgaria, Canada, China, Croatia, Czech Republic, Denmark, Egypt, France, Germany, Hungary, India, Israel, Italy, Japan, Kenya, Mexico, Netherlands, Nigeria, Norway, Portugal, Peru, Saudi Arabia, Serbia, Singapore, Slovenia, South Korea, Spain, Switzerland, Taiwan, Thailand, UK, Uruguay, USA, Vietnam	<i>C. braakii</i> , <i>C. freundii</i> , <i>C. koseri</i> , <i>E. aerogenes</i> , <i>E. cloacae</i> , <i>E. coli</i> , <i>H. parasuis</i> , <i>K. pneumoniae</i> , <i>Kluyvera</i> spp., <i>L. hongkongensis</i> , <i>M. morgani</i> , <i>P. aeruginosa</i> , <i>P. luteola</i> , <i>P. oryzae</i> , <i>P. mirabilis</i> , <i>Raoultella</i> spp., <i>S. enterica</i> , <i>S. marcescens</i> , <i>S. odorifera</i> , <i>S. flexneri</i> , <i>S. sonnei</i> , <i>Shigella</i> sp., <i>V. fluvialis</i>	IS26	(10, 14, 67, 73, 75, 100–102, 104, 110, 113, 116, 125, 127, 133, 138, 141, 145, 148, 150, 153, 155–157, 161, 167, 169, 172, 182, 184–187, 189, 190, 192, 194–196, 220, 226, 256, 261, 263, 266, 267, 270, 272, 275, 278, 284, 287, 288, 290, 296, 297, 301, 315, 316, 318, 319, 322, 336, 348–362)
<i>oqxAB</i>	Argentina, Czech Republic, China, Denmark, Hong Kong, Italy, Japan, Poland, Serbia, South Korea, Spain	<i>E. coli</i> , <i>K. oxytoca</i> , <i>K. pneumoniae</i> , <i>S. enterica</i>	IS26	(73, 101, 148, 156, 168, 202–206, 208, 339, 360, 363, 364)
<i>qepA1</i>	Belgium, Bolivia, Canada, China, Egypt, India, Japan, Mexico, Nigeria, South Korea, Spain, UK, USA, Vietnam	<i>E. cloacae</i> , <i>E. coli</i> , <i>K. pneumoniae</i> , <i>M. morgani</i> , <i>P. oryzae</i> , <i>S. enterica</i> , <i>S. flexneri</i>	IS26, ISCR3C	(11, 12, 75, 114, 138, 145, 150, 153, 156, 167, 187, 192, 196–198, 220, 261, 270, 272, 284, 285, 290, 321, 324, 361, 365–367)
<i>qepA2</i>	France	<i>E. coli</i>	ISCR3C	(199)

^a *Citrobacter* spp. contain both plasmid-mediated and chromosomal *qnr* genes. Genes for *qnrB22* and *qnrB23* were transferred by conjugation to *E. coli* and hence proved to be plasmid determined.

Table 3

Effect of different quinolone resistance mechanisms on quinolone susceptibility of *E. coli*.

<i>E. coli</i> Strain	MIC µg/ml		
	Ciprofloxacin	Levofloxacin	Nalidixic acid
J53	0.008	0.015	4
J53 <i>gyrA</i> (S83L)	0.25	0.5	256
J53 pMG252 (<i>qnrA1</i>)	0.25	0.5	16
J53 pMG298 (<i>qnrB1</i>)	0.25	0.5	16
J53 pMG306 (<i>qnrS1</i>)	0.25	0.38	16
J53 pMG320 (<i>aac(6')-Ib-cr</i>)	0.06	0.015	4
J53 pAT851 (<i>qepA</i>)	0.064	0.032	4
CLSI susceptibility breakpoint	1.0	2.0	16