CASE REPORT

Percutaneous breast implant herniation: a rare complication of miliary TB

Adam P Dale,¹ Martin J Dedicoat,² Tausif Saleem,³ Ed Moran²

¹Department of Medical Microbiology, Basingstoke and North Hampshire Hospitals, Basingstoke, UK

²Department of Infectious Diseases and Tropical Medicine, Birmingham Heartlands Hospital, Heart of England NHS Hospital Trust, Birmingham, UK

³Department of Histopathology, Birmingham Heartlands Hospital, Heart of England NHS Hospital Trust, Birmingham, UK

Correspondence to

Dr Adam P Dale,
adamdale@doctors.org.uk

Accepted 18 December 2014

SUMMARY

We describe the case of a 46-year-old female patient treated for disseminated tuberculosis (TB) infection involving the lungs, urinary tract and skin. Following initiation of antituberculous therapy, the patient's right breast implant eroded through the overlying skin and was seen to be herniating through the resulting defect. The breast implant was removed under local anaesthetic and histological analysis of the resected tissue demonstrated granuloma formation consistent with periprosthetic TB. Wound healing following implant removal was poor and future breast augmentation surgery was only considered following completion of 12 months anti-TB treatment. This case constitutes the first report in the literature of percutaneous breast implant herniation resulting from periprosthetic infection with TB. A high index of suspicion is required to ensure early detection and timely management of TB and, in cases where periprosthetic pus aspirate is sterile, mycobacterial infection must be actively excluded.

BACKGROUND

Periprosthetic breast implant infection presents as either an early or late postoperative complication. In individuals solely undergoing breast augmentation, the incidence of postoperative breast implant-associated infection ranges between 1.1% and 2.5%.¹ However, in patients undergoing breast reconstruction the incidence of infection can be substantially higher, ranging from 1.5% to 12% on analysis of studies spanning the past decade.¹⁻³

Direct inoculation of the breast implant with colonising skin or breast duct flora in the perioperative period is the underlying mechanism resulting in the majority of breast implant-associated infections.³ Consequently, commensal skin organisms, including *Staphylococcus aureus*, streptococci and coagulase negative staphylococci, are among the most frequently identified causative pathogens.¹ Less frequently, breast implant infections may result from haematogenous bacterial spread from a distant infective site and, in these cases, a wide variety of pathogenic Gram-positive and Gram-negative organisms may be isolated.³⁻⁶

Reports of breast implant infections secondary to environmental non-tuberculous mycobacteria (NTM) are increasingly common. *Mycobacterium fortuitum* is the most frequently reported NTM associated with breast implant infections with reports also describing infection with *Mycobacterium avium-intracellulare*⁷ and *Mycobacterium jaccuzii*.⁸ *Mycobacterium tuberculosis* (TB), unlike NTM, is not an environmental pathogen, and should therefore not be associated with intraoperative contamination.

Instead, TB-associated breast implant infections occur in individuals with pre-existing TB infection, resulting from either contiguous or haematogenous mycobacterial spread.

To the best of our knowledge, there is only one published case report to date that outlines a TB-associated breast implant infection.⁹ In the current report, we describe the first case of percutaneous breast implant herniation secondary to periprosthetic TB infection.

CASE PRESENTATION

In April 2013 a 46-year-old female patient of Indian origin was referred to the infectious diseases outpatient clinic by her dermatologist, after a lesion on her right hand was confirmed to be a tuberculous granuloma (figure 1). The granulomatous hand lesion, present since July 2012, was initially thought to be a fish tank granuloma, however, TB was isolated following culture of the skin biopsy in the laboratory. On further questioning, the patient reported a 6-week history of progressive dyspnoea, a cough productive of green sputum and a two stone loss in weight. It was also noted that a family member who resided in the same household had received treatment for pulmonary TB 5 years previously. Clinical assessment revealed the patient to be septic, therefore hospital admission was required for resuscitation and start of antituberculous therapy.

INVESTIGATIONS

Microscopic examination of repeated sputum smears, obtained during the inpatient stay, were negative for acid alcohol fast bacilli, however, following a period of incubation, fully sensitive TB was isolated on culture of early morning urine and sputum specimens. Chest radiography demonstrated changes consistent with pulmonary TB (figure 2) and, in view of disseminated infection, a diagnosis of miliary TB was made.

Figure 1 Right hand tuberculous granuloma.

CrossMark

To cite: Dale AP, Dedicoat MJ, Saleem T, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-207546

Figure 2 Chest radiograph demonstrating changes consistent with miliary *Mycobacterium tuberculosis* infection.

TREATMENT

The patient's clinical condition stabilised following initiation of standard four drug antituberculous therapy (rifampicin, isoniazid, pyrazinamide and ethambutol) and subsequently, 6 days following initial admission, the patient was discharged.

OUTCOME AND FOLLOW-UP

On follow-up assessment in the outpatients' clinic in May 2013 the patient was making good clinical progress and had noticed a significant increase in her exercise tolerance, along with shrinkage of the cutaneous tuberculoma. However, she reported that her right silicon breast implant, which had initially been inserted as part of an augmentation procedure for aesthetic reasons, had started to protrude through the overlying skin. Examination demonstrated no overt evidence of soft tissue infection, however, the breast implant had clearly eroded through the overlying skin and had begun to herniate through the skin defect.

The patient was referred for review by the plastic and reconstructive surgery team. Following consideration of the risks of general anaesthesia in a patient with significantly impaired

pulmonary function resulting from miliary TB, it was decided that the herniating implant was to be removed under local anaesthesia. Histological analysis of intraoperative soft tissue samples taken from the site of skin erosion was performed and demonstrated granulomatous tissue characteristic of TB infection (figure 3). Ziehl-Neelsen stain for acid-fast bacilli on the biopsy was examined but was negative, however, given known culture positivity, the histological findings were still consistent with a diagnosis of periprosthetic TB infection.

Wound healing was slow following breast implant removal and, on review in August 2013, the wound had still failed to heal fully. At this time, it was also noted that the patient's left breast implant was starting to erode through the overlying skin. The opportunity for the patient to undergo future breast augmentation surgery was discussed, however, the consensus was only to pursue this once the patient had completed her 12-month treatment course.

DISCUSSION

This case details the first report of percutaneous breast implant herniation resulting from periprosthetic TB infection. While the majority of non-mycobacterial and NTM-associated breast implant infections arise from intraoperative contamination, in this case it is likely that the infection resulted from haematogenous mycobacterial seeding. Although reports of breast implant-associated TB infection are rare, case reports describing tuberculous infections of other prosthetic devices and implants, for example, hip or knee replacements^{10–12} and heart valves,^{13 14} are increasingly common. As the prevalence of patients with breast implants increases, cases of periprosthetic breast implant TB may become more frequent, particularly among patient groups with a high burden of TB.

In the case described, the diagnosis of miliary TB was established prior to the development of the breast implant-associated infection. As a consequence, TB was considered as a potential cause from the outset. However, given the ability of TB to cause primary breast infection,^{9 15 16} it is possible that primary breast implant-associated TB infection could occur. In these cases a high index of suspicion would be required to establish the underlying diagnosis.

Practice relating to the management of periprosthetic breast implant infections varies substantially. There are no published studies comparing implant removal and antimicrobial therapy

Figure 3 Periprosthetic biopsy: Skin undermined by a palisade of epithelioid histiocytes with sporadic multinucleated giant cells accompanied by necrosis indicating granulomatous inflammation.

with antimicrobial therapy alone. In their recent review, Washer and Gutowski¹ outline an investigatory and management pathway for patients with breast implant infections. They suggest that in the majority of cases infected breast implants should be removed. However, in cases of superficial cellulitis responding to antimicrobial therapy, or in patients with culture negative non-purulent periprosthetic fluid aspirate (likely to be a seroma), a less aggressive approach may be taken.

Breast implant removal can prove distressing for the patient and has significant psychosocial implications.¹⁷ A number of reports detail successful salvage attempts utilising antimicrobial therapy in conjunction with irrigation methods and breast implant exchange.^{18 19} However, while these methods may be appropriate for individuals with mild cellulosic bacterial infection, evidence from one study suggests that implant salvage is likely to fail in the presence of atypical pathogens including fungi and NTM.²⁰

When suspected, TB should be actively excluded using direct detection and culture methods, particularly in cases where periprosthetic pus aspiration is sterile, or where the history reveals constitutional features consistent with a diagnosis of TB. Tissue specimens available following breast implant removal should be subject to histological analysis to exclude granulomatous findings consistent with a diagnosis of TB. The appropriate surgical strategy is likely to vary from case to case. A non-suppurative granuloma may respond to antituberculous therapy alone, whereas skin erosion and persistent discharge while on therapy, as in the case described here, clearly indicates the need for implant removal.

Before considering repeat breast augmentation surgery, the operator must be fully confident that the infective pathogen has been eradicated. Consequently, in cases of TB infection, the patient may be without prosthetic breast implants for several months. This stands in contrast to the shorter time period a

patient with a non-mycobacterial breast implant infection would experience.

This case highlights the importance of considering TB as a cause of periprosthetic or skin infection in patients known to have disseminated TB. If there is no history of TB, a high index of suspicion is required to ensure exclusion of this pathogen, or to allow initiation of appropriate treatment if TB is isolated.

Contributors APD was responsible for literature review and writing of the introduction, case report and discussion. MJD and EM were responsible for providing clinical information relating to case, providing expert opinion and reviewing, altering and finalising the manuscript prior to submission. TS provided expert histological analysis and histological slide images as well as reviewing and altering the manuscript prior to submission.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Washer LL, Gutowski K. Breast implant infections. *Infect Dis Clin North Am* 2012;26:111–25.
- 2 Nahabedian MY, Tsangaris T, Momen B, et al. Infectious complications following breast reconstruction with expanders and implants. *Plast Reconstr Surg* 2003;112:467–76.
- 3 Pittet B, Montandon D, Pittet D. Infection in breast implants. *Lancet Infect Dis* 2005;5:94–106.
- 4 Bernardi C, Saccomanno F. Late Klebsiella pneumoniae infection following breast augmentation: case report. *Aesthetic Plast Surg* 1998;22:222–4.
- 5 Memish ZA, Alazzawi M, Bannatyne R. Unusual complication of breast implants: Brucella infection. *Infection* 2001;29:291–2.
- 6 Hunter JG, Padilla M, Cooper-Vastola S. Late Clostridium perfringens breast implant infection after dental treatment. *Ann Plast Surg* 1996;36:309–12.
- 7 Wirth GA, Brenner KA, Sundine MJ. Delayed silicone breast implant infection with Mycobacterium avium-intracellulare. *Aesthet Surg J* 2007;27:167–71.
- 8 Rahav G, Pitlik S, Amitai Z, et al. An outbreak of Mycobacterium jaccuzii infection following insertion of breast implants. *Clin Infect Dis* 2006;43:823–30.
- 9 Miles G, Walters T, Shee C. Periprosthetic tuberculous breast infection. *J R Soc Med* 2003;96:556–7.
- 10 Shanbhag V, Kotwal R, Gaitonde A, et al. Total hip replacement infected with Mycobacterium tuberculosis. A case report with review of literature. *Acta Orthop Belg* 2007;73:268–74.
- 11 Marmor M, Parnes N, Dekel S. Tuberculosis infection complicating total knee arthroplasty: report of 3 cases and review of the literature. *J Arthroplasty* 2004;19:397–400.
- 12 Cansü E, Erdogan F, Ulusam AO. Incision infection with Mycobacterium tuberculosis after total hip arthroplasty without any primary tuberculosis focus. *J Arthroplasty* 2011;26:505.e1–3.
- 13 Liu A, Nicol E, Hu Y, et al. Tuberculous endocarditis. *Int J Cardiol* 2012;167:640–5.
- 14 Sultan FAT, Fatimi S, Jamil B, et al. Tuberculous endocarditis: valvular and right atrial involvement. *Eur J Echocardiogr* 2010;11:E13.
- 15 Sen M, Gorpelioglu C, Bozer M. Isolated primary breast tuberculosis: report of three cases and review of the literature. *Clinics (São Paulo)* 2009;64:607–10.
- 16 Green M, Millar E, Merai H, et al. Mammary tuberculosis in the young: a case report and literature review. *Breast Dis* 2012;34:39–42.
- 17 Roberts C, Wells KE, Daniels S. Outcome study of the psychological changes after silicone breast implant removal. *Plast Reconstr Surg* 1997;100:595–9.
- 18 Laveaux C, Pauchot J, Lory J, et al. [Acute periprosthetic infection after aesthetic breast augmentation. Report of three cases of implant “salvage”. Proposal of a standardized protocol of care]. *Ann Chir Plast Esthet* 2009;54:358–64.
- 19 Chun JK, Schulman MR. The infected breast prosthesis after mastectomy reconstruction: successful salvage of nine implants in eight consecutive patients. *Plast Reconstr Surg* 2007;120:581–9.
- 20 Spear SL, Seruya M. Management of the infected or exposed breast prosthesis: a single surgeon's 15-year experience with 69 patients. *Plast Reconstr Surg* 2010;125:1074–84.

Learning points

- Periprosthetic breast implant infections most commonly result from intraoperative bacterial contamination, however, less commonly, infections occur secondary to haematogenous bacterial spread.
- Tuberculosis (TB)-associated periprosthetic breast implant infections are extremely rare therefore a high index of suspicion is required to ensure early detection and timely management in these cases.
- When suspected, TB should be actively excluded using direct detection and culture methods, particularly in cases where periprosthetic pus aspiration is sterile, or where the history reveals constitutional features consistent with a diagnosis of TB.
- In cases where periprosthetic TB infection is confirmed, the surgical strategy is likely to vary depending on severity of symptoms. However, in cases where surgical excision is indicated, the operator must be confident that TB has been completely eradicated prior to consideration of reimplantation.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow