Published in final edited form as:

Clin Psychol Sci. 2015 January 1; 3(1): 126–139. doi:10.1177/2167702614560748.

Cognitive Bias Modification for Interpretation in Major Depression: Effects on Memory and Stress Reactivity

Jutta Joormann,

Northwestern University

Christian E. Waugh, and

Wake Forest University

Ian H. Gotlib

Stanford University

Abstract

Interpreting ambiguous stimuli in a negative manner is a core bias associated with depression. Investigators have used cognitive bias modification for interpretation (CBM-I) to demonstrate that it is possible to experimentally induce and modify these biases. This study extends previous research by examining whether CBM-I affects not only interpretation, but also memory and physiological stress response in individuals diagnosed with Major Depressive Disorder (MDD). We found that CBM-I was effective in inducing an interpretive bias. Participants also exhibited memory biases that corresponded to their training condition and demonstrated differential physiological responding in a stress task. These results suggest that interpretation biases in depression can be modified, and that this training can lead to corresponding changes in memory and to decreases in stress reactivity. Findings from this study highlight the importance of examining the relations among different cognitive biases in MDD and the possibility of modifying cognitive biases.

Keywords

Cognitive bias modification; training; interpretation; memory; depression

Cognition plays a critical role in human emotion. According to cognitive theories of emotion, appraisals or interpretations of a situation determine whether an emotion is experienced and which emotion that will be (Ellsworth & Scherer, 2003; Siemer, Mauss, & Gross, 2007). Importantly, therefore, cognition is also a primary mechanism through which emotions can be regulated. For example, cognitive reappraisal, or the reinterpretation of the emotion-eliciting situation, has been found to be a particularly effective emotion regulation strategy (Gross & John, 2002). Researchers have also demonstrated that memory functioning is critical for the regulation of negative affect (Joormann, Siemer, & Gotlib, 2007). Cognitive biases in interpretation or memory, specifically a tendency to interpret ambiguous material in a negative matter or to preferentially recall negative events, may

therefore impair emotion regulation and increase vulnerability to emotional disorders (Joormann, Yoon, & Siemer, 2010).

Indeed, cognitive theories of emotional disorders posit that depressed and anxious individuals, as well as people who are at risk for developing these disorders, exhibit cognitive biases in various aspects of information processing, including interpretation and memory (Mathews & MacLeod, 2005). These theories further propose that cognitive biases are not simply epiphenomenal in these disorders, but instead play an important role in increasing the risk for the onset, maintenance, and recurrence. Indeed, depressed individuals have been found to demonstrate increased elaboration of negative material and a tendency to interpret ambiguous material in a mood-congruent manner (e.g., Cowden Hindash & Amir, 2012). In addition, biased memory for negative, relative to positive, information is perhaps the most robust finding concerning cognitive biases associated with major depression (Matt, Vazquez, & Campbell, 1992).

Given the importance of cognition for everyday emotion regulation and risk for emotional disorders, modifying cognitive biases may alter individuals' reactivity to emotion-eliciting events and improve their ability to regulate negative affect. Furthermore, research examining how these biases can be changed may yield important insights into processes that underlie the onset and maintenance of disorders, and thus holds promise for efforts aimed at prevention and treatment. Indeed, researchers have investigated the effects of bias modification on various outcome measures including symptoms of depression and anxiety, and responses to acute stressors. Some of these studies have focused on manipulating attention by guiding participants to attend to either threatening or emotionally neutral stimuli (see Hallion & Ruscio, 2010 for a recent review). Biases in interpretation and memory, however, may have particularly detrimental effects on individuals' ability to regulate negative mood states and may serve to increase risk for the onset and maintenance of debilitating emotional disorders.

In a seminal study, Grey and Mathews (2000) demonstrated that biased interpretation of emotionally ambiguous homographs can be induced, and that participants are often not aware of this modification. In a related study, Mathews and Mackintosh (2000) used ambiguous scenarios to train individuals to make either non-anxious or anxious interpretations of ambiguous situations. In this and in later studies, participants who were trained to interpret ambiguity in a non-anxious manner had an attenuated self-reported anxious reaction to a subsequent stressor (e.g., Wilson, MacLeod, Mathews, & Rutherford, 2006; see also Mackintosh, Mathews, Yiend, Ridgeway, & Cook, 2006) and showed improvements in a subsequently induced negative mood state (Holmes, Lang, & Shah, 2009). Importantly, beneficial effects of interpretive training on levels of anxiety have been shown to be stable after a 24-hour delay between training and test (Yiend, Mackintosh, & Mathews, 2005). Not surprisingly, therefore, researchers have begun to examine the usefulness of cognitive bias modification for interpretation (CBM-I) as an intervention strategy. In a recent study, Bowler et al. (2012) demonstrated in a sample of participants with high levels of social anxiety symptoms that CBM-I led to symptom reduction, lower trait anxiety, and depression, as well as to improved self-reported attentional control.

Despite these successes when using CBM-I in anxiety disorders and in participants with elevated levels of anxiety, research in depression has been mostly lacking. Williams et al. (Williams, Blackwell, Mackenzie, Holmes, & Andrews, 2013) found that one week of CBM-I reduced the severity of symptoms in a sample of clinically depressed individuals and that these changes were at least partially mediated by changes in interpretation biases. This study, however, did not include a control group, which makes it difficult to attribute the symptom change to the intervention. In a study of depressed adolescents and young adults (14–21 years of age), Micco, Henin, and Hirshfeld-Becker (2013) included a control group in a four-session training. The authors reported that both groups, the active training and the control group, exhibited a reduced interpretation bias at follow-up with no significant between-group difference. When the authors limited their analysis to participants with negative biases at baseline, however, the intervention group exhibited greater improvements in interpretation bias. Still no group differences were found in depression or anxiety symptom change. Only one study thus far has examined whether CBM-I in a depressed sample affects stress reactivity and recovery. Yiend et al. (2013) reported that a single session of CBM-I was effective in changing interpretive biases in depressed adults. These authors did not, however, find a reduction in self-reported stress responding. This lack of transfer of training to stress responding may be due to the use of a single session of training or to the exclusive reliance on self-reported distress. Thus, it is clear that additional studies are needed in depressed adults using control groups and CBM-I training protocols that encompass more than a single session. Moreover, no study to date has examined whether CBM-I reduces physiological stress reactivity in depression. The present study was designed in part to test this possibility.

Modifying interpretation biases may be a promising approach for the treatment of depression not only because it affects the interpretation of novel situations and responses to stressors, but because changes in interpretation may also affect memory. Hirsch, Clark, and Mathews (2006) and Everaert, Koster and Derakshan (2012) noted that cognitive biases do not operate in isolation, but rather, influence and interact with one another. If individuals tend to interpret situations in a negative manner, they may also be more likely to remember them in a way that is consistent with this initial interpretation. Thus, memory biases may be due in part to earlier biases in interpretation (Hertel, 2004; Hertel & Brozovich, 2010), although very few studies have investigated this possibility empirically. Among the few studies that have done so is one conducted by Hertel, Brozovich, Joormann, and Gotlib (2008), who examined the association between interpretation and memory in individuals diagnosed with social phobia. When asked to recall details from previously presented ambiguous scenarios, socially anxious tended to produce more distortions than did non-anxious participants that reflected their initial negative interpretations of the scenarios. This result suggests that memory biases can be produced by biased interpretations and that modifying the initial interpretation of events could prevent subsequent biases in memory. Indeed, in a previous study using CBM-I in a non-clinical sample, Tran, Hertel, and Joormann (2011) reported a relation between interpretation and memory biases. In this study, the authors induced positive and negative interpretation biases in a student sample using a single session of CBM training and found effects of this training on subsequent

recall. This association between interpretation and memory biases, however, has not been examined in clinically depressed participants.

In the present study we examined whether two sessions of CBM-I in participants diagnosed with Major Depressive Disorder (MDD) affect not only participants' interpretation of novel scenarios, but also their memory for these scenarios. We also examined the effects of CBM-I on physiological responses to an acute stressor. In addition, we included several control groups. We compared participants diagnosed with MDD to a group of healthy control participants. We expected training effects in both groups, with a stronger training effect in the MDD group. In addition, to examine whether we could replicate Tran et al.'s (2011) findings, all participants were randomly assigned to receive either positive or negative interpretation training. While we expected that the negative training would induce a negative interpretation bias in the CTL group, we did not expect any effect of this training in the MDD group given that MDD participants should exhibit a negative interpretation bias even before the training. Finally, to examine whether training affects interpretation biases in the MDD group, we recruited a separate group of MDD participants who did not undergo training but participated in the follow-up assessment; this procedure allowed us to compare the effects of a no-training condition to a positive-training and a negative-training condition within the MDD sample.

Participants diagnosed with current MDD and non-disordered controls were randomly assigned to a positive or a negative interpretation training condition using ambiguous social scenarios as stimuli. We assessed the transfer of this training to the interpretation of subsequently presented novel ambiguous scenarios. We predicted that training would transfer to these novel scenarios; this transfer is indexed by whether participants judge positive or negative interpretations of these scenarios to be more or less similar to the novel scenarios. Thus, compared with participants in the negative training condition, participants in the positive training condition were predicted to judge positive interpretations to be more similar to the ambiguous scenarios. We also predicted that the training would be effective in both the MDD and the CTL participants, but that the MDD participants would show greater improvement than will the CTL participants. Furthermore, to investigate the relation between interpretation and memory biases, we also asked participants to recall these novel scenarios, guided by the hypothesis that CBM-I will not only change participants' interpretations of novel scenarios, but will also induce a training-congruent memory bias. Thus, we predicted that CBM-I would lead to a training-congruent memory bias for the ambiguous scenarios, such that participants will be more likely to report memory distortions (or intrusions) that are congruent with their training condition. Again, we predicted these training effects across MDD and CTL participants, but expected the effects to be stronger in the MDD than in the CTL group. Participants were also exposed to acute stressor while we assessed psychophysiological arousal. We predicted that participants who completed positive bias training would show reduced physiological arousal to the stressor compared to participants who received negative training. We again expected this effect in both the MDD and CTL groups, with a stronger effect in the MDD group. Finally, to further examine the effects of negative and positive interpretation training within the MDD group, we compared MDD participants in the two training conditions to MDD participants in the no-training condition. We expected to find no differences in any of our measures when comparing MDD

participants in the negative versus the no-training condition; however, we expected that MDD participants in the positive-training condition would exhibit more positive interpretations, better recall of positive material, and lower stress reactivity than would MDD participants in the no-training condition.

Method

Participants

Participants were recruited from a variety of sources. Clinical participants were solicited from two outpatient psychiatry clinics in a university teaching hospital, as well as through advertisements posted in numerous locations within the local community (e.g., internet bulletin boards, university kiosks, supermarkets). Healthy controls were recruited from the community through advertisements posted in the same locations. Participants' responses to a telephone interview provided initial selection information. This phone screen established that participants were fluent in English and were between 18 and 60 years of age. Participants were excluded for severe head trauma, learning disabilities, as well as for current psychotic symptoms, bipolar disorder, and alcohol or substance abuse within the past six months. This telephone interview was also used to identify individuals who were likely to meet criteria for one of the two diagnostic groups; these individuals were invited to come to the laboratory for a more extensive interview. Participants were paid \$25 per hour for participation in this study.

Trained interviewers administered the Structured Clinical Interview for the DSM-IV (SCID; First, Spitzer, Gibbon, & Williams, 1995) to these individuals during their first session in the study. The SCID has demonstrated good reliability for the majority of the disorders covered in the interview (J. B. Williams et al., 1992). All interviewers had extensive training in the use of the SCID, as well as previous experience in administering structured clinical interviews with psychiatric patients prior to beginning the current study. Our team of interviewers achieved excellent interrater reliability. The κ coefficients were .93 for the MDD diagnosis and .92 for the "nonpsychiatric control" diagnosis (i.e., the absence of current or lifetime psychiatric diagnoses, according to the DSM-IV criteria).

Participants were included in the depressed group if they met the Diagnostic and Statistical Manual of Mental Disorders, 4th ed. (DSM-IV; American Psychiatric Association, 1994) criteria for MDD. The never-disordered CTL group consisted of individuals with no current diagnosis and no history of any Axis I disorder. Participants also completed the Beck Depression Inventory-II (see below). Participants were scheduled for a second session of "computer tasks," usually within two weeks after the interview.

Seventy-six individuals (48 MDD, 28 CTL) participated in this study. The MDD participants were randomly assigned to the positive- ($N = 16$), the negative- ($N = 15$) or the no- training condition ($N = 17$). The CTL participants was randomly assigned to either the positive- ($N = 14$) or the negative-training condition ($N = 14$).

Questionnaires

Beck Depression Inventory (BDI-II; Beck & Steer, 1993; Beck, Steer, & Garbin, 1988)—Severity of current depressive symptoms was assessed using this 21-item self-report measure. The BDI-II is a widely used depression scale with good psychometric properties (Beck et al, 1988), and correlations with clinician ratings of depression of .62 to .66 (Foa, Riggs, Dancu, & Rothbaum, 1993).

Mood Ratings—To assess current mood, participants rated multiple mood items before and after each training session by indicating the degree to which the items described how they felt right now, on a scale of 1 (*not at all*) to 11 (*extremely*). Responses were averaged across items to yield positive (*interested, excited, enthusiastic*) and negative (*distressed, upset, nervous*) affect scores.

Cognitive Bias Modification Training

In this training, participants are exposed to ambiguous scenarios and are asked to complete word fragments that solve the ambiguity in either a positive or negative direction. The rationale behind the training is that participants develop implicit expectations that ambiguous situations are more likely to end positively or negatively depending on the training condition. Participants in the positive- and negative-training conditions took part in two training sessions that were scheduled within one week. Participants in the no-training condition participated only in the test phase (see below). In each training session, participants were presented with 10 blocks of 13 scenarios each, in which they were asked to imagine themselves as the central character. For example, in a scenario titled “Meeting a Friend,” participants were asked to picture themselves waiting at an empty bar to meet an old friend. Instructions and scenarios were taken from training studies reported by Mathews and Mackintosh (2000). Each scenario consisted of a title and two to three sentences and ended with a word fragment for participants to complete. Within each of the ten training blocks, eight training scenarios, two probe scenarios, and three filler scenarios were presented. The filler scenarios were identical in both training conditions and thus helped to disguise the training condition. For the training scenarios, each word fragment could be completed to produce only one possible solution, which disambiguated the meaning of the scenario according to the assigned training condition. Thus, for the positive-training group, the completed fragment produced a positive outcome for the scenario, whereas for the negative-training group, the completed fragment produced a negative outcome. In the probe scenarios, the word fragment disambiguated the sentence in the same way (either positive or negative) for all participants, regardless of training group. In the filler scenarios, completion of the word fragments produced a neutral meaning. Examples can be found in previous training studies (e.g., Mathews & Mackintosh, 2000).

Prior to training, participants were told that for each trial, they would be shown a brief scenario that ends with an incomplete word fragment. They were instructed to press the spacebar as soon as they were able to solve the word fragment. Following the key press, the scenario disappeared from the screen and participants then typed the completed word on the next screen. In the final segment of the trial, participants were presented with a simple comprehension question and responded by pressing “Y” (yes) or “N” (no). This

comprehension question served to emphasize the valence of each scenario, as well as test participants' understanding of the text. Participants were given the opportunity to complete three practice trials prior to beginning the training to ensure their understanding of the task. The order of the training blocks and the order of the scenarios within the blocks were newly randomized for each participant.

Test Phase

In the test phase, participants were presented with 20 novel ambiguous scenarios, identical in structure to those they had viewed in the training phase. The main difference between the training and test scenarios was the fact that, although there was only one possible solution for the word fragment, the resolved word maintained the ambiguity of the preceding text and did not resolve the scenario in a positive or a negative way. The scenarios were presented in random order. Participants were shown each scenario on the computer screen again and were instructed to press any key as soon as they were able to solve the word fragment. After doing so, the scenario disappeared from the screen and they were asked to enter the completed word. Next, to maintain consistency, participants were presented with a comprehension question and were asked to indicate their answers with a "Y" or "N" response. Following their response, they were presented with the next test scenario. Unlike the comprehension questions during the training phase, these questions did not draw attention to the emotional implications of the scenario.

Interpretation test—Following a brief filler task, participants completed a computer task in which they were presented with the titles of the previously presented test scenarios. Each title was presented one at a time on the screen in random order. For each scenario title, participants were instructed to rate four sentences according to their similarity to the original scenario with the corresponding title (1: very different, 2: fairly different, 3: fairly similar, 4: very similar). In each case, two of these sentences were target sentences - a positive target that matched a positive interpretation of the original scenario (e.g., *You arrange to meet in a bar and your friend arrives late*) and a negative target that matched a negative interpretation (e.g., *You arrange to meet in a bar but your friend doesn't turn up*). The other two sentences were positive and negative foils -- sentences with a generally positive or negative meaning that were unrelated to the ambiguous concept in the original scenario. These foil statements were included to assess broader valence effects of the training. The positive foil sentence assigned a generally positive meaning to the scenario, but was not based on any information provided by the original text (e.g., *Your friend wants to meet again but you don't have time*); similarly, the negative foil sentence assigned a generally negative meaning but was not based on the original text (e.g., *Your friend tells you that she does not want to meet you*).

To assess interpretations of the test scenarios, each of the four sentences (per scenario) was presented one at a time under the scenario title. When each sentence appeared on the screen, participants were instructed to type the number corresponding to their rating of similarity to the meaning of the scenario. Once they entered their similarity rating, they were presented with the next sentence until they had provided ratings for all four sentences corresponding to that scenario title. The order of these sentences was newly randomized for each participant.

Recall task—Finally, participants were asked to recall details from the 20 test scenarios. The titles of the scenarios were presented in random order on the computer screen. When presented with each title, participants were instructed to recall aloud as many details as they could from the corresponding scenario. A key press initiated the next trial. Responses were audio-taped and later transcribed. Two independent raters were trained to code the transcriptions for whether each recalled scenario contained intrusions or new ideas (yes/no), and if so, the valence of each intrusion (positive, negative, neutral). Once adequate levels of inter-rater agreement in the number and valence of intrusions were achieved ($r = 0.85$) using independent training scenarios, two raters who were blind to the experimental conditions categorized intrusions in each recalled scenario as belonging to one of three groups: negative (e.g., *everyone stares and laughs at you at the bar*), positive (e.g., *your friend arrives right as you walk in the door and is happy to see you*), or neutral (e.g. *you have a seat and wait at the bar*).

Stress Task

The stress induction was based on a procedure developed by Waugh, Panage, Mendes, and Gotlib (2010). After completing the test phase, participants sat and rested for five minutes. Participants were then instructed that they might have to give a five-minute speech and would have two minutes to prepare the speech. They were told that their speech would be recorded and judged by evaluators on clarity, coherence, and persuasiveness. They were then told that at the end of the two-minute preparation period, the experimenter would flip a coin to determine whether they actually had to give a speech. The experimenter then told participants the speech topic was “Why are you a good friend?”- a topic used successfully in previous studies to induce anticipatory stress responses (Fredrickson, Mancuso, Branigan & Tugade, 2000; Waugh, et al., 2010) - and left them alone to prepare the speech for two minutes. The experimenter then flipped a double-tailed coin to ensure that the participants would not have to give the speech. No participants reported being suspicious about this fixed coin flip. After anticipating giving the speech, participants sat and rested for five minutes. This anticipatory stress procedure has been shown to induce increases in both heart rate and negative affect that are comparable to stress procedures that have the participants actually give the speech. An additional advantage of this anticipatory stress procedure that is less true of speech stressors is that anticipatory stress-induced negative affect and heart rate are positively correlated, making it an effective test of whether changes in negative interpretation biases can influence physiological responses to stress (Waugh et al., 2010).

Self-reported distress

We used average scores on three items to assess self-reported distress during the stress task (*distressed, upset, nervous*). All three items were given three times during the stress task: at baseline before the onset of the stressor (Pre-stressor), after the anticipation phase of the stressor when participants were finished preparing the speech (Anticipation), and after a 5-min recovery period when they had been told that they did not have to give the speech (Recovery).

Heart rate

Immediately upon entering the lab for either their second training session plus test session (training conditions) or just their test session (no training condition), disposable snap electrodes were attached to the participants using a lead II configuration. During the stress task, cardiovascular responses were recorded at 1kHz sampling rate with the electrocardiogram (ECG) module of an integrated physiological acquisition system (Biopac MP150; Biopac Systems, Goleta, CA), and then high- (.5 Hz) and low-pass filtered (40 Hz). We inspected the ECG data for artifacts and missing R-peaks (based on improbable inter-beat intervals) using a physiological analysis software package (ANSLAB; Wilhelm & Peyk, 2005). If a single R-peak was missing, an R-peak was inserted at a time-point halfway in between the two neighboring R-peaks. If more than one R-peak was missing in a row, those data were not scored. Heart rate (BPM) was then calculated from the scored inter-beat intervals for each time period (baseline – 5 mins, anticipation – 2 mins, recovery – 5 mins).

Overall Procedure

All participants took part in the phone interview and the SCID, which took approximately two hours. Within one week after the SCID interview, participants in the training conditions took part in the first training session, which was followed by the second training session within three days. After the second training session, participants completed a 15-min word scramble filler task which was followed by the test phase. During the test phase, participants first read the set of 20 emotionally ambiguous scenarios and then completed another brief filler task, the Reverse Digit Span task from the Wechsler Adult Intelligence Scale-IV (WAIS-IV; Wechsler, 2008) before performing the interpretation and recall tests, in that order. The test phase was followed by the stress task. All participants (training and no training conditions) completed the test phase and the stress task. The training sessions lasted about 45–60 minutes, the test phase about 30 minutes, and the stress task 12 minutes. All study procedures were approved by the Institutional Review Board.

Results

Demographic Characteristics and Mood Ratings

Demographic characteristics and mood ratings throughout the training are presented in Table 1. Participants in the five groups did not differ in gender, age, or education. As expected, there was a main effect of diagnostic group for BDI-II scores, $F(1,55) = 213.28, p < .001, \eta^2 = .79$, there was no main effect of training condition, $F(1,55) < 1$, and no interaction of group and condition, $F(1,55) < 1$. About half of the MDD participants in each group presented with a comorbid condition, mostly anxiety disorders, $\chi^2(2, N = 48) = 3.86, p > .10$. We also compared mood ratings at three time-points during the training-- prior to the interpretation training (Time 1), after the first training session (Time 2), and after the second training session (Time 3) -- using a mixed-design analysis of variance (ANOVA) with group (MDD vs. CTL) and training condition (positive vs. negative) as the between-subjects factors and time-point and valence (positive vs. negative affect) as the within-subjects factors. The analysis yielded main effects of time, $F(2,110) = 20.37; p < .001, \eta^2 = .27$, valence, $F(1,55) = 104.6, p < .001, \eta^2 = .66$, and of group, $F(1,55) = 6.95, p = .018, \eta^2 = .11$, and an interaction of group and valence, $F(1,55) = 69.15, p < .001, \eta^2 = .56$. No main or

interaction effects involving training condition were found. Participants in both training conditions improved in their negative affect but MDD participants reported more negative affect throughout the course of the study. In addition, we found a greater decline in negative affect in the MDD compared to the CTL group. Importantly, however, training condition did not affect mood ratings.

Interpretation Training

Means and standard deviations for all main study variables are presented in Table 2. We first compared MDD and CTL participants in the positive- versus negative-training conditions. We predicted that the training should affect all participants but that training effects should be larger in the MDD than in the CTL group. To examine the effects of CBM-I on participants' interpretations of novel scenarios, similarity ratings for the test scenarios were analyzed by a mixed-model ANOVA with group (MDD vs. CTL) and training condition (positive vs. negative) as the between-subjects factors and valence of the statements (positive vs. negative) and type of statement (target vs. foil) as within-subjects factors. This analysis yielded a three-way interaction of training condition, valence, and type, $F(1, 52) = 11.19, p = .002, \eta^2 = .18$. Contrary to expectations, no effect of diagnostic group emerged. Follow-up analyses examining this three-way interaction were conducted separately for foil and target statements. Results of the follow-up tests revealed no significant training-condition differences in similarity ratings for positive foils, $t(54) = 1.64, p = .107$, or negative foils, $t(54) = 0.78, p = .436$. There were, however, significant training-condition differences for the target statements (see Figure 1). Compared to participants in the negative-training condition, participants in the positive-training condition chose higher similarity ratings for positive target statements, $t(54) = 3.39, p = .001$, and lower similarity ratings for negative target statements, $t(54) = 2.45, p = .018$. Overall, there was no significant main effect of group and no significant interactions with group. Thus, the training was effective in all participants, and the effects did not differ as a function of diagnostic group. Finally, we conducted ANOVAs to compare MDD participants in the positive- and negative-training conditions to those in the no-training condition. The results of these analyses yielded no training-condition differences for negative targets, $F(2,43) < 1$, for negative foils, $F(2,43) < 1$, or for positive foils, $F(2,43) = 1.34, p = .274$. MDD participants in the three training conditions did differ significantly, however, in their responses to the positive targets, $F(2,43) = 4.09, p < .024, \eta^2 = .16$. Depressed participants in the positive-training condition endorsed higher similarity ratings for positive target statements than did depressed participants in both the negative-, $t(27) = 2.46, p = .021$, and the no-training conditions, $t(30) = 2.64, p = .031$, who did not differ in their endorsement of positive statements, $t(29) < 1$. In summary, participants endorsed higher similarity ratings for target statements that corresponded to their training condition. No differences were found between MDD and CTL participants. Within the sample of MDD participants, those in the negative-training condition did not differ from those in the no-training condition. In contrast, MDD participants in the positive-training condition endorsed higher similarity ratings for positive target statements than did MDD participants in both the negative-training and the no-training conditions.

Recall Task

We also examined whether CBM-I affects participants' memory for the test scenarios. In this context, we predicted that CBM-I will result in a corresponding memory bias in the free recall task and that MDD and CTL participants will differ in the strength of this training effect. We assessed the number and valence of memory distortions or intrusions (i.e., new details that the participants "recalled" that had never been part of the original scenarios). Similar to the analytic approach reported above, we first conducted a mixed-model ANOVA with group (MDD vs. CTL) and training condition (positive vs. negative) as the between-subjects factors and valence of the intrusion (positive vs. negative vs. neutral) as the within-subjects factor to compare the training conditions on number of intrusions. This analysis yielded a significant interaction of training condition and intrusion valence, $F(2, 110) = 7.90, p = .001, \eta^2 = .13$; the effect of diagnostic group was not significant. Follow-up tests showed that participants in the two training conditions did not differ in the number of reported neutral intrusions, $t(57) < 1$; there were significant differences between the positive- and negative-training conditions, however, in the number of both positive intrusions, $t(57) = 2.48, p = .016$, and negative intrusions, $t(57) = 2.47, p = .017$. As illustrated in Figure 2, participants in the positive-training condition "recalled" more positive intrusions than did participants in the negative-training condition; similarly, participants in the negative-training condition "recalled" more negative intrusions than did participants in the positive-training condition. These results indicate that participants in each training condition reported more memory intrusions that corresponded to the valence of their training.

Finally, we conducted an ANOVA to compare the positive- and negative-training conditions to the no-training condition within the MDD sample. Although not reaching conventional levels of statistical significance, there were trends for both positive, $F(2, 45) = 2.60, p = .08$, and negative intrusions, $F(2, 45) = 2.24, p = .11$, that were in the predicted direction (neutral intrusions, $F(2, 45) = 1.1, p = .338$). As illustrated in Figure 2, MDD participants in the positive-training condition reported more positive intrusions than did MDD participants in the negative-training condition, $t(29) = 2.81, p = .037$; similarly, MDD participants in the negative-training condition reported more negative intrusions than did MDD participants in the positive-training condition, $t(29) = 2.51, p = .040$. MDD participants in both training conditions, however, did not differ significantly from participants in the no-training condition. Further supporting the formulation that interpretation and memory biases are related, significant correlations among number of intrusions and similarity ratings for target statements were obtained across all participants. These correlations were as follows: positive intrusions and positive targets, $r(73) = .41, p < .001$; negative intrusions and negative targets, $r(73) = .40, p < .001$. There were no significant correlations among intrusions and similarity ratings for target statements when examining each training condition separately.

Stress Task

We examined heart rate and self-reported distress to assess whether the training conditions differentially affected stress reactivity and recovery. We conducted ANOVAs with time (baseline, anticipation and recovery) as the within-subject variable and diagnostic group and training condition as the between-subjects variables. The pre-stressor questionnaire was not

given to participants in the no-training condition; therefore, the analyses for self-reported distress examine only the two training conditions. The ANOVA examining self-reported distress (see Table 1) yielded a main effect of time, $F(2,110) = 54.07, p < .001, \eta^2 = .49$, and a main effect of group, $F(1,55) = 22.43, p < .001, \eta^2 = .29$, which were qualified by a significant interaction of group and time, $F(2,110) = 4.14, p = .018, \eta^2 = .07$. These results indicate that the stressor was perceived as stressful and that the MDD and CTL groups differed in their perception of the stressfulness. There was, however, no effect of training on self-reported level of distress. The ANOVA for heart rate yielded a significant main effect of time, $F(2,116) = 27.80, p = .000, \eta^2 = .32$, which was qualified by a significant interaction of time and condition, $F(2,116) = 2.95, p = .005, \eta^2 = .05$; there was no effect of diagnostic group on heart rate. An inspection of Figure 3 shows that participants in the two training conditions differed significantly in their increase in heart rate from baseline to anticipation: participants in the negative-training condition exhibited a significantly higher increase in heart rate than did participants in the positive-training condition, $t(55) = 2.18, p = .033$. Participants in the negative- and positive-training conditions did not differ in heart rate at recovery, $t(55) < 1$. Finally, we compared MDD participants in the positive- and negative-training conditions to MDD participants in the no-training condition. This ANOVA yielded a significant interaction of time and condition, $F(1,43) = 2.78, p = .032, \eta^2 = .12$. MDD participants in the no-training condition did not differ from MDD participants in the positive-training condition with respect to increase in heart rate, $t(29) = 1.08, p = .286$, but they exhibited a smaller increase in heart rate than did MDD participants in the negative-training group condition, $t(27) = 2.38, p = .025$. These results should be interpreted with caution, however. As Figure 3 shows, the MDD participants in the no-training condition had elevated heart rate at baseline; therefore, the lack of reactivity in participants in this condition may be due to a ceiling effect.

Discussion

The present study was conducted to examine whether CBM-I in participants diagnosed with MDD affects interpretation of novel ambiguous scenarios, memory, and physiological stress responding. The results indicate that MDD and CTL participants in the positive-training condition demonstrated an interpretation bias that corresponded to their training condition. Specifically, in both diagnostic groups, participants in the positive-training condition were more likely to rate positive than negative target sentences as similar to the test scenario and were also more likely to do so than were participants in the negative-training condition. The study included a sample of MDD participants who did not undergo training. MDD participants in the positive-training condition were significantly more likely to endorse positive interpretations than were MDD participants in this no-training condition; in contrast, MDD participants who were assigned to the negative-training condition did not differ from MDD participants in the no-training condition. Importantly, we found training effects on memory and stress responding. Specifically, compared to the participants in the negative-training condition, those in the positive-training condition were more likely to report positive memory intrusions and showed a smaller increase in heart rate in response to the stress induction.

The current findings replicate past research demonstrating effects of CBM-I (e.g., Mackintosh et al., 2006; Mathews & Mackintosh, 2000; Salemink, van den Hout, & Kindt, 2007; Yiend et al., 2005). Most CBM-I studies, however, have focused on anxiety disorders. This study shows that CBM-I is beneficial in MDD participants which supports findings from recent studies that have used this training in MDD (Bowler et al., 2013). Like Bowler et al. (2013) and Yiend et al. (2013) we found effects of the training on interpretation biases in MDD. In contrast to the findings reported by Yiend et al., however, our training also had a positive impact on stress reactivity. The discrepant findings obtained in these two studies may be due to the difference in number of training sessions (2 session in our study versus 1 in the study by Yiend et al.) and/or may be due to the use of psychophysiological measures of stress responding. Indeed, in the current study we found no indication of changes in self-reported stress responding but a significant difference when examining heart rate. Heart rate and affective responses during stress are only very loosely correlated with each other (Cohen et al., 2010), and only under certain circumstances (Waugh et al., 2010), suggesting that the heart rate changes in this study may have been due to training-induced effects in some process other than negative affect, such as in the perceived effort of the task (Peters et al., 1998). More studies are clearly needed that examine stress reactivity using a multi-method approach and that add more training sessions to corroborate these findings.

In addition, previous studies have focused only on one cognitive bias, despite researchers' suggestion that cognitive biases interact and influence each other (e.g., Hirsch et al., 2006). Indeed, in the current study, our measures of interpretation and memory bias were significantly correlated across groups. These findings concerning memory bias are consistent with research demonstrating that how events are initially interpreted affects how they are subsequently recalled (Hertel & Brozovich, 2010). Given the evidence for memory biases in depression (Matt et al., 1992; Watkins, Mathews, Williamson, & Fuller, 1992) and the potential benefits on emotion regulation and stress reactivity from training positive biases, investigating the association between interpretation and memory is a critical area for further exploration. For example, the tendency to respond to negative events and mood states with rumination is an important risk factor for the onset of depression (Nolen-Hoeksema, Wisco, & Lyubomirsky, 2008). People who ruminate repetitively think about past events and interpret these events negatively. Indeed, rumination has been shown to strengthen mood-congruent memory biases (Lyubomirsky, Caldwell, & Nolen-Hoeksema, 1998). In contrast, the benefits of reappraisal as a more adaptive emotion regulation strategy may lie in its resemblance to positive interpretation training. Our findings suggest that the habitual use of rumination or reappraisal may have important consequences not only for acute emotional responding but also for how the emotion-eliciting event will be remembered and thereby affect emotional well-being in the future.

The present study also replicates previous results by Salemink, Hertel, and Mackintosh (2010), who trained biases in order to examine whether the recall of prior events could be modified by subsequently acquired biases. Participants in this study exhibited biased recall of their own interpretation of the scenarios, but not of the scenarios themselves. This lack of a bias in recall of the scenarios was predicted on the basis that recall intrusions arise from imagery processes that operate during initial interpretations (see Hertel et al., 2008; Hirsch, et al., 2006).

Although previous training studies have reported effects of both positive and negative training (Mathews & Mackintosh, 2000; Salemink et al., 2007; Yiend et al., 2005), our findings suggest that positive training had stronger effects than did negative training. MDD participants in the negative training group did not differ from the no-training group in their performance on any of the outcome measures. Indeed, our findings show that MDD compared to CTL participants show a negative bias from the outset and that whereas training induced a negative bias in CTL participants, it did not make MDD participants worse.

Findings concerning training effects on stress response in previous studies that have examined CBM-I in depression have been equivocal. The present study is the first to examine physiological stress responding; our results indicate that training affects physiological arousal in response to an acute stressor. This result has important clinical implications. Depression has been associated with cardiovascular dysregulation (Salomon, Clift, Karlsdottir, & Rottenberg, 2009) and has been shown to be risk factor for heart disease (Musselman, Evans, & Nemeroff, 1998). The finding that CBM-I training affects cardiovascular stress responding is promising, but it is clear that these findings require replication. Unfortunately, the fact that participants in the no-training condition were elevated in their heart rate at baseline precluded us from being able to compare the positive- and negative-training conditions to the no-training condition. It is likely that this initial heart-rate elevation is due to this being their first visit to the laboratory.

We should note three limitations of this study. First, we did not assess pre-training biases in interpretation or memory. We did, however, include a no-training control condition that allowed us to compare baseline responding to training-induced responding; our results show clear differences between the no-training and the positive-training conditions on all of our measures. We should point out, however, that we included a no-training condition only for the MDD group; we did not recruit a group of CTL participants who did not undergo training – including this group was not central to the main aim of this study, to examine CBM-I in clinical depression. It will be helpful in future work to include both CTL and MDD groups who do not undergo the training. Second, the training consisted of only two sessions. Two training sessions is an improvement over the many studies in this area that have used a single training session and clearly was sufficient to demonstrate that biases can be modified. Using a larger number of training sessions will be important in studies examining the longer-term clinical impact of these trainings. Indeed, recent studies examining CBM-I as an intervention for depression have examined interventions lasting 10 weeks or more. It is encouraging that the two-session training used in this study yielded significant findings, but future research is required to extend the training time in order to examine the effects of more prolonged interventions on memory and on stress reactivity. Finally, we used the training scenarios developed by Mathews and Mackintosh (2000) that describe situations that are most centrally relevant to socially anxious participants. Other researchers have now developed training protocols that are more specific to the concerns of depressed participants (Lester, Mathews, Davison, Burgess & Yiend, 2011; Yiend et al., 2014) and the use of such stimuli could result in stronger training effects than those documented in this study. Indeed, it is possible that the lack of group differences is due to

the use of scenarios that are not depression-specific. Future studies are needed to examine this possibility more explicitly and systematically.

Alternative explanations of our results involve mood and the order of our tasks. Mood affects memory and it is therefore possible that training-related changes in mood contributed to the changes in interpretation and memory that we found in this study. This account does not appear to be viable, however, given that mood did not change as a consequence of training, and that mood was not related to measures of memory or interpretation (see Footnote 1). Second, because we necessarily assessed interpretation bias before recall bias, it is possible that the assessment itself played a role in changes in our dependent measures. Participants in both training conditions were exposed to the same four sentences (2 targets and 2 foils); hence, condition-associated differences in recall cannot be explained by differences in exposure to statements in the interpretation task. In fact, we would have encountered several problems if we had measured recall first. During recall, participants are given more time to elaborate on their memory of the previously presented scenarios; therefore, their subsequent similarity ratings would primarily be assessing how participants recalled the descriptions during recall, rather than how they interpreted the situation initially. In the design of the current study, we presented the interpretation task as a brief exercise in which participants are asked to quickly indicate similarity ratings, and we allowed less time for elaboration. We believe that the main constraint that test order places on these results involves external validity. It may be that explicit interpretation is necessary to obtain biases in remembering the scenarios. This possibility should be investigated in future research.

Biases in interpretation and memory affect emotion regulation in various ways. They can influence judgments of the frequency of positive and negative events, and of the likelihood that these types of events will occur in the future. Memory biases have also been associated with lowered self-esteem and with increased levels of negative affect and hopelessness (MacLeod & Campbell, 1992). Increased accessibility of negative material and difficulties recalling mood-incongruent material may interfere with the use of effective mood regulation strategies. These difficulties in effectively managing negative mood may be an important mechanism by which memory biases affect the maintenance and recurrence of depressive episodes (Nolen-Hoeksema, 2000; Nolen-Hoeksema & Larson, 1999; Roberts, Gilboa, & Gotlib, 1998). Increased accessibility and recall of positive material, on the other hand, is related to higher levels of well-being (Charles, Mather & Carstensen, 2003). Reducing negative biases in memory and interpretation therefore, may improve mood and decrease emotional vulnerability following stressful experiences, thereby reducing the risk for the onset and recurrence of emotional disorders. Future studies are needed to examine underlying mechanisms and the durability of the training effects. Given the critical role of memory biases in the development and maintenance of emotional disorders, the ability to manipulate interpretation biases and thereby change individuals' memory for events could hold great promise for the prevention and treatment of these disorders and provide greater insight into the functional role of cognitive biases in emotional disorders.

¹No significant correlations were found among mood ratings assessed after the training and any of the interpretation or memory measures (i.e., similarity ratings for positive and negative targets and positive, negative, and neutral intrusions) in the two training groups.

Acknowledgments

The authors thank Brooke Gilbert for her help in recruiting and running the participants in this study. This research was supported by National Institute of Mental Health grant MH59259 awarded to IHG.

References

- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4. Washington, D.C: Author; 1994.
- Beck, AT.; Steer, RA. Beck Depression Inventory-Manual. San Antonio: Psychological Corp; 1993.
- Beck AT, Steer RA, Garbin MG. Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clinical Psychology Review*. 1988; 8(1):77–100.
- Bowler JO, Mackintosh B, Dunn BD, Mathews A, Dalgleish T, Hoppitt L. A comparison of cognitive bias modification for interpretation and computerized cognitive behavior therapy: Effects on anxiety, depression, attentional control, and interpretive bias. *Journal of Consulting and Clinical Psychology*. 2012; 80(6):1021–1033. [PubMed: 22963595]
- Charles S, Mather M, Carstensen LL. Aging and emotional memory: The forgettable nature of negative images for older adults. *Journal of Experimental Psychology: General*. 2003; 132(2):310–324. [PubMed: 12825643]
- Cohen S, Hamrick N, Rodriguez MS, Feldman PJ, Rabin BS, Manuck SB. The stability of and intercorrelations among cardiovascular, immune, endocrine, and psychological reactivity. *Annals of Behavioral Medicine*. 2000; 22(3):171–179. [PubMed: 11211850]
- Cowden Hindash AH, Amir N. Negative interpretation bias in individuals with depressive symptoms. *Cognitive Therapy and Research*. 2012; 36(5):502–511.
- Ellsworth, PC.; Scherer, KR. Appraisal processes in emotion. In: Davidson, RJ.; Goldsmith, H.; Scherer, KR., editors. *Handbook of Affective Sciences*. New York: Oxford University Press; 2003.
- Everaert J, Koster EW, Derakshan N. The combined cognitive bias hypothesis in depression. *Clinical Psychology Review*. 2012; 32(5):413–424. [PubMed: 22681914]
- First, MB.; Spitzer, MB.; Gibbon, M.; Williams, JBW. Structured Clinical Interview for DSM-IV Axis I Disorders - Patient Edition (SCID-I/P, Version 2.0). Biometrics Research Department, New York State Psychiatric Institute; 1995.
- Foa EB, Riggs D, Dancu CVS, Rothbaum BO. Reliability and validity of a brief instrument for assessing Post Traumatic Stress Disorder. *Journal of Traumatic Stress*. 1993; 6(4):459–473.
- Fredrickson BL, Mancuso RA, Branigan C, Tugade MM. The undoing effect of positive emotions. *Motivation and Emotion*. 2000; 24(4):237–258. [PubMed: 21731120]
- Grey S, Mathews A. Effects of training on interpretation of emotional ambiguity. *Quarterly Journal of Experimental Psychology*. 2000; 53A:1143–1162. [PubMed: 11131817]
- Gross, JJ.; John, OP. Wise emotion regulation. In: Barrett, LF.; Salovey, P., editors. *The wisdom in feeling: Psychological processes in emotional intelligence*. New York, NY: Guilford Press; 2002. p. 297–319.
- Hallion LS, Ruscio A. A meta-analysis of the effect of cognitive bias modification on anxiety and depression. *Psychological Bulletin*. 2011; 137(6):940–958. [PubMed: 21728399]
- Hertel, PT. Habits of thought produce memory biases in anxiety and depression. In: Yiend, J., editor. *Cognition, emotion, and psychopathology*. Cambridge, UK: Cambridge University Press; 2004. p. 109–129.
- Hertel PT, Vasquez E, Benbow A, Hughes M. Recollection is impaired by the modification of interpretation bias. *Journal of Abnormal Psychology*. 2011; 120(4):902–910. [PubMed: 21668083]
- Hertel PT, Brozovich F. Cognitive habits and memory distortions in anxiety and depression. *Current Directions in Psychological Science*. 2010; 19(3):155–160.
- Hertel PT, Brozovich F, Joormann J, Gotlib IH. Biases in interpretation and memory in generalized social phobia. *Journal of Abnormal Psychology*. 2008; 117(2):278–288. [PubMed: 18489204]

- Hirsch CR, Clark DM, Mathews A. Imagery and interpretations in social phobia: Support for the combined cognitive biases hypothesis. *Behavior Therapy*. 2006; 37:223–236. [PubMed: 16942974]
- Holmes EA, Lang TJ, Shah DM. Developing interpretation bias modification as a ‘cognitive vaccine’ for depressed mood: Imagining positive events makes you feel better than thinking about them verbally. *Journal of Abnormal Psychology*. 2009; 118(1):76–88. [PubMed: 19222316]
- Joormann J, Siemer M, Gotlib IH. Mood regulation in depression: Differential effects of distraction and recall of happy memories on sad mood. *Journal of Abnormal Psychology*. 2007; 116:484–491. [PubMed: 17696704]
- Joormann, J.; Yoon, K.L.; Siemer, M. Cognition and emotion regulation. In: Kring, A.; Sloan, D., editors. *Emotion Regulation and Psychopathology*. New York, NY: Guilford Press; 2010. p. 174–203.
- Lester KJ, Mathews A, Davison PS, Burgess JL, Yiend J. Modifying cognitive errors promotes cognitive well being: A new approach to bias modification. *Journal Of Behavior Therapy And Experimental Psychiatry*. 2011; 42(3):298–308. [PubMed: 21352718]
- Lyubomirsky S, Caldwell ND, Nolen-Hoeksema S. Effects of ruminative and distracting responses to depressed mood on retrieval of autobiographical memories. *Journal of Personality and Social Psychology*. 1998; 75(1):166–177. [PubMed: 9686457]
- Mackintosh B, Mathews A, Yiend J, Ridgeway V, Cook E. Induced biases in emotional interpretation influence stress vulnerability and endure despite changes in context. *Behavior Therapy*. 2006; 37(3):209–222. [PubMed: 16942973]
- MacLeod C, Campbell L. Memory accessibility and probability judgments: An experimental evaluation of the availability heuristic. *Journal of Personality and Social Psychology*. 1992; 63(6): 890–902. [PubMed: 1460558]
- Mathews A, Mackintosh B. Induced emotional interpretation bias and anxiety. *Journal of Abnormal Psychology*. 2000; 109:602–615. [PubMed: 11195984]
- Mathews A, MacLeod C. Cognitive vulnerability to emotional disorders. *Annual Review of Clinical Psychology*. 2005; 1:167–95.
- Matt GE, Vazquez C, Campbell WK. Mood-congruent recall of affectively toned stimuli: A meta-analytic review. *Clinical Psychology Review*. 1992; 12:227–255.
- Micco JA, Henin A, Hirshfeld-Becker DR. Efficacy of interpretation bias modification in depressed adolescents and young adults. *Cognitive Therapy and Research*. 2014; 38(2):89–102. [PubMed: 24729643]
- Musselman DL, Evans DL, Nemeroff CB. The relationship of depression to cardiovascular disease: Epidemiology, biology, and treatment. *Archives of General Psychiatry*. 1998; 55(7):580–592. [PubMed: 9672048]
- Nolen-Hoeksema S. The role of rumination in depressive disorders and mixed anxiety/depressive symptoms. *Journal of Abnormal Psychology*. 2000; 109:504– 511. [PubMed: 11016119]
- Nolen-Hoeksema, S.; Larson, J. *Coping with loss*. Mahwah, NJ: Erlbaum; 1999.
- Nolen-Hoeksema S, Wisco BE, Lyubomirsky S. Rethinking rumination. *Perspectives on Psychological Science*. 2008; 3(5):400–424.
- Peters ML, Godaert GLR, Ballieux RE, van Vliet M, Willemsen JJ, Sweep FCGJ, et al. Cardiovascular and endocrine responses to experimental stress: Effects of mental effort and controllability. *Psychoneuroendocrinology*. 1998; 23(1):1–17. [PubMed: 9618748]
- Salemink E, Hertel P, Mackintosh B. Interpretation training influences memory for prior interpretations. *Emotion*. 2010; 10:903–907. [PubMed: 21171760]
- Salemink E, van den Hout M, Kindt M. Trained interpretive bias and anxiety. *Behaviour Research and Therapy*. 2007; 45(2):329–340. [PubMed: 16859638]
- Salomon K, Clift A, Karlsdóttir M, Rottenberg J. Major depressive disorder is associated with attenuated cardiovascular reactivity and impaired recovery among those free of cardiovascular disease. *Health Psychology*. 2009; 28(2):157–165. [PubMed: 19290707]
- Siemer M, Mauss IB, Gross JJ. Same situation -- different emotions: How appraisals shape our emotions. *Emotion*. 2007; 7:592–600. [PubMed: 17683215]

- Tran TB, Hertel PT, Joormann J. Cognitive bias modification: Induced interpretive biases affect memory. *Emotion*. 2011; 11(1):145–152. [PubMed: 21401234]
- Watkins PC, Mathews A, Williamson DA, Fuller RD. Mood-congruent memory in depression: Emotional priming or elaboration? *Journal of Abnormal Psychology*. 1992; 101(3):581–586. [PubMed: 1500617]
- Waugh CE, Panage S, Mendes WB, Gotlib IH. Cardiovascular and affective recovery from anticipatory threat. *Biological Psychology*. 2010; 84:169–175. [PubMed: 20096747]
- Wechsler, D. Wechsler Adult Intelligence Scale. 4. San Antonio, TX: The Psychological Corporation; 2008. (WAIS-IV)
- Wilhelm, FH.; Peyk, P. ANSLAB: Autonomic Nervous System Laboratory (Version 2.4). 2005. Retrieved from the SPR Software Repository: <http://www.sprweb.org>
- Williams AD, Blackwell SE, Mackenzie A, Holmes EA, Andrews G. Combining imagination and reason in the treatment of depression: A randomized controlled trial of internet-based cognitive-bias modification and internet-CBT for depression. *Journal of Consulting and Clinical Psychology*. 2013; 81(5):793–799. [PubMed: 23750459]
- Williams JB, Gibbon M, First MB, Spitzer RL, Davis M, Borus J, et al. The Structured Clinical Interview for DSM-III-R (SCID): Multisite test-retest reliability. *Archives of General Psychiatry*. 1992; 49:630–636. [PubMed: 1637253]
- Wilson EJ, MacLeod C, Mathews A, Rutherford EM. The causal role of interpretive bias in anxiety reactivity. *Journal of Abnormal Psychology*. 2006; 115:103–111. [PubMed: 16492101]
- Yiend J, Lee J, Tekes S, Atkins L, Mathews A, Vrinten M, Shergill S. Modifying interpretation in a clinically depressed sample using ‘cognitive bias modification-errors’: A double blind randomised controlled trial. *Cognitive Therapy and Research*. 2014; 38(2):146–159.
- Yiend J, Mackintosh B, Andrews M. Enduring consequences of experimentally induced biases in interpretation. *Behaviour Research and Therapy*. 2005; 43:779–797. [PubMed: 15890169]

Figure 1.

Mean similarity ratings for positive and negative target statements when rating novel test scenarios separated by training group (PosTrain = Positive Training, NegTrain = Negative Training, NoTrain = No Training) and diagnostic group (CTL = control participants, MDD = participants diagnosed with Major Depressive Disorder). Similarity ratings ranged from 1 (*very different*) to 4 (*very similar*). PosTarget = positive target, NegTarget = negative target. Error bars represent 1 standard error.

Figure 2.

Mean number of positive, negative, and neutral memory intrusions (i.e., material that was not originally presented) made by each training group (PosTrain = Positive Training, NegTrain = Negative Training, NoTrain = No Training) and diagnostic group (CTL = control participants, MDD = participants diagnosed with Major Depressive Disorder) during the recall task. Error bars represent 1 standard error.

Figure 3.

Mean heartrate at baseline and after anticipation in the three training groups (PosTrain = Positive Training, NegTrain = Negative Training, NoTrain = No Training).

Table 1

Demographic Characteristics, Mood Ratings and Self-reported Distress in Stress Task

Group Training Condition	CTL		MDD		No training
	positive	negative	positive	negative	
N/N(female)	14/9	14/8	16/10	15/11	17/12
Age M(SD)	40.93 (11.24)	42.07 (12.77)	38.50 (10.77)	39.40(12.18)	33.88 (11.25)
% College	64	92	57	60	48
% Comorbid	0	0	43	53	70
Disorders					
BDI	3.07(3.61)	1.71 (2.61)	32.88 (8.18)	30.93 (12.07)	36.41 (10.77)
NA – time 1	1.12 (0.28)	1.16 (0.31)	1.81 (0.93)	2.02 (0.82)	
NA – time 2	1.14 (0.31)	1.11 (0.28)	1.66 (0.66)	1.65 (0.47)	
NA – time 3	1.02 (0.09)	1.04 (0.12)	1.35 (0.61)	1.64 (0.62)	
PA – time 1	3.57 (0.74)	3.33 (0.89)	1.93 (0.87)	2.20 (0.81)	
PA - time 2	3.11 (1.08)	2.76 (0.88)	1.75 (0.82)	2.00 (0.77)	
PA – time 3	3.07 (0.82)	2.62 (1.01)	1.81 (0.79)	1.67 (0.62)	
Stress Task: Self-reported Distress					
Baseline	1.28 (0.69)	1.14 (0.31)	1.75 (0.68)	1.93 (0.96)	
Anticipation	2.00 (1.15)	2.19 (0.95)	3.04 (0.94)	3.26 (1.11)	3.41 (1.15)
Recovery	1.19 (0.33)	1.29 (0.55)	1.39 (0.51)	1.71 (0.65)	2.15 (1.19)

Note. CTL = Control participants; MDD = Participants diagnosed with Major Depressive Disorder; BDI = Beck Depression Inventory; NA= negative affect, PA = positive affect

Table 2
Means and Standard Deviations (SD) of Recognition Ratings, Intrusions, and Heart Rate during Stress Task

Group Training Condition	CTL		MDD		No training	
	positive	negative	positive	negative	positive	negative
Recognition						
Foil-pos	1.56 (0.41)	1.44 (0.35)	1.97 (0.74)	1.61 (0.65)	1.68 (0.50)	
Foil-neg	1.34 (0.33)	1.52 (0.33)	1.76 (0.59)	1.85 (0.75)	1.86 (0.59)	
Target-pos	2.82 (0.38)	2.49 (0.35)	2.99 0.48)	2.50 (0.58)	2.48 (0.60)	
Target-neg	2.09 (0.46)	2.65 (0.39)	2.59 (0.55)	2.76 (0.58)	2.58 (0.54)	
Recall						
Intrusion-pos	5.89 (5.97)	2.96 (3.97)	4.21 (4.23)	1.60 (1.96)	2.88 (2.94)	
Intrusion-neg	4.42 (4.13)	6.32 (4.14)	3.75 (2.25)	8.20 (7.97)	6.29 (6.07)	
Intrusion-neu	9.64 (4.88)	10.93 (6.99)	8.21 (4.31)	9.54 (4.78)	7.26 (2.85)	
Stress Task: Heart Rate						
Baseline	68.33 (10.86)	69.28 (7.93)	72.26 (9.06)	69.32 (10.77)	79.74 (11.84)	
Anticipation	70.68 (11.64)	73.73 (11.10)	74.77 (8.84)	74.66 (11.20)	80.76 (12.35)	
Recovery	68.05 (11.44)	70.58 (8.34)	73.03 (9.82)	69.66 (9.15)	80.30 (12.11)	

Note. CTL = Control participants; MDD = Participants diagnosed with Major Depressive Disorder; pos = positive, neg = negative, neu = neutral