Published in final edited form as:

Psychol Med. 2011 January ; 41(1): 195–206. doi:10.1017/S0033291710000255.

A latent profile analysis of the typology of bulimic symptoms in an indigenous Pacific population: Evidence of cross-cultural variation in phenomenology

Jennifer J. Thomas, Ph.D.^{1,2}, Ross D. Crosby, Ph.D.^{3,4}, Stephen A. Wonderlich, Ph.D.^{3,4}, Ruth H. Striegel-Moore, Ph.D.⁵, and Anne E. Becker, M.D., Ph.D., Sc.M.^{1,6}

¹Eating Disorders Clinical & Research Program, Massachusetts General Hospital

²Klarman Eating Disorders Center, McLean Hospital

³Neuropsychiatric Research Institute

⁴University of North Dakota School of Medicine

⁵Department of Psychology, Montana State University

⁶Department of Global Health and Social Medicine, Harvard Medical School

Abstract

Background—Previous efforts to derive empirically based eating disorder (ED) typologies through latent structure modeling have been limited by ethnic and cultural homogeneity of their study populations and their reliance on DSM-IV ED signs and symptoms as indicator variables.

Methods—Ethnic Fijian schoolgirls ($N = 523$) responded to a self-report battery assessing ED symptoms, herbal purgative use, comorbid psychopathology, clinical impairment, cultural orientation, and peer influences. Participants who endorsed self-induced vomiting or herbal purgative use in the past 28 days ($n = 222$) were included in a latent profile analysis (LPA) to identify unique subgroups of bulimic symptomatology.

Results—LPA identified a bulimia nervosa (BN)-like class ($n = 86$) characterized by high rates of binge eating and self-induced vomiting; and an herbal purgative class ($n = 136$) characterized primarily by the use of indigenous Fijian herbal purgatives. Both ED classes endorsed greater eating pathology and general psychopathology than non-purging participants, and the herbal purgative class endorsed greater clinical impairment than either BN-like or non-purging participants. Cultural orientation did not differ between the two ED classes.

Conclusions—The inclusion of study populations underrepresented in mental health research and broadening the scope of relevant signs and symptoms in latent structure models may increase the generalizability of ED nosological schemes to encompass greater cultural diversity.

Correspondence: Jennifer J. Thomas, Eating Disorders Clinical and Research Program, Massachusetts General Hospital, WAC 816D, 15 Parkman St., Boston, MA 02114 or jjthomas@partners.org.

Previous Presentation: Portions of this paper were presented at the NIMH-sponsored R-13 meeting on Eating Disorder Classification in Arlington, Virginia on March 16, 2009; and the American Psychological Association Convention in Toronto, Canada on August 6, 2009.

Declaration of Interest: The authors report no competing financial interests.

Keywords

eating disorders; classification; latent profile analysis; Fiji; herbal purgative

The clinical utility of the DSM-IV eating disorder classification system has been critiqued insofar as the residual category, eating disorder not otherwise specified (EDNOS), is the most common eating disorder diagnosis in clinical settings. In contrast to the diagnostic categories of anorexia nervosa (AN) and bulimia nervosa (BN), EDNOS is phenomenologically heterogeneous (Thomas, *et al.*, 2009) and few clinical trials support relevant treatment strategies. As a result, the upcoming publication of DSM-V has inspired a wave of research on empirical approaches to reconsider optimal eating disorder nosology. To date, ten published studies (Sullivan *et al.*, 1998; Bulik *et al.*, 2000; Keel *et al.*, 2004; Striegel-Moore *et al.*, 2005; Wade *et al.*, 2006; Duncan *et al.*, 2007; Mitchell *et al.*, 2007; Pinheiro *et al.*, 2008; Striegel-Moore *et al.*, 2008; Eddy *et al.*, 2009) have utilized latent class analysis (LCA) or latent profile analysis (LPA) to derive eating disorder subtypes empirically based on the co-occurrence of attitudes and behaviors in symptomatic samples. These analyses have proved invaluable in highlighting the strengths and weaknesses of DSM-IV diagnoses. For example, while several LCA and LPA studies have confirmed the existence of latent classes resembling AN (Bulik *et al.*, 2000; Keel *et al.*, 2004) and BN (Sullivan *et al.*, 1998; Bulik *et al.*, 2000; Keel *et al.*, 2004; Striegel-Moore *et al.*, 2005; Pinheiro *et al.*, 2008; Eddy *et al.*, 2009), others have identified subgroups not yet recognized as formal diagnostic categories in DSM-IV, including binge-eating disorder (Bulik *et al.*, 2000; Striegel-Moore *et al.*, 2005; Mitchell *et al.*, 2007; Pinheiro *et al.*, 2008; Eddy *et al.*, 2009), purging disorder (Striegel-Moore *et al.*, 2005; Mitchell *et al.*, 2007; Pinheiro *et al.*, 2008), and night eating syndrome (Striegel-Moore *et al.*, 2008).

The generalizability of previous LCA and LPA findings to populations underrepresented in mental health research may be limited, however, by the demographic homogeneity of study samples. Specifically, only three studies have included participants from outside the United States, including Canada (Keel *et al.*, 2004), Europe (Keel *et al.*, 2004; Pinheiro *et al.*, 2008), and Australia (Wade *et al.*, 2006), and only one study has included a large proportion of non-Caucasian participants (Striegel-Moore *et al.*, 2005). To date, no LCA or LPA studies have investigated the latent structure of eating pathology among participants outside high-income countries. This lack of broad ethnic, social, and cultural representation is consistent with meta-trends in the broader psychiatric literature, in which just 6% percent of articles published in the field's top journals originate from the global regions outside of Western Europe, North America, Australia, or New Zealand that represent more than 90% of the world's population (Patel & Sumathipala, 2001).

Designated a priority area for adolescent mental health by the World Health Organization (WHO, 2003), eating disorders have global distribution and public health significance. Because eating pathology may present differently outside of North America and Europe, empirical typologies derived from previous eating disorder LCA and LPA studies may not capture the full range of possible presentations. For example, epidemiological and case-finding investigations have found AN to be rare among majority blacks in Curacao (Hoek *et*

al., 2005), Kenya (Njenga & Kangethe, 2004), and Ghana (Bennett *et al.*, 2004). However, a non-fat-phobic variant of AN has been identified in Hong Kong (Lee *et al.*, 2001), Singapore (Ong *et al.*, 1982), and Ghana (Bennett *et al.*, 2004). Interestingly, while two LCA studies (Bulik *et al.*, 2000; Keel *et al.*, 2004) have identified latent classes resembling AN, and five LCA and LPA studies have identified subgroups of low to normal weight participants exhibiting moderate weight concerns (Bulik *et al.*, 2000; Keel *et al.*, 2004; Wade *et al.*, 2006; Mitchell *et al.*, 2007; Eddy *et al.*, 2009), none has identified a class clearly resembling non-fat-phobic AN. This pattern of findings is consistent with etiologic theories that link the core psychopathology of eating disorders—i.e., weight and shape concerns—to specific cultural and historical contexts (Lee *et al.*, 2001; Becker *et al.*, 2009a).

The purpose of the present study was to conduct an LPA of eating disorder phenotypes in an ethnic Fijian study sample to examine potential ethnic and cultural variation in optimal categorization. Ethnic Fijians, a small-scale indigenous Pacific Islander population, are ethnically and culturally distinct from Western populations utilized in previous LCA studies. Although there is no indigenous Fijian nosologic category for eating disorders, weight management strategies are common among Fijian youth (McCabe *et al.*, 2009), and ethnic Fijian cultural traditions support attention to appetite, eating, and weight (Becker, 1995; Mavoa & McCabe, 2008). However, in the setting of globalizing economic and cultural influences on the nutritional environment (Hughes & Lawrence, 2005), urban migration, and social norms for appearance, eating pathology (Becker *et al.*, 2002) and obesity (Becker *et al.*, 2005; Mavoa & McCabe, 2008) have become more common in ethnic Fijian females. Weight management strategies include use of an extensive traditional herbal pharmacopoeia (*dranu*) both to prevent weight loss (Becker, 1995) and also to prevent weight gain (Becker *et al.*, 2009b), and suggest the potential for cross-cultural plasticity in eating disorder presentation.

Therefore, our primary hypothesis was that an LPA utilizing both conventional DSM-IV and uniquely Fijian eating disorder symptoms as indicators would identify novel and culture-specific subgroups with possibly greater local relevance than those identified in previous eating disorder LCAs and LPAs. Our secondary hypothesis was that symptomatic latent classes would differ from asymptomatic classes on external validators including eating pathology, clinical impairment, and general psychopathology. Finally, a *post-hoc* aim was to explore whether symptomatic latent classes were associated with differential social and cultural characteristics.

Methods

Sample Characteristics

Ethnic Fijian schoolgirls ($N = 523$) enrolled in 12 secondary schools registered within one sector of the Fiji Ministry of Education participated in this study, which was part of a larger investigation on social transition and psychopathology risk. Girls aged 15 to 20 years were eligible for inclusion. All of the invited schools agreed to participate, and 71% of eligible students enrolled in the study. Participants' mean age was 16.67 ($SD = 1.09$) years, and mean body mass index (BMI) was 23.96 ($SD = 3.35$) kg/m². Participants were evenly

divided between rural (50%, $n = 262$) and peri-urban (50%, $n = 261$) locations. Less than half (41%, $n = 216$) lived in relative material affluence (i.e., operationalized as having household access to electricity, a gas stove, a refrigerator, and running water); the majority (59%, $n = 307$) lived in relative material poverty (i.e., lacking one or more of these goods or services). Furthermore, 41% ($n = 216$) reported having sometimes gone hungry in the past month due to insufficient food in the home.

Procedure

The study utilized data originally collected as part of a two-stage design described previously (Becker *et al.*, 2009b; Becker *et al.*, 2009c). Briefly, all participants completed a self-report battery at Stage 1. Assessments were offered in either English (the language of formal instruction) or the vernacular (Fijian) language. For LPA indicators, we drew from Stage 1 self-report and anthropometric data available for the full sample ($n = 523$). Stage 2 interview data were available for a subsample ($n = 215$) selected by eligibility criteria specified in a related study identifying symptomatic ($n = 178$) and asymptomatic ($n = 37$) groups (described in Becker *et al.*, 2009b; Becker *et al.*, 2009c). The Partners Human Research Committee (PHRC), Harvard Medical School Human Subjects Committee, and Fiji National Research Ethical Review Committee approved the data collection. The PHRC approved the secondary data analyses.

Measures

Eating pathology—We assessed eating pathology with the Eating Disorder Examination Questionnaire (EDE-Q) version 5.2, adapted to include an item assessing herbal purgative use. The EDE-Q is a self-report measure evaluating the domains of Restraint, Eating Concern, Shape Concern, and Weight Concern (Fairburn & Beglin, 1994). All participants rated the frequency eating disorder symptoms over the past 28 days on a Likert scale ranging from “no days” to “every day.” Internal consistency, retest reliability, and construct validity were adequate in our sample (Becker *et al.*, 2009c).

Herbal purgative use—We evaluated herbal purgative use as a potential LPA indicator with the following item incorporated into the EDE-Q: “Over the past 28 days, how many times have you taken traditional Fijian *dranu* to cause diarrhea, clean out your stomach, or suppress your appetite as a means of controlling your shape or weight?” We have previously reported that self-reported herbal purgative use was prevalent in our sample, and significantly associated with both eating pathology (Becker *et al.*, 2009c) and eating-related clinical impairment (Becker *et al.*, 2009b).

Evaluation of herbal purgative use to qualify it as a potential indicator of eating pathology—Because the phrasing of this single assessment item could conceivably encompass non-pathological (i.e., culturally-sanctioned medicinal) use of *dranu*, we sought to ascertain that herbal purgative use was rationalized as a mode of weight management by at least some participants in order to justify its inclusion as an LPA indicator. In order to interrogate the rationales for herbal purgative use and confirm its local cultural relevance as a potential sign of eating pathology, we examined portions of Stage 2 interview transcripts from a convenience sample of 19 respondents who had endorsed this behavior on the EDE-

Q. Transcripts were from a semi-structured interview based upon the Eating Disorder Examination interview version 16.0 (Fairburn & Beglin, 1994) and adapted for this population by adding culturally and developmentally appropriate probe questions addressing the presence, motivation, and context of herbal purgative use. Of the 19, 74% ($n = 14$) confirmed using herbal purgatives specifically in order to influence their shape or weight, whereas 26% ($n = 5$), exclusively provided rationales less clearly linked to—or even independent of—eating pathology, such as the alleviation of postprandial discomfort after overeating ($n = 1$), the prevention of appetite loss ($n = 2$), or the treatment of physical illness ($n = 2$). Table 1 displays interview transcript excerpts from six participants who affirmed that their herbal purgative use was linked to weight management and purging, although some excerpts illustrate plural rationales. Indeed, herbal purgative use was compensatory for specific overeating episodes (e.g., cases 2, 3, 4), as well as non-compensatory, but still motivated by weight management goals (e.g., cases 1, 5, 6). Notably, we found evidence that family members sometimes encouraged or enabled herbal purgative use (e.g., cases 1, 3, 4, 6). Consistent with our study aim to encompass a broad range of behaviors that might reflect culture-specific variation in eating pathology, we conceptualized herbal purgative use as a possible eating disorder symptom and utilized it as an indicator variable in our LPA.

Body mass index (BMI)—BMI was calculated as weight in kg divided by height in meters squared, using height and weight estimates generated from measured values at Stage 1. We measured weight with an electronic scale to the nearest 0.2 kg and height to the nearest millimeter with a portable stadiometer in light clothing without shoes. Weight estimates were corrected for clothing by subtracting 0.5 kg, and height estimates were rounded to the nearest centimeter.

Impairment and comorbid psychopathology—The Clinical Impairment Assessment (CIA; Bohn *et al.*, 2008) evaluates the extent to which eating disorder symptoms negatively impact mood, cognitive functioning, and relationships, with 4-point response options ranging from “not at all” to “a lot.” The CIA was adapted and administered to Stage 2 participants ($n = 215$) as a structured interview. Internal consistency was adequate in our sample (Becker *et al.*, 2009b). The Center for Epidemiologic Studies Depression Scale (CES-D) is a self-report measure of depressive symptoms (e.g., low mood, hopelessness) over the past seven days (Radloff, 1977) and was used to measure psychopathology that is often comorbid with disordered eating. The items, which are scored on a 4-point Likert scale ranging from < 1 day to ≥ 5 days, have been used successfully in non-Western adolescent populations (Ghubash *et al.*, 2000; Yang *et al.*, 2004). For this study, participants received scores as long as they had responded to at least 90% of CES-D items.

Cultural orientation—We assessed cultural orientation with composite measures derived from Likert items relating multiple dimensions of both Western/global and ethnic Fijian cultural orientation. Development, translation, and psychometric evaluation of these measures is described elsewhere (Becker *et al.*, in press). For this study, participants received scores as long as they had answered at least 80% of scale items.

Indigenous practices: kava and dranu use—As an additional proxy for traditional Fijian orientation, we queried the use of indigenous health and social practices. We evaluated use of *kava* (a ceremonial and social traditional beverage prepared from dried *Piper methysticum* root) in the past 30 days on a Likert scale ranging from 0 (no days) to 5 (everyday). We also assessed whether participants had used traditional herbal medicine to prevent and/or treat an indigenous illness, *macake* (characterized by loss of appetite; see Becker, 1995), in the past month.

Peer eating pathology—We evaluated the perceived prevalence of peer eating pathology in participants' social networks by asking them to estimate the proportion of "your five or so closest friends" who had engaged in dieting, fasting, and vomiting/laxative abuse in the past 30 days on a 4-point Likert scale ranging from "none of them" to "all of them." These items exhibited "moderate" test-retest reliability ($\kappa = .41$ for dieting; $\kappa = .48$ for fasting; $\kappa = .53$ for vomiting/laxative use) (Landis & Koch, 1977).

Latent Profile Analysis

LPA is a technique that utilizes maximum likelihood estimation to assign participants to mutually exclusive populations called latent classes. Classes are latent because membership is not directly observable, but instead can be inferred by evaluating the pattern of inter-correlations among indicator variables. The purpose of LPA is to identify the smallest number of classes that could account for these inter-correlations, thus minimizing the associations among indicator variables within each latent class and achieving a state of conditional independence. In contrast to LCA, which accepts only dichotomous indicator variables, LPA utilizes nominal, ordinal, or continuous indicators.

Consistent with previous LCA and LPA studies utilizing community samples (Sullivan *et al.*, 1998; Bulik *et al.*, 2000; Striegel-Moore *et al.*, 2005), we included only symptomatic participants in our LPA in order to enhance statistical power to identify multiple pathological groups. Because the base rate of AN symptoms was quite low in our sample (i.e., only 1% had a BMI < 17.5 kg/m² and 4% endorsed amenorrhea), we restricted our LPA to symptoms characterizing a diagnosis of BN. Therefore, we included only those participants who self-reported vomiting or herbal purgative use in the past 28 days via EDE-Q ($n = 222$). LPA indicator variables reflected the frequency of EDE-Q bulimic symptoms, with the addition of herbal purgative use. Because both symptom frequencies and Likert endorsement of our indicator variables were positively skewed, we transformed indicators into dichotomies or ordered categories based on natural breaks in the distributions. Our LPA indicator variables were as follows: (1) overvaluation of weight and shape ("not at all" [mean of EDE-Q overvaluation items < 1], "slightly" [mean of EDE-Q overvaluation items 1, < 3], "moderately" to "markedly" [mean of EDE-Q overvaluation items ≥ 3]), (2) binge eating (0, 1–3, 4 or more episodes), (3) vomiting (0 versus ≥ 1 episode), (4) laxative use (0 versus ≥ 1 episode), (5) fasting (0 versus ≥ 1 episode), (6) driven exercise (0, 1–4, 5 or more episodes), and (7) herbal purgative use (0 versus ≥ 1 episode).

We conducted our LPA using Latent Gold 4.5 (Vermunt & Magidson, 2005). We evaluated conditional independence by identifying the class solution associated with the lowest values

of the Bayesian information criterion (BIC) and the consistent Akaike information criterion (cAIC), as well as the minimization of cross-classification probabilities. We examined bivariate residuals (indices of the remaining correlations among indicator variables within latent classes) to ensure that none was greater than 5. Because no more than 2% of participants had missing values on any indicator variable, all 222 participants who endorsed self-induced vomiting or herbal purgative use in the past 28 days were submitted to LCA, and the remaining 301 participants served as a non-purging control group in subsequent validation analyses.

Validation Analyses

We hypothesized that, if purging is a valid marker of psychopathology among ethnic Fijian girls, then the latent classes would score significantly higher than non-purging participants with regard to (1) eating pathology (EDE-Q); (2) clinical impairment associated with disordered eating symptoms (CIA); and (3) depressive symptoms (CES-D). Furthermore, given the potential confounding effect of impoverished food environment on the cross-cultural assessment of eating pathology (le Grange *et al.*, 2004), we wanted to ensure that the classes did not differ from one another or from the non-purging group on the one-month prevalence of hunger due to lack of food in the home. To evaluate these hypotheses, we conducted a series of ANOVAs and χ^2 tests. To reduce Type I error, we set α to .01 for each omnibus test. To follow up statistically significant F tests, we utilized Fisher's LSD *post-hoc* comparisons. To follow up statistically significant omnibus χ^2 tests, we utilized a Bonferroni correction by dividing $\alpha = .05$ by the number of unique pair-wise comparisons. We conducted our validation analyses on SPSS version 16.0 (SPSS, 2007).

Post-hoc Class Comparisons of Potential Etiological Variables

A third aim of the study was to generate hypotheses about the social and cultural characteristics that could provide insights into the potentially distinct etiologies of symptomatic latent classes. Therefore, we used t -tests and χ^2 analyses to compare levels of cultural orientation, indigenous health practices, and peer eating pathology, across classes. We again set α to .01 to reduce Type I error, and conducted these *post-hoc* analyses on SPSS version 16.0 (SPSS, 2007).

Results

Latent Profile Analysis

We evaluated models with one to five classes. In order to meet the assumptions of the LPA model, we allowed for the conditional dependence of three pair-wise correlations of indicators in which bivariate residuals were greater than 5: (1) herbal purgative use/laxative use; driven exercise/overvaluation; and laxative use/overvaluation. BIC and cAIC were lowest for the two-class model with 21 parameters (model fit indices available upon request

from the first author)¹. Table 2 displays the relative endorsement of each indicator variable in the two latent classes, and Figure 1 provides a graphical depiction of these data.

The first class comprised 39% ($n = 86$) of the purging sample and was characterized primarily by high prevalence of self-induced vomiting (100%) and binge eating (90%). Members of this “BN-like class” also had moderate prevalence of herbal purgative use (58%) and laxative use (39%). In addition, the majority endorsed exercise for weight control (87%). Over half endorsed fasting (56%), and approximately one third affirmed that weight and shape figured “moderately” to “markedly” into their self-evaluation (35%).

The second class comprised 61% of the purging sample ($n = 136$). All members of this “herbal purgative class” endorsed herbal purgative use. Other forms of purging, such as vomiting (11%) and laxative use (10%), were infrequently endorsed. Participants defined by this class also endorsed binge eating (58%) and exercise for weight control (60%) less frequently than members of the BN-like class. However, the prevalences of fasting (47%) and “moderate” to “marked” overvaluation of shape and weight (43%) were comparable to those observed in the BN-like class. Notably, had we not added this culturally relevant item to the EDE-Q, 71% of participants in the herbal purgative class would have been classified as non-purging because of their negative responses to the items probing self-induced vomiting and laxative use.

Validation Analyses

Results of the validation analyses are presented in Table 3; the p -values presented below represent Fisher’s LSD *post-hoc* pair-wise comparisons, which we conducted to follow-up omnibus tests that already met our $\alpha = .01$ criterion for statistical significance.

Eating pathology—Participants in the BN-like and herbal purgative classes endorsed similar levels of eating pathology on the EDE-Q. As hypothesized, the BN-like and herbal purgative classes both endorsed significantly greater EDE-Q Global, Restraint, Eating Concern, Shape Concern, and Weight Concern scale scores than the non-purging group (all p ’s $< .05$). The BN-like class, however, endorsed significantly greater Eating Concern than the herbal purgative class ($p = .03$).

Impairment and comorbid psychopathology—Contrary to our expectations, the herbal purgative class endorsed significantly greater clinical impairment related to disordered eating than the BN-like class ($p = .03$), and the BN-like class did not differ from the non-purging group with regard to clinical impairment. In line with predictions, the herbal purgative class endorsed greater clinical impairment than the non-purging group ($p < .001$). As expected, both the BN-like ($p = .005$) and herbal purgative ($p = .01$) classes reported higher levels of comorbid depressive symptoms than the nonpurging group.

¹Due to our recruitment strategy, participants could be considered nested within school. Class membership was associated with school, $\chi^2(22) = 34.92$, $p = .04$, but the effect size was small (Cramer’s $V = .18$). We therefore re-ran our LPA with school as a covariate. Results were similar with regard to number of classes, proportion of participants in each class, and characteristics of each class, so we chose to present the final results of our LPA without controlling for school.

Demographics—The classes two did not differ from one another or from the non-purging group on demographic characteristics including age, BMI, peri-urban school location, relative material poverty, and hunger.

Post-hoc Class Comparisons of Potential Etiological Variables

No significant socio-demographic or cultural differences were identified between BN-like and herbal purgative classes, including cultural orientation, indigenous practices, or peer eating pathology (Table 4).

Discussion

To our knowledge, this study is the first LCA or LPA of eating disorder symptoms in a small-scale indigenous population and outside a high-income country. Our LPA of ethnic Fijian schoolgirls identified two latent classes associated with comparable levels of eating and general psychopathology, which we have characterized as “BN-like” and “herbal purgative” classes, respectively. The BN-like class, with its high rates of binge eating and vomiting, is phenomenologically similar to the bulimic classes described in previous North American and European LCAs and LPAs (Sullivan *et al.*, 1998; Bulik *et al.*, 2000; Keel *et al.*, 2004; Striegel-Moore *et al.*, 2005; Pinheiro *et al.*, 2008; Eddy *et al.*, 2009). In contrast, the herbal purgative class, which was characterized primarily by the use of indigenous herbal purgatives, has not been observed in previous LCAs and LPAs, nor is it represented as a distinct subtype in DSM-IV. Although the two classes were distinguished by self-reported purging modalities, they appeared to share core pathology by exhibiting remarkably similar profiles of eating and general psychopathology. Our results highlight the importance of including as LPA indicators locally meaningful behaviors that might be culture-specific, such as herbal purgative use, in order to enhance the generalizeability of empirically derived nosologic schemes.

Contrary to expectations, our *post-hoc* analyses of social and cultural characteristics did not find evidence that the herbal purgative class differed from the BN-like class on either Fijian or Western/global cultural orientation or perceived disordered eating among peers. However, our study was only sufficiently powered to detect medium to large effects. Moreover, the multiple dimensions of acculturation likely have heterogeneous impacts on health outcomes (Guarnaccia *et al.*, 2009; Becker *et al.*, in press) and the two classes may differ on a facet of cultural orientation that we did not assess. An alternative interpretation is that ethnic Fijian girls view the two types of purging as interchangeable. Indeed, given that herbal purgatives are used to induce vomiting or diarrhea, it is conceivable that there is more overlap in purging modality than the self-report responses suggest, reflecting a response bias related to the perception that herbal purgative use is more socially acceptable than other modes of purging. Relatedly, it is possible that within-class variation with respect to purging confounds the relation between socio-cultural variables and class membership. For example, the BN-like class contains respondents who also use herbal purgatives, and the herbal purgative class may include respondents who do not explicitly link their herbal purgative use to weight management, or who espouse plural rationales for herbal purgative use. Moreover, the herbal purgative class we identified may not be culturally unique; surveys of North

American eating disorder patients indicate that 13% (Trigazis *et al.*, 2004) to 64% (Steffen *et al.*, 2006) have used herbal products to induce weight loss or vomiting. Of these products, the ingestion of ipecac syrup to induce emesis in Western populations (Silber, 2005) bears the greatest phenomenological similarity to the herbal purgative use observed in Fiji. Indeed, because no previous eating disorder LCA or LPA has used ipecac or herbal product use as an indicator variable, we cannot be certain whether the emergence of an herbal purgative class in our ethnic Fijian sample reflects actual culture-specific phenomenologic heterogeneity, or our more inclusive operational definition of disordered eating behaviors.

The substantial prevalence of herbal purgative use in our sample suggests local vulnerability to disordered eating that could benefit from additional research attention and clinical resources. However, there are several additional implications of our findings, beyond their immediate public health relevance for the local populace. First, these results suggest that the application of a “universal” criteria set for eating disorders developed from a largely Euro-American evidence base may be insufficient to characterize the full range of phenomenologic heterogeneity across populations outside of these regions; indeed, a broader—and more culturally informed—scope of symptom phenomena should be considered (cf. Kleinman, 1977). Second, indigenous nosologic categories may not identify all symptom profiles as culturally salient, even if they are associated with distress and impairment. Third, so-called “unsupervised learning techniques” such as latent class analysis (Magidson & Vermunt, 2002; p. 38), in which group size and composition is not known *a priori*, may mitigate some of the limitations inherent to the application of solely *etic* or *emic* perspectives in cross-cultural evaluation by identifying novel symptom patterns. Fourth, this study provides further evidence that social environment may promote unique symptom presentations. It would be misleading, however, to interpret these results to mean that cultural context results only in superficial epiphenomenal variants on universal core pathology. Additional data are necessary to establish whether the herbal purgative class represents a clinical phenomenon with a course and outcome similar to or distinct from eating disorders described in high-income countries. For example, our preliminary qualitative data analysis suggests that characteristics of Fijian girls’ herbal purgative use—insofar as it is sometimes abetted by parents or temporally unrelated to overeating—may be distinctive from purging typical of BN.

Also of note, the BN-like class in our study endorsed EDE-Q scores substantially lower than those reported for BN-like classes identified in an American LPA (Eddy *et al.*, 2009) and European LCA (Pinheiro *et al.*, 2009), but comparable to the population median for American female undergraduates (Luce *et al.*, 2008). Notably, more than half of participants in both classes endorsed the use of traditional herbal therapy to prevent an indigenous illness—*macake*—which is characterized by poor appetite. The juxtaposition of a weight loss prevention strategy with purgative use suggests that weight and shape concerns may not be central to all eating disorder presentations in Fiji. Eating disorders resembling AN, but presenting in the absence of weight concerns, have been also been described in non-Western populations (Ong *et al.*, 1982; Lee *et al.*, 2001; Bennett *et al.*, 2004). Taken together, this cultural variation in the presentation of eating pathology supports the possibility that cultural

context may attenuate or exacerbate the cognitive symptoms associated with disordered eating.

Our findings should be interpreted in light of design strengths and limitations. With regard to strengths, first, we developed and adapted assessments to be culturally appropriate and in the local vernacular language, drawing from foundational ethnographic data and expert local knowledge. Second, we considered and excluded potential confounding by impoverished food environment (le Grange *et al.*, 2004). Third, the mean CIA score for the herbal purgative class approached the clinical cut-off of 16 that separated eating disorder cases from non-cases in the original validation sample (Bohn *et al.*, 2008), which supports our interpretation of herbal purgative as an eating disorder symptom.

A limitation is the use of primarily self-report data to identify and validate the latent classes. Moreover, our characterization of the heterogeneity and complexity of rationales for herbal purgative use in this sample is preliminary and incomplete. Indeed, in our evaluation of herbal purgative use via narrative data, a sizeable minority of respondents did not explicitly attribute their herbal purgative use to shape and weight. Therefore, it is likely that at least some of the herbal purgative use observed in the present study may have been motivated by alternative socially sanctioned indications (e.g., medicinal usage). On the other hand, it is also conceivable that individuals may not formulate herbal purgative use as motivated by weight concerns if they do not regard weight control as a socially legitimate pursuit. Relatedly, other investigators have commented on the validity of evaluating eating pathology with self-report versus interview assessment, highlighting the potentially greater candor evidenced in self-reports of socially stigmatized behaviors (Keel *et al.*, 2002; Mond *et al.*, 2007). A second limitation is that our selection of LPA indicators and validators—although informed by extensive ethnographic work with this population—necessarily reflect our Western conceptual models of disordered eating and related distress. A third limitation is that, because we did not observe the two respective classes longitudinally, we could not evaluate crossover between classes or the directionality of the relation between class membership and external validators. Lastly, LPA cannot distinguish differences in kind versus differences in degree. Indeed, there were more similarities than differences between the BN-like and herbal purgative classes, and future taxometric work is needed to ascertain the nature of the boundary between groups.

This study augments empirical support for cultural variation in patterning of eating disorder symptom presentation (Lee *et al.*, 2001; Franko *et al.*, 2007) and also has implications for clinical practice. For example, without appropriate breadth to identify clinically relevant variation, clinical and research assessment of eating pathology can result in misclassification of true cases as non-cases (Lee *et al.*, 2001; Bennett *et al.*, 2004) and non-cases as cases (le Grange *et al.*, 2004). Indeed, the majority of participants in the herbal purgative class would have been classified as non-purging by the EDE-Q had we not included an item referencing a culture-specific means of purging. In sum, the inclusion of populations with culturally distinctive traditions and symptoms in latent structure modeling may enhance the clinical utility of statistically derived nosological schemes in both diverse social contexts and multi-cultural populations.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement

We gratefully acknowledge the assistance of Dr. Lepani Waqatakirewa, CEO - Fiji Ministry of Health and his team; the Fiji Ministry of Education; Joana Rokomatu, the late *Tui Sigatoka*; Dr. Jan Pryor, Chair of the FN-RERC; Dr. Tevita Qorimasi; Professor Bill Aalbersberg; A. Nisha Khan; Asenaca Bainivualiku; Professor Jane Murphy; Lauren Richards; Aliyah Shivji; Amy Heberle; Kesaia Navara; and members of the Senior Advisory Group for the HEALTHY Fiji Study. Finally, we thank all the Fiji-based principals and teachers who facilitated this study.

This project was funded by NIMH K23 MH 068575 (AEB), a Harvard Research Enabling Grant (AEB), and a Klarman Foundation Post-Doctoral Fellowship (JJT).

References

- Becker, AE. Body, self, and society: The view from Fiji. Philadelphia: University of Pennsylvania Press; 1995.
- Becker AE, Gilman SE, Burwell RA. Changes in prevalence of overweight and in body image among Fijian women between 1989 and 1998. *Obesity Research*. 2005; 13:110–117. [PubMed: 15761169]
- Becker AE, Burwell RA, Gilman SE, Herzog DB, Hamburg P. Eating behaviours and attitudes following prolonged exposure to television among ethnic Fijian adolescent girls. *British Journal of Psychiatry*. 2002; 180:509–514. [PubMed: 12042229]
- Becker, AE.; Fay, K.; Agnew-Blais, J.; Guarnaccia, PM.; Striegel-Moore, RH.; Gilman, SE. Transcultural Psychiatry. Development of a measure of 'acculturation' for ethnic Fijians: Methodologic and conceptual considerations for application to eating disorders research. (in press).
- Becker AE, Thomas JJ, Bainivualiku A, Richards L, Navara K, Roberts AL, Gilman SE, Striegel-Moore RH. Adaptation and evaluation of the Clinical Impairment Assessment to assess disordered eating related distress in an adolescent female ethnic Fijian population. *International Journal of Eating Disorders*. 2009b Published online: 23 March 2009.
- Becker AE, Thomas JJ, Bainivualiku A, Richards L, Navara K, Roberts AL, Gilman SE, Striegel-Moore RH. Validity and reliability of a Fijian translation and adaptation of the Eating Disorder Examination Questionnaire. *International Journal of Eating Disorders*. 2009c Published online: 23 March 2009.
- Becker AE, Thomas JJ, Pike KM. Should non-fat-phobic anorexia nervosa be included in DSM-V? *International Journal of Eating Disorders*. 2009a Published Online: 4 August 2009.
- Bennett D, Sharpe M, Freeman C, Carson A. Anorexia nervosa among female secondary school students in Ghana. *British Journal of Psychiatry*. 2004; 185:312–317. [PubMed: 15458991]
- Bohn K, Doll HA, Cooper Z, O'Connor M, Palmer RL, Fairburn CG. The measurement of impairment due to eating disorder psychopathology. *Behaviour Research and Therapy*. 2008; 46:1105–1110. [PubMed: 18710699]
- Bulik CM, Sullivan PF, Kendler KS. An empirical study of the classification of eating disorders. *American Journal of Psychiatry*. 2000; 157:886–895. [PubMed: 10831467]
- Duncan AE, Bucholz KK, Neuman RJ, Agrawal A, Madden PAF, Heath AC. Clustering of eating disorder symptoms in a general population twin sample: A latent class analysis. *Psychological Medicine*. 2007; 37:1097–1107. [PubMed: 17472759]
- Eddy KT, Crosby RD, Keel PK, Wonderlich SA, le Grange D, Hill L, Powers P, Mitchell JE. Empirical identification and validation of eating disorder phenotypes in a multisite clinical sample. *Journal of Nervous and Mental Disease*. 2009; 197:41–49. [PubMed: 19155809]
- Fairburn CG, Beglin SJ. The assessment of eating disorders: Interview or self-report questionnaire? *International Journal of Eating Disorders*. 1994; 16:363–370. [PubMed: 7866415]
- Franko DL, Becker AE, Thomas JJ, Herzog DB. Cross-ethnic differences in eating disorder symptoms and related distress. *International Journal of Eating Disorders*. 2007; 40:156–164. [PubMed: 17080449]

- Guarnaccia PJ, Pincay IM, Alegria M, Shrout PE, Lewis-Fernandez R, Canino GJ. Assessing diversity among Latinos: Results from the NLAAS. *Hispanic Journal of Behavioral Sciences*. 2007; 29:510–534. [PubMed: 19672330]
- Ghubash R, Daradkeh TK, Al Naseri KS, Al Bloushi NBA, Al Daher AM. The performance of the Center for Epidemiologic Study Depression Scale (CES-D) in an Arab female community. *International Journal of Social Psychiatry*. 2000; 46:241–249. [PubMed: 11201346]
- Hoek HW, Van Harten PN, Hermans KME, Katzman MA, Matroos GE, Susser ES. The incidence of anorexia nervosa on Curacao. *American Journal of Psychiatry*. 2005; 162:748–752. [PubMed: 15800148]
- Hughes RG, Lawrence MA. Globalisation, food and health in Pacific Island countries. *Asia Pacific Journal of Clinical Nutrition*. 2005; 14:298–306. [PubMed: 16326635]
- Keel PK, Crow S, Davis TL, Mitchell JE. Assessment of eating disorders: Comparison of interview and questionnaire data from a long-term follow-up study of bulimia nervosa. *Journal of Psychosomatic Research*. 2002; 53:1043–1047. [PubMed: 12445594]
- Keel PK, Fichter M, Quadflieg N, Bulik CM, Baxter MG, Thornton L, Halmi KA, Kaplan AS, Strober M, Woodside DB, Crow SJ, Mitchell JE, Rotondo A, Mauri M, Cassano G, Treasure J, Goldman D, Berrettini WH, Kaye WH. Application of a latent class analysis to empirically define eating disorder phenotypes. *Archives of General Psychiatry*. 2004; 61:192–200. [PubMed: 14757596]
- Kleinman AM. Depression, somatization, and the “new cross-cultural psychiatry”. *Social Science & Medicine*. 1977; 11:3–10. [PubMed: 887955]
- Landis J, Koch G. The measurement of observer agreement for categorical data. *Biometrics*. 1977; 33:159–174. [PubMed: 843571]
- le Grange D, Louw J, Breen A, Katzman MA. The meaning of ‘self-starvation’ in impoverished black adolescents in South Africa. *Culture, Medicine, and Psychiatry*. 2004; 28:439–461.
- Lee S, Lee AM, Ngai E, Lee DT, Wing YK. Rationales for food refusal in Chinese patients with anorexia nervosa. *International Journal of Eating Disorders*. 2001; 29:224–229. [PubMed: 11429985]
- Luce KH, Crowther JH, Pole M. Eating Disorder Examination Questionnaire (EDE-Q) norms for undergraduate women. *International Journal of Eating Disorders*. 2008; 41:273–276. [PubMed: 18213686]
- Magidson J, Vermunt JK. Latent class models for clustering: A comparison with K-means. *Canadian Journal of Marketing Research*. 2002; 20:37–44.
- Mavoa HM, McCabe M. Sociocultural factors relating to Tongans’ and Indigenous Fijians’ patterns of eating, physical activity and body size. *Asia Pacific Journal of Clinical Nutrition*. 2008; 17:375–384. [PubMed: 18818156]
- McCabe MP, Ricciardelli L, Waqa G, Goundar R, Fotu K. Body image and body change strategies among adolescent males and females from Fiji, Tonga and Australia. *Body Image*. 2009 Published online: 8 July 2009.
- Mitchell JE, Crosby RD, Wonderlich SA, Hill L, le Grange D, Powers P, Eddy KT. Latent profile analysis of a cohort of patients with eating disorders not otherwise specified. *International Journal of Eating Disorders*. 2007; 40:S95–S98. [PubMed: 17886266]
- Mond JM, Hay PJ, Rodgers B, Owen C. Self-report versus interview assessment of purging in a community sample of women. *European Eating Disorders Review*. 2007; 15:403–409. [PubMed: 17960776]
- Njenga FG, Kangethe RN. Anorexia nervosa in Kenya. *East African Medical Journal*. 2004; 81:188–193. [PubMed: 15884284]
- Ong YL, Tsoi WF, Chea JS. A clinical and psychosocial study of seven cases of anorexia nervosa in Singapore. *Singapore Medical Journal*. 1982; 23:255–261. [PubMed: 7157004]
- Patel V, Sumathipala A. International representation in psychiatric literature: Survey of six leading journals. *British Journal of Psychiatry*. 2001; 178:406–409. [PubMed: 11331553]
- Pinheiro AP, Bulik CM, Sullivan PF, Machado PPP. An empirical study of the typology of bulimic symptoms in young Portuguese women. *International Journal of Eating Disorders*. 2008; 41:251–258. [PubMed: 18095310]

- Radloff LS. The CES-D scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*. 1977; 1:385–401.
- Silber TJ. Ipecac syrup abuse, morbidity, and mortality: Isn't it time to repeal its over-the-counter status? *Journal of Adolescent Health*. 2005; 37:256–260. [PubMed: 16109351]
- SPSS. Statistical Package for the Social Sciences, 16.0 for Macintosh. Chicago, IL: SPSS, Inc.; 2007.
- Steffen KJ, Roerig JL, Mitchell JE, Crosby RD. A survey of herbal and alternative medication use among participants with eating disorder symptoms. *International Journal of Eating Disorders*. 2006; 39:741–746. [PubMed: 16941628]
- Striegel-Moore RH, Franko DL, Thompson D, Barton B, Schreiber GB, Daniels SR. An empirical study of the typology of bulimia nervosa and its spectrum variants. *Psychological Medicine*. 2005; 35:1563–1572. [PubMed: 16219114]
- Striegel-Moore RH, Franko DL, Thompson D, Affenito S, May A, Kraemer HC. Exploring the typology of night eating syndrome. *International Journal of Eating Disorders*. 2008; 41:411–418. [PubMed: 18306340]
- Sullivan PF, Bulik CM, Kendler KS. The epidemiology and classification of bulimia nervosa. *Psychological Medicine*. 1998; 28:599–610. [PubMed: 9626716]
- Thomas JJ, Vartanian LR, Brownell KD. The relationship between eating disorder not otherwise specified (EDNOS) and officially recognized eating disorders: Meta-analysis and implications for DSM. *Psychological Bulletin*. 2009; 135:407–433. [PubMed: 19379023]
- Trigasiz L, Tennankore D, Vohra S, Katzman DK. The use of herbal remedies by adolescents with eating disorders. *International Journal of Eating Disorders*. 2004; 35:223–228. [PubMed: 14994361]
- Vermunt, JK.; Magidson, J. *Latent Gold 4.5*. Belmont, MA: Statistical Innovations; 2005.
- Wade TD, Crosby RD, Martin NG. Use of latent profile analysis to identify eating disorder phenotypes in an adult Australian twin cohort. *Archives of General Psychiatry*. 2006; 63:1377–1384. [PubMed: 17146012]
- World Health Organization. *Caring for children and adolescents with mental disorders: Setting WHO directions*. Geneva, Switzerland: World Health Organization; 2003.
- Yang H-J, Soong W-T, Kuo P-H, Chang H-L, Chen W-J. Using the CES-D in a two-phase survey for depressive disorders among nonreferred adolescents in Taipei: A stratum-specific likelihood ratio analysis. *Journal of Affective Disorders*. 2004; 82:419–430. [PubMed: 15555693]

Figure 1.

Relative endorsement of eating disorder symptom indicators in the BN-like class (represented by the solid line) and herbal purgative class (represented by the dashed line)
Note. For dichotomous variables (vomiting, laxative use, fasting, and herbal purgative use), relative endorsement is plotted as the percentage of participants in each class who endorsed that symptom. For ordinal variables (overvaluation, binge eating, and driven exercise), relative endorsement is plotted as the mean of the three ordered categories (assigned point values of 0, 1, and 2 for increasingly greater frequencies) divided by two, so as to fit on a 0–1 scale.

Excerpted EDE interview transcript data illustrating a rationale for herbal purgative use to compensate for episodes of overeating and/or control shape or weight among six study participants who self-reported herbal purgative use in the past 28 days

Table 1

Case number	Frequency of herbal purgative use in past 28 days reported on the EDE-Q	Peri-urban school	Age	BMI	EDE-Q global score	Excerpts from participant responses to the initial EDE probe item: "Over the past four weeks, have you taken <i>dramu</i> as a means of controlling your shape or weight?"
1	1	No	18	24.14	2.48	"Yes, madam. [INT] [...] It's [called] <i>kura</i> . [INT] [...] [It causes] vomiting [INT] [...] and sometimes I get diarrhea. [INT] [...] It's, uh, Fijian medicine [...] [My parents] think that, uh, inside their body can come out and have the body to lose weight. [INT]: And they want you to lose weight?" Yes, ma'am."
2	1	Yes	18	28.12	4.19	"Yes, I tried to vomit. [INT] I drank to my mom's, like, herbal medicine I don't know for a cough, and I just tried it to take care of things, and I just drank it and then I started feeling sick and started vomiting. [INT]: Did you do it because you wanted to get rid of the food?" Yes. [...] [INT]: Did you do it after you felt like you had eaten too much, or did you just do it?" No, after I ate too much."
3	2	Yes	16	23.93	3.08	"Yes, ma'am. [INT] [...] Fijian <i>dramu</i> . [INT] [...] My mother gave it to me [INT] [...] so it, um, because my stomach, to clean it out. [INT]: Why did you feel like you needed to clean your stomach?" Because I had a lot to, I ate a lot. [INT]: So you ate a lot?" Yes ma'am. [INT]: And you wanted to get rid of it? But not to change your shape or weight?" Yes. [INT]: Or to stay the same?" Yes ma'am. [INT]: Was it to stay the same shape and weight?" Yes ma'am."
4	5	Yes	17	22.11	3.36	"Yes. [INT] A Fijian one. [INT] It was in liquid form [INT] from a plant. [INT] We usually drink it as a whole family [INT] [...] to cut down on my eating habits and to lose weight. [INT] [...] It cleans out my stomach and makes me go to the loo most of the time. [INT]: Did you do this after you felt you had eaten too much, like those times we talked about, or did you do it when you weren't eating too much?" When I was eating too much [INT] [...] so that all the unwanted stuff could come out and that you could lose weight, be fit and all."
5	5	No	17	22.77	2.28	"Yeah. [INT] It was, uh, it was before, when my father told me that I was like overweight. [...] I was, like, um, very, um, not satisfied with my weight, so I tried Fijian medicine [INT] [...] I took it and I tried to like make me sick so that I could, uh, vomit and like [...] [have] diarrhea [...] so that it could wash out. [...] [INT]: This was to control your shape and weight?" Shape and weight. [INT]: Because your father told you that you were too big?" Yeah, that's why I took it."

Case number	Frequency of herbal purgative use in past 28 days reported on the EDE-Q	Peri-urban school	Age	BMI	EDE-Q global score	Excerpts from participant responses to the initial EDE probe item: “Over the past four weeks, have you taken <i>dranu</i> as a means of controlling your shape or weight?”
6	28	Yes	18	24.82	3.24	“Yes. [INT] [...] It was a liquid. [INT] My mother [gave it to me]. [INT] Um, she says it’s for girls [INT] to clean the stomach from, uh, everything. [INT] [...] It helps. [INT] It makes me go to the, use the toilet. [INT: And did you do this to lose weight, too? Did you use it to control your weight?] Yes. [INT] Sometimes I take it, like, every day. [INT] [...] Just for keeping safe and, uh, control something, control anything that can happen. [...] [INT: Do you also do it to lose weight?] Yes. [INT: To be slim?] Uhhmm.”

Note. BMI = Body mass index; EDE-Q = Eating Disorder Examination Questionnaire; *Dranu* = a generic term for indigenous Fijian medicinal herbal preparations; *kuru* = a specific Fijian herbal preparation; [INT] = Interviewer question omitted; [...] = Participant comment omitted.

Table 2

Prevalence of eating disorder symptom endorsement in the LPA-derived BN-like class and herbal purgative class

	BN-like (n = 86)	Herbal Purgative (n = 136)
Overvaluation of weight and shape		
0 ("Not at all"/Mean of EDE-Q overvaluation items <1)	8 (9%)	21 (15%)
1 ("Slightly"/Mean of EDE-Q overvaluation items = 1, <3)	48 (56%)	57 (42%)
2 ("Moderately" to "Markedly"/Mean of EDE-Q overvaluation items ≥ 3)	30 (35%)	58 (43%)
Binge eating		
0 (None)	9 (10%)	57 (42%)
1 (1–3 episodes)	60 (70%)	62 (46%)
2 (≥ 4 episodes)	17 (20%)	16 (12%)
Self-induced vomiting		
0 (None)	0 (0%)	119 (89%)
1 (≥ 1 episode)	86 (100%)	15 (11%)
Laxative use		
0 (None)	51 (61%)	121 (90%)
1 (≥ 1 episode)	33 (39%)	13 (10%)
Fasting		
0 (None)	38 (44%)	72 (53%)
1 (≥ 1 episode)	48 (56%)	64 (47%)
Driven exercise		
0 (None)	11 (13%)	53 (40%)
1 (1–4 episodes)	51 (59%)	51 (39%)
2 (≥ 5 episodes)	24 (28%)	28 (21%)
Herbal purgative use		
0 (None)	36 (42%)	0 (0%)
1 (≥ 1 episode)	50 (58%)	136 (100%)

Note. Within each cell, the bolded percentage indicates the modal response for that variable within that latent class. Numbers in some cells do not add up to the total number of participants in that latent class due to missing data on that variable.

Table 3

Comparison of external validators in the LPA-derived BN-like and herbal purgative classes versus the non-purgating group

<i>M (SD) or n (%)</i>						
	BN-like (<i>n</i> = 86)	Herbal Purgative (<i>n</i> = 136)	Non-Purging (<i>n</i> = 301)	<i>F(df)</i> or $\chi^2(df)$	<i>p</i>	Effect size
Eating Pathology						
EDE-Q Global	2.08 (0.93) ^a	1.90 (1.11) ^a	1.49 (1.05) ^b	<i>F</i> (2, 519)=13.60	<.001	$\eta^2_p=.050$
EDE-Q Restraint	1.28 (1.09) ^a	1.20 (1.24) ^a	0.71 (0.88) ^b	<i>F</i> (2, 519)=16.39	<.001	$\eta^2_p=.059$
EDE-Q Eating Concern	1.84 (1.01) ^a	1.51 (1.20) ^b	1.28 (1.09) ^c	<i>F</i> (2, 519)=9.14	<.001	$\eta^2_p=.034$
EDE-Q Shape Concern	2.62 (1.28) ^a	2.50 (1.45) ^a	2.09 (1.47) ^b	<i>F</i> (2, 520)=6.77	.001	$\eta^2_p=.025$
EDE-Q Weight Concern	2.55 (1.33) ^a	2.39 (1.39) ^a	1.90 (1.42) ^b	<i>F</i> (2, 520)=10.18	<.001	$\eta^2_p=.038$
Impairment and Comorbid Psychopathology						
CIA θ	11.47 (10.43) ^a	15.39 (11.57) ^b	8.69 (10.15) ^a	<i>F</i> (2, 212)=6.34	.002	$\eta^2_p=.056$
CES-D	21.48 (7.14) ^a	20.85 (8.41) ^a	18.85 (7.25) ^b	<i>F</i> (2, 514)=5.76	.003	$\eta^2_p=.022$
Demographics						
Age	16.70 (1.14)	16.74 (1.14)	16.65 (1.06)	<i>F</i> (2, 520)=0.33	<i>ns</i>	$\eta^2_p=.001$
BMI	24.03 (3.12)	24.11 (3.29)	23.88 (3.45)	<i>F</i> (2, 520)=0.23	<i>ns</i>	$\eta^2_p=.001$
Peri-urban school location	38 (44%)	62 (46%)	161 (54%)	$\chi^2(2)=3.69$	<i>ns</i>	<i>V</i> =.08
Relative material poverty	54 (63%)	79 (58%)	174 (58%)	$\chi^2(2)=0.71$	<i>ns</i>	<i>V</i> =.04
Hunger due to insufficient food (past 30 days)	41 (48%)	55 (41%)	120 (40%)	$\chi^2(2)=1.71$	<i>ns</i>	<i>V</i> =.06

Note. Within each row, means and percentages with different superscripts (*a*, *b*, *c*) differ significantly from one another (utilizing Fisher's LSD *post-hoc* comparisons to follow up significant omnibus *F* tests, and Bonferroni corrections to follow up significant omnibus χ^2 tests). EDE-Q = Eating Disorder Examination Questionnaire; CIA = Clinical Impairment Assessment; CES-D = Center for Epidemiologic Studies Depression Scale; BMI = Body mass index; η^2_p =partial eta squared; *V* = Cramer's *V*.

θ Data available for Stage 2 participants only (*n* = 215).

Table 4
Class comparisons of selected potential etiological variables in the LPA-derived BN-like and herbal purgative classes

	BN-like (<i>n</i> = 86)	Herbal Purgative (<i>n</i> = 136)	<i>t</i> (<i>df</i>) or χ^2 (<i>df</i>)	<i>p</i>	Effect size
Cultural Orientation					
Global/Western orientation scale	4.54 (0.90)	4.44 (0.78)	<i>t</i> (220)=0.88	<i>ns</i>	<i>d</i> =.12
Ethnic Fijian orientation scale	4.99 (1.48)	4.74 (1.58)	<i>t</i> (220)=1.04	<i>ns</i>	<i>d</i> =.16
Indigenous Health Practices					
<i>Kava</i> use (Likert frequency in past month)	1.30 (0.51)	1.45 (0.62)	<i>t</i> (220)=1.91	<i>ns</i>	<i>d</i> =.25
Indigenous herbal medicine use to prevent <i>macake</i> (yes in past month)	44 (56%)	73 (60%)	χ^2 (1)=0.34	<i>ns</i>	<i>V</i> =.04
Peer Eating Pathology: Close Friends					
Dieting	1.99 (0.75)	1.93 (0.85)	<i>t</i> (219)=0.49	<i>ns</i>	<i>d</i> =.07
Fasting	1.74 (0.83)	1.63 (0.77)	<i>t</i> (219)=1.05	<i>ns</i>	<i>d</i> =.15
Vomiting/laxative use	1.48 (0.71)	1.32 (0.62)	<i>t</i> (219)=1.69	<i>ns</i>	<i>d</i> =.24

Note. *Macake* = indigenous illness characterized by appetite loss. *d* = Cohen's *d*; *V* = Cramer's *V*.