Letter to the Editor

Quantitative Serial T2 Relaxometry and Cerebral Cysticercosis

BEUY JOOB¹, VIROJ WIWANITKIT²

¹Sanitation 1 Medical Academic Center; Bangkok Thailand

²Hainan Medical University; China - Faculty of Medicine, University of Nis; Serbia - Joseph Ayobabalola University; Nigeria - Dr DY Patil Medical College; India

Key words: quantitative serial T2 relaxometry, cerebral cysticercosis

Sir,

the report on “Quantitative serial T2 relaxometry and cerebral cysticercosis” is very interesting¹. Nalini et al. described possible values of quantitative serial T2 relaxometry and cerebral cysticercosis on serial MRI in patients with a solitary cerebral cysticercal lesion¹. This observation might imply a role for quantitative serial T2 relaxometry in cerebral cysticercosis. However, there are many considerations

in this report. First, without a control group, it might not be possible to mention that the observed serial change is due to the response to the antiparasitic drug. Second, quantitative serial T2 relaxometry is not routinely used and available.

Routinely, the follow-up of a case is simple neuroimaging with the need to use quantitative serial T2 relaxometry². The use of radiation-free MRI might be more appropriate for follow-up³.

References

- 1 Nalini A, de Souza A, Saini J, et al. Quantitative serial T2 relaxometry: a prospective evaluation in solitary cerebral cysticercosis. *Neuroradiol J.* 2014; 27 (3): 339-349. doi: 10.15274/NRJ-2014-10054
- 2 Nogales-Gaete J, Arias-Quezada P, Jurado-Díaz F. Neurocysticercosis: five years follow-up of 21 patients. *Med Chil.* 2013; 141 (10): 1353-1356. doi: 10.4067/S0034-98872013001000020.

- 3 Verma R, Lalla R. Why MRI of brain is superior to CT in multiple neurocysticercosis? *BMJ Case Rep.* 2012; 2012. doi: 10.1136/bcr.03.2012.6046.

Beuy Joob
Sanitation 1 Medical Academic Center - Bangkok Thailand
E-mail: beuyjoob@hotmail.com