Specific Inactivation of Herpes Simplex Virus by Silver Nitrate at Low Concentrations and Biological Activities of the Inactivated Virus

FUMIO SHIMIZU,* YOSHINOBU SHIMIZU, AND KATSUO KUMAGAI

Department of Microbiology, Tohoku University School of Dentistry, Sendai, Japan

Received for publication 29 January 1976

The infectivities of herpes simplex virus types 1 and 2 were inactivated by silver nitrate at concentrations of 30 μ M or less, which did not affect at all the infectivities of hemagglutinating virus of Japan, vesicular stomatitis virus, poliovirus, vaccinia virus, and adenovirus. The inactivated virus retained the capability of adsorbing to the cell, with an adsorption kinetics quite similar to that of intact virus, and of inducing the concanavalin A agglutinability in the infected cells, whereas it lost completely the capability of producing viral antigens and other cytopathic changes.

Salts of heavy metal ions such as silver, mercury, or copper are bactericidal at very low concentrations, under which conditions they act by combining with sulfhydryl groups. Such sulfhydryl reagents have also been shown to inactivate the infectivity of many viruses. This reaction has been used by many previous investigators to obtain information on the superficial structure of viruses and the early stages of virus multiplication (1, 2, 4, 7).

It has been reported that the infectivity of herpes simplex virus (HSV) is almost completely diminished when the virus is treated with the compounds at concentrations as low as 10^{-4} to 10^{-5} M (7, 10). Thus, in vitro and in vivo effects of compounds such as silver nitrate (5) or silver sulfadiazine (3) on HSV infection have recently been investigated to evaluate them as local chemotherapeutic agents.

The present communication is concerned with further studies on the in vitro effect of silver nitrate on the infectivity of HSV. The inactivation kinetics of HSV by the compound and the specific inactivation of HSV at low concentrations of the compound, which did not affect the infectivity of other viruses, are described. It is also shown that the inactivated virus retained the capability to interact with the cellular membranes, although they did not undergo a growth cycle.

MATERIALS AND METHODS

Viruses. Two types of HSV (type 1, Miyama strain; type 2, UW 268 strain), vaccinia virus (IHD strain), adenovirus (type 5), hemagglutinating virus of Japan (HVJ; Fushimi strain), vesicular stomatitis virus (Indiana strain), and poliovirus (type 1, Mahoney strain) were used. Both types of HSV were

propagated in African green monkey kidney cells and centrifuged at $1,400 \times g$ for 15 min. Titers of the stock HSV-1 and -2 were 2×10^7 and 10^7 plaque-forming units/ml, respectively. Partially purified HSV-1 was prepared by centrifugation of the crude virus onto a cushion of 60% (wt/wt) sucrose in tris(hydroxymethyl)aminomethane (Tris) - hydrochloride buffer through a layer of 5% sucrose in Tris-hydrochloride at $64,000 \times g$ for 90 min at 4 C. The viral preparation at the interface between 5 and 60% sucrose layers was rinsed at once and resuspended in the buffer. Adenovirus was propagated in HEP-2 cells, and the titer was $10^{6.8}$ 50% tissue culture infectious doses (TCID₅₀)/ml. Vesicular stomatitis virus and vaccinia virus were propagated in rabbit kidney cells (RK13), and their titers were 10^7 and $10^{7.5}$ TCID₅₀/ml, respectively. Poliovirus was prepared in human embryonic lung cells, and the titer was $10^{7.2}$ TCID₅₀/ml. HVJ was prepared in the chorioallantoic cavity of 11-day-old embryonated chicken eggs, and the titer was $10^{9.6}$ 50% egg infectious doses/ml. All the viruses were stored at -80°C until used.

Tissue culture cells. Monolayer cultures of chicken embryo, RK13, African green monkey kidney, HEP-2, and human embryonic lung cells were grown in Eagle minimum essential medium (MEM) supplemented with 10% heat-inactivated bovine serum, penicillin (100 U/ml), and streptomycin (100 μ g/ml). The growth medium for monolayer cultures of 3T3 mouse cells was MEM containing 10% calf serum, penicillin, and streptomycin.

Infectivity assay. All the viruses other than HVJ were assayed by counting plaque-forming units or TCID₅₀ per milliliter, using African green monkey kidney or human embryonic lung cells. HVJ was assayed by hemagglutinin titration.

Reagents and radioisotopes. The reagents used were obtained from the following sources. Silver nitrate (AgNO₃), silver iodide (AgI), and reduced glutathione (R-glutathione) were from Wako Chemical Industries, Tokyo. Potassium iodide (KI) was

from Junsei Chemicals, Tokyo. AgNO_3 was dissolved in distilled water at 100 $\mu\text{g/ml}$, and lower concentrations were prepared by diluting the solution with MEM. KI and AgI were diluted with MEM to the desired concentrations. R-glutathione was dissolved in Tris buffer, and the pH was adjusted to 7.4. [^3H]thymidine (specific activity, 10 Ci/mmol) was obtained from New England Nuclear Corp.

Inactivation of viruses by silver nitrate. In most experiments, inactivation of viruses with AgNO_3 was performed by adding 1.0 ml of 2×10^6 plaque-forming units or 2×10^5 50% egg infectious doses (HVJ) of viruses per ml to 1.0 ml of a 60 μM AgNO_3 solution and agitating the mixture at 37 C for varying times. For controls, 1.0 ml of phosphate-buffered saline (PBS), pH 7.2, was added to the virus suspension before agitation at 37 C. At intervals, serial dilutions into MEM were made, and their virus titers were assayed by counting plaque-forming units or TCID_{50} .

Preparation of labeled HSV. Monolayer cultures of 2×10^7 African green monkey kidney cells were infected with HSV-1 at an input multiplicity of 6. After a 2.5-h adsorption period at 37 C, cell sheets were washed three times with PBS to remove the unadsorbed virus and flooded with 30 ml of fresh medium. After 4 h of incubation, [^3H]thymidine in MEM was then added to monolayer cultures to a concentration of 4 $\mu\text{Ci/ml}$. The cultures were incubated at 37 C until the cytopathic effect became evident (about 24 h). After centrifugation of the culture medium at $4,000 \times g$ at 4 C for 15 min to remove cell debris, the supernatant fluid was harvested. The virus was purified by the cushion ultracentrifugation method in a sucrose gradient (20 to 70%), and 1.0-ml fractions of virus were collected from the top to the bottom. The fractions were assayed for infectivity and radioactivity.

ConA agglutination tests. The methods used were essentially those of Inbar and Sacks (6). For agglutination assays, chicken embryo cells were infected with HSV inactivated by 30 μM AgNO_3 or with noninactivated HSV. After an adsorption period of 90 min at 37 C, the cell sheets were washed three times with culture medium to remove the unadsorbed virus and flooded with 6 ml of fresh medium. Six hours after infection, the cell sheets were washed with Ca- and Mg-free PBS, prewarmed to 37 C, and then brought into single cell suspensions with 0.02% disodium versenate solution (ethylenediaminetetraacetate). The cells were then washed with saline and suspended with PBS at a concentration of 10^6 cells/ml, and 0.3-ml samples of cell suspension were transferred to 15-mm plastic trays to which was added either 0.3 ml of concanavalin A (ConA) or 0.3 ml of saline. The mixture was rocked gently several times by hand at room temperature and then examined for cell agglutination under an inverted microscope after 45 min.

Tests for hemadsorption of sensitized erythrocytes. Sensitization of sheep erythrocytes with rabbit anti-sheep erythrocyte serum and hemadsorption on cover slip cultures were carried out as described by Watkins (12). Briefly, a 1% suspension of erythrocytes was mixed with an equal volume of anti-

erythrocyte serum (1 hemagglutination unit), and the mixture was incubated at 37 C for 30 min. The erythrocytes were washed three times with glucose veronal buffer and used as a 0.5% suspension for hemadsorption tests. After removing the culture medium, the cover slip cultures, in petri dishes, were gently rinsed three times with glucose veronal buffer. Two milliliters of a suspension of sensitized erythrocytes was then added. After incubation for 2 h at room temperature, the cover slip cultures were gently washed two times with glucose veronal buffer to remove nonadsorbed erythrocytes, and hemadsorption was then examined microscopically.

Fluorescent staining. The antibodies to HSV used were rabbit antisera from rabbits immunized with partially purified virus preparations of HSV-1, Miyama strain. The fluorescein isothiocyanate-conjugated sera used were sheep anti-rabbit gamma globulin (Behringwerke, Uppsala, Sweden).

Production of HSV antigens in the cells was detected by the indirect method. Smear preparations of infected cells fixed with acetone were first covered with an appropriate dilution of antiserum and incubated for 60 min in a moist chamber at 37 C. At this time the preparations were removed, washed with PBS, covered with fluorescein isothiocyanate-conjugated serum, and then reincubated at 37 C for 60 min.

To detect HSV-induced surface antigens, infected monolayer cultures were washed three times with glucose veronal buffer and incubated with the antiserum for 30 min at 4 C. The antiserum was removed by washing, and the cells were stained with a 1:8 dilution of fluorescein isothiocyanate-conjugated serum.

RESULTS

Kinetics of inactivation of HSV by AgNO_3 . The first experiment was performed to investigate the rate of inactivation of HSV-1 during exposure to AgNO_3 . A total of 10^5 TCID_{50} of HSV per ml was treated with 30 μM AgNO_3 for up to 120 min at 37 C in a water bath and tested for the titers of residual virus. HSV was rapidly inactivated with time (Fig. 1). After 60 min, the virus was almost completely inactivated.

To investigate whether the inactivation of HSV by AgNO_3 was dependent upon the amounts of test virus, we exposed different infectivities of HSV to 30 μM AgNO_3 for 60 min at 37 C and tested for the titers of residual virus (Fig. 2). When the unpurified virus preparation was used as a test virus, less than 10^6 TCID_{50} of HSV per ml was completely inactivated upon exposure to the compound, whereas 10^7 TCID_{50} of the virus per ml lost no virus titers. Although the inactivation by AgNO_3 of unpurified HSV thus appeared to be dependent on the virus titers, partially purified virus at 10^7 $\text{TCID}_{50}/\text{ml}$ was completely inactivated by the same concentration of the compound (Fig. 2). The protein concentrations of unpurified and purified prep-

FIG. 1. Effect of exposure time on inactivation of HSV-1 by silver nitrate.

arations, both of which contained 10^7 TCID₅₀/ml, were 2.7 and 0.2 mg/ml, respectively, as determined by the method of Lowry et al. (8). These results, therefore, suggest that lack of inactivation by AgNO_3 of the unpurified virus at 10^7 TCID₅₀/ml may be due to the presence of the proteins at concentrations that would inhibit the effect of AgNO_3 on the virus. When the purified virus at 10^7 TCID₅₀/ml, after addition of bovine serum albumin at a final concentration of 0.8%, was tested for inactivation by AgNO_3 , no inactivation was observed (Fig. 2). Thus, it was concluded that AgNO_3 at $30 \mu\text{M}$ can inactivate HSV independently of the virus titers, at least up to 10^7 TCID₅₀/ml.

Effect of AgNO_3 on different viruses. After studying the inactivation kinetics of HSV-1 by AgNO_3 , the compound was tested for its effect on different viruses, including HSV-2. A total of 10^5 TCID₅₀ or 10^5 50% egg infectious doses (HVJ) of viruses per ml were treated with AgNO_3 at different concentrations (less than $30 \mu\text{M}$) for 60 min at 37°C and then tested for the titers of residual virus. There was a marked contrast between the results for HSV (types 1 and 2) and those for the other viruses (Fig. 3). Both HSV-1 and -2 underwent at least a 10^4 -fold loss of titer when treated with 15 or $30 \mu\text{M}$ AgNO_3 . By contrast, the other viruses, irrespective of whether they were enveloped or non-enveloped and deoxyribonucleic acid or ri-

bonucleic acid viruses, were not reduced in titer by treatment with the same concentrations of this reagent.

The protein concentrations contained in the test preparations were calculated for each virus by the estimation of the concentrations of undiluted samples and were found to be as low as 27, 28, 0.2, 70, 97, 83, and $140 \mu\text{g/ml}$ for HSV-1 and -2, HVJ, vesicular stomatitis virus, vaccinia virus, and polio- and adenoviruses, respectively. These results indicate that non-inactivation by AgNO_3 of the viruses other than HSV may be due to the insensitivity to the compound of the virus itself, in contrast to the case of an undiluted preparation of HSV-1 at 10^7 TCID₅₀/ml, in which non-inactivation of HSV is possibly due to the diminution of the AgNO_3 effect by the nonviral proteins. Thus, it is concluded that AgNO_3 at a concentration of $30 \mu\text{M}$ was specific to herpesviruses and could not inactivate other test viruses.

Inhibition of AgNO_3 effect by different reagents. An experiment was performed to test whether the inactivation effect of AgNO_3 is due to the binding of ionized Ag to the sulfhydryl groups in the virus (Table 1). The results clearly show that no inactivation of HSV was obtained with AgNO_3 after it was reacted with a sulfhydryl compound, R-glutathione, or with KI, the reaction of which with AgNO_3 results in formation of AgI. In addition, AgI per se, which is an extremely low-ionizing compound, could not inactivate HSV at all. These results indicate that this specific inactivation of HSV by

FIG. 2. Loss of infectivity of HSV of various infectivities by silver nitrate. Symbols: ●, crude virus; ○, purified virus; △, purified virus after addition of bovine serum albumin.

FIG. 3. Effect of silver nitrate on various viruses.

TABLE 1. Inhibition by different reagents of the AgNO_3 effect on HSV-1

Compounds added to HSV ^a	Residual virus ^b	
	PFU/ml	% of control
AgNO_3	<10	<0.02
AgNO_3 + R-glutathione ^c	4.8×10^4	88.9
AgNO_3 + KI ^c	4.3×10^4	79.5
AgI	3.9×10^4	72.2
MEM	5.4×10^4	

^a Final concentrations of AgNO_3 , R-glutathione, KI, and AgI in the virus preparations were 30, 10^3 , 300, and $0.006 \mu\text{M}$, respectively.

^b Residual virus was assayed after exposure of the virus to the compound at 37 C for 60 min. PFU, Plaque-forming units.

^c AgNO_3 and R-glutathione or AgNO_3 and KI were incubated at room temperature for 5 min after mixing and then were added to the virus preparation.

AgNO_3 may be due to the binding of ionizing Ag to the sulfhydryl groups in the virus.

Biological activities retained in AgNO_3 -inactivated HSV. To investigate the steps by which replication of the AgNO_3 -treated virus was prevented, we tested whether the inactivated virus retained some biological activities to the cells.

First, we tested the ability of the virus to adsorb to the cell surface. [^3H]thymidine-labeled HSV (12,000 counts/min per ml), which possessed an infectivity of 7×10^5 , was divided into two equal portions. Infectivity of one por-

tion of the virus was completely inactivated by exposure to $30 \mu\text{M}$ AgNO_3 at 37 C for 60 min. The other portion was incubated in PBS at 37 C for 60 min and served as an intact virus. Both virus preparations were diluted fourfold in MEM, and 1 ml of each was then inoculated into approximately 4×10^5 3T3 cells and incubated at 37 C. At intervals, radioactivities in the supernatant fluids and the cells were counted in a liquid scintillation spectrometer (Fig. 4). The results clearly show that the radioactivity of inactivated virus decreased in the supernatant fluids and increased in the cells, with time, at rates almost identical to those of intact virus. Simultaneously, the infectivity residues in the supernatant fluids were assayed in the cultures infected with intact virus (Fig. 4). The results show that the rates of decrease in radioactivities of both the intact and inactivated viruses were similar to that of the decrease in infectivity. This indicates that the HSV inactivated with AgNO_3 retained almost fully the activity of adsorbing to the cells.

A further attempt was made to test whether infection with the inactivated virus can induce the early surface changes and synthesis of viral antigens observed in the cells infected with intact HSV. The results clearly show that neither a hemadsorption phenomenon nor cell fusion and polykaryocytes occurred in the cells infected with the inactivated virus (Table 2). Neither surface nor intracellular viral antigens, as detected by immunofluorescence tests, were observed (Table 2).

Tevethia et al. have reported that, soon after

FIG. 4. Adsorption of AgNO_3 -inactivated HSV to the cells. Symbols: Circles represent the radioactivity in the cells—●, cells infected with AgNO_3 -inactivated HSV; ○, cells infected with intact virus. Squares represent the radioactivity in the culture fluids—■, AgNO_3 -inactivated HSV; □, intact virus. Triangles represent the infectivity in the supernatant fluids of the cultures infected with intact virus.

infection, the cells infected with HSV showed agglutinability by ConA before production of infectious virus (11). We thus examined the agglutinability by ConA of cells infected with the AgNO_3 -inactivated virus. The results clearly show that the chicken embryo cells infected with inactivated virus, as well as the cells infected with intact virus, were agglutinated in the presence of $250 \mu\text{g}$ of ConA per ml, by which the noninfected cells tested as a control were not agglutinated at all (Table 3). Hence, it would seem that infection of the inactivated virus underwent the early steps that produce certain cell surface changes responsible for ConA agglutinability, but not other surface changes and viral antigens.

DISCUSSION

It is well known that many agents exist that have potent antiviral properties against extracellular HSV (7) and that mercaptide-forming sulphydryl reagents are a group of potent inactivators (9). The present report has shown that the inactivation of HSV by AgNO_3 is also probably attributable to the formation of mercaptides of Ag with the SH base of the virion components. Previous investigators have re-

TABLE 2. Surface changes and antigen production in cells infected with intact and AgNO_3 -inactivated HSV^a

Surface changes and antigen production	HSV ^b			
	Intact		AgNO_3 -inactivated	
	6 h	20 h	6 h	20 h
Hemadsorption	1+	3+	0	0
Surface antigens	1+	4+	0	0
Intracellular antigens	1+	4+	0	0
Cell fusion and polykaryocytes	0	3+	0	0

^a HSV-1 containing 5×10^5 TCID₅₀/ml, and its preparation inactivated by $30 \mu\text{M}$ AgNO_3 were diluted to one-fifth in MEM and inoculated into 10^5 3T3 cells. The infected cells were cultivated at 37 C and tested for their surface changes and production of viral antigens 6 and 20 h after infection.

^b Positive reactions were graded from 1+ to 4+ depending upon the degree of reaction: 4+, almost all the cells were positive; 1+, less than one-fourth of the total cells were positive.

ported that the infectivities of many viruses other than HSV were also inactivated by 10^{-3} to 10^{-5} M mercuric compounds such as mercuric

TABLE 3. Agglutination by ConA of chicken embryo cells infected with AgNO₃-inactivated HSV-1

Cells infected with:	Agglutination by ConA (μ g/ml)		
	500	250	0
Intact HSV ^a	+++	++	—
AgNO ₃ -inactivated HSV ^a	++	++	—
Not infected (7.5 μ M AgNO ₃) ^b	—	—	—
Not infected	—	—	—

^a A 1-ml amount of HSV-1 containing 2×10^6 TCID₅₀/ml and its preparation inactivated by 30 μ M AgNO₃ were diluted to one-fourth in MEM. Two milliliters of each was inoculated into the monolayers of 10⁶ chicken embryo cells in plastic petri dishes. The infected cells were then tested for agglutination by ConA as described in the text.

^b To test whether AgNO₃ per se, of which trace amounts probably are residue in the AgNO₃-treated virus preparation, could induce ConA agglutinability, the cells were incubated in the presence of 7.5 μ M AgNO₃ for 6 h at 37 C and then tested for their ConA agglutinability.

chloride, *p*-chloromercuribenzoate, or merthiolate (9). We have, however, demonstrated in this paper that 10⁻⁵ M AgNO₃ could inactivate specifically the infectivity of HSV at exposure times that did not affect the infectivity of other deoxyribonucleic acid or enveloped viruses or of ribonucleic acid or non-enveloped viruses tested so far. The biochemical basis for the differential inactivation of herpesviruses and other viruses is unknown. However, it was concluded that the sulfhydryl groups of the HSV virion may play an important role, more so than those in other viruses, in keeping the superficial structure required for the infectivity of the virus.

The present experiments have also demonstrated that AgNO₃ did not affect the rate of attachment of HSV to host cells. Philipson and Choppin (4, 9) have demonstrated that treatment with mercuric chloride and *p*-chloromercuribenzoate prevented the hemagglutination by many Echo- and Coxsackie viruses and the attachment of Echo-7 virus to host cells. In contrast, treatment with such sulfhydryl reagents have been shown not to have any effect on hemagglutination by fowl plague virus and other myxoviruses (9) and on attachment of fowl plague and vaccinia viruses to host cells (1). These results and our findings thus strongly suggest the possibility that the sulfhydryl groups in the enveloped viruses, in contrast to those in the non-enveloped viruses, are not involved in the adsorption of the virus to host cells.

It appears that AgNO₃ may prevent the multiplication of HSV at a certain step(s) of infection soon after attachment. The present paper does not indicate a precise step(s) after attachment responsible for the abortion of infection. Allison has presented some evidence indicating that after treatment with mercurials the uncoating of the vaccinia virus may not occur, although the virus has been taken up by the cells (1). Whether a similar mechanism exists for the inactivation of HSV by AgNO₃ is under investigation.

It is interesting to note that infection of the cells with AgNO₃-inactivated virus could induce ConA agglutinability, whereas it produced no virus-determined surface changes. Tevethia et al. have reported that the surface change leading to the agglutination of HSV-infected cells is caused by a virus-determined function but not by a nonspecific change due to virus-cell interaction (11). This may not be the case here, since it seems unlikely that the inactivated virus functioned as a genome determined only for the induction of ConA agglutinability and not for the other virus-specific changes. It seems more likely that certain nonspecific changes occurred in the steps of virus adsorption or penetration that were responsible for the induction of the ConA agglutinability obtained in our study.

ACKNOWLEDGMENT

We wish to thank Shoko Tanabe for her excellent assistance in preparation of the manuscript.

LITERATURE CITED

1. Allison, A. C. 1962. Observations on the inactivation of viruses by sulfhydryl reagents. *Virology* 17:176-183.
2. Allison, A. C., F. E. Buckland, and C. H. Andrewes. 1962. Effects of sulfhydryl reagents on infectivity of some viruses. *Virology* 17:171-175.
3. Chang, Te-Wen, and L. Weinstein. 1975. In vitro activity of silver sulfadiazine against *Herpesvirus hominis*. *J. Infect. Dis.* 132:79-81.
4. Choppin, P. W., and L. Philipson. 1960. The inactivation of enterovirus infectivity by the sulfhydryl reagent *p*-chloromercuribenzoate. *J. Exp. Med.* 113:713-734.
5. Coleman, V. R., J. Wilkie, W. E. Levinson, T. Stevens, and E. Jawetz. 1973. Inactivation of *Herpesvirus hominis* types 1 and 2 by silver nitrate in vitro and in vivo. *Antimicrob. Agents Chemother.* 4:259-262.
6. Inbar, M., and L. Sachs. 1969. Interaction of the carbohydrate-binding protein concanavalin A with normal and transformed cells. *Proc. Natl. Acad. Sci. U.S.A.* 63:1418-1425.
7. Levinson, W., V. Coleman, B. Woodson, A. Rabson, J. Lanier, J. Whitchee, and C. Dawson. 1974. Inactivation of herpes simplex virus by thiosemicarbazones and certain cations. *Antimicrob. Agents Chemother.* 5:398-402.
8. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193:265-275.

9. Philipson, L., and P. W. Choppin. 1960. On the role of virus sulfhydryl groups in the attachment of enteroviruses to erythrocytes. *J. Exp. Med.* 112:455-478.
10. Sery, T. W., and F. P. Furguieles. 1961. The inactivation of herpes-simplex virus by chemical agents. *Am. J. Ophthalmol.* 51:42-57.
11. Tevethia, S. S., S. Lowry, W. E. Rawls, J. L. Melnick, and V. McMillan. 1972. Detection of early cell surface changes in herpes simplex virus infected cells by agglutination with concanavalin A. *J. Gen. Virol.* 15:93-97.
12. Watkins, J. F. 1964. Adsorption of sensitized sheep erythrocytes to HeLa cells infected with herpes simplex virus. *Nature (London)* 202:1364-1365.