Published in final edited form as:

Circ Res. 2014 October 10; 115(9): e26–e27. doi:10.1161/CIRCRESAHA.114.304975.

Angiotensin II Type-2 Receptor Effects: Author Reply

Brand A. Kemp¹, Nancy L. Howell¹, John J. Gildea², Susanna R. Keller¹, Shetal H. Padia¹, and Robert M. Carey¹

¹Division of Endocrinology and Metabolism, Department of Medicine, University of Virginia Health System, Charlottesville, VA

²Department of Pathology, University of Virginia Health System, Charlottesville, VA

The authors thank Drs. Maiolino, Naso and Calo for their kind recognition of our study (1). We are delighted to learn of the evidence from the human model of Bartter's and Gietelman's syndromes that AT₂ receptors (AT₂R) are activated via angiotensin signaling. Indeed, in both of these human syndromes there is marked activation of the renin-angiotensin system (RAS) associated with hypotension, to which AT₂Rs may contribute by direct vasodilation and/or natriuresis.

Numerous studies in experimental animals have shown that AT₂Rs oppose the pressor actions of angiotensin II (Ang II) via AT₁ receptors (AT₁Rs) and lower blood pressure, especially in the presence of AT₁R blockade (2-9). Indeed, AT₂R stimulation lowers blood pressure predominantly under circumstances in which the RAS is activated (10-13) by increasing pressure-natriuresis (11,13). It is likely that the depressor and hypotensive effects of AT₂R activation described in experimental animals translate to human diseases such as Bartter's and Gietelman's syndromes, as evidenced by the above-cited work of Drs. Maiolino, Naso and Calo.

Our recent study (1) demonstrates that selective renal AT₂R activation induces natriuresis in experimental animals by inducing proximal tubule sodium transporter internalization and inactivation. The mechanism by which sodium transporters are inhibited was identified as increased AT₂R-induced renal production of bradykinin, nitric oxide and cyclic GMP, similar to the signaling pathway mediating most, if not all, of the described AT₂R functions, including vasodilation (14).

In contrast to the actions of Ang II to internalize and desensitize AT₁Rs, our studies show that AT₂R activation recruits receptors from intracellular compartments to the apical plasma membranes of proximal tubule cells (1). AT₂R recruitment may therefore serve as a reinforcing positive feedback mechanism which sustains natriuresis in response to activation of the RAS, opposing AT₁R-mediated reduction in sodium excretion. Studies to date show that AT₂R recruitment to the apical plasma membrane is requisite to inhibition of proximal tubule sodium reabsorption (1). The mechanism for translocation of AT₂Rs from cytosol to

apical membrane in response to AT₂R activation has been identified by our group as likely to be increased renal cyclic AMP production (15).

It is important to point out that the primary/preferred agonist for endogenous AT₂R activation in experimental animals is not Ang II, but des¹-aspartyl-Ang II (Ang III) (16). Ang II seems to work primarily or exclusively after conversion to Ang III by the action of aminopeptidase A (16,17). Whether Ang III acts as an endogenous AT₂R agonist in human disorders, such as the Bartter's and Gietelman's syndromes, awaits further study.

Recent studies have pointed out the importance of the intrarenal RAS as a local tissue system functioning independently from the systemic system in the pathophysiology of diseases such as hypertension. These studies have shown that the intrarenal RAS is important in the regulation of blood pressure and renal function in experimental animals. This concept is highly likely to apply also to humans. Overactivation of the intrarenal RAS leads to increased Ang II levels, which unlike the systemic RAS, can induce auto-amplification of Ang II production (18-22). AT₁R-dependent Ang II cellular internalization and up-regulated renin synthesized in the distal nephron may provide a sufficient source of continuing intrarenal Ang II formation, to maintain vasoconstriction, antinatriuresis and hypertension (20-22). The precise role of the AT₂R as a counter-regulatory receptor in modulating these intrarenal responses is currently unknown and may be an important area of future investigation.

We anticipate that AT₂R activation might be applied to human conditions of sodium retention such as congestive heart failure and/or hypertension. Our observations suggest that chronic systemic administration of Compound 21, a potent, selective non-peptide AT₂R agonist, counteracts sodium retention and lowers blood pressure in the angiotensin II-infusion model of experimental hypertension in rats (1). Since there is no available diuretic for clinical use in humans that effectively reduces sodium reabsorption at the proximal tubule, it is possible that AT₂R activation in that nephron segment would complement actions of available diuretic/natriuretic agents acting in the ascending limb of Henle, distal tubule and/or collecting duct. This exciting possibility awaits further study.

References

1. Kemp BA, Howell NL, Gildea JJ, Keller SR, Padia SH, Carey RM. AT₂ receptor activation induces natriuresis and lowers blood pressure. *Circ Res*. 2014; 115:388–399. [PubMed: 24903104]
2. Tsutsumi Y, Matsubara H, Masake H. Angiotensin type2 receptor overexpression activates the vascular kinin system and causes vasodilation. *J Clin Invest*. 1999; 104:925–935. [PubMed: 10510333]
3. Moore AF, Heiderstedt NT, Huang E, Howell NL, Wang ZQ, Siragy HM, Carey RM. Selective inhibition of the renal angiotensin type 2 receptor increases blood pressure in conscious rats. *Hypertension*. 2001; 37:1285–1291. [PubMed: 11358942]
4. Carey RM, Howell NL, Jin XH, Siragy HM. Angiotensin type-2 (AT₂) receptor-mediated hypotension in type-1 (AT₁) receptor-blocked rats. *Hypertension*. 2001; 38:1272–1277. [PubMed: 11751702]
5. Li XC, Widdop RE. AT₂ receptor-mediated vasodilation is unmasked by AT₁ receptor blockade in conscious SHR. *Br J Pharmacol*. 2004; 142:821–830. [PubMed: 15197103]

6. Duke LM, Evans RG, Widdop RE. AT₂ receptors contribute to acute blood pressure-lowering and vasodilator effects of AT₁ receptor antagonism in conscious normotensive but not hypertensive rats. *Am J Physiol Heart Circ Physiol*. 2005; 288:H2289–H2297. [PubMed: 15615839]
7. Bosnyak S, Welungoda IK, Hallberg A, Alterman M, Widdop RE, Jones ES. Stimulation of angiotensin AT₂ receptors by the non-peptide agonist, Compound 21, evokes vasodepressor effects in conscious spontaneously hypertensive rats. *Br J Pharmacol*. 2010; 159:709–716. [PubMed: 20128808]
8. Jones ES, Del Borgo MP, Kirsch JF, Clayton D, Bosnyak S, Welungoda I, Hausler N, Unabia S, Perlmutter P, Thomas WG, Aguilar MI, Widdop RE. A single β -amino acid substitution to angiotensin II confers AT₂ receptor selectivity and vascular function. *Hypertension*. 2011; 57:570–576. part 2. [PubMed: 21300665]
9. Ali Q, Wu Y, Hussain T. Chronic AT₂ receptor activation increases renal ACE2 activity, attenuates AT₁ receptor function and blood pressure in obese Zucker rats. *Kidney Int*. 2013 e-pub ahead of print.
10. Siragy HM, Carey RM. Protective role of the angiotensin AT₂ receptor in renal vascular hypertension in conscious rats. *Hypertension*. 1999; 33:1237–1242. [PubMed: 10334818]
11. Siragy HM, Inagami T, Ichiki T, Carey RM. Sustained hypersensitivity to angiotensin II and its mechanism in mice lacking the subtype-2 (AT₂) angiotensin receptor. *Proceedings of the National Academy of Sciences USA*. 1999; 96:6506–6510.
12. Siragy HM, deGasparo M, Carey RM. Angiotensin type 2 receptor mediates valsartan-induced hypotension. *Hypertension*. 2000; 35:1074–1077. [PubMed: 10818067]
13. Gross V, Schunck WF, Honek H, Milia AF, Kargel E, Walther T, Bader M, Inagami T, Schneider W, Luft FC. Inhibition of pressure natriuresis in mice lacking the AT₂ receptor. *Kidney Int*. 2000; 57:191–202. [PubMed: 10620200]
14. Padia SH, Carey RM. AT₂ receptors: beneficial counter-regulatory role in cardiovascular and renal function. *Pflügers Arch*. 2013; 120:476–479.
15. Padia SH, Kemp BA, Howell NL, Keller SR, Gildea JJ, Carey RM. Mechanisms of dopamine D(1) and angiotensin type 2 receptor interaction in natriuresis. *Hypertension*. 2012; 59:437–445. [PubMed: 22203736]
16. Padia SH, Kemp BA, Howell NL, Fournie-Zaluski MC, Roques BP, Carey RM. Conversion of renal angiotensin II to angiotensin III is critical for AT₂receptor-mediated natriuresis in rats. *Hypertension*. 2008; 51:460–465. Part 2. [PubMed: 18158338]
17. Kemp BA, Bell JF, Rottkamp DM, Howell NL, Shao W, Navar LG, Padia SH, Carey RM. Intrarenal angiotensin III is the predominant agonist for proximal tubule angiotensin type 2 receptors. *Hypertension*. 2012; 60:387–395. [PubMed: 22689743]
18. Kobori H, Harrison-Bernard LM, Navar LG. Enhancement of angiotensinogen expression in angiotensin II-dependent hypertension. *Hypertension*. 2001; 37:1329–1335. [PubMed: 11358949]
19. Kobori H, Nangaku M, Navar LG, Nishiyama A. The intrarenal renin-angiotensin system: from physiology to pathobiology of hypertension and kidney disease. *Pharmacol Rev*. 2007; 59:251–287. [PubMed: 17878513]
20. Navar LG, Kobori H, Prieto MC, Gonzalez-Villalobos RA. Intratubular renin-angiotensin system in hypertension. *Hypertension*. 2011; 57:355–362. [PubMed: 21282552]
21. Prieto-Carrasquero MC, Kobori H, Ozawa Y, Gutierrez A, Seth D, Navar LG. AT₁ receptor-mediated enhancement of collecting duct renin in angiotensin II-dependently hypertensive rats. *Am J Physiol Renal Physiol*. 2005; 289:F632–F637. [PubMed: 15870381]
22. Prieto-Carrasquero MC, Williams DE, Liu L, Kavanaugh KI, Mullins JJ, Mitchell KD. Enhancement of renin and prorenin receptor in collecting duct of Cyp11a1-Ren 2 rats may contribute to development and progression of malignant hypertension. *Am J Physiol Renal Physiol*. 2011; 300:F581–F588. [PubMed: 21068087]