REFERENCES

1. Raghavan D, Khan S. The use of otoacoustic emission to detect subclinical inner ear damage in divers of the Indian Navy. *Med J Armed Forces India*. 2014;70:344–348.
2. Skogstad M, Eriksen T, Skare O. A twelve-year longitudinal study of hearing thresholds among professional divers. *Undersea Hyperb Med*. 2009;36:25–31.
3. Taylor DM, Lippmann, Smith D. The absence of hearing loss in otologically asymptomatic recreational scuba divers. *Undersea Hyperb Med*. 2006 Mar-Apr;33:135–141.
4. Goplen FK, Aasen T, Grønning M, Molvær OI, Nordahl SHG. Hearing loss in divers: a 6 year prospective study. *Eur Arch Otorhinolaryngol*. 2011;268:979–985.
5. Smith PF. Towards a standard for hearing conservation for underwater and hyperbaric environments. *J Aud Res*. 1985 Oct;25:221–238.
6. Hausmann D, Laabing S, Hoth S, Plinkert PK, Klingmann C. Assessment of the peripheral hearing system of sport divers. *Undersea Hyperb Med*. 2011 Nov–Dec;38:515–526.
7. Salami A, Dellepiane M, Barbierato M, et al. The effects of active sonar for the protection of moored and anchored warships on the human hearing. *Eur Arch Otorhinolaryngol*. 2010 Feb;267:207–211.
8. Hizel SB, Muluk NB, Budak B, Budak G. Does scuba diving cause hearing loss? *J Otolaryngol*. 2007 Aug;36:247–252.

Lt Col K. Manoj Kumar*
Assistant Professor, Dept of ENT, Armed Forces Medical College,
Pune 411040, India

Col N. Ramakrishnan
Professor & Head of Department, Dept of ENT,
Armed Forces Medical College, Pune 411040, India

Col S.K. Singh
Associate Professor, Dept of ENT, Armed Forces Medical College,
Pune 411040, India

*Corresponding author.
E-mail address: manojkanzhuly@gmail.com (K. Manoj Kumar)

Available online 27 December 2014

DOI of original article: <http://dx.doi.org/10.1016/j.mjafi.2014.04.004>

<http://dx.doi.org/10.1016/j.mjafi.2014.12.005>
0377-1237/© 2014, Armed Forces Medical Services (AFMS). All rights reserved.

Reply

Dear Editor,

I would like to thank the readers for the interest shown in our article. We are grateful for the comments. As a whole, we by and large agree with, and appreciate the comments in the above letter regarding our article. As brought out in the article and in the above Letter to the Editor, the damage to the Inner Ear is probably multi-factorial and cumulative.

It is because of the likely cumulative nature of the damage to the Inner Ear that we studied the possible incidence of increased Inner Ear Damage with rise in Diving Years. However our study did not find a correlation between the length of the Diving Career and incidence of Inner Ear Damage as measured by OAE. This does not however exclude the possibility; a much larger study may shed further light on this subject.

Noise as a cause of Inner Ear Damage has been mentioned in the article and in the Letter to the Editor. However with the present facilities and equipment available, it may not be possible to conclusively prove what proportion of the Inner Ear Damage is due to Noise. However it is pertinent to mention that Abalone Divers who are not exposed to Noise also have Inner Ear Damage and SNHL as brought out in our article. Stratification of Dive Profiles and the Dive Objective especially with respect to Noise Exposure in a future study may be required to quantify the contribution of the various risk factors towards Inner Ear Damage.

As pointed out in the Letter to the Editor, we agree that ideally, OAE measurements and Pure Tone Audiometry should be done immediately prior to and after a dive. However given the difficulty in achieving sound treated/sound proof conditions to conduct audiometry at the dive site and limitations of logistically covering dispersed Diving Teams with the limited equipment and manpower available, the same could not be done; hence the method of checking for cumulative damage was employed.

Further studies on this subject will probably clear the picture.

Surg Capt Dilip Raghavan
Senior Advisor (ENT), INHS Asvini, Colaba, Mumbai 400005, India
E-mail address: dilipraghavan@hotmail.com

Available online 27 December 2014

DOI of original article: <http://dx.doi.org/10.1016/j.mjafi.2014.11.015>

<http://dx.doi.org/10.1016/j.mjafi.2014.11.016>
0377-1237/© 2014, Armed Forces Medical Services (AFMS). All rights reserved.