ORIGINAL RESEARCH COMMUNICATION

Association of *Nrf2* Polymorphism Haplotypes with Acute Lung Injury Phenotypes in Inbred Strains of Mice

Hye-Youn Cho,¹ Anne E. Jedlicka,² Wesley Gladwell,¹ Jacqui Marzec,¹ Zackary R. McCaw,¹ Rachelle J. Bienstock,³ and Steven R. Kleeberger¹

Abstract

Aims: *Nrf2* is a master transcription factor for antioxidant response element (ARE)-mediated cytoprotective gene induction. A protective role for pulmonary *Nrf2* was determined in model oxidative disorders, including hyperoxia-induced acute lung injury (ALI). To obtain additional insights into the function and genetic regulation of *Nrf2*, we assessed functional single nucleotide polymorphisms (SNPs) of *Nrf2* in inbred mouse strains and tested whether sequence variation is associated with hyperoxia susceptibility. **Results:** *Nrf2* SNPs were compiled from publicly available databases and by re-sequencing DNA from inbred strains. Hierarchical clustering of *Nrf2* SNPs categorized the strains into three major haplotypes. Hyperoxia susceptibility was greater in haplotypes 2 and 3 strains than in haplotype 1 strains. A promoter SNP –103 T/C adding an Sp1 binding site in haplotype 2 diminished promoter activation basally and under hyperoxia. Haplotype 3 mice bearing nonsynonymous coding SNPs located in (1862 A/T, His543Gln) and adjacent to (1417 T/C, Thr395Ile) the Neh1 domain showed suppressed nuclear transactivation of pulmonary *Nrf2* relative to other strains, and overexpression of haplotype 3 *Nrf2* showed lower ARE responsiveness than overexpression of haplotype 1 *Nrf2* in airway cells. Importantly, we found a significant correlation of *Nrf2* haplotypes and hyperoxic lung injury phenotypes. **Innovation and Conclusion:** The results indicate significant influence of *Nrf2* polymorphisms and haplotypes on gene function and hyperoxia susceptibility. Our findings further support *Nrf2* as a genetic determinant in ALI pathogenesis and provide useful tools for investigators who use mouse strains classified by *Nrf2* haplotypes to elucidate the role for *Nrf2* in oxidative disorders. *Antioxid. Redox Signal.* 22, 325–338.

Introduction

EXCESS PRODUCTION OF REACTIVE OXYGEN SPECIES (ROS) during oxidative stress has been implicated in the pathogenesis of many acute and chronic pulmonary disorders (3, 12, 36). Acute lung injury (ALI) or its severe form, acute respiratory distress syndrome (ARDS), is a major lung disease that affects millions worldwide. ALI is characterized by disrupted epithelial-vascular barrier function, resulting in increased permeability of alveoli to water, proteins, and other solutes and intra-alveolar accumulation of inflammatory cells and erythrocytes, which leads to abnormal gas exchange and

variable late phase responses, including fibrosis in the airways (53). Release of ROS and proteases, as well as increased synthesis of chemokine and cytokines from leukocytes, is known to play a key role in the pathogenesis. Hyperoxia exposure to laboratory rodents causes pulmonary damage that resembles ALI/ARDS phenotypes.

An inbred mouse screen and genetic linkage analysis with *F₂* and recombinant inbred mice from hyperoxia-susceptible (C57BL/6J, B6) and -resistant (C3H/HeJ, C3) strains identified nuclear factor, erythroid derived 2, like 2 (*Nfe2l2* or NF-E2-related factor 2, *Nrf2*) as a susceptibility gene for hyperoxia-induced lung injury (7, 20). *Nrf2* is a ubiquitous

¹Laboratory of Respiratory Biology, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, North Carolina.

²Department of Molecular Biology and Immunology, Johns Hopkins University Bloomberg School of Public Health, Baltimore, Maryland.

³Laboratory of Structural Biology, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, North Carolina.

Innovation

Nrf2 is a key transcription factor for antioxidant defense. While the protective role for Nrf2 has been widely investigated in model oxidative airway disorders using mice with targeted deletion of *Nrf2*, genetic variation of *Nrf2* and its functional relevance to disease susceptibility has been rarely studied. Single nucleotide polymorphisms (SNPs) of murine *Nrf2* classified inbred strains into three major haplotypes. We found a significant correlation of haplotypes with hyperoxia-induced acute lung injury (ALI) phenotypes, and determined functional consequences of SNPs within the haplotypes that provide important determinants of ALI susceptibility. The results suggest that mouse strains classified by haplotypes are useful tools to elucidate the role of Nrf2 in various oxidative disorders.

transcription factor that binds to a *cis*-acting antioxidant response element (ARE) for a variety of cytoprotective proteins, including phase 2 detoxifying enzymes, drug efflux pumps, and ROS scavengers (21, 47). Using mice genetically deficient in *Nrf2* (*Nrf2*^{-/-}), we demonstrated a protective role for the Nrf2-ARE pathway in hyperoxia-induced lung injury (6, 39, 40).

In order to obtain additional insights into the function and genetic regulation of *Nrf2* beyond that which was obtained with the simple gene deletion, we systematically evaluated sequence variation and structure of *Nrf2* and tested whether variants and haplotypes are associated with hyperoxia susceptibility in inbred strains of mice. First, *Nrf2* single nucleotide polymorphisms (SNPs) in 72 inbred strains were compiled from publicly available databases and three major SNP haplotypes were identified. We also re-sequenced *Nrf2* exons and 1000 bp upstream promoter of 16 strains from the three haplotypes to further elucidate *Nrf2* SNPs. Next, the 16 strains were exposed to hyperoxia to determine whether the susceptibility phenotypes were associated with the haplotypes. Functional relevance of SNP haplotypes was also assessed by comparing lung mRNA expression and Nrf2-ARE activation among the strains. Finally, we tested *in vitro* the roles of potentially informative SNPs for their influence on Nrf2 function in response to hyperoxia.

This novel approach extends our understanding of the genetic regulation and expression of Nrf2 beyond what was known in B6 and C3 mice to include additional functional variants in the mouse genome, and may provide greater insights into variation in human NRF2 function. The investigation also further characterized the role of genetic variation in *Nrf2* and its role in susceptibility to oxidant-induced lung injury that could not be understood using gene-specific knockout mice (6). The current model with genetically diverse, and differentially susceptible, strains of mice may provide an additional understanding of the role of NRF2 in clinical ALI and ARDS (32).

Results

Nrf2 SNPs and haplotypes

A total of 173 SNPs were identified in *Nrf2* including the 5000 bp upstream promoter and 3402 bp downstream regions

(chr2:75547698–75510174 bp) from the mouse phenome database (MPD, <http://phenome.jax.org/SNP>) for 72 inbred strains (16, 49, 51, 55). Hierarchical clustering identified three major *Nrf2* haplotypes: haplotype 1 (28 strains, in blue); haplotype 2 (23 strains, in red); and haplotype 3 (15 strains, in green) (Fig. 1A and Supplementary Table S1; Supplementary Data are available online at www.liebertpub.com/ars). Six strains (in purple), including three wild derived, are distinct from each other and haplotypes 1–3. Additional *Nrf2* SNPs were collected from NCBI dbSNP (www.ncbi.nlm.nih.gov/SNP) and the recent update of the MPD (5). We also found novel SNPs by re-sequencing the *Nrf2* promoter and coding regions of 16 strains from the three haplotype groups (Supplementary Table S2 and Table 1). Overall, 1114 polymorphisms were identified in *Nrf2* spanning 36,123 bp of mouse chromosome 2 (GRCm38.p1 Primary Assembly; Ch2.75709641-75673519), which included 179 in the 5 kb upstream region, 49 in exons 1–5, 806 in introns, and 80 in 2000 bp downstream regions (Supplementary Table S2). This indicates a SNP frequency of 1 per every 32 bp. While most of the SNP alleles are bi-allelic, 13 tri-allelic SNPs were also identified (Supplementary Table S2). Ten percent of SNPs vary between haplotypes 1 and 2, while 39% and 35% are polymorphic between haplotypes 2 and 3 and haplotypes 1 and 3, respectively (Supplementary Table S2 and Fig. 1B).

The distal promoter region of *Nrf2* overlaps with a neighboring RIKEN gene (E030042O20Rik, Ch2:75704770-75710024 bp) that is located from 129 bp upstream of the *Nrf2* TSS. Among SNPs within the 1000 bp upstream region (Table 1), four SNPs at positions –921, –887/–884, and –103 [formerly –336 (7)] in haplotype 2 potentially alter transcription factor binding function relative to haplotypes 1 or 3 as determined by TFSEARCH (www.cbrc.jp/research/db/TFSEARCH.html; Fig. 2 and Table 1). Among 11 SNPs in coding sequences (Table 1), 2 nonsynonymous coding SNPs (Cns) are located in (c.1862A/T, p.His543Gln) and adjacent to (c.1417T/C, p.Thr395Ile) the ECH homology I (Neh1) domain (Fig. 2), which is essential for nuclear translocation and DNA (ARE) binding (5, 22).

Effect of SNP haplotypes on hyperoxia susceptibility

When 16 mouse strains selected from three haplotypes were exposed to hyperoxia ($\geq 95\%$ O₂, 72 h), total bronchoalveolar lavage (BAL) cell numbers were significantly increased in many strains from haplotypes 2 (C57BL/10J or B10, SJL/J, AKR/J, and B6) and 3 (SM/J, P/J, SEA/GnJ) and C3 mice from haplotype 1, relative to those of strain-matched air controls (Fig. 3A). Compared with haplotype 1 mice, hyperoxia-induced total BAL cell numbers were significantly greater in mice from haplotypes 2 ($p = 0.02$) and 3 ($p = 0.01$), and those from haplotypes 2 and 3 were not significantly different from each other. Mean total BAL cells and SNP haplotypes were significantly correlated (Pearson correlation coefficient = 0.658, $p = 0.006$; Spearman Rank Order coefficient = 0.635, $p = 0.008$). Significant differences between haplotype groups were also found for total BAL protein (data not shown). Hyperoxia exposure significantly decreased body weight in haplotype 2 strains (72 h) and SM/J and P/J of haplotype 3 (48 and 72 h); body weight of the majority of haplotype 1 strains was not significantly changed by hyperoxia (Fig. 3B).

FIG. 1. *Nrf2* haplotypes in inbred mice. (A) Hierarchical clustering of 72 inbred strains of mice based on 173 *Nrf2* SNPs from the MPD (<http://phenome.jax.org/SNP>). Numbers represent the uncertainty in genotype changes after clustering (Φ), where $\Phi = 1 - \text{cluster entropy/crude entropy}$. If $\Phi = 1$, after clustering all strains in the cluster have the same genotype; if $\Phi = 0$, the genotype distribution within the cluster is as variable as the genotype distribution across all strains. Asterisks indicate strains of mice that were phenotyped for their response to hyperoxia exposure. Clusters of the strains are arranged by color for haplotypes (haplotype 1, blue; haplotype 2, red; haplotype 3, green), and six strains (in purple) are distinct from each other and haplotypes 1–3. (B) Frequency of sequence variation on the chromosomal loci of *Nrf2* (Ch2:75704641–75675519 bp) and flanking regions (5 kb upstream and 2 kb downstream) between haplotypes is indicated as percent polymorphic. A total of 945 SNPs were identified between haplotypes 1 and 2 strains, while only 239 SNPs of them were found in haplotype 3 mice (details in Supplementary Table S2). MPD, mouse phenome database; SNP, single nucleotide polymorphism. To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars

Functional roles of *Nrf2* SNPs in vitro

Promoter SNP. The effects of a promoter SNP –103T/C on the regulation of *Nrf2* expression was determined by transient transfection of airway epithelial cells with luciferase vectors bearing the 5' upstream region (–612 to +228) from haplotype 1 (p*Nrf2*_{C3}) or 2 (p*Nrf2*_{B6}) mice. Basal promoter activity as well as hyperoxia (85% O_2 , 4 h)-enhanced reporter gene activity in cells was significantly lower with p*Nrf2*_{B6} transfection than with p*Nrf2*_{C3} transfection (Fig. 4A, top). Effects of the –103T/C promoter SNP on potential Sp1 binding activity was further analyzed by gel shift assays, and nuclear protein binding activity on the B6 (haplotype 2)-like promoter probe, including the –103T/C SNP, was relatively more intense than that on the C3 (haplotype 1)-like promoter probe (Fig. 4A, bottom).

Coding SNP. Binding activities of nuclear proteins to either ARE or NF-E2 consensus sequences determined by gel shift analysis (Fig. 4B, top) were greater in airway epithelial cells with overexpressed haplotype 1 *Nrf2* (pFlag-*Nrf2*_{Hap1}) than in cells with overexpressed haplotype 3 *Nrf2* (pFlag-*Nrf2*_{Hap3}) after hyperoxia (85% O_2 , 6 h). Similarly, luciferase-ARE reporter gene activity with pFlag-*Nrf2*_{Hap1}

transfection was significantly higher than that with pFlag-*Nrf2*_{Hap3} transfection after hyperoxia and at baseline (Fig. 4B, bottom).

Functional role of *Nrf2* SNPs in vivo. Effects of sequence variations on *Nrf2* gene expression were determined in mouse lungs. The amount of *Nrf2* message did not vary between the strains after air exposure (Fig. 5A). Relative to respective air controls, hyperoxia ($\geq 95\% \text{O}_2$, 48 h) induced *Nrf2* mRNA levels in 16 strains. When the ratio of hyperoxia-induced *Nrf2* mRNA expression and total BAL cells for each haplotype strain were evaluated together, we found a statistically significant inverse relationship between *Nrf2* mRNA/total cells and the three haplotypes ($\chi^2 = 13.206$; 6 d.f.; $p = 0.040$); that is, the ratio of *Nrf2* mRNA and total cells was greater in haplotype 1 strains compared with haplotype 2 strains ($\chi^2 = 10.305$; 3 d.f.; $p = 0.016$). We also elucidated, relative to haplotype 1 and 2 strains, marked attenuation of lung *Nrf2*-ARE binding ability at baseline and after hyperoxia in haplotype 3 mice bearing a Cns SNP (His543Gln or H543Q) in the DNA binding Neh1 domain and another (Thr395Ile or T395I) in the adjacent region (Fig. 5B). Mean mRNA expression levels of pulmonary ARE-responsive antioxidant enzymes, NAD(P)H:quinone oxidoreductase 1 or

TABLE 1. SINGLE NUCLEOTIDE POLYMORPHISMS IN MOUSE *Nrf2* (OR *Nfe2l2*) VARIED BETWEEN HAPLOTYPES 1 AND 3 IN 1 KB 5' UPSTREAM AND EXONS (NC_000068.7, GI: 372099108)

SNP ID	Chr2 map position (B6)	SNP location	<i>Nrf2</i> map position	Haplotype 1	Haplotype 2	Haplotype 3	SNP allele ^a	Change of SNP function (vs. C3) ^b
rs47274959	75705562	Upstream	-921	A ^c	T	A	T/A	(-) Oct-1
rs46461765	75705528	Upstream	-887	C ^c	A	C	A/C	(+) TATA-box-binding protein, (+) SRY, (-) GATA-binding factor 1, 2
rs51449853	75705525	Upstream	-884	T ^c	G	T	G/T	Unknown
rs45651867	75705498	Upstream	-857	C ^c	A	C	A/C	Unknown
rs27978306	75705449	Upstream	-808	T	A	A	A/T	Unknown
rs27978307	75705400	Upstream	-759	G	G	A	G/A	Unknown
rs247519047	75704922	Upstream	-281	-C	C	-C	Deletion	Unknown
rs27978309	75704887	Upstream	-246	T	C	T	C/T	Unknown
(rs51915758)								
rs217197904	75704744	Upstream	-103	T	C	T	C/T	(+) Sp1
NA	75704727-33	Upstream	-86: -92	NA	NA	+G	Insertion	Unknown
rs27978310	75704617	Exon 1	25	T	T	C	T/C	Unknown
rs27978312	75704605	-5' UTR	37	T	C	C	C/T	Unknown
rs27978313	75704497	-5' UTR	145	G	C	G	C/G	Unknown
rs13460861	75679262	Exon 2	446	C	T	C	T/C	Cs (F71)
rs13460859	75679202	Exon 2	506	T	C	C	C/T	Cs (H91)
rs27978436	75679187	Exon 2	521	C	C	T	C/T	Cs (T96)
rs226131070	75679178	Exon 2	530	C	C	T	C/T	Cs (S99)
rs27978444	75678500	Exon 3	623	A	A	T	A/T	Cs (V130)
rs13459064	75676732	Exon 5	1256	A	G	G	G/A	Cs (T341)
rs13459064	756767328	Exon 5	1417	C	C	T	C/T	Cns (T395I)
rs225593319	75676126	Exon 5	1862	T	T	A	T/A	Cns (H543Q)

SNPs acquired from re-sequencing or from mouse phenotype database (<http://phenome.jax.org/SNP>) and NCBI SNP Database (www.ncbi.nlm.nih.gov/SNP).

^aAll SNPs and alleles presented are complement sequences, as *Nrf2* locus lies reverse in 44.75 cM of chromosome 2. A full list of SNPs, including those in the <5 kb promoter, introns, and downstream 2 kb, are in Supplementary Tables S2 and S3.

^bTranscription factor binding site determined TFSEARCH (www.cbrc.jp/research/db/TFSEARCH.html) indicated as deletion (-) or addition (+). Amino acid F-phenylalanine, H-histidine, T-threonine, V-valine, I-isoleucine, Q-glutamine; Oct-1, octamer-binding protein; SRY, sex-determining region Y protein.

^cGenotyped by Celera whole genome shotgun sequencing using mixed strain DNA (A/J, DBA/2J, 129Sv1/SvJ, and 129S1/SvImJ). B6, C57BL/6J.

Shaded SNPs differ from those of haplotype 1.

Cs, coding-synonymous; Cns, coding-nonsynonymous; NA, not available; SNP, single nucleotide polymorphism.

FIG. 2. Informative SNPs in mice with three *Nrf2* haplotypes and predicted consequences of the mutations. Promoter (≤ 1 kb) and exon SNPs that are polymorphic among three haplotypes are mapped on mouse *Nrf2* spanning 75704641–75675519 bp on chromosome 2. Functional SNPs that may affect promoter activity and amino-acid codons are specified. Chromosome map locations are referenced to GRCm38.p1 (C57BL/6J, B6, haplotype 2), and the comparison template is C3H/HeJ (C3, haplotype 1). Red and green triangles indicate haplotypes 2 and 3 alleles, respectively. Arrows indicate SNPs predicting changes in transcription factor binding sites (promoter) or protein amino-acid sequences (exons). To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars

quinone reductase 1 (NQO1 or QR1), glutathione peroxidase 2 (GPx-2), and glutathione *S*-transferase A4 (GSTA4) were significantly lower in strains from haplotypes 2 and 3 compared with haplotype 1 after 48–72 h of hyperoxia (Fig. 5C, left). Consistent with these results, chromatin immunoprecipitation (ChIP) assays followed by semi-quantitative and quantitative polymerase chain reactions (PCRs) determined that specific binding activity of pulmonary Nrf2-Maf proteins in ARE sequences on *Nqo1* and *Gpx2* promoters was increased by hyperoxia in all three haplotype mice, while the abundance of ARE-bound protein was relatively lower in haplotype 2 (AKR/J) and 3 (SEA/GnJ) strains compared with haplotype 1 (DBA/2J) mice (Fig. 5C, middle and right). Lung protein levels of QR1, GPx-2, and GSTA1 after hyperoxia varied between haplotypes, which was consistent with their differential mRNA expression profiles (Fig. 5D).

Protein structural modeling of murine Nrf2. A three-dimensional molecular model of the murine Nrf2 DNA binding basic region/leucine zipper (bZIP) domain (amino-acid residues 490–550), and adjacent sequences bearing two Cns SNPs (T395I, H543Q) from the three haplotypes were developed based on published crystal structures of other bZIP transcription factor family members (Fig. 6A). The template bZIP structures used to develop the homology model were the SKN-1 binding domain in complex with DNA [protein data bank, PDB:1SKN (41)], which shares 36% sequence identity with murine Nrf2 (residues 444–512), the MafG-DNA complex [PDB:3A5T (28)], which shares 22% sequence identity with murine Nrf2 (residues 455–549), and CCAAT/enhancer-binding protein (C/EBP) beta bZip dimeric structure bound to DNA [PDB:1GU4], which shares 22% sequence identity with murine Nrf2 (residues 491–550). A

molecular homology model was also developed for the bZIP region of human NRF2 protein (residues 502–557) based on the crystal structure of heterodimeric human c-Fos:c-Jun complex [PDB:1FOS (18)] with which it shares 34% sequence identity (Fig. 6B). The model for human NRF2 (in yellow) is shown superimposed on the complete human c-Fos:c-Jun structure solved with DNA that suggests potential human NRF2-DNA interaction (Fig. 6B). This proposed structural model indicates that H543Q is located in the long helical segment which packs against the other helix of the bZIP structure, while T395I is not a part of this helical structure interaction within the bZIP domain. The sizes (mean volume of buried residue) of the H and Q residues are similar (167.6 Å³ vs. 161.1 Å³, respectively), and the change in size of the residue would not have a large impact on the protein structure. However, Q is a hydrophilic amino-acid residue with a terminal extended NH₂ group available for hydrogen bonding, while the imidazole group of H is the only ionizable one of all amino acids at neutral pH (pKa 6.3–6.6). Within the coiled-coil region of the Nrf2 protein, H could thus produce charged and electrostatic interactions with other residues. Therefore, it is postulated that the H543Q substitution of the side chain structure may have a potential steric and charge effect on the overall DNA binding ability (Fig. 6C). PolyPhen-2 (<http://genetics.bwh.harvard.edu/pph2/>) also determined that H543Q is predicted to be possibly damaging with a score of 0.895 (sensitivity: 0.82, specificity: 0.94). A Cns SNP L451V detected in strains such as SPRET/EiJ (Supplementary Tables S2 and S3) is located within the helical domain that organizes DNA-contact elements based on structural modeling. While it is difficult from the modeling to suggest the direct effect of the L451V SNP, steric implications on a binding partner are also plausible due to the size difference between leucine and valine.

FIG. 3. Association of hyperoxia susceptibility and *Nrf2* SNP haplotypes. (A) Relative to haplotype 1 (black symbols), mean numbers of total cells were significantly greater in haplotype 2 ($p=0.02$, gray symbols) and haplotype 3 ($p=0.01$, white symbols) mice (one-way ANOVA) after hyperoxia exposure. Pulmonary cellular injury and *Nrf2* haplotypes were significantly correlated across the inbred strains (Pearson correlation coefficient=0.658) ($p=0.006$). Mean $\pm$ SEM ($n=3-5$ /group for air, 4-7/group for hyperoxia) presented. * $p \leq 0.05$ between strain-matched air- and hyperoxia-exposed mice. Bars (air) or symbols (hyperoxia) in black are for haplotype 1, in dark gray for haplotype 2, and white for haplotype 3. Dotted lines separate strains by haplotypes. (B) Percent body weight changes of hyperoxia-exposed mice relative to strain-matched air controls within haplotypes. *, significantly different from strain-matched air-exposed mice ($p \leq 0.05$). Color and patterns of the bar graphs are coordinated by exposure and haplotypes, and dotted lines separate strains by haplotypes. ANOVA, analysis of variance; SEM, standard error of the mean.

Discussion

High-density mapping of mammalian genomes has supported a wide range of genetic investigations. SNP discovery is particularly useful in genetic studies of model organisms such as mice. Tsang *et al.* (49) first compiled 673 SNPs in 55

mouse strains and constructed a phylogenetic tree to correlate and clarify the origins of strains based on the assembled mouse genome sequence and SNP data (51, 54, 55). Currently, millions of mouse SNPs and haplotypes from more than 120 strains have been published and provide a valuable reference for dissecting the genetic basis of complex traits (16, 27, 31).

Use of animals bearing targeted deletion of genes can provide important information regarding their roles in normal and disease status, although it may not contribute to understanding gene regulation. Knowledge on allelic variation of individual genes and association with phenotypes has enhanced understanding of their function in mouse disease models (15). However, *Nrf2* polymorphisms in inbred strains and their correlation to disease risk have received little attention. In this study, we assembled variations in *Nrf2* from public databases and new sequencing analysis, and determined genotype-phenotype associations in a model of oxidative ALI. Our results indicate that murine *Nrf2* sequences vary widely across strains (~ 1 SNP per every 32 bp). Importantly, *Nrf2* SNP haplotypes influenced its function and, at least in part, associated with differential pulmonary susceptibility to hyperoxia. We also found intra-haplotype variations in *Nrf2* expression and transactivation *in vivo*, indicating that polymorphisms in *Nrf2* do not account for all of the variation in susceptibility to hyperoxia in inbred strains. This is not surprising, as ALI is a complex disease with multiple phenotypes that are likely controlled by many genes.

The $-103T/C$ (formerly $-336T/C$) *Nrf2* SNP and simple sequence length polymorphism markers in *Nrf2* (D2Mit248 and D2Mit94) co-segregated with hyperoxia susceptibility in B6C3F₂ mice (7). This promoter SNP added an additional Sp1 binding site or at least increased Sp1 binding affinity in the hyperoxia-susceptible haplotype 2 strains of mice (including B6) compared with hyperoxia-resistant haplotype 1 strains (including C3; see Fig. 4A). In addition, it affected promoter activity in response to hyperoxia.

Although SNPs in exon 5 did not cause complete loss of function, we found an inverse relationship between the severity of injury and *Nrf2* transactivation activity in resistant haplotype 1 strains and susceptible haplotypes 2 and 3. While we did not determine directly whether hyperoxia susceptibility can be attributed in part to the exon 5 SNPs of haplotype 3, the H543Q substitution in the ARE-binding Neh1 domain in haplotype 3 strains has the potential to influence *Nrf2* function. Based on our protein structural modeling, we postulate that the H to Q substitution impacts charge (H can be positively charged depending on pKa), which may influence steric hindrance and side chain helix structure, ultimately impeding on *Nrf2*-DNA binding interactions (Fig. 6). The SNP-function analysis program PolyPhen-2 also predicted that the H543Q SNP is possibly damaging. In support of this possibility, previous studies showed that substitution of amino acid H by Q or *vice versa* modifies structural organization and affects protein activities and disease risk (23, 24, 56).

More than 502 SNPs have been discovered in human *NRF2*, and some of these genetic mutations have been shown to be associated with disease risks (5). *NRF2* promoter SNPs, including -617 (rs6721961), -651 (rs6706649), -653 (rs35652124), and $-684/686$ positions, or in introns (*e.g.*,

FIG. 4. Functional assessment of informative *Nrf2* SNPs *in vitro*. (A) Transactivating activity of the *Nrf2* promoter region bearing the –103 SNP was compared by co-transfection of A549 cells with a vector containing either haplotype 1 (pNrf2_{C3}) or haplotype 2 (pNrf2_{B6}) promoter sequences. The cells were then exposed to either air or hyperoxia (4 h), and luciferase activity was determined and normalized to β -gal. Mean $\pm$ SEM presented ($n=8$ /group). $^*p \leq 0.05$ between hyperoxia and air exposure under same transfection condition; $^+p \leq 0.05$ compared with hyperoxia-exposed pNrf2_{C3}. Aliquots of lung nuclear protein were subjected to gel shift analysis on haplotype 1 (C3 –103) or haplotype 2 (B6 –103) promoter probes, including the –103 SNP to determine the effect of the –103 T/C SNP on potential addition of Sp1 binding site in the haplotype 2 promoter. Nuclear protein binding activities were compared with those on Sp1 consensus sequence for specificity. Representative band image from replicates is presented. (B) Airway epithelial cells (BEAS-2B) overexpressing haplotype 1- or haplotype 3-originated Nrf2 were prepared by transient transfection with either BALB/cJ-Nrf2 or SM/J-Nrf2 expression vector (pFlag-Nrf2_{Hap1} or pFlag-Nrf2_{Hap3}, respectively). ARE and NF-E2 binding activity after hyperoxia exposure was determined by gel shift analysis on nuclear proteins from BEAS-2B cells overexpressing haplotype 1 (pFlag-Nrf2_{Hap1})- or haplotype 3 (pFlag-Nrf2_{Hap3})-originated Nrf2 using a γ P³²-end labeled ARE consensus sequence. FP, free probes; SB, shifted bands. Representative band image from replicates is presented. Vector indicates the binding activity of nuclear proteins transfected with empty vector (pFlag). BEAS-2B cells overexpressing haplotypes 1- or 3-originated Nrf2 were co-transfected with *Nqo1* ARE-luciferase reporter constructs, and exposed to either air or hyperoxia (6 h). Reporter gene activity was measured by a dual luciferase detection system. Vector indicates the activity of cells co-transfected with *Nqo1* ARE-luciferase reporter constructs and empty vector (pFlag). Mean $\pm$ SEM presented ($n=8$ /group). $^*p \leq 0.05$ between hyperoxia and air exposure under same transfection condition; $^+p \leq 0.05$ compared with exposure-matched pFlag-Nrf2_{C3}. ARE, antioxidant response element; NF-E2, nuclear factor, erythroid-derived 2.

rs2364722, rs10183914) were found to be associated with ALI, chronic obstructive pulmonary disorder, asthma, systemic lupus erythematosus, gastric cancer, and Parkinson's disease (2, 13, 32, 33, 50). Interestingly, a unique triplet repeat polymorphism was also identified in the promoter region (between –20 and –6) of Japanese cohorts (57). Non-synonymous coding SNPs have also been found in at least 17 amino-acid residues (Supplementary Table S4), including P233S (17), although they have not been associated with disease susceptibility. Considering the essential role of *Nrf2* in various model diseases, further population studies of *NRF2* in oxidative lung disorders are warranted (10, 11, 38, 48).

Recent investigations have identified somatic (missense) mutations in *NRF2* and Kelch-like ECH-associated protein 1 gene (*KEAP1*), a cytosolic suppressor for NRF2, in human lung adenocarcinoma cells known to be associated with increased risk of neoplasia and chemoresistance (43, 45, 47, 58). Somatic mutations clustered in DLG/ETGE motifs (Neh2 domain) for KEAP1 binding in *NRF2* led to NRF2 accumulation in cancer cells by suppressing its ubiquitination or KEAP1 binding (26, 44). Epigenetic changes on CpG islands resulting in constitutive NRF2 activation (19, 25, 34, 52, 59) also support “pathogenic mutations” of *KEAP1* and *NRF2* in various cancers.

FIG. 5. Functional assessment of SNP haplotypes *in vivo*. (A) Association between lung *Nrf2* mRNA abundance (y axis) assessed by semi-quantitative RT-PCR after air or 48 h hyperoxia exposure and cellular injury susceptibility (x axis) assessed by BAL total cell counts after air or 72 h hyperoxia exposure in 16 strains. Mean $\pm$ SEM ($n=3$ /group) is presented. Dot graphs for individual strains indicate *Nrf2* SNP haplotypes (black-1, gray-2, and white-3). (B) Gel shift analysis demonstrates differential ARE binding activity of nuclear Nrf2 from three haplotypes of mice after air and hyperoxia (48 h) exposure. SB indicates ARE binding activity of lung nuclear proteins (5 μ g), and SSB indicates specific ARE-Nrf2 binding activity. Representative band image from replicates is presented. All lanes were run on the same gel but were noncontiguous. (C) Left; Differential mRNA induction of Nrf2 effector antioxidants (QR1, GPx-2, and GSTA4) in 16 strains of mice from three *Nrf2* haplotypes was determined by quantitative RT-PCR. Average ($n=3$ /group) mRNA abundance of each strain was normalized to that of air-exposed A/J mice. Dot graphs for individual strains indicate Nrf2 SNP haplotypes (black-1, gray-2, and white-3). Middle and right; ChIP analysis followed by semi-quantitative and/or quantitative PCR determined differential binding activity of lung Nrf2-small Maf proteins to AREs of *Nqo1* and *Gpx-2* promoter after hyperoxia (48 h). Mean $\pm$ SEM ($n=3$ /group) presented. * $p \leq 0.05$ different between hyperoxia and strain-matched air control; + $p \leq 0.05$ compared with exposure-matched haplotype 1 mice. (D) Differential lung protein levels of antioxidants in 16 strains of mice were compared after air or hyperoxia (48 h) exposure by Western blotting analysis using an aliquot (60 μ g) of lung lysates and mouse-specific antibodies against QR1, GPx-2, and GSTA1. Actin level was detected as a protein loading control. Representative band image from replicates is presented. All lanes were run on the same gel but were noncontiguous. ChIP, chromatin immunoprecipitation; GPx-2, glutathione peroxidase 2; GSTA4, glutathione S-transferase A4; QR1, quinone reductase 1; RT-PCR, reverse transcriptase-polymerase chain reaction; SSB, super-shifted bands.

FIG. 6. Modeling of three-dimensional structures for bZIP domain of Nrf2 protein. (A) Putative protein structure of the Neh1 domain, including bZIP region (amino-acid residues 490–550 for mouse Nrf2), was modeled based on the published crystal structures of similar bZIP transcription factors, including SKN-1, MafG, and C/EBP beta. Amino-acid residue H543 is predicted to be located in the long helical segment that packs against the other helix of the bZIP structure side chain. (B) bZIP transcription factor domain (amino-acid residues 498–558) of human NRF2 (in yellow) was superimposed on the heterodimeric complex structure of human c-Fos:c-Jun bound to DNA. (C) Close-up images illustrate side chains with H543 or Q543 residues, predicting potential steric/charge effects of the H543Q side chain substitution overall on DNA binding ability of Nrf2 in haplotype 3. bZIP, basic region/leucine zipper; C/EBP, CCAAT/enhancer-binding protein. To see this illustration in color, the reader is referred to the web version of this article at www.liebertpub.com/ars

It is important to note that, in addition to genetic background, other extrinsic (*e.g.*, age) and intrinsic (*e.g.*, gender) factors may influence the function of Nrf2. We selected adult, male mice for this study, because the original investigation that identified *Nrf2* as a susceptibility gene for hyperoxia-induced ALI used B6 and C3 mice with this age and gender (7). However, a role of gender in model ALI induced by hyperoxia has been demonstrated as indicated by differential disease susceptibility (29, 30, 37). We have also found that, although Nrf2 has an important protective role in the neonatal pulmonary response to hyperoxia (11), it was not associated with neonatal susceptibility to hyperoxia as determined in a genome-wide association study with 30 inbred strains of mice (35). Additional investigations are necessary to understand potential gene-by-gene, gene-by-gender, and gene-by-age interactions that may affect *Nrf2* expression and function.

In conclusion, we identified distinct genetic variation in *Nrf2* across multiple inbred strains of mice and SNP haplotypes that were correlated with differential susceptibility to oxidant-induced lung injury. We also used *in vitro* genomic

sequencing and protein structural modeling approaches to characterize novel genetic mutations in *Nrf2* that have functional effects on Nrf2 activity and function, and are likely to contribute to differential susceptibility to oxidative lung injury. An important strength of the current model is that, by investigating variation in *Nrf2* across many genetically diverse strains, it should provide insights into the genetic variation of the human *NRF2* homolog and its role in ALI. Understanding the genetic basis of murine *Nrf2* function and strains bearing *Nrf2* haplotypes, thus, provides a novel means to investigate the role and the regulation of Nrf2 in oxidative disease predisposition.

Materials and Methods

Animals

C3, A/J, DBA/2J, BALB/cJ, LP/J, CBA/J, B6, C57BL/10J, C57L/J, SWR/J, SJL/J, NZW/LACJ, AKR/J, SM/J, P/J, and SEA/GnJ strains of mice (5–7 weeks, male) were purchased from Jackson Laboratory (Bar Harbor, ME). The mice were provided food (NIH-31) and water *ad libitum*. All animal

use was approved by the NIEHS Animal Care and Use Committee.

Inhalation exposure and BAL analysis

Mice were placed individually in a Hazleton M60 battery inside a Hazleton 1000 chamber (Lab Products, Maywood, NJ). After acclimation in the chambers for 24 h/day for 2–3 days, mice were exposed to $\geq 95\%$ O₂ (UHP grade, minimum purity 99.994% O₂ tanks; National Welders, Durham, NC) 24 h/day for 48 or 72 h. Water and feed were provided *ad libitum* during the exposure. The temperature ($72 \pm 3^\circ\text{F}$) and humidity ($50\% \pm 15\%$) of the chambers were monitored. Mice for room-air exposure were placed in a Hazleton M60 battery located on a countertop with food and water provided *ad libitum* for the exposure duration. The chambers were opened for ~ 30 min at the same time each morning to allow animal health checks (morbidity and mortality), water and feed checks, and excreta paper change. Immediately after the end of designated exposure, mice were euthanized by sodium pentobarbital overdose (104 mg/kg). The right lung of each mouse was lavaged *in situ*, four consecutive times with HBSS (0.5 ml/25 g body weight), and the BAL fluid returns were then centrifuged (1000 g, 10 min at 4°C). Supernatants from the first lavage return were analyzed for total protein content to assess lung permeability change, and cells from pooled lavage returns were re-suspended and counted to assess inflammatory cell influx and epithelial cell exfoliation as previously described (6).

Nrf2 SNP discovery and compiling

SNPs in murine *Nrf2*, including the 5 kb upstream promoter region, were compiled for a total of 72 inbred strains from publicly available sources, including the MPD (<http://phenome.jax.org/SNP>) and NCBI dbSNP (www.ncbi.nlm.nih.gov/SNP), which presents SNP records from Celera, ENSEMBL, ABI, etc. (16, 49, 51, 55). *Nrf2* from 16 inbred strains was re-sequenced using mouse *Nrf2*-specific primers. Briefly, genomic DNA was isolated from individual mouse kidney of 16 inbred strains using a DNeasy Blood & Tissue Kit (Qiagen, Inc., Valencia, CA). New SNPs were discovered by re-sequencing of the 1 kb 5'-upstream and coding region of *Nrf2* using a high-fidelity DNA amplification system (Epicentre Technologies, Madison, WI), mouse *Nrf2*-specific primers (4), and ABI-Prism fluorescent methods and reagents on an ABI377 Automated DNA sequencer (Applied Biosystems, Foster City, CA).

RNA isolation and RT-PCR

Total lung RNA was isolated from homogenates using the RNeasy Mini Kit (Qiagen, Inc.). One microgram of total lung RNA was reverse transcribed into cDNA in a 50 μl volume (6). For semi-quantitative PCR of *Nrf2*, cDNA (1.5 or 2.5 μl) equivalent to 50 ng of RNA was amplified in a 25 μl reaction containing 240 nM of forward and reverse primers in a Gene Amp 9700 PCR System (Applied Biosystems), and expression levels were quantified with 18s ribosomal RNA as an internal control following the procedures described elsewhere (6). For quantitative PCR, cDNA equivalent to 20–50 ng of RNA was amplified in a 20 μl volume containing gene-specific primers for GPx-2 (381 forward 5'-tgcaaccagt tcggacatc-3',

531 reverse 5'-aggcaaagacaggatgctc-3'), GSTA4 (Real Time Primers, LLC, Elkins Park, PA), QR1 (1141 forward 5'-agcgagctgg aaaatactct-3', 1310 reverse 5'-ggccattgtta ctttgagc-3'), or 18s (324 forward 5'-tacctggttgatcctgccag-3', 507 reverse 5'-ccgtcgcatgtattagctc-3') and 2 $\times$ Master Mix (Applied Biosystems) using an ABI Prism 7700 Sequence Detection System (Applied Biosystems). The relative quantification of gene expression was calculated from the threshold cycle (C_T) values for each sample and normalized in relation to the expression of 18s rRNA using the comparative C_T method.

Protein isolation

Total proteins were prepared from lung homogenates or cell extracts in radioimmunoprecipitation buffer, including phenylmethanesulfonyl fluoride (PMSF, 10 $\mu\text{g}/\text{ml}$) and protease/phosphatase inhibitor cocktail (Sigma-Aldrich, St. Louis, MO). A Polytron homogenizer was used for lung tissues. Nuclear proteins were prepared from pulverized lung tissues ($n=3/\text{group}$) or harvested cells ($n=3/\text{group}$) following procedures described elsewhere (8). Proteins were quantified and stored in aliquots at -70°C until needed.

Electrophoretic mobility shift assay

Nuclear DNA binding activities were determined by gel shift analyses of nuclear protein aliquots (5 μg , pooled for each group) on 3×10^4 cpm [γ - ^{32}P] ATP end-labeled double-stranded oligonucleotides containing an ARE consensus binding sequence for Nrf2 or Sp1 consensus sequence following the procedure previously described (6). Specific binding activity of Nrf2 was determined by preincubation of nuclear proteins with anti-Nrf2 antisera (9) for 2 h in ice followed by electrophoretic mobility shift assay. The gel was autoradiographed using an intensifying screen at -70°C , and bands were scanned using a Bio-Rad Gel Doc 2000 system (Hercules, PA). Representative band images from two analyses are presented.

Western blot analyses

Proteins (20–100 μg , pooled for each group) were separated on SDS-phage gels and analyzed by Western blotting against Flag (F3165; Sigma-Aldrich), QR1 (sc-16464; Santa Cruz Biotechnology, Inc., Santa Cruz, CA), GPx-2 (ab4501; Abcam, Cambridge, MA), GSTA1 (a gift from Dr. C.C. Reddy, Pennsylvania State University), or pan actin (sc-1615; Santa Cruz Biotechnology, Inc.). Representative band images from two independent blots are presented. The bands were scanned using a Bio-Rad Gel Doc 2000 system.

Plasmid construction

The distal promoter region of the *Nrf2* gene (spanning from -612 to $+228$, GI:149338249) from haplotypes 1 (C3) and 2 (B6), including the -103 SNP, was amplified by PCR using primers bearing restriction enzyme sites (*Sal*I–*Sac*I or *Nhe*I–*Xba*I) 5'-gcgtcgacgagctAGCAACTGAATGGGACC TGG-3' and 5'-gctctagagctagcAGGGCGGACGCTGTG GTA-3'. Each promoter segment was processed for cloning into the *Sac*I and *Nhe*I sites of a pGL3-basic vector (Promega, Madison, WI) containing the SV40 endogenous promoter using Lipofectamine 2000 Reagent (Invitrogen, Grand Island, NY) to create p*Nrf2*-C3 and p*Nrf2*-B6. A mammalian

overexpression vector for murine *Nrf2* was generated by PCR amplification of haplotypes 1 (BALB/cJ) and 3 (P/J) mouse cDNAs using oligonucleotide primers 5'-gggtaccacATGATGGACTTGGGAGTTGCCA-3' and 5'-gctctagaCTCCATCCTCCCGAACCTAGT-3'. cDNAs encoding haplotypes 1 or 3 *Nrf2* were then cloned into the *Kpn1* and *Xba1* sites of p3XFLAG-CMV-10 expression vector (Sigma-Aldrich), respectively (pFlag-*Nrf2*_{HAP1}, pFlag-*Nrf2*_{HAP3}). All plasmid sequences were confirmed by sequencing analysis performed in the NIEHS Sequencing Core facility.

Transfection, in vitro hyperoxia exposure, and reporter gene analyses

Transient transfection was performed in airway epithelial cells with 60%–70% confluency following standard procedures. For promoter function analysis, cells were co-transfected with 100 ng of either p*Nrf2*_{C3} or p*Nrf2*_{B6} along with 1 ng of an internal control vector (pSV- β -galactosidase; Promega). After 4 h exposure to hyperoxia (85 $\pm$ 5%), cells were harvested and aliquots of cell extracts were assayed for luciferase activity. For ARE activation analysis of coding SNPs in exon 5, 100 ng of overexpressing plasmid for *Nrf2* haplotype 1 or 3 (pFlag-*Nrf2*_{HAP1} or pFlag-*Nrf2*_{HAP3}) was co-transfected into BEAS-2B cells (ATCC, Manassas, VA) with 600 ng of an ARE-bearing luciferase expression vector (p*Nqo1*-ARE, a gift from Dr. S.R. Reddy, University of Illinois at Chicago) and *Renilla* luciferase (*Rluc*) vector (pRL-TK; Promega) as an internal normalization control. After exposure to air or hyperoxia (85 $\pm$ 5%, 6 h), aliquots of cell extracts were assayed for firefly and *Rluc* activities using the Dual-Luciferase Reporter Assay System (Promega) in a microplate luminometer (Fluoroskan Ascent FL; Thermo Scientific, Inc., Waltham, MA) according to the manufacturer's protocol. Promoter activity of each group was normalized to that of air-exposed p*Nqo1*-ARE group with no *Nrf2* overexpression, which was assigned a value of 1. All transfections were performed in triplicate, and each experiment was repeated.

ChIP assay

Frozen lung tissues from individual mice were pooled for each experiment group ($n = 3$ lungs/group) and pulverized in liquid nitrogen. Pooled tissues (18 mg/group) were processed for chromatin isolation, immunoprecipitation of protein-DNA cross-links, and DNA purification following the manufacturer's protocol (Magna ChIP G Tissue Kit; EMD Millipore Corporation, Billerica, MA). Antibodies used in immunoprecipitation included anti-*Nrf2* (4 μ g, sc-722x; Santa Cruz Biotechnology, Inc.) and anti-MafF/G/K (4 μ g, sc-22831x; Santa Cruz Biotechnology, Inc.). Normal rabbit IgG (2 or 4 μ g) was used in immunoprecipitation for negative controls. Target DNA (2 μ l, 4% of target) isolated through ChIP or input DNA (2 μ l, 0.25% of total chromatin DNA) was subjected to semi-quantitative (FailSafe PCR Kit; Epicentre) or quantitative PCR (Power SYBR-Green 2 $\times$ Master Mix, ABI) using primers designed for *Gpx-2*-ARE (GI:5305215, –1050 forward: 5'-aacaatcatgacccgga-3', –851 reverse: 5'-ataatgccatccacggt-3') and *Nqo1* ARE (GI:37694057, –516 forward: 5'-gcagtttctaagagcagaac-3', –341 reverse: 5'-gtagattagctcactcagccg-3').

Protein structure modeling

Comparative protein structure modeling was done by combining multiple templates and optimizing sequence-to-structure alignments using ITASSER (60), SwissModel (42), and M4T (14) to develop models for human NRF2/murine *Nrf2* and the optimum consensus models selected for presentation. HHpred (46) was used to provide secondary structure prediction as a check for the molecular structures. Sequence alignments and similarity were checked with the BLAST server at NCBI (1).

Statistics

Hamming distance between each pair of strains was calculated as the number of polymorphic positions within *Nrf2* at which the strains differed. Hierarchical clustering on the distance matrix using Ward's method (<http://stat.ethz.ch/R-manual/R-patched/library/stats/html/hclust.html>) partitioned the strains into haplotype groups. A minimum of three strains were required to form a haplotype group for statistical analyses. Experimental data were expressed as the group mean $\pm$ standard error of the mean. When data were normally distributed and homoscedastic, analysis of variance (ANOVA) was used for all the experimental data, and the Student–Newman–Keuls or Holm–Sidak tests were used for *a posteriori* comparisons of means ($p \leq 0.05$). When data did not meet the normality and skedasticity conditions, we assigned individual data for all strains to quartiles and analyzed for haplotype effects by χ^2 analysis. Correlation analysis (Pearson correlation, Spearman Rank Order) was performed for SNP haplotypes and total BAL cells. All statistical analyses were performed using SigmaStat 3.0 or SigmaPlot 11.0 software program (SPSS Science, Inc., Chicago, IL).

Acknowledgments

This research was supported, in part, by the Intramural Research Program of the National Institute of Environmental Health Sciences. Hyperoxia exposures were conducted at the NIEHS Inhalation Facility under contract to Alion Science and Technology, Inc. The authors thank Dr. Daniel Morgan and Mr. Herman Price for coordinating the exposures. Drs. Don Cook and Michael Fessler at the NIEHS provided an excellent critical review of this article.

Author Disclosure Statement

The authors declare that no competing financial interests exist.

References

1. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, and Lipman DJ. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389–3402, 1997.
2. Arisawa T, Tahara T, Shibata T, Nagasaka M, Nakamura M, Kamiya Y, Fujita H, Yoshioka D, Arima Y, Okubo M, Hirata I, and Nakano H. The influence of promoter polymorphism of nuclear factor-erythroid 2-related factor 2 gene on the aberrant DNA methylation in gastric epithelium. *Oncol Rep* 19: 211–216, 2008.

3. Bowler RP, Barnes PJ, and Crapo JD. The role of oxidative stress in chronic obstructive pulmonary disease. *COPD* 1: 255–277, 2004.
4. Chan K, Lu R, Chang JC, and Kan YW. NRF2, a member of the NFE2 family of transcription factors, is not essential for murine erythropoiesis, growth, and development. *Proc Natl Acad Sci U S A* 93: 13943–13948, 1996.
5. Cho HY. Genomic structure and variation of nuclear factor (erythroid-derived 2)-like 2. *Oxid Med Cell Longev* 2013: 286524, 2013.
6. Cho HY, Jedlicka AE, Reddy SP, Kensler TW, Yamamoto M, Zhang LY, and Kleeberger SR. Role of NRF2 in protection against hyperoxic lung injury in mice. *Am J Respir Cell Mol Biol* 26: 175–182, 2002.
7. Cho HY, Jedlicka AE, Reddy SP, Zhang LY, Kensler TW, and Kleeberger SR. Linkage analysis of susceptibility to hyperoxia. Nrf2 is a candidate gene. *Am J Respir Cell Mol Biol* 26: 42–51, 2002.
8. Cho HY, Morgan DL, Bauer AK, and Kleeberger SR. Signal transduction pathways of tumor necrosis factor—mediated lung injury induced by ozone in mice. *Am J Respir Crit Care Med* 175: 829–839, 2007.
9. Cho HY, Reddy SP, Debiase A, Yamamoto M, and Kleeberger SR. Gene expression profiling of NRF2-mediated protection against oxidative injury. *Free Radic Biol Med* 38: 325–343, 2005.
10. Cho HY, Reddy SP, Yamamoto M, and Kleeberger SR. The transcription factor NRF2 protects against pulmonary fibrosis. *FASEB J* 18: 1258–1260, 2004.
11. Cho HY, van Houten B, Wang X, Miller-Degraff L, Fostel J, Gladwell W, Perrow L, Panduri V, Kobzik L, Yamamoto M, Bell DA, and Kleeberger SR. Targeted deletion of Nrf2 impairs lung development and oxidant injury in neonatal mice. *Antioxid Redox Signal* 17: 1066–1082, 2012.
12. Chow CW, Herrera Abreu MT, Suzuki T, and Downey GP. Oxidative stress and acute lung injury. *Am J Respir Cell Mol Biol* 29: 427–431, 2003.
13. Cordova EJ, Velazquez-Cruz R, Centeno F, Baca V, and Orozco L. The NRF2 gene variant, –653G/A, is associated with nephritis in childhood-onset systemic lupus erythematosus. *Lupus* 19: 1237–1242, 2010.
14. Fernandez-Fuentes N, Madrid-Aliste CJ, Rai BK, Fajardo JE, and Fiser A. M4T: a comparative protein structure modeling server. *Nucleic Acids Res* 35: W363–W368, 2007.
15. Flint J and Eskin E. Genome-wide association studies in mice. *Nat Rev Genet* 13: 807–817, 2012.
16. Frazer KA, Eskin E, Kang HM, Bogue MA, Hinds DA, Beilharz EJ, Gupta RV, Montgomery J, Morenzoni MM, Nilsen GB, Pethiyagoda CL, Stuve LL, Johnson FM, Daly MJ, Wade CM, and Cox DR. A sequence-based variation map of 8.27 million SNPs in inbred mouse strains. *Nature* 448: 1050–1053, 2007.
17. Fukushima-Uesaka H, Saito Y, Maekawa K, Kamatani N, Kajio H, Kuzuya N, Noda M, Yasuda K, and Sawada J. Genetic variations and haplotype structures of transcriptional factor Nrf2 and its cytosolic reservoir protein Keap1 in Japanese. *Drug Metab Pharmacokinet* 22: 212–219, 2007.
18. Glover JN and Harrison SC. Crystal structure of the heterodimeric bZIP transcription factor c-Fos-c-Jun bound to DNA. *Nature* 373: 257–261, 1995.
19. Hanada N, Takahata T, Zhou Q, Ye X, Sun R, Itoh J, Ishiguro A, Kijima H, Mimura J, Itoh K, Fukuda S, and Saijo Y. Methylation of the KEAP1 gene promoter region in human colorectal cancer. *BMC Cancer* 12: 66, 2012.
20. Hudak BB, Zhang LY, and Kleeberger SR. Inter-strain variation in susceptibility to hyperoxic injury of murine airways. *Pharmacogenetics* 3: 135–143, 1993.
21. Itoh K, Chiba T, Takahashi S, Ishii T, Igarashi K, Katoh Y, Oyake T, Hayashi N, Satoh K, Hatayama I, Yamamoto M, and Nabeshima Y. An Nrf2/small Maf heterodimer mediates the induction of phase II detoxifying enzyme genes through antioxidant response elements. *Biochem Biophys Res Commun* 236: 313–322, 1997.
22. Itoh K, Wakabayashi N, Katoh Y, Ishii T, Igarashi K, Engel JD, and Yamamoto M. Keap1 represses nuclear activation of antioxidant responsive elements by Nrf2 through binding to the amino-terminal Neh2 domain. *Genes Dev* 13: 76–86, 1999.
23. Jacob CO, Eisenstein M, Dinanuer MC, Ming W, Liu Q, John S, Quismorio FP, Jr., Reiff A, Myones BL, Kaufman KM, McCurdy D, Harley JB, Silverman E, Kimberly RP, Vyse TJ, Gaffney PM, Moser KL, Klein-Gitelman M, Wagner-Weiner L, Langefeld CD, Armstrong DL, and Zidovetzki R. Lupus-associated causal mutation in neutrophil cytosolic factor 2 (NCF2) brings unique insights to the structure and function of NADPH oxidase. *Proc Natl Acad Sci U S A* 109: E59–E67, 2012.
24. Joosten PH, Hol FA, van Beersum SE, Peters H, Hamel BC, Afink GB, van Zoelen EJ, and Mariman EC. Altered regulation of platelet-derived growth factor receptor- α gene-transcription in vitro by spina bifida-associated mutant Pax1 proteins. *Proc Natl Acad Sci U S A* 95: 14459–14463, 1998.
25. Khor TO, Huang Y, Wu TY, Shu L, Lee J, and Kong AN. Pharmacodynamics of curcumin as DNA hypomethylation agent in restoring the expression of Nrf2 via promoter CpGs demethylation. *Biochem Pharmacol* 82: 1073–1078, 2011.
26. Kim YR, Oh JE, Kim MS, Kang MR, Park SW, Han JY, Eom HS, Yoo NJ, and Lee SH. Oncogenic NRF2 mutations in squamous cell carcinomas of oesophagus and skin. *J Pathol* 220: 446–451, 2010.
27. Kirby A, Kang HM, Wade CM, Cotsapas C, Kostem E, Han B, Furlotte N, Kang EY, Rivas M, Bogue MA, Frazer KA, Johnson FM, Beilharz EJ, Cox DR, Eskin E, and Daly MJ. Fine mapping in 94 inbred mouse strains using a high-density haplotype resource. *Genetics* 185: 1081–1095, 2010.
28. Kurokawa H, Motohashi H, Sueno S, Kimura M, Takagawa H, Kanno Y, Yamamoto M, and Tanaka T. Structural basis of alternative DNA recognition by Maf transcription factors. *Mol Cell Biol* 29: 6232–6244, 2009.
29. Lingappan K, Jiang W, Wang L, Couroucli XI, Barrios R, and Moorthy B. Sex-specific differences in hyperoxic lung injury in mice: implications for acute and chronic lung disease in humans. *Toxicol Appl Pharmacol* 272: 281–290, 2013.
30. Lingappan K, Srinivasan C, Jiang W, Wang L, Couroucli XI, and Moorthy B. Analysis of the transcriptome in hyperoxic lung injury and sex-specific alterations in gene expression. *PLoS One* 9: e101581, 2014.
31. Maddatu TP, Grubb SC, Bult CJ, and Bogue MA. Mouse phenotype database (MPD). *Nucleic Acids Res* 40: D887–E894, 2012.
32. Marzec JM, Christie JD, Reddy SP, Jedlicka AE, Vuong H, Lanken PN, Aplenc R, Yamamoto T, Yamamoto M, Cho HY, and Kleeberger SR. Functional polymorphisms in the transcription factor NRF2 in humans increase the risk of acute lung injury. *FASEB J* 21: 2237–2246, 2007.

33. Masuko H, Sakamoto T, Kaneko Y, Iijima H, Naito T, Noguchi E, Hirota T, Tamari M, and Hizawa N. Lower FEV1 in non-COPD, nonasthmatic subjects: association with smoking, annual decline in FEV1, total IgE levels, and TSLP genotypes. *Int J Chron Obstruct Pulmon Dis* 6: 181–189, 2011.
34. Muscarella LA, Barbano R, D'Angelo V, Copetti M, Coco M, Balsamo T, la Torre A, Notarangelo A, Troiano M, Parisi S, Icolaro N, Catapano D, Valori VM, Pellegrini F, Merla G, Carella M, Fazio VM, and Parrella P. Regulation of KEAP1 expression by promoter methylation in malignant gliomas and association with patient's outcome. *Epi-genetics* 6: 317–325, 2011.
35. Nichols JL, Gladwell W, Verhein KC, Cho HY, Wess J, Suzuki O, Wiltshire T, and Kleeberger SR. Genome-wide association mapping of acute lung injury in neonatal inbred mice. *FASEB J* 28: 2538–2550, 2014.
36. Paz-Elizur T, Krupsky M, Elinger D, Schechtman E, and Livneh Z. Repair of the oxidative DNA damage 8-oxoguanine as a biomarker for lung cancer risk. *Cancer Biomark* 1: 201–205, 2005.
37. Prows DR, Hafertepen AP, Gibbons WJ, Jr., Winterberg AV, and Nick TG. A genetic mouse model to investigate hyperoxic acute lung injury survival. *Physiol Genomics* 30: 262–270, 2007.
38. Rangasamy T, Guo J, Mitzner WA, Roman J, Singh A, Fryer AD, Yamamoto M, Kensler TW, Tudor RM, Georas SN, and Biswal S. Disruption of Nrf2 enhances susceptibility to severe airway inflammation and asthma in mice. *J Exp Med* 202: 47–59, 2005.
39. Reddy NM, Kleeberger SR, Kensler TW, Yamamoto M, Hassoun PM, and Reddy SP. Disruption of Nrf2 impairs the resolution of hyperoxia-induced acute lung injury and inflammation in mice. *J Immunol* 182: 7264–7271, 2009.
40. Reddy NM, Suryanarayana V, Kalvakolanu DV, Yamamoto M, Kensler TW, Hassoun PM, Kleeberger SR, and Reddy SP. Innate immunity against bacterial infection following hyperoxia exposure is impaired in NRF2-deficient mice. *J Immunol* 183: 4601–4608, 2009.
41. Rupert PB, Daughdrill GW, Bowerman B, and Matthews BW. A new DNA-binding motif in the Skn-1 binding domain-DNA complex. *Nat Struct Biol* 5: 484–491, 1998.
42. Schwede T, Kopp J, Guex N, and Peitsch MC. SWISS-MODEL: an automated protein homology-modeling server. *Nucleic Acids Res* 31: 3381–3385, 2003.
43. Shibata T, Kokubu A, Gotoh M, Ojima H, Ohta T, Yamamoto M, and Hirohashi S. Genetic alteration of Keap1 confers constitutive Nrf2 activation and resistance to chemotherapy in gallbladder cancer. *Gastroenterology* 135: 1358–1368, 1368.e1351–e1354, 2008.
44. Shibata T, Ohta T, Tong KI, Kokubu A, Odogawa R, Tsuta K, Asamura H, Yamamoto M, and Hirohashi S. Cancer related mutations in NRF2 impair its recognition by Keap1-Cul3 E3 ligase and promote malignancy. *Proc Natl Acad Sci U S A* 105: 13568–13573, 2008.
45. Singh A, Misra V, Thimmulappa RK, Lee H, Ames S, Hoque MO, Herman JG, Baylin SB, Sidransky D, Gabrielson E, Brock MV, and Biswal S. Dysfunctional KEAP1-NRF2 interaction in non-small-cell lung cancer. *PLoS Med* 3: e420, 2006.
46. Soding J, Biegert A, and Lupas AN. The HHpred interactive server for protein homology detection and structure prediction. *Nucleic Acids Res* 33: W244–W248, 2005.
47. Taguchi K, Motohashi H, and Yamamoto M. Molecular mechanisms of the Keap1-Nrf2 pathway in stress response and cancer evolution. *Genes Cells* 16: 123–140, 2011.
48. Thimmulappa RK, Scollick C, Traore K, Yates M, Trush MA, Liby KT, Sporn MB, Yamamoto M, Kensler TW, and Biswal S. Nrf2-dependent protection from LPS induced inflammatory response and mortality by CDDO-Imidazole. *Biochem Biophys Res Commun* 351: 883–889, 2006.
49. Tsang S, Sun Z, Luke B, Stewart C, Lum N, Gregory M, Wu X, Subleski M, Jenkins NA, Copeland NG, and Munroe DJ. A comprehensive SNP-based genetic analysis of inbred mouse strains. *Mamm Genome* 16: 476–480, 2005.
50. von Otter M, Landgren S, Nilsson S, Celajevic D, Bergstrom P, Hakansson A, Nissbrandt H, Drozdziak M, Bialecka M, Kurzawski M, Blennow K, Nilsson M, Hammarsten O, and Zetterberg H. Association of Nrf2-encoding NFE2L2 haplotypes with Parkinson's disease. *BMC Med Genet* 11: 36, 2010.
51. Wade CM, Kulbokas EJ, 3rd, Kirby AW, Zody MC, Mullikin JC, Lander ES, Lindblad-Toh K, and Daly MJ. The mosaic structure of variation in the laboratory mouse genome. *Nature* 420: 574–578, 2002.
52. Wang R, An J, Ji F, Jiao H, Sun H, and Zhou D. Hypermethylation of the Keap1 gene in human lung cancer cell lines and lung cancer tissues. *Biochem Biophys Res Commun* 373: 151–154, 2008.
53. Ware LB and Matthay MA. The acute respiratory distress syndrome. *N Engl J Med* 342: 1334–1349, 2000.
54. Waterston RH, Lindblad-Toh K, Birney E, Rogers J, Abril JF, Agarwal P, Agarwala R, Ainscough R, Alexandersson M, An P, Antonarakis SE, Attwood J, Baertsch R, Bailey J, Barlow K, Beck S, Berry E, Birren B, Bloom T, Bork P, Botcherby M, Bray N, Brent MR, Brown DG, Brown SD, Bult C, Burton J, Butler J, Campbell RD, Carninci P, Cawley S, Chiaromonte F, Chinwalla AT, Church DM, Clamp M, Clee C, Collins FS, Cook LL, Copley RR, Coulson A, Couronne O, Cuff J, Curwen V, Cutts T, Daly M, David R, Davies J, Delehaunty KD, Deri J, Dermitzakis ET, Dewey C, Dickens NJ, Diekhans M, Dodge S, Dubchak I, Dunn DM, Eddy SR, Elnitski L, Emes RD, Eswara P, Eyraas E, Felsenfeld A, Fewell GA, Flicek P, Foley K, Frankel WN, Fulton LA, Fulton RS, Furey TS, Gage D, Gibbs RA, Glusman G, Gnerre S, Goldman N, Goodstadt L, Grafham D, Graves TA, Green ED, Gregory S, Guigo R, Guyer M, Hardison RC, Haussler D, Hayashizaki Y, Hillier LW, Hinrichs A, Hlavina W, Holzer T, Hsu F, Hua A, Hubbard T, Hunt A, Jackson I, Jaffe DB, Johnson LS, Jones M, Jones TA, Joy A, Kamal M, Karlsson EK, et al. Initial sequencing and comparative analysis of the mouse genome. *Nature* 420: 520–562, 2002.
55. Wiltshire T, Pletcher MT, Batalov S, Barnes SW, Tarantino LM, Cooke MP, Wu H, Smylie K, Santrosyan A, Copeland NG, Jenkins NA, Kalush F, Mural RJ, Glynne RJ, Kay SA, Adams MD, and Fletcher CF. Genome-wide single-nucleotide polymorphism analysis defines haplotype patterns in mouse. *Proc Natl Acad Sci U S A* 100: 3380–3385, 2003.
56. Winkler DD, Schuermann JP, Cao X, Holloway SP, Borchelt DR, Carroll MC, Proeschler JB, Culotta VC, and Hart PJ. Structural and biophysical properties of the pathogenic SOD1 variant H46R/H48Q. *Biochemistry* 48: 3436–3447, 2009.
57. Yamamoto T, Yoh K, Kobayashi A, Ishii Y, Kure S, Koyama A, Sakamoto T, Sekizawa K, Motohashi H, and

- Yamamoto M. Identification of polymorphisms in the promoter region of the human NRF2 gene. *Biochem Biophys Res Commun* 321: 72–79, 2004.
58. Yoo NJ, Kim HR, Kim YR, An CH, and Lee SH. Somatic mutations of the KEAP1 gene in common solid cancers. *Histopathology* 60: 943–952, 2012.
59. Zhang P, Singh A, Yegnasubramanian S, Esopi D, Kom-bairaju P, Bodas M, Wu H, Bova SG, and Biswal S. Loss of Kelch-like ECH-associated protein 1 function in prostate cancer cells causes chemoresistance and radio-resistance and promotes tumor growth. *Mol Cancer Ther* 9: 336–346, 2010.
60. Zhang Y. I-TASSER server for protein 3D structure prediction. *BMC Bioinformatics* 9: 40, 2008.

Address correspondence to:

Dr. Hye-Youn Cho

Laboratory of Respiratory Biology

National Institute of Environmental Health Sciences

National Institutes of Health

111 TW Alexander Drive

Building 101, MD D-201

Research Triangle Park, NC 27709

E-mail: cho2@niehs.nih.gov

Date of first submission to ARS Central, April 7, 2014; date of final revised submission, September 9, 2014; date of acceptance, September 29, 2014.

Abbreviations Used

ALI = acute lung injury
 ANOVA = analysis of variance
 ARDS = acute respiratory distress syndrome
 ARE = antioxidant response element
 BAL = bronchoalveolar lavage
 bp = base pair
 bZIP = basic region/leucine zipper
 C/EBP = CCAAT/enhancer-binding protein
 ChIP = chromatin immunoprecipitation
 Cns = coding-nonsynonymous
 Cs = coding-synonymous
 GPx-2 = glutathione peroxidase 2
 GSTA4 = glutathione S-transferase A4
 HBSS = Hank's balanced salt solution
 kb = kilobase
 KEAP1 = Kelch-like ECH-associated protein 1
 MPD = mouse phenome database
 NF-E2 = nuclear factor-erythroid 2
 NQO1 or QR1 = NAD(P)H:quinone oxidoreductase 1 or quinone reductase 1
 PCRs = polymerase chain reactions
 PDB = protein data bank
 PMSF = phenylmethanesulfonyl fluoride
 ROS = reactive oxygen species
 RT-PCR = reverse transcriptase-polymerase chain reaction
 SDS = sodium dodecyl sulfate
 SEM = standard error of the mean
 SNPs = single nucleotide polymorphisms