Published in final edited form as:

Neuroscience. 2015 February 12; 0: 97–108. doi:10.1016/j.neuroscience.2014.11.042.

Methamphetamine differentially affects BDNF and cell death factors in anatomically defined regions of the hippocampus

Melissa H. Galinato^{1,2,*}, Laura Orio^{3,*}, and Chitra D. Mandyam^{1,2}

¹Committee on the Neurobiology of Addictive Disorders, The Scripps Research Institute, La Jolla, CA, 92037 USA

²Department of Neurosciences, University of California San Diego, La Jolla, CA, 92037, USA

³Dpto. Psicobiología, Facultad Psicología, Universidad Complutense de Madrid, Campus Somosaguas, 28223 Pozuelo de Alarcón, Madrid

Abstract

Methamphetamine exposure reduces hippocampal long-term potentiation (LTP) and neurogenesis and these alterations partially contribute to hippocampal maladaptive plasticity. The potential mechanisms underlying methamphetamine-induced maladaptive plasticity were identified in the present study. Expression of brain-derived neurotrophic factor (BDNF; a regulator of LTP and neurogenesis), and its receptor tropomyosin-related kinase B (TrkB) were studied in the dorsal and ventral hippocampal tissue lysates in rats that intravenously self-administered methamphetamine in a limited access (1 h/day) or extended access (6 h/day) paradigm for 17 days post baseline sessions. Extended access methamphetamine enhanced expression of BDNF with significant effects observed in the dorsal and ventral hippocampus. Methamphetamine-induced enhancements in BDNF expression were not associated with TrkB receptor activation as indicated by phospho (p)-TrkB-706 levels. Conversely, methamphetamine produced hypophosphorylation of NMDA receptor subunit 2B (GluN2B) at Tyr-1472 in the ventral hippocampus, indicating reduced receptor activation. In addition, methamphetamine enhanced expression of anti-apoptotic protein Bcl-2 and reduced pro-apoptotic protein Bax levels in the ventral hippocampus, suggesting a mechanism for reducing cell death. Analysis of Akt, a pro-survival kinase that suppresses apoptotic pathways and pAkt at Ser-473 demonstrated that extended access methamphetamine reduces Akt expression in the ventral hippocampus. These data reveal that alterations in Bcl-2 and Bax levels by methamphetamine were not associated with enhanced Akt expression. Given that hippocampal function and neurogenesis vary in a subregion-specific fashion, where dorsal hippocampus regulates spatial processing and has higher levels of neurogenesis, whereas ventral hippocampus regulates anxiety-related behaviors, these data suggest that methamphetamine self-

© 2014 IBRO. Elsevier Ltd. All rights reserved.

Correspondence: Chitra D. Mandyam, Ph.D., Committee on the Neurobiology of Addictive Disorders, The Scripps Research Institute, 10550 North Torrey Pines Road, SP30-2400, La Jolla, CA 92037 USA; Tel: (858) 784-9039; Fax: (858) 784-2086; cmandyam@scripps.edu.

*equal contribution

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

administration initiates distinct allostatic changes in hippocampal subregions that may contribute to the altered synaptic activity in the hippocampus, which may underlie enhanced negative affective symptoms and perpetuation of the addiction cycle.

Keywords

Self-administration; ventral hippocampus; GluN2B; TrkB; Bcl-2; Bax

Introduction

Methamphetamine abuse takes severe emotional and financial tolls on society, cutting across ages, races, ethnicities, and genders, and causes significant damage to self by producing cognitive impairment (Thompson et al., 2004, SAMHSA, 2008, Price et al., 2011, Weber et al., 2012). Hippocampus-dependent cognitive impairments in methamphetamine-abusing subjects can be attributed to methamphetamine-induced alterations in structural and functional plasticity of hippocampal neurons. For example, human imaging studies show reduced hippocampal volume, particularly gray matter volume, and decreased hippocampal responsiveness in chronic methamphetamine users (Thompson et al., 2004, Kim et al., 2010, Schwartz et al., 2010, Daumann et al., 2011, Nakama et al., 2011, Orikabe et al., 2011, Morales et al., 2012), indicating maladaptive hippocampal networking in methamphetamine-exposed individuals. Postmortem analyses in human brain tissue confirms that chronic methamphetamine use produces neurotoxicity in the hippocampus (Kitamura, 2009, Kitamura et al., 2010), which suggests an association between hippocampal dysfunction and toxicity in methamphetamine addicts.

The behavioral deficits observed in methamphetamine addicts have been demonstrated in preclinical models of binge methamphetamine exposure and methamphetamine self-administration with extended access to the drug. For example, these experimental paradigms produce cognitive dysfunction and memory impairments dependent on the hippocampus (Itoh et al., 1984, Yoshikawa et al., 1991, Yamamoto, 1997, Friedman et al., 1998, Rogers et al., 2008, Recinto et al., 2012). Neurodegeneration and neurotoxicity in the hippocampus is also observed in animal models of binge methamphetamine exposure and methamphetamine self-administration, suggesting a positive correlation between methamphetamine-induced toxicity and methamphetamine-induced behavioral deficits (Commins and Seiden, 1986, Eisch et al., 1996, Schmued and Bowyer, 1997, Mandym et al., 2008). Additional mechanistic studies show that methamphetamine exposure alters the functional and structural plasticity of hippocampal neurons. For example, acute and systemic methamphetamine treatment reduces long-term potentiation (LTP) of CA1 pyramidal neurons through activation of D1 receptors and increases baseline excitatory synaptic transmission (Swant et al., 2010). Alternatively, self-administration of methamphetamine produces robust and long-lasting morphological changes in CA1 neurons (Onaivi et al., 2002, Crombag et al., 2005, Swant et al., 2010). Acute methamphetamine exposure reduces excitability of dentate gyrus neurons, whereas repeated exposure to methamphetamine increases excitability of these neurons (Criado et al., 2000). Furthermore, emerging studies in adult rats self-administering methamphetamine under extended access conditions (6h of

drug access per day) demonstrate that methamphetamine self-administration reduces dentate gyrus neurogenesis, and these effects were relative to the amount of methamphetamine consumed (Mandyam et al., 2008, Yuan et al., 2011). Withdrawal from methamphetamine self-administration produces compensatory changes in dentate gyrus neurogenesis (i.e., the enhanced survival of newly born neurons born after methamphetamine self-administration), and the hypothesis is that these changes in the hippocampus during withdrawal may regulate relapse to methamphetamine seeking (Recinto et al., 2012). These studies demonstrate that methamphetamine exposure (via either experimenter-delivered or self-administration paradigms) produces synaptic maladaptation in the hippocampus that may mediate some of the addiction behaviors dependent on the hippocampus.

In this context, brain-derived neurotrophic factor (BDNF), via interactions with its receptor tropomyosin related kinase B (TrkB), regulates the function of neurons within the mesolimbic dopamine system and other reward regions to modulate the motivation to take drugs, while BDNF also independently affects hippocampal neurogenesis (Lee et al., 2000, Pierce and Bari, 2001, Katoh-Semba et al., 2002, Bolanos and Nestler, 2004, Scharfman et al., 2005, Donovan et al., 2008, Russo et al., 2010). Enhanced BDNF levels in the dentate gyrus positively correlate with increased number of newly born neurons in the dentate gyrus in non-pathological conditions; however, enhancing BDNF in an ischemic brain reduces ischemia-induced neurogenesis (Lee et al., 2000, Katoh-Semba et al., 2002, Larsson et al., 2002, Scharfman et al., 2005), suggesting a critical relationship between the neurogenic niche, BDNF, and neurogenesis in the hippocampus. In the context of addiction, cocaine exposure produces a widespread, but transient, induction of BDNF protein in the nucleus accumbens, prefrontal cortex, ventral tegmental area, and the central and basolateral nuclei of the amygdala (Grimm et al., 2003, Le Foll et al., 2005, Graham et al., 2007), whereas fewer studies have been conducted to examine BDNF protein levels after exposure to amphetamines. For example, BDNF protein expression in the hippocampus is enhanced in juvenile animals after forced administration of neurotoxic doses of amphetamines and the findings demonstrate that these changes in the neurotrophin may relate to the cognitive deficits in methamphetamine experienced animals (Skelton et al., 2007, Grace et al., 2008). In contrast to these studies using noncontingent methamphetamine injections, there are no published studies on the effect of methamphetamine self-administration on hippocampal BDNF levels, and therefore, was the focus of the current study. We also investigated the receptor systems (tropomyosin-related kinase B (TrkB), NMDA receptor subunit 2B (GluN2B)), cell death factors (Bcl2 and Bax) and downstream signaling proteins (Akt, a pro-survival kinase) modulated by BDNF (Lin et al., 1998, Chao, 2003, Almeida et al., 2005).

Methods

Materials and Methods

Animals—Surgical and experimental procedures were carried out in strict adherence to the National Institutes of Health Guide for the Care and Use of Laboratory Animals (NIH publication number 85–23, revised 1996) and approved by the Institutional Animal Care and Use Committee of The Scripps Research Institute. Nineteen adult, male Wistar rats (Charles

River), weighing 200–250 g at the start of the experiment, were housed two per cage in a temperature-controlled vivarium under a reverse light/dark cycle (lights off 8:00 AM–8:00 PM) for at least one week.

Surgery—Twenty-two rats underwent surgery for catheter implantation for intravenous self-administration. Rats were anesthetized with 2–3% of isoflurane mixed in oxygen and implanted with a sterilized silastic catheter (0.64 ID × 1.19 OD mm; Dow Corning Co.) into the right jugular vein under aseptic conditions. The distal end of the catheter was threaded under the skin to the back of the rat and exited the skin via a metal guide cannula (22G, Plastic One, Inc.). Immediately after surgery, Flunixin® (2.5 mg/kg, s.c.; Bimeda – MTC Animal Health Inc) was given as analgesic. The rats were subjected to antibiotic therapy with Timentin® (20 mg, i.v, once daily; SmithKline Beecham) during 10 days after the surgeries, and sulfamethoxazoletrimethoprim oral suspension (TMS, Hi-Tech Pharmacal Co., Inc.) in the drinking water (0.48 mg/ml) when health complications appeared during the study. Catheters were flushed daily with heparinized saline (30USPunits/ml) and tested eventually for patency using methohexital sodium (Brevital®, 10mg/ ml, 2 mg/rat; King Pharmaceutical Inc.). Eight rats did not experience methamphetamine. Three of the eight rats were trained to self-administer saline (under FR 1 schedule with each active lever delivering 90–100 ul of sterile saline) and experienced saline self-administration for 1 h for the same number of sessions as methamphetamine rats. Fourteen rats experienced methamphetamine self-administration.

Baseline training sessions and maintenance on an extended access schedule

—Four to five days after surgery rats ($n = 14$) were trained to press a lever according to an FR 1 schedule of methamphetamine reinforcement (0.05 mg/kg/injection of methamphetamine hydrochloride, generously provided by the National Institute on Drug Abuse) in operant boxes (Med Associates) under baseline (acquisition) conditions (1h access per day for six-seven days). During daily sessions, a response on the active lever resulted in a 4 second infusion (90–100 μ l volume), followed by a 20 second time-out period to prevent overdose. Each infusion was paired for 4 seconds with white stimulus light over the active lever (conditioned stimulus [CS]). Response during the time-out or on the inactive lever was recorded but resulted in no programmed consequences. All animals were housed on a reverse cycle (lights off at 8 am) and were transferred from their home cages to their operant chambers between 9 and 10 am. Training on the first and second day was initiated with two-three priming (non-contingent) infusions of methamphetamine during the first ten minutes. Rats were allowed to respond for the remaining fifty minutes without any additional priming. Acquiring methamphetamine self-administration was defined as maintenance of similar number of infusions over three consecutive days during baseline training sessions. All animals acquired methamphetamine self-administration (data not shown). After baseline (acquisition) training, the rats were subjected to a short (ShA, $n = 8$, 1h per day under an FR1 schedule) or long (LgA, $n = 6$, 6h per day under an FR1 schedule) access schedule of methamphetamine reinforcement. Methamphetamine self-administration was performed for 17 days.

Western blotting—Procedures optimized for measuring levels of both phosphoproteins and total proteins were employed (Graham et al., 2007, Orio et al., 2009, Edwards et al., 2013). Methamphetamine-naïve rats ($n = 8$), ShA ($n = 8$) and LgA ($n = 6$) rats were killed via rapid decapitation under light isoflurane anesthesia 16–20 h after the last self-administration session. Brains were quickly removed and flash-frozen. Tissue punches enriched in dorsal hippocampus (−3.12 to −4.44 mm from bregma) or ventral hippocampus (−5.40 to −6.12 mm from bregma) from 500 μ m thick sections were homogenized on ice by sonication in buffer (320 mM sucrose, 5 mM HEPES, 1 mM EGTA, 1 mM EDTA, 1% SDS, with Protease Inhibitor Cocktail and Phosphatase Inhibitor Cocktails II and III diluted 1:100; Sigma), heated at 100 degrees C for five minutes, and stored at −80 degrees C until determination of protein concentration by a detergent-compatible Lowry method (Bio-Rad).

Mature BDNF protein levels were determined in 20 μ g protein samples (mixed 1:1) with a Tricine sample buffer containing β -mercaptoethanol) subjected to SDS-PAGE (15% acrylamide) using a Tris-Tricine-SDS buffer (Bio-Rad), followed by electrophoretic transfer to polyvinylidene fluoride membranes (PVDF pore size 0.2 μ m). TrkB, pTrkB, GluN2B, pGluN2B, Bcl-2 and Bax protein levels were determined in 20–30 μ g protein samples (mixed 1:1) with a Laemmli sample buffer containing β -mercaptoethanol) subjected to SDS-PAGE (8–12% acrylamide) using a Tris-Glycine-SDS buffer (Bio-Rad), followed by electrophoretic transfer to PVDF membranes. Blots were blocked with 5% milk (w/v) in TBST (25 mM Tris-HCl (pH 7.4), 150 mM NaCl and 0.1% Tween 20 (v/v)) for 1 h at room temperature and were incubated with the primary antibody for 16–20 h at 4 °C: antibody to BDNF (1:200, Santa Cruz cat. no. sc-546, predicted molecular weight 14 kDa, observed band between 15–20 kDa), TrkB (1:200, Santa Cruz cat. no. sc-8316, predicted molecular weight 95–145 kDa, observed band ~130 kDa), pTrkB Tyr-706 (1:200, Santa Cruz cat. no. sc-8316, predicted molecular weight 95–145 kDa, observed band ~95 kDa), GluN2B (1:200, Santa Cruz cat. no. sc-9057, predicted molecular weight 178 kDa, observed band ~180 kDa), antibody to pGluN2B Tyr-1472 (1:200, Cell Signaling cat. no. 4208S, predicted molecular weight 190 kDa, observed band ~180 kDa), Bcl-2 (1:500, R&D Systems cat. no. MAB8272, predicted molecular weight 24 kDa, observed band ~25 kDa), Bax (1:500, Santa Cruz cat. no. sc-493, predicted molecular weight 23 kDa, observed band ~20 kDa), antibody to Akt (1:500, Cell Signaling cat. no. 4691S, predicted molecular weight 60 kDa, observed band ~60 kDa), antibody to pAkt Ser-473 (1:500, Cell Signaling cat. no. 4060S, predicted molecular weight 60 kDa, observed band ~60 kDa). Blots were then washed three times for 15 min in TBST, and then incubated for 1 h at room temperature (24 °C), with horseradish peroxidase-conjugated goat antibody to rabbit (1:2,000, BioRad) in TBST. After another three washes for 15 min with TBST, immunoreactivity was detected using SuperSignal West Dura chemiluminescence detection reagent (Thermo Scientific) and collected using HyBlot CL Autoradiography film (Denville Scientific) and a Kodak film processor. Net intensity values were determined using the Image Studio Lite (version 3.1). For normalization purposes, membranes were incubated with 0.125% coomassie stain for 5 minutes and washed three times for 5–10 minutes in destain solution.

Statistical analysis—The methamphetamine and saline self-administration data is expressed as active lever presses and methamphetamine data is also expressed as mean

mg/kg methamphetamine self-administered per session. The effect of session duration on methamphetamine self-administration during the 1h session, 6h session and during the first hour of the 6h session was examined over the 17 escalation sessions using a two-way repeated-measures analysis of variance (ANOVA; session duration $\times$ daily session) followed by the Fisher's Least Significant Difference (LSD) post hoc test or Student-Newman-Keuls *post hoc* test. The pattern of responding for methamphetamine is expressed as the mean mg/kg per hour over 6 h sessions in LgA rats and were compared between the first and $> 10^{\text{th}}$ escalation sessions. Differences in the rate of responding between the first and other escalation sessions were evaluated using the paired *t*-test. Differences in the rate of responding between the first and other saline sessions were evaluated using one-way repeated-measures ANOVA. Western blotting analysis was conducted separately for four groups, nonsaline controls, saline controls, limited access and extended access methamphetamine. Because protein levels were not significantly different between nonsaline and saline controls they were combined and were analyzed as methamphetamine naïve controls. Differences in density of proteins were analyzed by split-plot ANOVAs with methamphetamine access as a between-subjects factor and hippocampal subregion as a within-subject factor. Following significant omnibus effects ($p < 0.05$), pairwise comparisons used Student's *t*-tests for split-plot ANOVAs. Data are expressed as mean $\pm$ SEM and were analyzed using SPSS or GraphPad Prism. Values of $p < 0.05$ were considered statistically significant. Graphs were generated using GraphPad Prism 5.0 software.

Results

Extended access to methamphetamine self-administration resulted in escalation of methamphetamine intake

Rats experienced saline or methamphetamine for 17 days (Figure 1a). Repeated measures ANOVA did not detect a change in saline intake in control animals (Figure 1c). Repeated measures two-way ANOVA detected a significant Days of Self-Administration $\times$ Methamphetamine Access interaction (Figure 1c, d, $F_{16,192} = 6.323$, $p < 0.0001$) and an increase in methamphetamine intake across 17 sessions when analyzing both the first hour of the session (Figure 1d, $F_{16,80} = 2.684$, $p < 0.01$) and the total 6 h session ($F_{16,80} = 4.136$, $p < 0.0001$) in LgA animals. *Post hoc* analysis revealed a significant increase in methamphetamine intake during the 6h session in LgA animals during Day 13–17 compared with Days 1–2. Repeated measures ANOVA did not detect a change in methamphetamine intake for ShA animals (Figure 1c, d).

Limited and extended access methamphetamine self-administration increase BDNF expression but do not alter expression of TrkB or the phosphorylation of TrkB at Tyr-706

To determine whether limited access and extended access methamphetamine self-administration alters BDNF expression in the dorsal and ventral hippocampus, Western blot analyses were conducted on hippocampal protein lysates for methamphetamine naïve control animals, LgA, and ShA animals sacrificed 16–20h after the last methamphetamine self-administration session. Protein levels of BDNF (mature form) were analyzed in the dorsal and ventral hippocampus-enriched regions in control, LgA and ShA animals (Figure

1b). Data were subjected to ANOVA to examine the effects of treatment (control vs. methamphetamine groups) across hippocampal subregion (dorsal vs. ventral). ANOVA detected a significant main effect of methamphetamine access on BDNF expression ($F_{2,38} = 3.7$, $p = 0.03$), indicating that methamphetamine experience enhanced BDNF protein expression in the hippocampus. Unpaired $-t$ -test demonstrated significant increases in BDNF protein levels in the LgA animals in dorsal and ventral hippocampus compared with drug naïve controls (p 's = 0.02; Figure 2).

Additionally, we determined whether methamphetamine self-administration altered total TrkB receptor expression and pTrkB expression in the dorsal and ventral hippocampus. Protein levels of TrkB (total and pTrkB at Tyr-706) were analyzed in the dorsal and ventral hippocampus-enriched regions in drug naïve controls, LgA and ShA animals. Data was subjected to ANOVAs to examine the effects of treatment (control vs. methamphetamine groups) across hippocampal subregion (dorsal vs. ventral) on total TrkB and pTrkB/TrkB expression. ANOVA did not detect a significant main effect of methamphetamine or hippocampal region on total TrkB and pTrkB/TrkB expression (Figure 3).

Extended access methamphetamine self-administration results in hypophosphorylation of GluN2B at Tyr-1472 in the ventral hippocampus

To determine whether methamphetamine self-administration altered total GluN2B receptor expression and pGluN2B expression in the dorsal and ventral hippocampus, protein levels of GluN2B (total and pGluN2B at Tyr-1472) were analyzed in the dorsal and ventral hippocampus-enriched regions in drug naïve control, LgA and ShA animals. Data was subjected to ANOVAs to examine the effects of treatment (control vs. methamphetamine groups) across hippocampal subregion (dorsal vs. ventral) on total and pGluN2B/GluN2B expression. ANOVA did not detect a significant main effect of methamphetamine or hippocampal region on total GluN2B expression. ANOVA showed significant main effect of methamphetamine access on pGluN2B/GluN2B ratio ($F_{2,38} = 3.2$, $p = 0.05$). Unpaired $-t$ -test demonstrated significant reduction in pGluN2B/GluN2B ratio in the ventral hippocampus in LgA animals compared with drug naïve controls ($p = 0.008$; Figure 4).

Both limited and extended access methamphetamine self-administration increased anti-apoptotic protein levels and reduced pro-apoptotic protein levels in the ventral hippocampus

We next determined whether methamphetamine self-administration alters expression of pro-apoptotic and anti-apoptotic factors in the dorsal and ventral hippocampus. Protein levels of Bcl-2 (anti-apoptotic) and Bax (pro-apoptotic) were analyzed in the dorsal and ventral hippocampus-enriched regions in controls, LgA and ShA animals. Data were subjected to two-way ANOVAs to examine the effects of treatment (control vs. methamphetamine groups) across hippocampal subregion (dorsal vs. ventral) on pro-and anti-cell death factor expression. ANOVA detected a significant effect of hippocampal subregion on Bcl-2 expression ($F_{1,34} = 4.8$, $p = 0.03$; Figure 5). Unpaired $-t$ -test detected an increase in Bcl-2 protein levels in the ventral hippocampus of LgA rats ($p = 0.04$) compared with controls. ANOVA detected a significant access $\times$ hippocampal subregion interaction ($F_{2,34} = 6.1$, $p = 0.001$), effect of hippocampal subregion ($F_{1,34} = 24.7$, $p < 0.001$), and effect of

methamphetamine ($F_{2,34} = 4.4$, $p = 0.02$; Figure 5) on Bax expression. Post hoc analysis revealed a significant decrease in Bax protein levels in the ventral hippocampus of LgA animals ($p < 0.0001$) and ShA animals ($p = 0.01$) compared with controls.

Extended access methamphetamine self-administration reduces the expression of Akt in the ventral hippocampus

To determine whether methamphetamine self-administration altered total Akt expression and pAkt expression in the ventral hippocampus, protein levels of Akt (total and pAkt at Ser-473) were analyzed in the ventral hippocampus-enriched regions in drug naïve control, LgA and ShA animals. Data was subjected to one-way ANOVAs to examine the effects of treatment (control vs. methamphetamine groups) on total and pAkt/tAkt expression. ANOVA showed significant main effect of methamphetamine access on tAkt ($F_{2,21} = 4.29$, $p = 0.02$). Dunnett's multiple comparison test demonstrated significant reduction in tAkt in the ventral hippocampus in LgA animals compared with drug naïve controls ($p < 0.05$; Figure 6). ANOVA did not detect a significant main effect of methamphetamine on pAkt or p/tAkt expression.

Discussion

Functional dissociation exists along the dorsal-ventral gradient in the rat hippocampus. For example, the ventral hippocampus when compared with the dorsal hippocampus has greater output connections with the prefrontal cortex and bed nucleus of the stria terminalis, and also has extensive reciprocal connections with the amygdala (Henke, 1990, Pitkanen et al., 2000, Ishikawa and Nakamura, 2006), suggesting that neuroadaptations in the ventral hippocampus may be strongly associated with negative affective symptoms that promote drug dependence (Koob, 2003). Indeed, partial lesion and regional inactivation studies have proposed that the dorsal hippocampus is vital for spatial learning, whereas the ventral hippocampus is involved in regulating fear and anxiety (Moser et al., 1995, Kjelstrup et al., 2002, Bannerman et al., 2004, Pothuizen et al., 2004, Pentkowski et al., 2006). In the context of drug-seeking behaviors, the dorsal hippocampus is particularly critical for acquisition of drug memories and reconsolidation of reactivated drug-related associative memories (Meyers et al., 2006, Wells et al., 2011), suggesting that the act of drug taking and seeking produces neuroadaptive changes in the dorsal hippocampus that allow for long-term memory storage of drug-context memories, and allow the recall of associative memories during relapse. The ventral hippocampus, however, is important for reinstatement of drug-seeking behavior triggered by drug contexts, drug cues, or the drug itself (Rogers and See, 2007, Lasseter et al., 2010, Deschaux et al., 2012), suggesting that neuroadaptive changes in the ventral hippocampus contribute to the motivational state that drives drug-seeking behavior during abstinence. Given the distinct, yet supporting roles of the dorsal and ventral hippocampal regions in addiction, we examined the specific proteins implicated in hippocampal neuroplasticity in dorsal and ventral hippocampus-enriched tissue lysates.

Synaptic plasticity, and consequently memory formation, requires new protein synthesis (Kandel, 2001, Bruehl-Jungerman et al., 2007, Costa-Mattioli et al., 2009). Endogenous pro-BDNF is the precursor for mature BDNF and undergoes intracellular processing to be

rapidly converted to the mature form, indicating that pro-BDNF is a transient biosynthetic intermediate (Matsumoto et al., 2008). BDNF, in its mature form is a prominent neurotrophin, mediating many enduring changes in neuroplasticity and synaptogenesis; it has been shown to play a key role as mediator of activity-induced LTP in the hippocampus (Bramham and Messaoudi, 2005, Lu et al., 2008, Minichiello, 2009, Park and Poo, 2013, Leal et al., 2014). Upon high frequency stimulation, endogenous hippocampal BDNF (is generated and) is secreted from axon terminals (presynaptic site) and dendrites (postsynaptic site) of glutamatergic principal neurons (hippocampal granule cells and pyramidal cells; (Lu, 2003), in a calcium-dependent and GluN2B-dependent manner (Hartmann et al., 2001, Balkowiec and Katz, 2002, Gartner and Staiger, 2002, Aicardi et al., 2004). Emerging evidence demonstrates that elevated endogenous BDNF is also required for the maintenance of high frequency stimulation-associated late-phase hippocampal LTP (Lu et al., 2008), suggesting that BDNF levels are low under normal conditions and are elevated by stimuli that enhance hippocampal activity. In this context, chronic methamphetamine exposure enhances baseline hippocampal CA1 activity, increases excitability of dentate gyrus neurons, and increases paired pulse facilitation, and these alterations result in occlusion of hippocampal LTP development (Criado et al., 2000, Onaivi et al., 2002, Ishikawa et al., 2005, Hori et al., 2010, Swant et al., 2010, North et al., 2013). Therefore, it can be hypothesized that methamphetamine induces the same neurobiological changes in the hippocampus that are required for generation and maintenance of hippocampal synaptic activity and additional activation as a consequence of tetanic stimulations in the hippocampus of methamphetamine experienced animals may not produce LTP (Hope, 1998, Schulz et al., 1999, Onaivi et al., 2002).

In the context of the above hypothesis, our findings demonstrate that extended access methamphetamine self-administration enhances BDNF expression in the dorsal and ventral hippocampus that may result in a modification of the characteristics of synaptic transmission in the dorsal and ventral hippocampus (Scharfman et al., 2005). This finding is consistent with previous studies in amphetamine-experienced animals, in which drug exposure (experimenter-delivered) significantly enhanced BDNF protein levels in the hippocampus (Skelton et al., 2007, Grace et al., 2008), but significantly extend the findings with methamphetamine self-administration, a paradigm that produces much lower levels of drug intake compared with high dose experimenter delivered paradigms. Upon release into the synaptic cleft, BDNF can bind to its receptor TrkB, localized at both pre- and postsynaptic sites of glutamatergic synapses in the hippocampus, and BDNF binding to TrkB induces receptor dimerization and phosphorylation at the autophosphorylation site Tyr-706 (Drake et al., 1999, Chao, 2003). Further analysis revealed that enhanced BDNF expression in the hippocampus in extended access animals did not concurrently alter expression of total TrkB or pTrkB-706. While the lack of effect of methamphetamine on total TrkB is consistent with previous studies using amphetamine treatments (Meredith et al., 2002, McGinty et al., 2011), a possible explanation for no effect on pTrkB-706 could be that BDNF-mediated TrkB activation occurs in a time-dependent manner following BDNF release. For example, TrkB phosphorylation was not evident 15 min after a single amphetamine injection but was detected 2 h later, suggesting that there is a delicate relationship between BDNF release and TrkB activation (McGinty et al., 2011). Thus, it is possible that TrkB activation occurs at a

different time point compared with the one used for the current study and should be further investigated. It is also possible that neuroadaptations have occurred in the hippocampus following methamphetamine self-administration, such that the typical neuronal activity-induced secretion of pro-BDNF and conversion of pro-BDNF to mature BDNF, and TrkB phosphorylation via BDNF release is hampered and therefore hinders hippocampal synaptic activity (LTP) that is dependent on TrkB activation (Aloyz et al., 1999, Binder et al., 1999, Pang et al., 2004, Nagappan and Lu, 2005).

BDNF release in the hippocampus also enhances tyrosine phosphorylation of GluN2B subunits at Tyr-1472 of the NMDA receptor via TrkB-mediated phosphorylation, and this constitutes a critical step in mediating BDNF stimulation of synaptic transmission (Lin et al., 1998, Caldeira et al., 2007). Furthermore, the increase in GluN2B tyrosine phosphorylation elicited by BDNF is comparable to that seen after hippocampal LTP induction (Rosenblum et al., 1996, Rostas et al., 1996), suggesting that BDNF-stimulated phosphorylation of GluN2B may provide a mechanism for trophic modulation of hippocampal LTP. The present results demonstrate that extended access methamphetamine self-administration reduces phosphorylation of GluN2B at Tyr-1472 in the ventral hippocampus. Functional changes in NMDARs are significantly linked with phosphorylation and dephosphorylation of GluN2B subunits (Ali and Salter, 2001, Nakazawa et al., 2001, Nakazawa et al., 2006), with phosphorylation of GluN2B at Tyr-1472 regulating internalization of NMDARs via clathrin-mediated endocytosis, such that phosphorylation suppresses NMDAR endocytosis and dephosphorylation enhances NMDAR endocytosis (Lavezzari et al., 2003, Prybylowski et al., 2005). Dephosphorylation of GluN2B by methamphetamine may thus prevent neuronal activity-induced potentiation of LTP in the hippocampus. Our observations add to the growing body of evidence indicating that modification of neuronal networks in the hippocampus by methamphetamine may play a role in altering hippocampal plasticity, which may partly underlie dysfunctional hippocampal-dependent behaviors.

Emerging evidence supports the beneficial role of BDNF against excitotoxicity-induced cell death in the hippocampus (Almeida et al., 2005). For example, BDNF prevents nitrous oxide (NO)-induced cell death in hippocampal neurons (Mattson et al., 1995, Kume et al., 1997), and these effects are mediated via BDNF-induced activation of extracellular signal-regulated kinase (ERK) 1/2 and phosphatidylinositol-3-kinase (PI3K)/Akt pathways (Skaper et al., 1998, Hetman et al., 1999, Han and Holtzman, 2000, Almeida et al., 2005). Notably, hypophosphorylation of GluN2B at Tyr-1472 also reduces excitotoxicity-induced cell death in cortical neurons and these effects could be mediated via Akt activation (Chalecka-Franaszek and Chuang, 1999, Hashimoto et al., 2002). Furthermore, Akt activation inhibits neuronal apoptosis by reducing the expression of pro-apoptotic protein Bax and enhancing the expression of anti-apoptotic protein Bcl-2 (Datta et al., 1997, del Peso et al., 1997), suggesting that Akt activation by BDNF is critical for BDNF-induced increases in Bcl-2 and reduction in Bax (Almeida et al., 2005). In this context, methamphetamine has been reported to increase Akt activity in cultured hippocampal neurons and reduce apoptosis in these neurons (Rau et al., 2011), suggesting enhancement of neuronal survival in the hippocampus. Our findings extend this *in vitro* study to demonstrate that limited and extended access methamphetamine self-administration increases Bcl-2 and reduces Bax

expression in the ventral hippocampus, suggesting enhancement of survival of hippocampal neurons in the ventral hippocampus. These data also suggest that alterations in cell death proteins by methamphetamine are not predicted by the amount of methamphetamine consumed. Analysis of Akt expression and Akt activity via pAkt expression suggests that methamphetamine self-administration induced increases in Bcl-2 and reduction in Bax expression in the ventral hippocampus did not correlate with increases in Akt expression, whereas, these changes in the extended access animals were associated with reduced Akt expression. One of the most compelling explanations for the differences observed in our study versus the *in vitro* study is that we examined the effects of methamphetamine under self-administration conditions. It is intriguing to consider the possibility that the effects of methamphetamine on Akt expression observed here are neurotoxic and are not associated with neuroprotection. Indeed, the Akt signaling cascade via phosphorylation at Ser-473 has been identified as being crucial to the induction of protein synthesis-dependent synaptic plasticity required for hippocampus-dependent learning and memory processes (Opazo et al., 2003, Shehadah et al., 2010, Zhang et al., 2010), and a reduction in Akt expression in extended access animals suggests reduction in synaptic plasticity (Swant et al., 2010). Therefore, the ability of methamphetamine to enhance BDNF expression, produce hypophosphorylation of GluN2B, together with increases in Bcl-2 and reductions in Bax and Akt, suggests that methamphetamine-induced activation of signaling in the hippocampus may function to transduce the BDNF-initiated survival signal in the ventral hippocampal neurons to assist with maladaptive hippocampal plasticity that contributes to the cognitive and motivational states that drive methamphetamine addiction. Future mechanistic studies with these candidate proteins will determine their involvement in the behavioral deficits that are associated with extended access methamphetamine self-administration.

Acknowledgements

The study was supported by funds from the National Institute on Drug Abuse (DA022473 and DA034140 to CDM), National Institute on Alcoholism and Alcohol Abuse (AA020098 and AA06420 to CDM) and Alcohol Beverage Medical Research Foundation to CDM, National Science Foundation (DGE-1144086) to MG, and Fulbright Award (FU-2006-0200) to LO. We appreciate the technical support of Robert Lintz, Ilham Polis and Yanabel Grant for assistance with intravenous self-administration studies. We thank Mathew Soleiman from the independent study program at UCSD for assistance with tissue processing. We thank Drs. Scott Edwards (LSUHSC-New Orleans) and Miranda Staples and Ms. Rosa Ojeda (NIDA summer intern) for assistance with brain tissue punching, tissue processing and immunoblotting. We thank Drs. Scott Edwards and Sucharita Somkuwar for critical reading of the manuscript and McKenzie Fannon for editorial assistance. This is publication number 27060 from The Scripps Research Institute.

References

- Aicardi G, Argilli E, Cappello S, Santi S, Riccio M, Thoenen H, Canossa M. Induction of long-term potentiation and depression is reflected by corresponding changes in secretion of endogenous brain-derived neurotrophic factor. *Proc Natl Acad Sci U S A*. 2004; 101:15788–15792. [PubMed: 15505222]
- Ali DW, Salter MW. NMDA receptor regulation by Src kinase signalling in excitatory synaptic transmission and plasticity. *Curr Opin Neurobiol*. 2001; 11:336–342. [PubMed: 11399432]
- Almeida RD, Manadas BJ, Melo CV, Gomes JR, Mendes CS, Graos MM, Carvalho RF, Carvalho AP, Duarte CB. Neuroprotection by BDNF against glutamate-induced apoptotic cell death is mediated by ERK and PI3-kinase pathways. *Cell Death Differ*. 2005; 12:1329–1343. [PubMed: 15905876]
- Aloyz R, Fawcett JP, Kaplan DR, Murphy RA, Miller FD. Activity-dependent activation of TrkB neurotrophin receptors in the adult CNS. *Learn Mem*. 1999; 6:216–231. [PubMed: 10492004]

- Balkowiec A, Katz DM. Cellular mechanisms regulating activity-dependent release of native brain-derived neurotrophic factor from hippocampal neurons. *J Neurosci*. 2002; 22:10399–10407. [PubMed: 12451139]
- Bannerman DM, Rawlins JN, McHugh SB, Deacon RM, Yee BK, Bast T, Zhang WN, Pothuisen HH, Feldon J. Regional dissociations within the hippocampus--memory and anxiety. *Neurosci Biobehav Rev*. 2004; 28:273–283. [PubMed: 15225971]
- Binder DK, Routbort MJ, McNamara JO. Immunohistochemical evidence of seizure-induced activation of trk receptors in the mossy fiber pathway of adult rat hippocampus. *J Neurosci*. 1999; 19:4616–4626. [PubMed: 10341259]
- Bolanos CA, Nestler EJ. Neurotrophic mechanisms in drug addiction. *Neuromolecular Med*. 2004; 5:69–83. [PubMed: 15001814]
- Bramham CR, Messaoudi E. BDNF function in adult synaptic plasticity: the synaptic consolidation hypothesis. *Prog Neurobiol*. 2005; 76:99–125. [PubMed: 16099088]
- Bruel-Jungerman E, Davis S, Laroche S. Brain plasticity mechanisms and memory: a party of four. *Neuroscientist*. 2007; 13:492–505. [PubMed: 17901258]
- Caldeira MV, Melo CV, Pereira DB, Carvalho RF, Carvalho AL, Duarte CB. BDNF regulates the expression and traffic of NMDA receptors in cultured hippocampal neurons. *Mol Cell Neurosci*. 2007; 35:208–219. [PubMed: 17428676]
- Chalecka-Franaszek E, Chuang DM. Lithium activates the serine/threonine kinase Akt-1 and suppresses glutamate-induced inhibition of Akt-1 activity in neurons. *Proc Natl Acad Sci U S A*. 1999; 96:8745–8750. [PubMed: 10411946]
- Chao MV. Neurotrophins and their receptors: a convergence point for many signalling pathways. *Nat Rev Neurosci*. 2003; 4:299–309. [PubMed: 12671646]
- Commings DL, Seiden LS. alpha-Methyltyrosine blocks methylamphetamine-induced degeneration in the rat somatosensory cortex. *Brain Res*. 1986; 365:15–20. [PubMed: 3947981]
- Costa-Mattioli M, Sossin WS, Klann E, Sonenberg N. Translational control of long-lasting synaptic plasticity and memory. *Neuron*. 2009; 61:10–26. [PubMed: 19146809]
- Criado JR, Gombart LM, Huitron-Resendiz S, Henriksen SJ. Neuroadaptations in dentate gyrus function following repeated methamphetamine administration. *Synapse*. 2000; 37:163–166. [PubMed: 10881037]
- Crombag HS, Gorny G, Li Y, Kolb B, Robinson TE. Opposite effects of amphetamine self-administration experience on dendritic spines in the medial and orbital prefrontal cortex. *Cereb Cortex*. 2005; 15:341–348. [PubMed: 15269111]
- Datta SR, Dudek H, Tao X, Masters S, Fu H, Gotoh Y, Greenberg ME. Akt phosphorylation of BAD couples survival signals to the cell-intrinsic death machinery. *Cell*. 1997; 91:231–241. [PubMed: 9346240]
- Daumann J, Koester P, Becker B, Wagner D, Imperati D, Gouzoulis-Mayfrank E, Tittgemeyer M. Medial prefrontal gray matter volume reductions in users of amphetamine-type stimulants revealed by combined tract-based spatial statistics and voxel-based morphometry. *Neuroimage*. 2011; 54:794–801. [PubMed: 20817105]
- del Peso L, Gonzalez-Garcia M, Page C, Herrera R, Nunez G. Interleukin-3-induced phosphorylation of BAD through the protein kinase Akt. *Science*. 1997; 278:687–689. [PubMed: 9381178]
- Deschaux O, Vendruscolo LF, Schlosburg JE, Diaz-Aguilar L, Yuan CJ, Sobieraj JC, George O, Koob GF, Mandyam CD. Hippocampal neurogenesis protects against cocaine-primed relapse. *Addict Biol*. 2012
- Donovan MH, Yamaguchi M, Eisch AJ. Dynamic expression of TrkB receptor protein on proliferating and maturing cells in the adult mouse dentate gyrus. *Hippocampus*. 2008; 18:435–439. [PubMed: 18240316]
- Drake CT, Milner TA, Patterson SL. Ultrastructural localization of full-length trkB immunoreactivity in rat hippocampus suggests multiple roles in modulating activity-dependent synaptic plasticity. *J Neurosci*. 1999; 19:8009–8026. [PubMed: 10479701]
- Edwards S, Baynes BB, Carmichael CY, Zamora-Martinez ER, Barrus M, Koob GF, Gilpin NW. Traumatic stress reactivity promotes excessive alcohol drinking and alters the balance of prefrontal cortex-amygdala activity. *Transl Psychiatry*. 2013; 3:e296. [PubMed: 23982628]

- Eisch AJ, O'Dell SJ, Marshall JF. Striatal and cortical NMDA receptors are altered by a neurotoxic regimen of methamphetamine. *Synapse*. 1996; 22:217–225. [PubMed: 9132989]
- Friedman SD, Castaneda E, Hodge GK. Long-term monoamine depletion, differential recovery, and subtle behavioral impairment following methamphetamine-induced neurotoxicity. *Pharmacol Biochem Behav*. 1998; 61:35–44. [PubMed: 9715805]
- Gartner A, Staiger V. Neurotrophin secretion from hippocampal neurons evoked by long-term-potential-inducing electrical stimulation patterns. *Proc Natl Acad Sci U S A*. 2002; 99:6386–6391. [PubMed: 11983920]
- Grace CE, Schaefer TL, Herring NR, Skelton MR, McCrea AE, Vorhees CV, Williams MT. (+)-Methamphetamine increases corticosterone in plasma and BDNF in brain more than forced swim or isolation in neonatal rats. *Synapse*. 2008; 62:110–121. [PubMed: 17992688]
- Graham DL, Edwards S, Bachtell RK, DiLeone RJ, Rios M, Self DW. Dynamic BDNF activity in nucleus accumbens with cocaine use increases self-administration and relapse. *Nat Neurosci*. 2007; 10:1029–1037. [PubMed: 17618281]
- Grimm JW, Lu L, Hayashi T, Hope BT, Su TP, Shaham Y. Time-dependent increases in brain-derived neurotrophic factor protein levels within the mesolimbic dopamine system after withdrawal from cocaine: implications for incubation of cocaine craving. *J Neurosci*. 2003; 23:742–747. [PubMed: 12574402]
- Han BH, Holtzman DM. BDNF protects the neonatal brain from hypoxic-ischemic injury in vivo via the ERK pathway. *J Neurosci*. 2000; 20:5775–5781. [PubMed: 10908618]
- Hartmann M, Heumann R, Lessmann V. Synaptic secretion of BDNF after high-frequency stimulation of glutamatergic synapses. *Embo J*. 2001; 20:5887–5897. [PubMed: 11689429]
- Hashimoto R, Hough C, Nakazawa T, Yamamoto T, Chuang DM. Lithium protection against glutamate excitotoxicity in rat cerebral cortical neurons: involvement of NMDA receptor inhibition possibly by decreasing NR2B tyrosine phosphorylation. *J Neurochem*. 2002; 80:589–597. [PubMed: 11841566]
- Henke PG. Hippocampal pathway to the amygdala and stress ulcer development. *Brain Res Bull*. 1990; 25:691–695. [PubMed: 2289157]
- Hetman M, Kanning K, Cavanaugh JE, Xia Z. Neuroprotection by brain-derived neurotrophic factor is mediated by extracellular signal-regulated kinase and phosphatidylinositol 3-kinase. *J Biol Chem*. 1999; 274:22569–22580. [PubMed: 10428835]
- Hope BT. Cocaine and the AP-1 transcription factor complex. *Ann N Y Acad Sci*. 1998; 844:1–6. [PubMed: 9668659]
- Hori N, Kadota MT, Watanabe M, Ito Y, Akaike N, Carpenter DO. Neurotoxic effects of methamphetamine on rat hippocampus pyramidal neurons. *Cell Mol Neurobiol*. 2010; 30:849–856. [PubMed: 20232135]
- Ishikawa A, Kadota T, Kadota K, Matsumura H, Nakamura S. Essential role of D1 but not D2 receptors in methamphetamine-induced impairment of long-term potentiation in hippocampal-prefrontal cortex pathway. *Eur J Neurosci*. 2005; 22:1713–1719. [PubMed: 16197511]
- Ishikawa A, Nakamura S. Ventral hippocampal neurons project axons simultaneously to the medial prefrontal cortex and amygdala in the rat. *J Neurophysiol*. 2006; 96:2134–2138. [PubMed: 16837666]
- Itoh K, Fukumori R, Suzuki Y. Effect of methamphetamine on the locomotor activity in the 6-OHDA dorsal hippocampus lesioned rat. *Life Sci*. 1984; 34:827–833. [PubMed: 6422178]
- Kandel ER. The molecular biology of memory storage: a dialogue between genes and synapses. *Science*. 2001; 294:1030–1038. [PubMed: 11691980]
- Katoh-Semba R, Asano T, Ueda H, Morishita R, Takeuchi IK, Inaguma Y, Kato K. Riluzole enhances expression of brain-derived neurotrophic factor with consequent proliferation of granule precursor cells in the rat hippocampus. *FASEB J*. 2002; 16:1328–1330. [PubMed: 12154010]
- Kim YT, Lee JJ, Song HJ, Kim JH, Kwon DH, Kim MN, Yoo DS, Lee HJ, Kim HJ, Chang Y. Alterations in cortical activity of male methamphetamine abusers performing an empathy task: fMRI study. *Hum Psychopharmacol*. 2010; 25:63–70. [PubMed: 20041477]
- Kitamura O. Detection of methamphetamine neurotoxicity in forensic autopsy cases. *Leg Med (Tokyo)*. 2009; 11(Suppl 1):S63–S65. [PubMed: 19269222]

- Kitamura O, Takeichi T, Wang EL, Tokunaga I, Ishigami A, Kubo S. Microglial and astrocytic changes in the striatum of methamphetamine abusers. *Leg Med (Tokyo)*. 2010; 12:57–62. [PubMed: 20110187]
- Kjelstrup KG, Tuvnes FA, Steffenach HA, Murison R, Moser EI, Moser MB. Reduced fear expression after lesions of the ventral hippocampus. *Proc Natl Acad Sci U S A*. 2002; 99:10825–10830. [PubMed: 12149439]
- Koob GF. Neuroadaptive mechanisms of addiction: studies on the extended amygdala. *Eur Neuropsychopharmacol*. 2003; 13:442–452. [PubMed: 14636960]
- Kume T, Kouchiyama H, Kaneko S, Maeda T, Akaike A, Shimohama S, Kihara T, Kimura J, Wada K, Koizumi S. BDNF prevents NO mediated glutamate cytotoxicity in cultured cortical neurons. *Brain Res*. 1997; 756:200–204. [PubMed: 9187333]
- Larsson E, Mandel RJ, Klein RL, Muzyczka N, Lindvall O, Kokaia Z. Suppression of insult-induced neurogenesis in adult rat brain by brain-derived neurotrophic factor. *Exp Neurol*. 2002; 177:1–8. [PubMed: 12429205]
- Lasseter HC, Xie X, Ramirez DR, Fuchs RA. Sub-region specific contribution of the ventral hippocampus to drug context-induced reinstatement of cocaine-seeking behavior in rats. *Neuroscience*. 2010; 171:830–839. [PubMed: 20870011]
- Lavezzi G, McCallum J, Lee R, Roche KW. Differential binding of the AP-2 adaptor complex and PSD-95 to the C-terminus of the NMDA receptor subunit NR2B regulates surface expression. *Neuropharmacology*. 2003; 45:729–737. [PubMed: 14529712]
- Le Foll B, Diaz J, Sokoloff P. A single cocaine exposure increases BDNF and D3 receptor expression: implications for drug-conditioning. *Neuroreport*. 2005; 16:175–178. [PubMed: 15671872]
- Leal G, Comprido D, Duarte CB. BDNF-induced local protein synthesis and synaptic plasticity. *Neuropharmacology*. 2014; 76(Pt C):639–656. [PubMed: 23602987]
- Lee J, Duan W, Long JM, Ingram DK, Mattson MP. Dietary restriction increases the number of newly generated neural cells, and induces BDNF expression, in the dentate gyrus of rats. *J Mol Neurosci*. 2000; 15:99–108. [PubMed: 11220789]
- Lin SY, Wu K, Levine ES, Mount HT, Suen PC, Black IB. BDNF acutely increases tyrosine phosphorylation of the NMDA receptor subunit 2B in cortical and hippocampal postsynaptic densities. *Brain Res Mol Brain Res*. 1998; 55:20–27. [PubMed: 9645956]
- Lu B. BDNF and activity-dependent synaptic modulation. *Learn Mem*. 2003; 10:86–98. [PubMed: 12663747]
- Lu Y, Christian K, Lu B. BDNF: a key regulator for protein synthesis-dependent LTP and long-term memory? *Neurobiol Learn Mem*. 2008; 89:312–323. [PubMed: 17942328]
- Mandyam CD, Wee S, Crawford EF, Eisch AJ, Richardson HN, Koob GF. Varied access to intravenous methamphetamine self-administration differentially alters adult hippocampal neurogenesis. *Biol Psychiatry*. 2008; 64:958–965. [PubMed: 18490002]
- Matsumoto T, Rauskolb S, Polack M, Klose J, Kolbeck R, Korte M, Barde YA. Biosynthesis and processing of endogenous BDNF: CNS neurons store and secrete BDNF, not pro-BDNF. *Nat Neurosci*. 2008; 11:131–133. [PubMed: 18204444]
- Mattson MP, Lovell MA, Furukawa K, Markesbery WR. Neurotrophic factors attenuate glutamate-induced accumulation of peroxides, elevation of intracellular Ca²⁺ concentration, and neurotoxicity and increase antioxidant enzyme activities in hippocampal neurons. *J Neurochem*. 1995; 65:1740–1751. [PubMed: 7561872]
- McGinty JF, Bache AJ, Coleman NT, Sun WL. The Role of BDNF/TrkB Signaling in Acute Amphetamine-Induced Locomotor Activity and Opioid Peptide Gene Expression in the Rat Dorsal Striatum. *Front Syst Neurosci*. 2011; 5:60. [PubMed: 21811445]
- Meredith GE, Callen S, Scheuer DA. Brain-derived neurotrophic factor expression is increased in the rat amygdala, piriform cortex and hypothalamus following repeated amphetamine administration. *Brain Res*. 2002; 949:218–227. [PubMed: 12213320]
- Meyers RA, Zavala AR, Speer CM, Neisewander JL. Dorsal hippocampus inhibition disrupts acquisition and expression, but not consolidation, of cocaine conditioned place preference. *Behav Neurosci*. 2006; 120:401–412. [PubMed: 16719704]

- Minichiello L. TrkB signalling pathways in LTP and learning. *Nat Rev Neurosci.* 2009; 10:850–860. [PubMed: 19927149]
- Morales AM, Lee B, Hellemann G, O'Neill J, London ED. Gray-matter volume in methamphetamine dependence: cigarette smoking and changes with abstinence from methamphetamine. *Drug Alcohol Depend.* 2012; 125:230–238. [PubMed: 22445480]
- Moser MB, Moser EI, Forrest E, Andersen P, Morris RG. Spatial learning with a minislab in the dorsal hippocampus. *Proc Natl Acad Sci U S A.* 1995; 92:9697–9701. [PubMed: 7568200]
- Nagappan G, Lu B. Activity-dependent modulation of the BDNF receptor TrkB: mechanisms and implications. *Trends Neurosci.* 2005; 28:464–471. [PubMed: 16040136]
- Nakama H, Chang L, Fein G, Shimotsu R, Jiang CS, Ernst T. Methamphetamine users show greater than normal age-related cortical gray matter loss. *Addiction.* 2011; 106:1474–1483. [PubMed: 21438934]
- Nakazawa T, Komai S, Tezuka T, Hisatsune C, Umemori H, Semba K, Mishina M, Manabe T, Yamamoto T. Characterization of Fyn-mediated tyrosine phosphorylation sites on GluR epsilon 2 (NR2B) subunit of the N-methyl-D-aspartate receptor. *J Biol Chem.* 2001; 276:693–699. [PubMed: 11024032]
- Nakazawa T, Komai S, Watabe AM, Kiyama Y, Fukaya M, Arima-Yoshida F, Horai R, Sudo K, Ebine K, Delawary M, Goto J, Umemori H, Tezuka T, Iwakura Y, Watanabe M, Yamamoto T, Manabe T. NR2B tyrosine phosphorylation modulates fear learning as well as amygdaloid synaptic plasticity. *Embo J.* 2006; 25:2867–2877. [PubMed: 16710293]
- North A, Swant J, Salvatore MF, Gamble-George J, Prins P, Butler B, Mittal MK, Heltsley R, Clark JT, Khoshbouei H. Chronic methamphetamine exposure produces a delayed, long-lasting memory deficit. *Synapse.* 2013; 67:245–257. [PubMed: 23280858]
- Onaivi ES, Ali SF, Chirwa SS, Zwiller J, Thiriet N, Akinshola BE, Ishiguro H. Ibogaine signals addiction genes and methamphetamine alteration of long-term potentiation. *Ann N Y Acad Sci.* 2002; 965:28–46. [PubMed: 12105083]
- Opazo P, Watabe AM, Grant SG, O'Dell TJ. Phosphatidylinositol 3-kinase regulates the induction of long-term potentiation through extracellular signal-related kinase-independent mechanisms. *J Neurosci.* 2003; 23:3679–3688. [PubMed: 12736339]
- Orikabe L, Yamasue H, Inoue H, Takayanagi Y, Mozue Y, Sudo Y, Ishii T, Itokawa M, Suzuki M, Kurachi M, Okazaki Y, Kasai K. Reduced amygdala and hippocampal volumes in patients with methamphetamine psychosis. *Schizophr Res.* 2011; 132:183–189. [PubMed: 21784619]
- Orio L, Edwards S, George O, Parsons LH, Koob GF. A role for the endocannabinoid system in the increased motivation for cocaine in extended-access conditions. *J Neurosci.* 2009; 29:4846–4857. [PubMed: 19369553]
- Pang PT, Teng HK, Zaitsev E, Woo NT, Sakata K, Zhen S, Teng KK, Yung WH, Hempstead BL, Lu B. Cleavage of proBDNF by tPA/plasmin is essential for long-term hippocampal plasticity. *Science.* 2004; 306:487–491. [PubMed: 15486301]
- Park H, Poo MM. Neurotrophin regulation of neural circuit development and function. *Nat Rev Neurosci.* 2013; 14:7–23. [PubMed: 23254191]
- Pentkowski NS, Blanchard DC, Lever C, Litvin Y, Blanchard RJ. Effects of lesions to the dorsal and ventral hippocampus on defensive behaviors in rats. *Eur J Neurosci.* 2006; 23:2185–2196. [PubMed: 16630065]
- Pierce RC, Bari AA. The role of neurotrophic factors in psychostimulant-induced behavioral and neuronal plasticity. *Rev Neurosci.* 2001; 12:95–110. [PubMed: 11392459]
- Pitkanen A, Pikkarainen M, Nurminen N, Ylinen A. Reciprocal connections between the amygdala and the hippocampal formation, perirhinal cortex, and postrhinal cortex in rat. A review. *Ann N Y Acad Sci.* 2000; 911:369–391. [PubMed: 10911886]
- Pothuizen HH, Zhang WN, Jongen-Relo AL, Feldon J, Yee BK. Dissociation of function between the dorsal and the ventral hippocampus in spatial learning abilities of the rat: a within-subject, within-task comparison of reference and working spatial memory. *Eur J Neurosci.* 2004; 19:705–712. [PubMed: 14984421]
- Price KL, DeSantis SM, Simpson AN, Tolliver BK, McRae-Clark AL, Saladin ME, Baker NL, Wagner MT, Brady KT. The impact of clinical and demographic variables on cognitive

- performance in methamphetamine-dependent individuals in rural South Carolina. *Am J Addict*. 2011; 20:447–455. [PubMed: 21838844]
- Prybylowski K, Chang K, Sans N, Kan L, Vicini S, Wenthold RJ. The synaptic localization of NR2B-containing NMDA receptors is controlled by interactions with PDZ proteins and AP-2. *Neuron*. 2005; 47:845–857. [PubMed: 16157279]
- Rau TF, Kothiwala A, Zhang L, Ulatowski S, Jacobson S, Brooks DM, Cardozo-Pelaez F, Chopp M, Poulsen DJ. Low dose methamphetamine mediates neuroprotection through a PI3K-AKT pathway. *Neuropharmacology*. 2011; 61:677–686. [PubMed: 21635908]
- Recinto P, Samant AR, Chavez G, Kim A, Yuan CJ, Soleiman M, Grant Y, Edwards S, Wee S, Koob GF, George O, Mandym CD. Levels of neural progenitors in the hippocampus predict memory impairment and relapse to drug seeking as a function of excessive methamphetamine self-administration. *Neuropsychopharmacology*. 2012; 37:1275–1287. [PubMed: 22205547]
- Rogers JL, De Santis S, See RE. Extended methamphetamine self-administration enhances reinstatement of drug seeking and impairs novel object recognition in rats. *Psychopharmacology (Berl)*. 2008; 199:615–624. [PubMed: 18493748]
- Rogers JL, See RE. Selective inactivation of the ventral hippocampus attenuates cue-induced and cocaine-primed reinstatement of drug-seeking in rats. *Neurobiol Learn Mem*. 2007; 87:688–692. [PubMed: 17337218]
- Rosenblum K, Dudai Y, Richter-Levin G. Long-term potentiation increases tyrosine phosphorylation of the N-methyl-D-aspartate receptor subunit 2B in rat dentate gyrus in vivo. *Proc Natl Acad Sci U S A*. 1996; 93:10457–10460. [PubMed: 8816822]
- Rostas JA, Brent VA, Voss K, Errington ML, Bliss TV, Gurd JW. Enhanced tyrosine phosphorylation of the 2B subunit of the N-methyl-D-aspartate receptor in long-term potentiation. *Proc Natl Acad Sci U S A*. 1996; 93:10452–10456. [PubMed: 8816821]
- Russo SJ, Dietz DM, Dumitriu D, Morrison JH, Malenka RC, Nestler EJ. The addicted synapse: mechanisms of synaptic and structural plasticity in nucleus accumbens. *Trends Neurosci*. 2010; 33:267–276. [PubMed: 20207024]
- SAMHSA. Results from the 2007 National Survey on Drug Use and Health: Detailed Tables. Substance Abuse and Mental Health Services Administration, Office of Applied Studies. 2008
- Scharfman H, Goodman J, Macleod A, Phani S, Antonelli C, Croll S. Increased neurogenesis and the ectopic granule cells after intrahippocampal BDNF infusion in adult rats. *Exp Neurol*. 2005; 192:348–356. [PubMed: 15755552]
- Schmued LC, Bowyer JF. Methamphetamine exposure can produce neuronal degeneration in mouse hippocampal remnants. *Brain Res*. 1997; 759:135–140. [PubMed: 9219871]
- Schulz S, Siemer H, Krug M, Holtt V. Direct evidence for biphasic cAMP responsive element-binding protein phosphorylation during long-term potentiation in the rat dentate gyrus in vivo. *J Neurosci*. 1999; 19:5683–5692. [PubMed: 10377374]
- Schwartz DL, Mitchell AD, Lahna DL, Lubner HS, Huckans MS, Mitchell SH, Hoffman WF. Global and local morphometric differences in recently abstinent methamphetamine-dependent individuals. *Neuroimage*. 2010; 50:1392–1401. [PubMed: 20096794]
- Shehadah A, Chen J, Zacharek A, Cui Y, Ion M, Roberts C, Kapke A, Chopp M. Niaspan treatment induces neuroprotection after stroke. *Neurobiol Dis*. 2010; 40:277–283. [PubMed: 20554037]
- Skaper SD, Floreani M, Negro A, Facci L, Giusti P. Neurotrophins rescue cerebellar granule neurons from oxidative stress-mediated apoptotic death: selective involvement of phosphatidylinositol 3-kinase and the mitogen-activated protein kinase pathway. *J Neurochem*. 1998; 70:1859–1868. [PubMed: 9572269]
- Skelton MR, Williams MT, Schaefer TL, Vorhees CV. Neonatal (+)-methamphetamine increases brain derived neurotrophic factor, but not nerve growth factor, during treatment and results in long-term spatial learning deficits. *Psychoneuroendocrinology*. 2007; 32:734–745. [PubMed: 17606327]
- Swant J, Chirwa S, Stanwood G, Khoshbouei H. Methamphetamine reduces LTP and increases baseline synaptic transmission in the CA1 region of mouse hippocampus. *PLoS ONE*. 2010; 5:e11382. [PubMed: 20614033]

- Thompson PM, Hayashi KM, Simon SL, Geaga JA, Hong MS, Sui Y, Lee JY, Toga AW, Ling W, London ED. Structural abnormalities in the brains of human subjects who use methamphetamine. *J Neurosci*. 2004; 24:6028–6036. [PubMed: 15229250]
- Weber E, Blackstone K, Iudicello JE, Morgan EE, Grant I, Moore DJ, Woods SP. Neurocognitive deficits are associated with unemployment in chronic methamphetamine users. *Drug Alcohol Depend*. 2012; 125:146–153. [PubMed: 22560676]
- Wells AM, Lasseter HC, Xie X, Cowhey KE, Reittinger AM, Fuchs RA. Interaction between the basolateral amygdala and dorsal hippocampus is critical for cocaine memory reconsolidation and subsequent drug context-induced cocaine-seeking behavior in rats. *Learn Mem*. 2011; 18:693–702. [PubMed: 22005750]
- Yamamoto J. Cortical and hippocampal EEG power spectra in animal models of schizophrenia produced with methamphetamine, cocaine, and phencyclidine. *Psychopharmacology (Berl)*. 1997; 131:379–387. [PubMed: 9226740]
- Yoshikawa T, Shibuya H, Kaneno S, Toru M. Blockade of behavioral sensitization to methamphetamine by lesion of hippocampo-accumbal pathway. *Life Sci*. 1991; 48:1325–1332. [PubMed: 2002758]
- Yuan CJ, Quiocho JM, Kim A, Wee S, Mandyam CD. Extended access methamphetamine decreases immature neurons in the hippocampus which results from loss and altered development of neural progenitors without altered dynamics of the S-phase of the cell cycle. *Pharmacol Biochem Behav*. 2011; 100:98–108. [PubMed: 21855565]
- Zhang L, Qu Y, Tang J, Chen D, Fu X, Mao M, Mu D. PI3K/Akt signaling pathway is required for neuroprotection of thalidomide on hypoxic-ischemic cortical neurons in vitro. *Brain Res*. 2010; 1357:157–165. [PubMed: 20705061]

Highlights

- Methamphetamine self-administration enhances BDNF expression in the hippocampus
- Enhanced BDNF expression is associated with hypophosphorylation of NR2B
- Enhanced BDNF expression is associated with altered expression of cell death factor

a) Schematic of experimental procedure

b)

c)

d)

Figure 1.

Rats that underwent extended access to methamphetamine self-administration escalated methamphetamine intake. (a) Schematic showing baseline training followed by 17 sessions of either 1 hr/day saline ($n = 3$), 1hr/day (ShA, $n = 8$; short vertical lines) or 6 hr/day (LgA, $n = 6$; long vertical lines) methamphetamine self-administration. (b) Schematic of brain regions indicating area of tissue punches used for Western blot analysis. Red circles, dorsal hippocampus; blue circles, ventral hippocampus. (c–d) Active lever presses for saline and methamphetamine (c) and methamphetamine intake in mg/kg per session from ShA and LgA animals (d). The first hour intake is also indicated for LgA animals. $*p < 0.01$ compared with initial sessions in the same animals by repeated measures two-way ANOVA followed by Student-Newman-Keuls *post hoc* test. Data are represented as mean $\pm$ SEM.

Figure 2.

Methamphetamine self-administration enhances BDNF expression in the dorsal and ventral hippocampus. (a): Quantitative analysis of BDNF expression (mature form) by Western blot analysis of dorsal and ventral hippocampus-enriched protein lysates. (b): Qualitative representation of BDNF immunoreactive bands used for quantitative data analysis from dorsal hippocampus; (c) corresponding Coomassie staining of the membrane is shown as loading control. Molecular weights of proteins are indicated in kDa. * $p < 0.05$ compared to naïve controls. Data are represented as mean + SEM.

Figure 3.

Methamphetamine self-administration does not alter TrkB expression or pTrkB levels in the hippocampus. **(a)**: Quantitative analysis of TrkB and pTrkB/TrkB expression by Western blot analysis of dorsal and ventral hippocampus-enriched protein lysates. **(b)**: Qualitative representation of phospho (p)TrkB and total (t)TrkB immunoreactive bands used for quantitative data analysis from dorsal hippocampus; **(c)** corresponding Coomassie staining of the membrane is shown as loading control. Molecular weights of proteins are indicated in kDa.

Figure 4.

Methamphetamine self-administration produces hypophosphorylation of GluN2B receptors at Tyr-1472 in the ventral hippocampus. **(a)**: Quantitative analysis of total (t)GluN2B and phospho (p)GluN2B/tGluN2B expression by Western blot analysis of dorsal and ventral hippocampus-enriched protein lysates. **(b)**: Qualitative representation of pGluN2B and tGluN2B immunoreactive bands used for quantitative data analysis. **(c)** corresponding Coomassie staining of the membrane is shown as loading control. Molecular weights of proteins are indicated in kDa. * $p < 0.05$ compared to naïve controls. Data are represented as mean + SEM.

Figure 5. Methamphetamine self-administration regulates cell death factor expression in the ventral hippocampus. **(a):** Quantitative analysis of Bcl-2 and Bax expression by Western blot analysis of dorsal and ventral hippocampus-enriched protein lysates. **(b–c):** Qualitative representation of Bcl-2 immunoreactive bands used for quantitative data analysis **(b)**; corresponding Coomassie staining of the membrane used as loading control **(c)**. **(d–e):** Qualitative representation of Bax immunoreactive bands used for quantitative data analysis **(d)**; corresponding Coomassie staining of the membrane used as loading control **(e)**.

Molecular weights of proteins are indicated in kDa. * $p < 0.05$ compared to naïve controls; ** $p < 0.01$ compared to naïve controls; # $p < 0.01$ main effect of hippocampal subregion. Data are represented as mean + SEM.

Figure 6.

Methamphetamine self-administration reduces Akt levels in the ventral hippocampus. **(a)**: Quantitative analysis of total (t)Akt and phospho (p)Akt/tAkt expression by Western blot analysis of ventral hippocampus-enriched protein lysates. **(b)**: Qualitative representation of pAkt and tAkt immunoreactive bands used for quantitative data analysis. **(c)** corresponding Coomassie staining of the membrane is shown as loading control. Molecular weights of proteins are indicated in kDa. * $p < 0.05$ compared to naïve controls. Data are represented as mean + SEM.