subsequently lead to an impaired lung function, a hallmark of COPD. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

K. de Jong, Ph.D.
J. M. Vonk, Ph.D.
University of Groningen, University Medical Center Groningen, Groningen
Research Institute for Asthma and COPD (GRIAC)
Groningen, The Netherlands

H. Kromhout, Ph.D.
R. Vermeulen, Ph.D.
University of Utrecht
The Netherlands

D. S. Postma, M.D., Ph.D.
H. M. Boezen, Ph.D.
University of Groningen, University Medical Center Groningen, Groningen
Research Institute for Asthma and COPD (GRIAC)
Groningen, The Netherlands

LifeLines Cohort Study

Investigators of the LifeLines Cohort Study: B. Z. Alizadeh, H. M. Boezen, H. L. Hillege, H. Snieder, and R. P. Stolk (Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands); R. A. de Boer, P. van der Harst, and H. L. Hillege (Department of Cardiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands); M. Bruinenberg (the LifeLines Cohort Study, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands); L. Franke and C. Wijmenga (Department of Genetics, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands); M. M. van der Klauw and B. H. R. Wolffenbuttel (Department of Endocrinology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands); G. Navis (Department of Internal Medicine, Division of Nephrology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands); J. Ormel (Interdisciplinary Center of Psychopathology of Emotion Regulation, Department of Psychiatry, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands); D. S. Postma (Department of Pulmonology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands); and J. P. Slaets (University Center for Geriatric Medicine, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands).

References

- Boezen HM. Genome-wide association studies: what do they teach us about asthma and chronic obstructive pulmonary disease? *Proc Am Thorac Soc* 2009;6:701–703.
- Hoppin JA, Valcin M, Henneberger PK, Kullman GJ, Umbach DM, London SJ, Alavanja MC, Sandler DP. Pesticide use and chronic bronchitis among farmers in the Agricultural Health Study. *Am J Ind Med* 2007;50:969–979.
- de Jong K, Boezen HM, Kromhout H, Vermeulen R, Postma DS, Vonk JM; LifeLines Cohort Study. Pesticides and other occupational exposures are associated with airway obstruction: the LifeLines cohort study. *Occup Environ Med* 2014;71:88–96.
- Hernández AF, Parrón T, Alarcón R. Pesticides and asthma. *Curr Opin Allergy Clin Immunol* 2011;11:90–96.
- Koutros S, Berndt SI, Hughes Barry K, Andreotti G, Hoppin JA, Sandler DP, Yeager M, Burdett LA, Yuenger J, Alavanja MC, et al. Genetic susceptibility loci, pesticide exposure and prostate cancer risk. *PLoS One* 2013;8:e58195.
- Dardiotis E, Xiromerisiou G, Hadjichristodoulou C, Tsatsakis AM, Wilks MF, Hadjigeorgiou GM. The interplay between environmental and genetic factors in Parkinson's disease susceptibility: the evidence for pesticides. *Toxicology* 2013;307:17–23.
- Stolk RP, Rosmalen JGM, Postma DS, de Boer RA, Navis G, Slaets JPJ, Ormel J, Wolffenbuttel BH. Universal risk factors for multifactorial diseases: LifeLines: a three-generation population-based study. *Eur J Epidemiol* 2008;23:67–74.
- van Diemen CC, Postma DS, Vonk JM, Bruinenberg M, Schouten JP, Boezen HM. A disintegrin and metalloprotease 33 polymorphisms and lung function decline in the general population. *Am J Respir Crit Care Med* 2005;172:329–333.
- Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, Maller J, Sklar P, de Bakker PI, Daly MJ, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;81:559–575.
- Swann K, Lai FA. PLC ζ and the initiation of Ca(2+) oscillations in fertilizing mammalian eggs. *Cell Calcium* 2013;53:55–62.
- Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–560.
- Hamilton SR, Li H, Wischniewski H, Prasad A, Kerley-Hamilton JS, Mitchell T, Walling AJ, Davidson RC, Wildt S, Gemgross TU. Intact alpha-1,2-endomannosidase is a typical type II membrane protein. *Glycobiology* 2005;15:615–624.
- Hancock DB, Martin ER, Vance JM, Scott WK. Nitric oxide synthase genes and their interactions with environmental factors in Parkinson's disease. *Neurogenetics* 2008;9:249–262.
- Aminuddin F, Hackett TL, Stefanowicz D, Saferali A, Paré PD, Gulsvik A, Bakke P, Cho MH, Litorjua A, Lomas DA, et al. Nitric oxide synthase polymorphisms, gene expression and lung function in chronic obstructive pulmonary disease. *BMC Pulm Med* 2013;13:64.
- Kobzik L, Bredt DS, Lowenstein CJ, Drazen J, Gaston B, Sugarbaker D, Stamler JS. Nitric oxide synthase in human and rat lung: immunocytochemical and histochemical localization. *Am J Respir Cell Mol Biol* 1993;9:371–377.
- Brindicci C, Kharitonov SA, Ito M, Elliott MW, Hogg JC, Barnes PJ, Ito K. Nitric oxide synthase isoenzyme expression and activity in peripheral lung tissue of patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2010;181:21–30.
- Morán JM, Ortiz-Ortiz MA, Ruiz-Mesa LM, Fuentes JM. Nitric oxide in paraquat-mediated toxicity: a review. *J Biochem Mol Toxicol* 2010;24:402–409.

Copyright © 2014 by the American Thoracic Society

Proximity to a Major Road, Vitamin D Insufficiency, and Severe Asthma Exacerbations in Puerto Rican Children

To the Editor:

Traffic-related air pollution (TRAP) causes asthma morbidity, and vitamin D insufficiency (a plasma 25[OH]D level < 30 ng/ml) is associated with severe asthma exacerbations in Puerto Rican (PR) children (1). Because vitamin D has antioxidant effects, we examined whether proximity to a major road (a marker of TRAP) is associated with increased risk for severe asthma exacerbations

The authors are supported by grants HL079966 and HL117191 from the National Institutes of Health and by an endowment from the Heinz Foundation.

Author Contributions: G.C. and J.C.C. participated in study design and implementation; F.R., J.M.B., E.F., K.S.K., and J.C.C. participated in data analysis; E.A.-P. participated in data generation; F.R. and J.C.C. wrote the initial draft of the manuscript; and all authors participated in the review of the manuscript and approved its final version.

This letter has an online supplement, which is accessible from this issue's table of contents at www.atsjournals.org

Table 1. Multivariate Analysis of Distance from a Major Roadway (by Quartiles [Model 1] and as a Continuous Variable [Model 2]) and Severe Asthma Exacerbations in Puerto Rican Children

Covariates	Model 1	Model 2
Residential distance from a major roadway	By quartiles	Per every 100 m
Quartile 1 (<172 m)	3.04* (1.43–6.45)	—
Quartile 2 (172–209 m)	2.02* (1.01–4.06)	—
Quartile 3 (210–441 m)	1.93 (0.94–3.99)	—
Quartile 4 (>441 m)	1.00 (reference)	—
Continuous	—	1.15* [†] (1.05–1.27)
Vitamin D insufficiency (plasma 25[OH]D level < 30 ng/dl)	2.46* (1.43–4.28)	2.46* (1.42–4.23)
Age, yr	0.84* (0.75–0.93)	0.87* (0.79–0.95)
Gender (male)	1.19 (0.71–1.98)	1.21 (0.73–2.01)
Annual household income > \$15,000	0.81 (0.45–1.47)	0.83 (0.48–1.45)
Inhaled corticosteroid use during the last 6 mo	3.94* (2.16–7.19)	3.97* (2.19–7.19)

Data are given as odds ratio (95% confidence interval).

* $P < 0.01$.

[†]Per every 100 m closer to a major roadway, the odds of a severe exacerbation increase by ~15%.

in PR children and whether coexisting vitamin D insufficiency further increases this risk.

From March 2009 to June 2010, 351 children with asthma were chosen from randomly selected households in San Juan, Puerto Rico, as previously described (1). Study participants completed a protocol including questionnaires, spirometry, and allergy skin testing. Plasma 25-hydroxy-vitamin D (hereafter referred to as vitamin D) was measured using tandem mass spectrophotometry (Waters Corporation, Milford, MA) (1). Written parental consent and child's assent was obtained for participants in the study, which was approved by the institutional review boards of the University of Puerto Rico in San Juan; Brigham and Women's Hospital in Boston, Massachusetts; and the University of Pittsburgh in Pennsylvania.

To measure residential distance from a major road, the home address of each participant was geocoded (linked) to a 15-digit 2000 US Census Federal Information Processing Standard + block code at the University of Puerto Rico. Using the program ArcMAP10.1 (ArcGIS 10.1, Esri, Redlands, CA), centroids were then created by obtaining X, Y coordinates for the center of individual census blocks based on a 2000 US Census map for Puerto Rico (see Figure E1 in the online supplement). The distance from the residential block centroid of a participant to the nearest major road (defined by Esri 2012 Data and Maps major road layer) was then calculated using the geo-processing proximity tool "near," which measures the nearest distance "as the crow flies" between two features. Because of the lack of a complete home address, one participant was excluded.

Figure 1. Categories of residential proximity to a major roadway (living > vs. ≤441 m away from a major road) and vitamin D insufficiency (plasma vitamin D level < 30 ng/dl) and severe asthma exacerbations. Multivariate analysis adjusted for age, sex, household income, and use of inhaled corticosteroids in the prior 6 months.

Our primary outcome was one or more severe asthma exacerbations (one or more hospitalizations or visits to the emergency department [ED] or urgent care requiring treatment with systemic corticosteroids, or one or more courses of systemic steroids [2]) in the previous year. Secondary outcomes included visits to the ED/urgent care and hospitalizations for asthma in the previous year.

Logistic multivariable regression was used for the analysis of TRAP and severe asthma exacerbations, adjusting for age, sex, household income, vitamin D insufficiency, and use of inhaled corticosteroids. To examine the joint effects of vitamin D insufficiency and proximity to a major road, we created four categories of exposure: vitamin D sufficiency and living furthest from a major road (in the upper or fourth quartile, based on results from a multivariate model), serving as the reference group; vitamin D sufficiency and living closer to a major road (first to third quartiles); vitamin D insufficiency and living furthest from a major road; and vitamin D insufficiency and living closer to a major road. Given nearly identical results for the two intermediate categories, they were collapsed, resulting in three categories for analysis: vitamin D sufficiency and living furthest from a major road, either living close to a major road or living farther from a road (but not both), and both living close to a major road and having vitamin D insufficiency.

Table E1 summarizes the characteristics of study participants: two-thirds or more lacked private/employer-based health insurance, nearly half were exposed to current or early-life environmental tobacco smoke (ETS), more than 40% had vitamin D insufficiency, slightly more than half had one or more severe exacerbations, and most were atopic. Proximity to a major road was significantly associated with one or more severe asthma exacerbations in the previous year.

Table E2 shows a comparison of children who either did or did not have one or more severe asthma exacerbations in the prior year by residential distance to a major road and other characteristics. Compared with children who had no severe exacerbations, those who did were significantly more likely to be younger and to have been exposed to ETS in early life, to have lower lung function and vitamin D insufficiency, and to not have used inhaled corticosteroids. There was also a trend for an inverse association between residential distance to a major road and one or more severe exacerbations ($P = 0.05$).

Table 1 shows the results of the multivariable analysis of residential distance to a major road and severe asthma exacerbations. Children living closer to a major road had two- to threefold increased odds of one or more severe exacerbation (Model 1). In this model, vitamin D insufficiency was also associated with 2.5 times increased odds of one or more severe exacerbations. Similar results were obtained when distance to a major road was treated as continuous (Model 2). Moreover, our findings were unchanged after additional adjustment for lung function measures or ETS in our multivariate models (data not shown).

Figure 1 and Table E3 show the findings from an analysis stratified by categories of proximity to a road and vitamin D insufficiency. In this analysis, children who had either vitamin D insufficiency or lived closest to a major road were 1.6 times more likely to have had one or more severe asthma exacerbations than those who were vitamin D sufficient and lived furthest from a major road. Compared with children who were both vitamin D sufficient and lived furthest from a major road, those who were both

vitamin D insufficient and lived closer to a major road were nearly five times more likely to have had one or more severe exacerbations. Findings for secondary outcomes were similar for ED/urgent care visits but less significant for hospitalizations for asthma, at least partly because of the small sample size (see Tables E4 and E5).

Distance from a centroid may under- or overestimate residential distance from a major road, but this misclassification would be modest and nondifferential. Although we lack data on specific pollutants, examining TRAP by residential proximity to a major road integrates exposure to a complex milieu of pollutants (3), some of which (e.g., nitrogen dioxide, particulate matter and ozone) have been implicated in the pathogenesis of asthma and severe asthma exacerbations (4, 5). Seasonal variation has no major effect on vitamin D levels in Puerto Rico, and our findings were unchanged after accounting for month of sample collection or other relevant variables measured in our study. Confounding by unmeasured factors is a plausible but improbable explanation for our results, given prior findings implicating vitamin D insufficiency or TRAP as a risk factor for severe asthma exacerbations in children.

To our knowledge, this is the first report of joint detrimental effects of vitamin D insufficiency and TRAP on severe asthma exacerbations in children. Our findings suggest that airway oxidative stress induced by air pollutants is exacerbated by vitamin D insufficiency (6–8). Alternatively, vitamin D insufficiency and TRAP may impair immune responses to viral infections, a major cause of severe asthma exacerbations in childhood (9). Clinical trials should determine whether vitamin D ameliorates the negative effects of TRAP on asthma. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Franziska Rosser, M.D.
John M. Brehm, M.D., M.P.H.
Erick Forno, M.D., M.P.H.
University of Pittsburgh School of Medicine
Pittsburgh, Pennsylvania

Edna Acosta-Pérez, Ph.D.
University of Puerto Rico
San Juan, Puerto Rico

Kristen Kurland, B.A.
Carnegie Mellon University
Pittsburgh, Pennsylvania

Gloria Canino, Ph.D.
University of Puerto Rico
San Juan, Puerto Rico

Juan C. Celedón, M.D., Dr.P.H.
University of Pittsburgh School of Medicine
Pittsburgh, Pennsylvania

References

1. Brehm JM, Acosta-Pérez E, Klei L, Roeder K, Barmada M, Boutaoui N, Forno E, Kelly R, Paul K, Sylvia J, *et al.* Vitamin D insufficiency and severe asthma exacerbations in Puerto Rican children. *Am J Respir Crit Care Med* 2012;186:140–146.
2. Reddel HK, Taylor DR, Bateman ED, Boulet LP, Boushey HA, Busse WW, Casale TB, Chané P, Enright PL, Gibson PG, *et al.*; American Thoracic Society/European Respiratory Society Task Force on Asthma Control and Exacerbations. An official American Thoracic Society/European Respiratory Society statement: asthma control and exacerbations: standardizing endpoints for clinical asthma trials and clinical practice. *Am J Respir Crit Care Med* 2009;180:59–99.

3. Pratt GC, Parson K, Shinoda N, Lindgren P, Dunlap S, Yawn B, Wollan P, Johnson J. Quantifying traffic exposure. *J Expo Sci Environ Epidemiol* 2014;24:290–296.
4. Salam MT, Islam T, Gilliland FD. Recent evidence for adverse effects of residential proximity to traffic sources on asthma. *Curr Opin Pulm Med* 2008;14:3–8.
5. Auerbach A, Hernandez ML. The effect of environmental oxidative stress on airway inflammation. *Curr Opin Allergy Clin Immunol* 2012;12:133–139.
6. Rafacho BP, Santos P, Assalin HB, Ardisson LP, Roscani MG, Polegato BF, Chiuso-Minicucci F, Fernandes AA, Azevedo PS, Minicucci MF, et al. Role of vitamin D in the cardiac remodeling induced by tobacco smoke exposure. *Int J Cardiol* 2012;155:472–473.
7. Sharifi N, Amani R, Hajiani E, Cheraghian B. Does vitamin D improve liver enzymes, oxidative stress, and inflammatory biomarkers in adults with non-alcoholic fatty liver disease? A randomized clinical trial. *Endocrine* 2014;47:70–80.
8. Li N, Hao M, Phalen RF, Hinds WC, Nel AE. Particulate air pollutants and asthma. A paradigm for the role of oxidative stress in PM-induced adverse health effects. *Clin Immunol* 2003;109:250–265.
9. Paul G, Brehm JM, Alcorn JF, Holguín F, Aujla SJ, Celedón JC. Vitamin D and asthma. *Am J Respir Crit Care Med* 2012;185:124–132.

Copyright © 2014 by the American Thoracic Society

Native and Active Vitamin D in Intensive Care: Who and How We Treat Is Crucially Important

To the Editor:

We read with great interest the results of the double-blind, placebo-controlled trial by Leaf and colleagues investigating the effects of the active vitamin D metabolite 1,25-dihydroxyvitamin D [calcitriol, 1,25(OH)₂D] on short-term biochemical outcomes in adult critically ill patients with severe sepsis (1). The authors report that a single dose of intravenous calcitriol had no effect on plasma human cathelicidin cationic antimicrobial protein (hCAP)-18 levels, but hCAP-18 and interleukin 10 mRNA expression was significantly higher in the intervention group after 24 hours. This study is very welcome, as it is one of only a few interventional trials among several observational studies aimed at understanding the role of vitamin D in critical care. Although the results are intriguing, several points need to be raised to fully appreciate the limitations of these findings:

1. Who was treated? Vitamin D treatment makes most sense in those with low vitamin D status. In analogy, well-known and powerful drugs such as insulin, antibiotics, or beta blockers are only useful in selected, not all, intensive care unit patients. In this study, low vitamin D status was not a prerequisite for study inclusion, and although a median 25-hydroxyvitamin D [25(OH)D] level of 16.5 ng/ml at baseline is reported in the intervention group, it is unclear what proportion of the study cohort was vitamin D deficient. It would have been interesting to analyze whether patients with very low 25(OH)D levels (i.e., below median levels) responded better to calcitriol than those with higher levels, despite the limitation of sample size. It is also worth noting that the reported liquid chromatography–mass spectrometry interassay coefficients of variation for 25(OH)D (17.5%) and calcitriol (14.5%) are surprisingly high and, as such, were a cause for some concern regarding the conclusions drawn on this data.

2. How were the patients treated? Calcitriol is the natural active vitamin D metabolite and has a half-life of only a few hours, in contrast to the much longer half-life of the native vitamin D metabolite cholecalciferol (2–3 weeks). The authors opted for a single dose of calcitriol and at the same time appreciated reasons why local 25(OH)D might be more relevant for immunomodulatory effects than circulating calcitriol levels (2). They eventually justified their choice to use calcitriol by suggesting impaired conversion of 25(OH)D to 1,25(OH)₂D in critical illness based on two studies (3, 4). One study investigated 30 patients who developed acute kidney injury (3), and the second looked at peritoneal dialysate monocytes from patients with end-stage renal disease (4). The current study design, however, excluded the only patient cohort that was likely to profit most from calcitriol treatment; namely, patients with end-stage renal disease. There is very limited evidence that 1 α -hydroxylase activity may indeed be reduced in critical illness. In fact, circulating levels of calcitriol and 25(OH)D have been shown to increase more than twofold within days of high-dose cholecalciferol supplementation in critically ill patients (5). Similar increases in 1,25(OH)₂D levels were found in a study in which 25(OH)D was given to intensive care unit patients, and the relative increase was even larger than in the group receiving a 2- μ g calcitriol dose every second day (6).

We strongly believe that native vitamin D, and in particular cholecalciferol (vitamin D₃), remains the best and most physiologic vitamin D metabolite to correct low 25(OH)D levels in critically ill patients with or without sepsis. Active vitamin D analogs may be useful as adjunctive agents in the small subgroup of patients with advanced acute or chronic renal failure. Regarding the question posed by Leaf and colleagues of whether calcitriol increases circulating hCAP-18 levels in humans with severe sepsis, it seems the data from this study are unable to provide adequate answers, because we doubt that either the chosen target population or the single calcitriol treatment regimen was optimal. To accurately determine whether vitamin D may play a beneficial role in the treatment of septic patients, future studies are needed, carefully choosing the target population, type of vitamin D supplementation, route/frequency/dose of administration, and methods used to quantify pertinent biomarkers. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Christian Schnedl, M.D.
Harald Dobnig, M.D.
Medical University of Graz
Graz, Austria

Sadeq A. Quraishi, M.D., M.H.A., M.Sc.
Massachusetts General Hospital-Harvard Medical School
Boston, Massachusetts

J. Dayre McNally, M.D., Ph.D.
Children's Hospital of Eastern Ontario
Ottawa, Ontario, Canada

Karin Amrein, M.D., M.Sc.
Medical University of Graz
Graz, Austria

References

1. Leaf DE, Raed A, Donnino MW, Ginde AA, Waikar SS. Randomized controlled trial of calcitriol in severe sepsis. *Am J Respir Crit Care Med* 2014;190:533–541.