Published in final edited form as:

Conf Proc IEEE Eng Med Biol Soc. 2014 August ; 2014: 4759–4762. doi:10.1109/EMBC.2014.6944688.

A Metabolic Signature of Colon Cancer Initiating Cells

Kai-Yuan Chen,

School of Electrical and Computer Engineering, Cornell University, Ithaca, NY 14853 USA

Xiaoqing Liu,

Division of Nutrition Sciences. Cornell University, Ithaca, NY 14853 USA

Pengcheng Bu,

Department of Biomedical Engineering, Cornell University, Ithaca, NY 14850 USA

Chieh-Sheng Lin,

Department of Biomedical Engineering, Cornell University, Ithaca, NY 14853 USA. Cornell University, Ithaca, NY 14850 USA

Nikolai Rakhilin,

School of Electrical and Computer Engineering Cornell University, Ithaca, NY 14850 USA. Cornell University, Ithaca, NY 14853 USA

Jason W. Locasale, and

Division of Nutrition Sciences. Cornell University, Ithaca, NY 14853 USA

Xiling Shen

School of Electrical and Computer Engineering and the Department of Biomedical Engineering, Cornell University, Ithaca, NY 14853 USA

Xiaoqing Liu: xl296@cornell.edu; Pengcheng Bu: pb345@cornell.edu; Chieh-Sheng Lin: cl979@cornell.edu; Nikolai Rakhilin: nr233@cornell.edu; Jason W. Locasale: locasale@cornell.edu; Xiling Shen: xs66@cornell.edu

Abstract

Colon cancer initiating cells (CCICs) are more tumorigenic and metastatic than the majority of colorectal cancer (CRC) cells. CCICs have also been associated with stem cell-like properties. However, there is a lack of system-level understanding of what mechanisms distinguish CCICs from common CRC cells. We compared the transcriptomes of CD133+ CCICs and CD133– CRC cells from multiple sources, which identified a distinct metabolic signature for CD133^{high} CCICs. High-resolution unbiased metabolomics was then performed to validate this CCIC metabolic signature. Specifically, levels of enzymes and metabolites involved in glycolysis, the citric acid (TCA) cycle, and cysteine and methionine metabolism are altered in CCICs. Analyses of the alterations further suggest an epigenetic link. This metabolic signature provides mechanistic insights into CCIC phenotypes and may serve as potential biomarkers and therapeutic targets for future CRC treatment.

I. Introduction

A rare population isolated from primary CRC tumors, CCICs play important roles in CRC tumorigenesis [1][2]. CCICs also possess certain stem cell-like traits, including self-renewal, differentiation, and asymmetric division[3]. CCICs were originally identified by the marker CD133[1][2][4]. Since then, they have also been associated with other markers such as CD44, ALDH1, and Lgr5[3][5].

However, it has remained largely unclear whether CCICs isolated from different CRC tumors indeed share common mechanisms that account for their phenotype, or alternatively they are completely different cells that were categorized simply by their tumorigenic capacity.

To address this question, we first analyzed 5 GEO microarray datasets that measured the transcriptomes of CD133+ versus CD133- CRC cells[6]. The transcriptome analysis suggested that CD133+ cells consistently regulate certain metabolic enzymes differentially from CD133- cells. Unbiased metabolomics by high-resolution mass spectrometry further corroborated the metabolic signature of CD133+ CCICs, which involve the glycolysis, TCA cycle, and cysteine/methionine metabolism pathways.

II. Material and Methods

A. Microarray Data From GEO

Five sets of microarray data from GEO were analyzed. These data sets include 28 FACS (Fluorescence-Activated Cell Sorting) sorted CD133+ vs. CD133- pairs from 3 CRC patient tumors and 4 CRC cell lines (Table I).

B. Statistical Analysis

An R-package, *Bioconductor*[7], was used to extract pre-analyzed GEO data and for post-processing. Genes that were significantly (p-value <0.05) up-regulated or down-regulated in CD133+ vs. CD133- cells were identified by differential analysis using t-test and fold-change analysis. Gene frequencies and Venn diagrams were further generated by the R-packages *limma*[8] and *vennDiagram*[9] to integrate analytical results from the 5 GEO datasets.

C. Metabolomics Data

We isolated CD133+ and CD133- populations from patient-derived CRC lines we have previously described. 6 samples were collected and FACS sorted using CD133 antibodies, and their metabolites levels were detected and measured by a high-resolution qExactive liquid chromatography-mass spectrometer (LC-MS).

D. Pathway Analysis

We performed pathway analyses on the CCIC-regulated genes identified from the GEO datasets using Gene Set Enrichment Analysis (GSEA)[10] and the Kyoto Encyclopedia of

Genes Genomes (KEGG)[11]. The metabolomics and transcriptomics data were then integrated by using the KEGG Pathway Online Module.

III. Results

A. Differential Transcriptional Profiling in CD133+ Cells

Transcriptional profiling of the five CRC sets showed that CD133+ and CD133- cells have on average 3178 significant-differential (P-value < 0.05) genes, of which 1628 genes are upregulated and 1550 genes are downregulated in CD133+ cells (Figure 1, Table II). To compare the gene expression profiling of the 5 sets of microarray data from 4 different platforms, we converted the probe IDs of each platform into unified Entrez format. A total of 5521 genes were identified as upregulated in CD133+ cells and 5527 genes as downregulated in CD133+ cells. Based on the frequencies of these identified genes across the 5 data sets, 946 genes were upregulated in more than one CRCs set and 718 genes were downregulated in more than one CRCs set (darker regions in Figure 2). However, only one gene (CD133) was consistently upregulated in all 5 datasets, and no gene was consistently downregulated in all 5 datasets.

Our differential analysis suggests that the transcriptional landscapes are consistently altered in CD133+ CCICs vs. CD133- CRC cells, and many genes are involved. However, CD133+ CCICs from different CRC tumor origins involve diverse genes, reflecting their various genetic backgrounds.

B. Pathway Enrichment Analysis

To find out what pathways these genes are involved with, we performed GSEA and KEGG pathway analyses on the identified list of differential genes. Pathway analyses showed that the curated gene sets of METABOLIC PATHWAYS, PATHWAY IN CANCER, and TRANSCRIPTIONAL MISREGULATION IN CANCER are most highly enriched. KEGG identified 282 enriched pathways, among which metabolic pathways scored highest gene hit rates – 50 curated metabolic pathways involve 614 differentially regulated metabolic genes. The metabolic alterations in CD133+ CCICs are visualized in the global metabolic map shown in Figure 3, wherein red lines represent upregulated enzymatic reactions and blue lines represent downregulated reactions. Darker color indicates a higher frequency of the metabolic gene across the 5 datasets. The global analysis suggests that metabolic pathways are highly reprogrammed in CD133+ CCICs.

C. Metabolomic Profiling

The gene and pathway enrichment analyses suggest that metabolic reprogramming may be a signature of CD133+ CCICs. To validate this hypothesis, we FACS sorted pure CD133+ and CD133- populations from patient-derived CRC lines we have previously published [3] and performed unbiased, high-resolution metabolomics to measure metabolite levels in triplicate samples using LC-MS. Differential analysis identified 54 metabolites that were differentially expressed in a statistically significant way (P-value < 0.05), among which 28 metabolites were upregulated in CD133+ CCICs and 26 metabolites were downregulated in CD133+ CCICs. The list of differential expressed metabolites is summarized in Table III.

D. Integrated Transcriptomic and Metabolomic Network Analysis

Above transcriptomic and metabolomic analyses suggest that metabolic pathways are altered in CD133+ CCICs across different CRC backgrounds. To investigate the connections between CCIC transcriptome and metabolome, we performed integrated KEGG pathway analysis on the list of differential genes and metabolites. The analysis identified carbohydrate metabolism (glycolysis, the TCA cycle) and cysteine and methionine metabolism as consistently altered in CCICs, with highly altered enzyme and metabolite levels (Figures 4, 5).

In most cells, glycolysis converts glucose into pyruvate, which then enters the TCA cycle. Compared to CD133- CRC cells, major glycolysis enzymes PGK1, BPGM, and ENO1 are significantly downregulated in CD133+ CCICs, indicating suppressed upstream activities (Figure 4). Moreover, ACSS1 (Acyl-CoA Synthetase Short-chain family member 1), a mitochondrial acetyl-CoA synthetase enzyme that catalyzes acetate to acetyl-CoA, is significantly upregulated in CD133+ CCICs, which potentially convert more upstream acetate into acetyl-CoA. On the other hand, downstream reactions to catalyze acetyl-CoA into malonyl-CoA and acetoacetyl-CoA are suppressed, hence forcing the extra acetyl-CoA to enter the TCA cycle. Consistently, the upstream metabolites in the TCA-cycle, i.e., citrate, aconitate, isocitrate, and α -ketoglutarate, are highly accumulated in CD133+ CCICs. At the end point of the TCA cycle, significantly upregulated PCK1 and PCK2 form a positive feedback loop to further facilitate the conversion of oxaloacetate into acetyl-CoA to fuel the TCA cycle. It has been shown that reprogramming of energy metabolism is a hallmark of pluripotent stem cells reprogram [14][15]. Stem cells influence epigenetic regulation such as histone acetylation and histone/DNA demethylation by controlling intermediate metabolite substrates Acetyl-CoA and α -ketoglutarate [16][17]. Therefore, altered energy metabolism and elevated Acetyl-CoA and α -ketoglutarate levels may explain some of the stem cell-like phenotypes observed in CD133+ CCICs.

Another consistently altered metabolic pathway is cysteine and methionine metabolism (Figure 5). In this pathway, enzyme CBS (Cystathionine- β -Synthase) combines homocysteine and serine to generate cystathionine and subsequently CTH catalyzes cystathionine into cysteine. Both CBS and CTH are significantly upregulated in CD133+ CCICs. Interestingly, cysteine has been reported to regulate neural stem cells through the CBS/H₂S pathway[18]. In methionine metabolism, DNMT1/3L, SRM, AMD1, and MTAP, downstream enzymes that catalyze S-AdenosylMethionine (SAM), are significantly downregulated in CD133+ CCICs. The metabolomics data confirm that the downregulated enzymatic reactions lead to accumulation of SAM in CD133+ CCICs. Among the downregulated enzymes, DNMT1/3L is an important DNA methyltransferase that catalyzes the transfer of methyl groups from SAM to DNA. Therefore, altered methionine metabolism might impact DNA methylation in CD133+ CCICs.

IV. Discussion

CD133+ CCICs from different CRC tumors are likely to have diverse mechanisms. However, by perform system-level transcriptomic and metabolomic analyses on various CRC sources, we identified a distinct metabolic signature of CD133+ CCICs that involve

glycolysis, the TCA cycle, and cysteine/methionine metabolism. The metabolite substrates involved in epigenetic regulations are highly altered, suggesting a potential epigenetic link. RNA-seq, metabolic flux analysis (MFA), and functional assays are currently being performed to further establish such links. The identified metabolic signature provides insights into reported stem cell-like properties of CD133+ CCICs. The involved metabolic enzymes and metabolites may provide biomarkers for CRC diagnosis and prognosis. New CRC treatments may also target them to suppress CCICs in the tumor population to reduce the risk of relapse and metastasis.

Acknowledgment

We thank the other members of the Shen lab and the Locasale lab for useful discussion and generous support.

Resrach supported by NIH NIGMS R01GM95990 and NSF 1137269.

References

1. Ricci-Vitiani L, Lombardi DG, Pilozzi E, Biffoni M, Todaro M, Peschle C, De Maria R. Identification and expansion of human colon-cancer-initiating cells. *Nature*. 2007; 445:111–115. [PubMed: 17122771]
2. O'Brien CA, Pollett A, Gallinger S, Dick JE. A human colon cancer cell capable of initiating tumour growth in immunodeficient mice. *Nature*. 2007; 445:106–110. [PubMed: 17122772]
3. Bu P, Chen K-Y, Chen JH, Wang L, Walters J, Shin YJ, Goerger JP, Sun J, Witherspoon M, Rakhilin N, Li J, Yang H, Milsom J, Lee S, Zipfel W, Jin MM, Gümü ZH, Lipkin SM, Shen X. A microRNA miR-34a-regulated bimodal switch targets notch in colon cancer stem cells. *Cell Stem Cell*. 2013 May; 12(5):602–615. [PubMed: 23642368]
4. Ren F, Sheng W-Q, Du X. CD133: a cancer stem cells marker, is used in colorectal cancers. *World J. Gastroenterol*. 2013; 19:2603–2611. [PubMed: 23674867]
5. Medema JP. Cancer stem cells: the challenges ahead. *Nat. Cell Biol*. 2013; 15:338–344. [PubMed: 23548926]
6. Barrett T, Troup DB, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, Marshall KA, Phillippy KH, Sherman PM, Muerter RN, Holko M, Ayanbule O, Yefanov A, Soboleva A. NCBI GEO: archive for functional genomics data sets--10 years on. *Nucleic Acids Res*. 2011; 39:D1005–D1010. [PubMed: 21097893]
7. Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, Ellis B, Gautier L, Ge Y, Gentry J, Hornik K, Hothorn T, Huber W, Iacus S, Irizarry R, Leisch F, Li C, Maechler M, Rossini AJ, Sawitzki G, Smith C, Smyth G, Tierney L, Yang JYH, Zhang J. Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol*. 2004; 5:R80. [PubMed: 15461798]
8. Davis JW. Bioinformatics and Computational Biology Solutions Using R and Bioconductor. *Journal of the American Statistical Association*. 2007; 102:388–389.
9. Chen H, Boutros PC. VennDiagram: a package for the generation of highly-customizable Venn and Euler diagrams in R. *BMC Bioinformatics*. 2011; 12:35. [PubMed: 21269502]
10. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A*. 2005 Oct; 102(43):15545–15550. [PubMed: 16199517]
11. Kanehisa M, Goto S, Kawashima S, Okuno Y, Hattori M. The KEGG resource for deciphering the genome. *Nucleic Acids Res*. 2004 Jan; 32(Database issue):D277–D280. [PubMed: 14681412]
12. Jaksch M, Múnera J, Bajpai R, Tersikh A, Oshima RG. Cell cycle-dependent variation of a CD133 epitope in human embryonic stem cell, colon cancer, and melanoma cell lines. *Cancer Res*. 2008; 68:7882–7886. [PubMed: 18829544]

13. Chao C, Carmical JR, Ives KL, Wood TG, Aronson JF, Gomez GA, Djukom CD, Hellmich MR. CD133+ colon cancer cells are more interactive with the tumor microenvironment than CD133- cells. *Laboratory Investigation*. 2012; 92:420–436. [PubMed: 22157717]
14. Zhang J, Nuebel E, Daley GQ, Koehler CM, Teitell Ma. Metabolic Regulation in Pluripotent Stem Cells during Reprogramming and Self-Renewal. *Cell Stem Cell*. 2012 Nov; 11(5):589–595. [PubMed: 23122286]
15. Shyh-Chang N, Zheng Y, Locasale JW, Cantley LC. Human pluripotent stem cells decouple respiration from energy production. *EMBO J*. 2011 Dec; 30(24):4851–4852. [PubMed: 22166995]
16. Lu C, Thompson CB. Metabolic regulation of epigenetics. *Cell Metab*. 2012; 16:9–17. [PubMed: 22768835]
17. Johnson C, Warmoes MO, Shen X, Locasale JW. Epigenetics and cancer metabolism. *Cancer Letters*. 2013
18. Wang Z, Liu D, Wang F, Zhang Q, Du Z, Zhan J, Yuan Q, Ling E-A, Hao A. L-cysteine promotes the proliferation and differentiation of neural stem cells via the CBS/H(2)S pathway. *Neuroscience*. 2013

Figure 1.

Differentially regulated genes in the 5 GEO datasets. Upper panels: Heatmaps of microarray datasets. Each row is one gene, and each column is a sample. Red and blue colors represent high and low expression levels respectively. Lower panels: Volcano plots of differentially regulated genes calculated by p-values and fold changes, Each dot represents a gene. The red dots represent significantly upregulated genes, and the blue dots represent significantly downregulated genes.

Figure 2.

Venn Diagrams of differentially regulated genes that are statistically significant. Each circle represents a GEO dataset. The number in each intersecting region represents the number of overlapping genes.

Figure 3.

A global metabolic map showing metabolic alterations in CD133+ CCICs. Nodes represent metabolites, and lines represent enzymatic reactions. Red color refers to significantly upregulated enzymatic genes, and blue color refers to significantly downregulated enzymatic genes. Color saturation represents the frequency of the gene across GEO datasets, from 5 (darkest) to 1 (lightest).

Figure 4.

Altered carbohydrate metabolism in CCIC. Arrows represent metabolic flows regulated by marked metabolic enzymes. Metabolites are shown as nodes. Upregulated genes and metabolites are shown in red, downregulated genes and metabolites are shown in blue.

Figure 5.

Altered cysteine and methionine metabolism in CCIC. The annotation is consistent with that of Figure 4.

TABLE I

GEO Microarray Data on CRCs.

Microarray Data List			
<i>GEO accession number</i>	<i>Bio marker</i>	<i>Cell Type</i>	<i>Sample Number</i>
GSE11757[12]	CD133	CACO-2	3 × CD133+ 3 × CD133–
GSE23295	CD133	SW620	2 × CD133+ 2 × CD133–
GSE24747	CD133	CACO-2	3 × CD133+ 3 × CD133–
GSE34053[13]	CD133	Patient specimen	3 × CD133+ 3 × CD133–
GSE38049	CD133	HCT116	3 × CD133+ 3 × CD133–

TABLE II

Summary of Differential Analysis

GEO accession number	Total Significant-Differential Genes	Significant- Differential Upregulated Genes	Significant- Differential Downregulated Genes
GSE11757	1676	861	815
GSE23295	6935	3089	3846
GSE24747	4011	2723	1288
GSE34053	2230	954	1276
GSE38049	1038	512	526
AVERAGE	3178	1628	1550

TABLE III

List of Significantly-Differential Metabolites

<i>54 differentially expressed metabolites (p-value < 0.05)</i>	
Upregulated Metabolites (28 metabolites)	Downregulated Metabolites (26 metabolites)
fructose-16-bisphosphate	serine
inosine	adenine
Phosphorylcholine	betaine aldehyde
UTP	2-keto-isovalerate
CDP-ethanolamine	sarcosine
malate	2-hydroxygluturate
ATP	Atrolactic acid
dGTP	Phenyllactic acid
S-methyl-5-thioadenosine	hydroxyphenylpyruvate
orotate	aspartate
uridine	Pyroglutamic acid
Maleic acid	histidine
octulose-monophosphate (O8P-O1P)	folate
NAD ⁺	2-Isopropylmalic acid
glutathione disulfide	glycolate
4-aminobutyrate	phenylpyruvate
dimethylglycine	Hydroxyisocaproic acid
S-adenosyl-L-methionine	glucono-D-lactone
UDP-N-acetyl-glucosamine	Pyridoxamine
citrulline	Ascorbic acid
aconitate	N-acetyl-glutamine
ADP	acetoacetate
dGDP	sn-glycerol-3-phosphate
citrate-isocitrate	Pyrophosphate
betaine	6-phospho-D-glucono-15-lactone
3-phosphoglycerate	2,3-dihydroxybenzoic acid
alanine	
a-ketoglutarate	