Published in final edited form as:

J Med Genet. 2009 September ; 46(9): 626–634. doi:10.1136/jmg.2008.062471.

Genomic duplication resulting in increased copy number of genes encoding the sister chromatid cohesion complex conveys clinical consequences distinct from Cornelia de Lange

Jiong Yan¹, Feng Zhang¹, Ellen Brundage¹, Angela Scheuerle², Brendan Lanpher³, Robert P. Erickson⁴, Zoe Powis⁴, Haynes B. Robinson⁵, Pamela L. Trapane⁶, Danuta Stachiw-Hietpas⁷, Kim M. Keppler-Noreuil⁸, Seema R. Lalani^{1,11}, Trilochan Sahoo^{1,11}, A. Craig Chinault¹, Ankita Patel^{1,11}, Sau Wai Cheung^{1,11}, and James R. Lupski^{1,9,10,11,*}

¹Department of Molecular and Human Genetics, Houston, Texas 77030

²Tesserae Genetics, Dallas, Texas 75230

³Department of Pediatrics, Vanderbilt University Medical Center, Nashville, TN 37232

⁴Department of Pediatrics, University of Arizona College of Medicine, Tucson, AZ 85724

⁵Department of Pathology, Akron Children's Hospital, Akron, OH 44308

⁶Children's Specialty Group, Medical College of Wisconsin, Milwaukee, WI 53201

⁷Children's Hospital of Wisconsin, Milwaukee, WI 53201

⁸Dept Pediatrics/Med Genetics, Univ Iowa Hosps & Clinics, Iowa City, IA 52242

⁹Department of Pediatrics, Baylor College of Medicine, Houston, Texas 77030

¹⁰Texas Children's Hospital, Houston, Texas 77030

¹¹Medical Genetics Laboratories (MGL), Houston, Texas 77030

Abstract

Cornelia de Lange Syndrome (CdLS) is a multisystem congenital anomaly disorder. Heterozygous point mutations in three genes (*NIPBL*, *SMC3* and *SMC1A*), encoding components of the sister chromatid cohesion apparatus, are responsible for ~ 50-60% of CdLS cases. Recent studies have revealed a high degree of genomic rearrangements (e.g. deletions and duplications) in the human genome, which result in gene copy number variations (CNV). CNVs have been associated with a wide range of both Mendelian and complex traits including disease phenotypes such as Charcot-

*Correspondence: James R. Lupski, M.D., Ph. D., Dept. of Molecular & Human Genetics, Baylor College of Medicine, Room 604B, One Baylor Plaza, Houston, TX 77030, U.S.A., Tel: 713-798 6530, FAX: 713-798 5073, E-mail: jlupski@bcm.tmc.edu.

The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive license (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in *Journal of Medical Genetics* and any other BMJPG products and sublicenses such use and exploit all subsidiary rights, as set out in our license (<http://jmg.bmj.com/misc/fora/licenceform.shtml>).

We have obtained the written consent from the patient or his/her legal guardian for publication of the images in print and online.

Publisher's Disclaimer: Online First contains unedited articles in manuscript form that have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Online First articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Online First articles must include the digital object identifier (DOIs) and date of initial publication.

Marie-Tooth type 1A, Pelizaeus-Merzbacher, Parkinson, Alzheimer, autism and schizophrenia. Increased versus decreased copy number of the same gene can potentially cause either similar or different clinical features. We identified duplications on chromosomes 5 or X using genome wide array Comparative Genomic Hybridization (aCGH). The duplicated regions contain either the *NIPBL* or the *SMC1A* genes. Junction sequences analyses revealed the involvement of three genomic rearrangement mechanisms. The patients share some common features including mental retardation, developmental delay, sleep abnormalities, and craniofacial and limb defects. The systems affected are the same as in CdLS, but clinical manifestations are distinct from CdLS; particularly the absence of the CdLS facial gestalt. Our results confirm the notion that duplication CNV of genes can be a common mechanism for human genetic diseases. Defining the clinical consequences for a specific gene dosage alteration represents a new “reverse genomics” trend in medical genetics that is reciprocal to the traditional approach of delineation of the common clinical phenotype preceding the discovery of the genetic etiology.

Keywords

Copy number changes; CNV; CdLS; *NIPBL*; cohesion complex

Introduction

The human genome has been found to contain a large number of CNVs [1-3], which encompass ~ 12% of the genome and 14.5% of the genes implicated in human diseases. Copy number alteration of dosage sensitive genes has long been linked to genetic diseases. Charcot-Marie-Tooth disease type 1A [CMT1A, MIM118220] and hereditary neuropathy with liability to pressure palsies [HNPP, MIM162500] result from reciprocal duplication and deletion of the dosage sensitive *PMP22* gene, respectively [4-7]. Deletions and duplications of *PLP1* (proteolipid protein gene) both result in Pelizaeus-Merzbacher disease [PMD, MIM312080], an X-linked dysmyelinating disorder [8-10]. Recent studies have linked CNVs to a broader range of diseases including complex traits, such as Parkinson [11], Alzheimer [12], autism [13-14] and schizophrenia [15-18]. These findings indicate that CNV assessment should be incorporated into the investigations of the genetic bases of diseases including inherited, complex, and sporadic traits [19].

Copy number variation (CNV) results from genomic rearrangements such as duplication and deletion. Three mechanisms have been implicated for the genomic rearrangements including nonallelic homologous recombination (NAHR), nonhomologous end joining (NHEJ) and the recently proposed DNA replication mechanism of Fork Stalling and Template Switching (FoSTeS) [20-24]. NAHR accounts for the majority of the genomic disorders [20] associated with recurrent rearrangements. It occurs mostly between low copy repeats (LCRs) and occasionally using repetitive sequences such as *Alu* or L1 as recombination substrates. NHEJ has been associated with several non-recurrent rearrangements [22]. Further study of the molecular mechanism for non-recurrent rearrangements, especially the complex rearrangements, has recently led to FoSTeS, a replication-based mechanism [24].

Cornelia de Lange Syndrome [CdLS, MIM122470] is a multiple congenital anomaly condition with growth and mental retardation, characteristic facial features, limb defects, neurobehavioral problems, and abnormalities in gastrointestinal, genitourinary and cardiovascular systems [25-27]. The prevalence is estimated to be 1/10,000 [28]. Marked phenotypic variability has been documented. According to the severity of the phenotype, patients have been classified as classical and mild types [25,26,29,30]. Classical facial features consist of synophrys, long eyelashes, long philtrum, thin upper lip and down turned mouth and provide a readily recognizable facial gestalt characteristic of the syndrome.

Heterozygous mutations in the *NIPBL* gene [MIM608667] have been found in ~ 50% of the CdLS cases [31-36]. Genotype-phenotype correlation analyses revealed that patients carrying *NIPBL* point mutations tend to have a more severe phenotype than mutation negative patients [31,34-36]. Furthermore, there is a trend that for some clinical features, such as growth delay and limb defects, individuals with missense mutations have a milder phenotype than those with truncating mutations. This suggests that *NIPBL* is a dosage sensitive gene. Recently, mutations in two additional cohesin subunit encoding genes, *SMC3* [MIM606062] and *SMC1A* [MIM300040], have been found to contribute to ~ 5% of CdLS cases [37-38]. These patients demonstrated a mild to moderate phenotype.

Our medical genetics diagnostic laboratory [39-40] identified six cases carrying duplications on chromosome 5 or on the X chromosome with the duplicated regions encompassing *NIPBL* or *SMC1A*. This presented a unique opportunity to study the dosage effect of the genes encoding the subunits of the cohesion complex using a “reverse genomics” approach [41]. These patients have mental retardation, developmental delay, neurobehavioral abnormalities, and craniofacial and limb dysmorphisms. The affected systems are the same as in CdLS. Genome wide 244K oligonucleotide aCGH analyses were performed to refine the size extent, and gene contents of the duplicated intervals and “genomotype” – phenotype correlations were analyzed. Junction sequences were obtained for four patients to study the mechanisms resulting in the duplications. Our results not only confirm that CNV of genes should be considered in searching for etiology of human genetic diseases, but also represent a new “reverse genomics” trend, which is defining the clinical consequences for CNV of a specific genomic interval and attempts to determine the specific gene(s) whose dosage alteration is potentially causative for the phenotype.

Materials and Methods

Subjects and clinical evaluation

Patients were recruited by their referring physicians after our medical genetics diagnostic laboratory initial genomic analyses were performed by the clinical implementation of array Comparative Genomic Hybridization (aCGH) using a targeted array. Clinical phenotype was assessed by experienced medical geneticists. Studies were approved by the Institutional Review Board (IRB) of Baylor College of Medicine and all participants or parental guardians volunteered after informed consent.

Genome wide 244k oligonucleotide aCGH analyses

To perform genome wide 244K oligonucleotide aCGH analyses, genomic DNA was isolated from peripheral blood with a Puregene kit (Gentra Systems). aCGH analyses were performed according to the manufacturer's protocol (Agilent Technologies). DNA samples (~ 2 µg) from patients and gender-matched normal individuals as reference were digested with *AluI* and *RsaI* and labeled with Cy5 or Cy3 using a BioPrime array CGH genomic labeling module (Invitrogen). After hybridization, slides were scanned and analyzed for relative gain or loss of fluorescent signals from hybridization of the patient and reference DNAs. Genomic region analyses were performed according to the human reference sequence build 36.1 (UCSC genome browser: <http://genome.ucsc.edu>) with the software provided by Agilent.

The junctional sequences analyses

Long range PCR primers were designed from the aCGH genomic coordinates to traverse the position of change from gain to “no gain”. Under the assumption of tandem duplication, outwardly facing primers were used to amplify breakpoint junctions. Obtained PCR products were sent for Sanger dideoxy DNA sequencing (SeqWright, Texas).

Results

Chromosomal Microarray Analyses (CMA)

Six unrelated patients were found to harbor gains of a gene encoding a cohesion subunit on either chromosome 5 (*NIPBL*) or the X chromosome (*SMC1A*) by CMA [39] (table 1). The gains were confirmed by interphase fluorescence *in situ* hybridization (FISH) analyses that revealed the extra copy mapped adjacent to the region consistent with interstitial duplication (data not shown). A genomic interval interrogated by BAC clones RP11-317I23, RP11-7M4 (contains *NIPBL*) and RP11-209M10 on chromosome 5 was found to be duplicated in patients 1 and 3. Patients 2 and 5 contain the duplications interrogated by RP11-7M4 and RP11-209M10. A region identified by clones RP11-317I23 and RP11-7M4 is duplicated in patient 4. Patient 6 has a gain in copy number in the proximal region of the short arm of chromosome X covered by BAC clones RP11-416B14, RP11-52N6 (contains *SMC1A*), RP11-70P16. The duplicated regions contain genes in which point mutations result in CdLS. The gains were not detected in the available parental samples (both parents for two patients and 3 maternal samples for three patients). These specific CNVs were not present in the CNV databases of genomic variants (<http://projects.tcag.ca/variation/>).

High resolution genomic analyses of duplications

In order to analyze the duplications at a higher resolution, we performed genome wide 244k oligonucleotide aCGH analyses (table 1, fig. 1). Patients 1 and 2 appear to have a similar sized 390 kb duplication on chromosome 5, which includes the *NIPBL* gene and exon 3 to the 3'UT region of the *FLJ13231* gene, a gene of unknown function. Patient 3 carries a larger (1.07 Mb) duplication on chromosome 5, which contains *SLC1A3*, *NUP155*, and exons 1 to 9 of the *WDR70* gene in addition to the *NIPBL* and *FLJ13231* genes. *NIPBL* is the only gene located within the 250 kb duplication in patient 4. A 330 kb duplication has

been found in patient 5, which includes exons 39 to 47 of *NIPBL*, *FLJ13231*, *NUP155*, and exons 1 and 2 of *WDR70*. Patient 6 has a 7.14 Mb duplication on X chromosome, which contains about 174 genes (fig. 2). Among them are the *WAS*, *CLCN5*, and *SMC1A* genes responsible for Wiskott-Aldrich syndrome, X-linked nephrolithiasis, and CdLS respectively [37,42,43]. This large duplication also includes the *HUWE1* gene; duplication and point mutation of which have recently been associated with X-linked mental retardation (XLMR) in males [44]. The high resolution 244k Agilent array results are consistent with the CMA findings and further refine the size, extent, and genomic content of duplication and breakpoint region. Additional CNVs identified are provided as a supplementary table.

Breakpoint junctional sequence analyses

To further fine map the duplications and potentially infer rearrangement mechanism(s) from the junctional sequences, we performed breakpoint mapping. The PCR amplification of breakpoint junctions were achieved in 4 out of 6 patients (Pt 2, 3, 4, and 5, fig. 3). Although the oligonucleotide aCGH analyses mapped the breakpoints of patients 1 and 2 to the same genomic interval, the primers that successfully amplified the breakpoint junction of patient 2 failed in patient 1. Therefore these two duplications were not identical. At the breakpoint junction of patient 2, only one base pair was shared between the proximal and distal reference sequences, suggesting the involvement of the non-homologous end joining (NHEJ) mechanism in this rearrangement. The rearrangement in patient 3 was complex. An additional small (176 bp) inverted duplication was observed between the two copies of the large genomic duplication. The distal breakpoint was hypothesized to be caused by non-allelic homologous recombination (NAHR) between two L1 repeats. However, since a 12-bp microhomology (ACAATTAAAAGA) was found, the distal breakpoint was also consistent with our recently proposed DNA replication based FoSTeS (replication Fork Stalling and Template Switching) mechanism while the microhomology could alternatively act in template switching and annealing between replication forks, restarting DNA replication, and generating the DNA rearrangement. No microhomology but the addition of a T was observed in the proximal breakpoint, which was consistent with NHEJ. Similar to patient 2, only one base pair was shared by reference sequences in the breakpoint of patient 4, suggesting that NHEJ caused the duplication. In patient 5, the microhomology of TTT was consistent with NHEJ or FoSTeS $\times$ 1. In aggregate, the above observations suggest different mechanisms (NHEJ, FoSTeS, and/or NAHR) are responsible for the nonrecurrent genomic duplications that can include the *NIPBL* gene.

Clinical phenotype

Clinical features are summarized in table 2. All patients had developmental delay and mental retardation. The majority of the patients had low birth weight and became overweight in adulthood. Hypotonia and seizures were commonly seen and periventricular leukomalacia (2/2) was observed in two head imaging studies (one CT and one MRI) available. Patients shared some craniofacial dysmorphic features including microbrachycephaly (3/6), frontal bossing (3/6), a short philtrum (4/6), low-set ears (3/6), with one patient having a significant skull deformity (patient 4, fig. 4G and 4H). The patients also had long-fingers (3/6) and neurobehavioral abnormalities including sleep

disturbances (3/6) (table 2). Available individual clinical details are provided as supplementary material.

Discussion

Both increased and decreased copy number of a dosage sensitive gene can cause congenital diseases. This is the case both for autosomal genes and genes on the X chromosome. Reciprocal duplication and deletion of the *PMP22* gene result in CMT1A and HNPP respectively [4-6], both are inherited neuropathies but with distinct clinical presentations. Williams -Beuren syndrome (WBS) and its reciprocal duplication syndrome are the results of deletion and duplication of a region on 7q11.23 [45-48]. WBS is featured by cognitive deficiency with relative preservation of linguistic abilities, neurobehavioral problems, recognizable facial dysmorphism, and cardiac and ophthalmologic abnormalities. Patients with the apparent reciprocal duplication of the WBS region have mild dysmorphisms and a severe delay in expressive language [46-48]. In most cases, the increased dosage from duplication causes a milder phenotype than deletion of the same region as in the Smith-Magenis syndrome [SMS, MIM182290] and its reciprocal duplication syndrome, Potocki-Lupski syndrome [PTLS, MIM610883] [49-50], as well as in the duplication and loss of function of the *MECP2* gene [51-53]. Patients with SMS have characteristic craniofacial, skeletal and neurobehavioral features as well as defects in multiple organ systems. PTLS is featured by mental retardation and behavioral problems including autistic features and sleep apnea. Sometimes both duplication and deletion result in similar phenotype as in Pelizaeus-Merzbacher disease (PMD), an X-linked dysmyelinating disorder [8-10].

Heterozygous mutations of the *NIPBL* or the *SMC1A* genes results in CdLS, which is characterized by mental retardation, developmental delay, neurobehavioral abnormalities, and craniofacial and limb defects [25,26,32,33,37]. We identified six patients with genomic duplications containing either *NIPBL* (N=5) or *SMC1A*. These individuals also have mental retardation, developmental delay, and neurobehavioral abnormalities such as hypotonia and sleep disturbances (table 2). Only a limited number of patients have been observed [five that have duplications including *NIPBL* and only one whose large > 7 Mb duplication includes *SMC1A*] and the clinical manifestations may be complicated by *in utero* exposures (maternal diabetes for patient 4 and ethanol during the pregnancy of patient 5) and different sized genomic duplications. Nevertheless, the patients clearly do not have CdLS and thus we can conclude that duplications harboring genes for the cohesion complex present with phenotypes distinct from those observed with haploinsufficiency effects for the same genes.

Common craniofacial features include frontal bossing, broad nasal root, low set ears, short philtrum and high palate. The shared facial defects make the patients appear similar to each other (fig. 4), but with this limited patient set from different ages no recognizable facial gestalt is apparent. The craniofacial features are different from those that are typical for CdLS, the latter of which include synophrys, long eyelashes, long philtrum, thin upper lip and down turned mouth [25-26]. However, some facial abnormalities have been frequently observed in patients with CdLS, such as low set ears, high palate, and ocular anomalies. Patients with duplication of the gene for cohesion complex proteins also have increased weight and long fingers, which are opposite to characteristics observed in CdLS.

NIPBL is the only gene duplicated in patient 4 (table 1). But it is hard to determine whether the severe skull deformity is the result of the duplication. The skull deformity complicated the assessment of the phenotype. Further follow up will hopefully reveal the effect of the duplication. Patients 1 and 2 have a similar sized genomic duplication, which results in the duplication of *NIPBL* and exon 3 and 3' untranslated region of *FLJ13231*, a gene of unknown function. Both *NIPBL* and *FLJ13231* are duplicated in patient 3, who has clinical features similar to patients 1 and 2. The duplication in patient 5 results in the duplication of *FLJ13231* and *Nup155*, as in patient 3. But the *NIPBL* gene is only partially duplicated, which most likely does not result in the increased expression of *NIPBL*. Indeed, the phenotype in patient 5 is the least similar to other patients, and the phenotypic analysis is likely complicated by the maternal ingestion of alcohol in the first trimester. Based on these observations, *NIPBL* is likely the major dosage sensitive gene for the phenotype, but more patients are necessary to further delineate the effects of the increased dosage of the *NIPBL* gene.

Patient 6 has a large duplication containing the *SMC1A* gene. Her phenotype is similar to patient 1 and 2; however, with only one patient available it is difficult to assess if other genes also contribute to the phenotype. The duplication contains the *WAS* gene, responsible for Wiskott-Aldrich syndrome, featured by recurrent infections and eczema [42], that have also been observed in patient 6. Whether this is the result of the duplication of *WAS* needs further investigation. This female patient is also duplicated for the *HUWE1* gene, duplication or point mutation of which has been recently associated with XLMR in males. Five of the families segregating XLMR in association with duplication of *HUWE1* also include the *SMC1A* within their duplications [44]. However, these males are described as having nonsyndromic MR [44]. For all six patients reported herein with duplications including genes encoding subunits of the cohesin complex, other genomic CNV were identified (supplementary table), but most are present in the database of genomic variants. Whether such variants can sometimes modify the ultimate phenotype is unclear at this time.

Three mechanisms have been implicated for the genomic rearrangements including NAHR, NHEJ and recently proposed FoSTeS [20-24]. NAHR occurs mostly between low copy repeats (LCRs) and occasionally through repetitive sequences such as *Alu* or *L1*. Sequence analysis surrounding the duplication breakpoints failed to detect any LCRs (data not shown) and our junction analyses revealed the involvement of all three mechanisms. Despite the similar duplication in aCGH analyses for patients 1 and 2, our results indicate that a more complicated rearrangement may have occurred in patient 1.

A tremendous amount of copy number variations (CNVs) has been identified in the human genome recently [1-3]. CNVs have been associated with multiple congenital diseases as well as some complex traits such as autism and schizophrenia [13-15]. Our findings confirm the notion that CNV of genes involved in dominant Mendelian traits could be a common mechanism for human genetic diseases. We further document that phenotypes conveyed by duplication of dosage sensitive gene may be milder than those caused by loss-of-function due to haploinsufficiency caused by deletion or point mutation. The distinct phenotype conveyed by duplication versus deletion may challenge discovery of duplication alleles at a given disease locus.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the patients and their families for their generous participation. This work was supported in part by a grant from the National Institute of Child Health and Human development (PO1 HD39420) and the Baylor College of Medicine Mental Retardation Research Center (MRRC) (HD 2406407).

References

1. Khaja R, Zhang J, MacDonald JR, He Y, Joseph-George AM, Wei J, Rafiq MA, Qian C, Shago M, Pantano L, Aburatani H, Jones K, Redon R, Hurles M, Armengol L, Estivill X, Mural RJ, Lee C, Scherer SW, Feuk L. Genome assembly comparison identifies structural variants in the human genome. *Nat Genet.* 2006; 38:1413–1418. [PubMed: 17115057]
2. Redon R, Ishikawa S, Fitch KR, Feuk L, Perry GH, Andrews TD, Fiegler H, Shaperro MH, Carson AR, Chen W, Cho EK, Dallaire S, Freeman JL, González JR, Gratacòs M, Huang J, Kalaitzopoulos D, Komura D, MacDonald JR, Marshall CR, Mei R, Montgomery L, Nishimura K, Okamura K, Shen F, Somerville MJ, Tchinda J, Valsesia A, Woodward C, Yang F, Zhang J, Zerjal T, Zhang J, Armengol L, Conrad DF, Estivill X, Tyler-Smith C, Carter NP, Aburatani H, Lee C, Jones KW, Scherer SW, Hurles ME. Global variation in copy number in the human genome. *Nature.* 2006; 444:444–454. [PubMed: 17122850]
3. Lupski JR. Structural variation in the human genome. *N Engl J Med.* 2007; 356:1169–1171. [PubMed: 17360997]
4. Lupski JR, de Oca-Luna RM, Slaugenhaupt S, Pentao L, Guzzetta V, Trask BJ, Saucedo-Cardenas O, Barker DF, Killian JM, Garcia CA, Chakravarti A, Patel PI. DNA duplication associated with Charcot-Marie-Tooth disease type 1A. *Cell.* 1991; 66:219–232. [PubMed: 1677316]
5. Chance PF, Abbas N, Lensch MW, Pentao L, Roa BB, Patel PI, Lupski JR. Two autosomal dominant neuropathies result from reciprocal DNA duplication/deletion of a region on chromosome 17. *Hum Mol Genet.* 1994; 3:223–228. [PubMed: 8004087]
6. Lupski JR. Charcot-Marie-Tooth polyneuropathy: duplication, gene dosage, and genetic heterogeneity. *Pediatr Res.* 1999; 45:159–165. [PubMed: 10022584]
7. Lupski JR, Wise CA, Kuwano A, Pentao L, Parke JT, Glaze DG, Ledbetter DH, Greenberg F, Patel PI. Gene dosage is a mechanism for Charcot-Marie-Tooth disease type 1A. *Nat Genet.* 1992; 1:29–33. [PubMed: 1301995]
8. Inoue K, Osaka H, Sugiyama N, Kawanishi C, Onishi H, Nezu A, Kimura K, Yamada Y, Kosaka K. A duplicated PLP gene causing Pelizaeus-Merzbacher disease detected by comparative multiplex PCR. *Am J Hum Genet.* 1996; 59:32–39. [PubMed: 8659540]
9. Siermans EA, de Coe RF, De Wijs IJ, Van Oost BA. Duplication of the proteolipid protein gene is the major cause of Pelizaeus-Merzbacher disease. *Neurology.* 1998; 50:1749–1754. [PubMed: 9633722]
10. Inoue K, Osaka H, Thurston VC, Clarke JT, Yoneyama A, Rosenbarker L, Bird TD, Hodes ME, Shaffer LG, Lupski JR. Genomic rearrangements resulting in *PLP1* deletion occur by nonhomologous end joining and cause different dysmyelinating phenotypes in males and females. *Am J Hum Genet.* 2002; 71:838–853. [PubMed: 12297985]
11. Singleton AB, Farrer M, Johnson J, Singleton A, Hague S, Kachergus J, Hulihan M, Peuralinna T, Dutra A, Nussbaum R, Lincoln S, Crawley A, Hanson M, Maraganore D, Adler C, Cookson MR, Muenter M, Baptista M, Miller D, Blancato J, Hardy J, Gwinn-Hardy K. alpha-Synuclein locus triplication causes Parkinson's disease. *Science.* 2003; 302:841. [PubMed: 14593171]
12. Rovelet-Lecrux A, Hannequin D, Raux G, Le Meur N, Laquerriere A, Vital A, Dumanchin C, Feuillette S, Brice A, Vercelletto M, Dubas F, Frebourg T, Campion D. APP locus duplication causes autosomal dominant early-onset Alzheimer disease with cerebral amyloid angiopathy. *Nat Genet.* 2006; 38:24–26. [PubMed: 16369530]

13. Sebat J, Lakshmi B, Malhotra D, Troke J, Lese-Martin C, Walsh T, Yamrom B, Yoon S, Krasnitz A, Kendall J, Leotta A, Pai D, Zhang R, Lee YH, Hicks J, Spence SJ, Lee AT, Puura K, Lehtimäki T, Ledbetter D, Gregersen PK, Bregman J, Sutcliffe JS, Jobanputra V, Chung W, Warburton D, King MC, Skuse D, Geschwind DH, Gilliam TC, Ye K, Wigler M. Strong association of de novo copy number mutations with autism. *Science*. 2007; 316:445–449. [PubMed: 17363630]
14. Weiss LA, Shen Y, Korn JM, Arking DE, Miller DT, Fossdal R, Saemundsen E, Stefansson H, Ferreira MA, Green T, Platt OS, Ruderfer DM, Walsh CA, Altshuler D, Chakravarti A, Tanzi RE, Stefansson K, Santangelo SL, Gusella JF, Sklar P, Wu BL, Daly MJ. Autism Consortium. Association between microdeletion and microduplication at 16p11.2 and autism. *N Engl J Med*. 2008; 358:667–675. [PubMed: 18184952]
15. Walsh T, McClellan JM, McCarthy SE, Addington AM, Pierce SB, Cooper GM, Nord AS, Kusenda M, Malhotra D, Bhandari A, Stray SM, Rippey CF, Roccanova P, Makarov V, Lakshmi B, Findling RL, Sikich L, Stromberg T, Merriman B, Gogtay N, Butler P, Eckstrand K, Noory L, Gochman P, Long R, Chen Z, Davis S, Baker C, Eichler EE, Meltzer PS, Nelson SF, Singleton AB, Lee MK, Rapoport JL, King MC, Sebat J. Rare structural variants disrupt multiple genes in neurodevelopmental pathways in schizophrenia. *Science*. 2008; 320:539–543. [PubMed: 18369103]
16. Stefansson H, Rujescu D, Cichon S, Pietiläinen OP, Ingason A, Steinberg S, Fossdal R, Sigurdsson E, Sigmundsson T, Buizer-Voskamp JE, Hansen T, Jakobsen KD, Muglia P, Francks C, Matthews PM, Gylfason A, Halldorsson BV, Gudbjartsson D, Thorgeirsson TE, Sigurdsson A, Jonasdottir A, Jonasdottir A, Bjornsson A, Mattiasdottir S, Blondal T, Haraldsson M, Magnusdottir BB, Giegling I, Möller HJ, Hartmann A, Shianna KV, Ge D, Need AC, Crombie C, Fraser G, Walker N, Lonnqvist J, Suvisaari J, Tuulio-Henriksson A, Paunio T, Touloupoulou T, Bramon E, Di Forti M, Murray R, Ruggeri M, Vassos E, Tosato S, Walshe M, Li T, Vasilescu C, Mühleisen TW, Wang AG, Ullum H, Djurovic S, Melle I, Olesen J, Kiemeny LA, Franke B, Sabatti C, Freimer NB, Gulcher JR, Thorsteinsdottir U, Kong A, Andreassen OA, Ophoff RA, Georgi A, Rietschel M, Werge T, Petursson H, Goldstein DB, Nöthen MM, Peltonen L, Collier DA, St Clair D, Stefansson K, Kahn RS, Linszen DH, van Os J, Wiersma D, Bruggeman R, Cahn W, de Haan L, Krabbendam L, Myin-Germeys I. Genetic Risk and Outcome in Psychosis (GROUP). Large recurrent microdeletions associated with schizophrenia. *Nature*. 2008; 455:232–236. [PubMed: 18668039]
17. The International Schizophrenia Consortium. Rare chromosomal deletions and duplications increase risk of schizophrenia. *Nature*. 2008; 455:237–241. [PubMed: 18668038]
18. Lupski JR. Schizophrenia: Incriminating genomic evidence. *Nature*. 2008; 455:178–179. [PubMed: 18784712]
19. Lupski JR. Genomic rearrangements and sporadic disease. *Nat Genet*. 2007; 39(7 Suppl):S43–47. [PubMed: 17597781]
20. Lupski JR. Genomic disorders: structural features of the genome can lead to DNA rearrangements and human disease traits. *Trends Genet*. 1998; 14:417–422. [PubMed: 9820031]
21. Stankiewicz P, Lupski JR. Genome architecture, rearrangements and genomic disorders. *Trends Genet*. 2002; 18:74–82. [PubMed: 11818139]
22. Shaw CJ, Lupski JR. Implications of human genome architecture for rearrangement-based disorders: the genomic basis of disease. *Hum Mol Genet*. 2004; 13(Spec No 1):R57–64. [PubMed: 14764619]
23. Lupski JR, Stankiewicz P. Genomic disorders: molecular mechanisms for rearrangements and conveyed phenotypes. *PLoS Genet*. 2005; 1:e49. [PubMed: 16444292]
24. Lee JA, Carvalho CM, Lupski JR. A DNA replication mechanism for generating nonrecurrent rearrangements associated with genomic disorders. *Cell*. 2007; 131:1235–1247. [PubMed: 18160035]
25. Ireland M, Donnai D, Burn J. Brachmann-de Lange syndrome. Delineation of the clinical phenotype. *Am J Med Genet*. 1993; 47:959–964. [PubMed: 8291539]
26. Jackson L, Kline AD, Barr MA, Koch S. de Lange syndrome: a clinical review of 310 individuals. *Am J Med Genet*. 1993; 47:940–946. [PubMed: 8291537]
27. Liu J, Krantz ID. Cohesin and Human Disease. *Annu Rev Genomics Hum Genet*. 2008; 9:303–320. [PubMed: 18767966]

28. Opitz JM. The Brachmann-de Lange syndrome. *Am J Med Genet.* 1985; 22:89–102. [PubMed: 3901753]
29. Van Allen MI, Filippi G, Siegel-Bartelt J, Yong SL, McGillivray B, Zuker RM, Smith CR, Magee JF, Ritchie S, Toi A, Reynolds JF. Clinical variability within Brachmann-de Lange syndrome: A proposed classification system. *Am J Med Genet.* 1993; 47:947–958. [PubMed: 8291538]
30. Allanson JE, Hennekam RCM, Ireland M. De Lange syndrome: subjective and objective comparison of the classical and mild phenotypes. *J Med Genet.* 1997; 34:645–650. [PubMed: 9279756]
31. Gillis LA, McCallum J, Kaur M, DeScipio C, Yaeger D, Mariani A, Kline AD, Li HH, Devoto M, Jackson LG, Krantz ID. *NIPBL* mutational analysis in 120 individuals with Cornelia de Lange syndrome and evaluation of genotype-phenotype correlations. *Am J Hum Genet.* 2004; 75:610–623. [PubMed: 15318302]
32. Krantz ID, McCallum J, DeScipio C, Kaur M, Gillis LA, Yaeger D, Jukofsky L, Wasserman N, Bottani A, Morris CA, Nowaczyk MJM, Toriello H, Bamshad MJ, Carey JC, Rappaport E, Kawauchi S, Lander AD, Calof AL, Li HH, Devoto M, Jackson LG. Cornelia de Lange syndrome is caused by mutations in *NIPBL*, the human homolog of *Drosophila melanogaster* *Nipped-B*. *Nat Genet.* 2004; 36:631–635. [PubMed: 15146186]
33. Tonkin ET, Wang TJ, Lisgo S, Bamshad MJ, Strachan T. *NIPBL*, encoding a homolog of fungal Scc2-type sister chromatid cohesion proteins and fly Nipped-B, is mutated in Cornelia de Lange syndrome. *Nat Genet.* 2004; 36:636–641. [PubMed: 15146185]
34. Bhuiyan ZA, Klein M, Hammond P, van Haeringen A, Mannens MMAM, Van Berckelaer-Onnes I, Hennekam RCM. Genotype-phenotype correlations of 39 patients with Cornelia de Lange syndrome: the Dutch experience. *J Med Genet.* 2006; 43:568–575. [PubMed: 16236812]
35. Yan J, Saifi GM, Wierzbica TH, Withers M, Bien-Willner GA, Limon J, Stankiewicz P, Lupski JR, Wierzbica J. Mutational and genotype-phenotype correlation analyses in 28 Polish patients with Cornelia de Lange syndrome. *Am J Med Genet A.* 2006; 140:1531–1541. [PubMed: 16770807]
36. Selicorni A, Russo S, Gervasini C, Castronovo P, Milani D, Cavalleri F, Bentivegna A, Masciadri M, Domi A, Divizia MT, Sforzini C, Tarantino E, Memo L, Scarano G, Larizza L. Clinical score of 62 Italian patients with Cornelia de Lange syndrome and correlations with the presence and type of *NIPBL* mutation. *Clin Genet.* 2007; 72:98–108. [PubMed: 17661813]
37. Musio A, Selicorni A, Focarelli ML, Gervasini C, Milani D, Russo S, Vezzoni P, Larizza L. X-linked Cornelia de Lange syndrome owing to *SMC1L1* mutations. *Nat Genet.* 2006; 38:528–530. [PubMed: 16604071]
38. Deardorff MA, Kaur M, Yaeger D, Rampuria A, Korolev S, Pie J, Gil-Rodriguez C, Arnedo M, Loeys B, Kline AD, Wilson M, Lillquist K, Siu V, Ramos FJ, Musio A, Jackson LS, Dorsett D, Krantz ID. Mutations in cohesin complex members SMC3 and SMC1A cause a mild variant of cornelia de Lange syndrome with predominant mental retardation. *Am J Hum Genet.* 2007; 80:485–494. [PubMed: 17273969]
39. Cheung SW, Shaw CA, Yu W, Li J, Ou Z, Patel A, Yatsenko SA, Cooper ML, Furman P, Stankiewicz P, Lupski JR, Chinault AC, Beaudet AL. Development and validation of a CGH microarray for clinical cytogenetic diagnosis. *Genet Med.* 2005; 7:422–432. [PubMed: 16024975]
40. Lu X, Shaw CA, Patel A, Li J, Cooper ML, Wells WR, Sullivan CM, Sahoo T, Yatsenko SA, Bacino CA, Stankiewicz P, Ou Z, Chinault AC, Beaudet AL, Lupski JR, Cheung SW, Ward PA. Clinical implementation of chromosomal microarray analysis: summary of 2513 postnatal cases. *PLoS ONE.* 2007; 2:e327. [PubMed: 17389918]
41. Bi W, Sapir T, Shchelochkov OA, Zhang F, Withers M, Hunter JV, Levy T, Shinder V, Peiffer DA, Gunderson K, Nezarati MM, Shotts VA, Amato SS, Savage SK, Harris DJ, Day-Salvatore D, Horner M, Lu X, Sahoo T, Yanagawa Y, Obata K, Beaudet AL, Cheung SW, Eichele G, Martinez S, Lupski JR, Reiner O. Duplications in the 17p13.3 Miller-Dieker syndrome region: LIS1 increased expression affects human and mouse brain development. *Nat Genet.* 2008 in press.
42. Derry JM, Ochs HD, Francke U. Isolation of a novel gene mutated in Wiskott-Aldrich syndrome. *Cell.* 1994; 78:635–644. [PubMed: 8069912]
43. Fisher SE, van Bakel I, Lloyd SE, Pearce SH, Thakker RV, Craig IW. Cloning and characterization of *CLCN5*, the human kidney chloride channel gene implicated in Dent disease (an X-linked hereditary nephrolithiasis). *Genomics.* 1995; 29:598–606. [PubMed: 8575751]

44. Froyen G, Corbett M, Vandewalle J, Jarvela I, Lawrence O, Meldrum C, Bauters M, Govaerts K, Vandeleur L, Van Esch H, Chelly J, Sanlaville D, van Bokhoven H, Ropers HH, Laumonnier F, Ranieri E, Schwartz CE, Abidi F, Tarpey PS, Futreal PA, Whibley A, Raymond FL, Stratton MR, Fryns JP, Scott R, Peippo M, Sipponen M, Partington M, Mowat D, Field M, Hackett A, Marynen P, Turner G, Géczy J. Submicroscopic duplications of the hydroxysteroid dehydrogenase *HSD17B10* and the E3 ubiquitin ligase *HUWE1* are associated with mental retardation. *Am J Hum Genet.* 2008; 82:432–443. [PubMed: 18252223]
45. Francke U. Williams-Beuren syndrome: genes and mechanisms. *Hum Mol Genet.* 1999; 8:1947–1954. [PubMed: 10469848]
46. Somerville MJ, Mervis CB, Young EJ, Seo EJ, del Campo M, Bamforth S, Peregrine E, Loo W, Lilley M, Pérez-Jurado LA, Morris CA, Scherer SW, Osborne LR. Severe expressive-language delay related to duplication of the Williams-Beuren locus. *N Engl J Med.* 2005; 353:1694–1701. [PubMed: 16236740]
47. Berg JS, Brunetti-Pierri N, Peters SU, Kang SH, Fong CT, Salamone J, Freedenberg D, Hannig VL, Prock LA, Miller DT, Raffalli P, Harris DJ, Erickson RP, Cunniff C, Clark GD, Blazo MA, Peiffer DA, Gunderson KL, Sahoo T, Patel A, Lupski JR, Beaudet AL, Cheung SW. Speech delay and autism spectrum behaviors are frequently associated with duplication of the 7q11.23 Williams-Beuren syndrome region. *Genet Med.* 2007; 9:427–441. [PubMed: 17666889]
48. Orellana C, Bernabeu J, Monfort S, Roselló M, Oltra S, Ferrer I, Quiroga R, Martínez-Garay I, Martínez F. Duplication of the Williams-Beuren critical region: case report and further delineation of the phenotypic spectrum. *J Med Genet.* 2008; 45:187–189. [PubMed: 18310268]
49. Greenberg F, Lewis RA, Potocki L, Glaze D, Parke J, Killian J, Murphy MA, Williamson D, Brown F, Dutton R, McCluggage C, Friedman E, Sulek M, Lupski JR. Multi-disciplinary clinical study of Smith-Magenis syndrome (deletion 17p11.2). *Am J Med Genet.* 1996; 62:247–254. [PubMed: 8882782]
50. Potocki L, Bi W, Treadwell-Deering D, Carvalho CM, Eifert A, Friedman EM, Glaze D, Krull K, Lee JA, Lewis RA, Mendoza-Londono R, Robbins-Furman P, Shaw C, Shi X, Weissenberger G, Withers M, Yatsenko SA, Zackai EH, Stankiewicz P, Lupski JR. Characterization of Potocki-Lupski syndrome (dup(17)(p11.2p11.2)) and delineation of a dosage-sensitive critical interval that can convey an autism phenotype. *Am J Hum Genet.* 2007; 80:633–649. [PubMed: 17357070]
51. Amir RE, Van den Veyver IB, Wan M, Tran CQ, Francke U, Zoghbi HY. Rett syndrome is caused by mutations in X-linked *MECP2*, encoding methyl-CpG-binding protein 2. *Nat Genet.* 1999; 23:185–188. [PubMed: 10508514]
52. Van Esch H, Bauters M, Ignatius J, Jansen M, Raynaud M, Hollanders K, Lugtenberg D, Bienvenu T, Jensen LR, Gecz J, Moraine C, Marynen P, Fryns JP, Froyen G. Duplication of the *MECP2* region is a frequent cause of severe mental retardation and progressive neurological symptoms in males. *Am J Hum Genet.* 2005; 77:442–453. [PubMed: 16080119]
53. del Gaudio D, Fang P, Scaglia F, Ward PA, Craigen WJ, Glaze DG, Neul JL, Patel A, Lee JA, Irons M, Berry SA, Pursley AA, Grebe TA, Freedenberg D, Martin RA, Hsich GE, Khera JR, Friedman NR, Zoghbi HY, Eng CM, Lupski JR, Beaudet AL, Cheung SW, Roa BB. Increased *MECP2* gene copy number as the result of genomic duplication in neurodevelopmentally delayed males. *Genet Med.* 2006; 8:784–792. [PubMed: 17172942]
54. Basile E, Villa L, Selicorni A, Molteni M. The behavioural phenotype of Cornelia de Lange Syndrome: a study of 56 individuals. *J Intellect Disabil Res.* 2007; 51(Pt 9):671–681. [PubMed: 17845236]

Figure 1. aCGH analyses for patients. Map of the *NIPBL* region on chromosome 5 (top) with the extent of the gains detected shown by horizontal bars. Below are the aCGH results from the Agilent 244K arrays. A gain, as evidenced by log₂ ratios > 0.5, is denoted by red dots with each signifying the relative signal from a single interrogating oligonucleotide. To the right are given the sizes of the individual duplications.

Figure 2. aCGH analyses for patient 6. Shown on top is the *SMC1A* region on chromosome X. The gain detected is shown by a horizontal bar. Below is the aCGH result with the size given at the right.

Figure 3.

Breakpoint sequence analyses. The amplification of breakpoint junctions were achieved in four patients (Pt 2, 3, 4, and 5). The duplication regions were shown in dash rectangles. The breakpoint coordinates (NCBI Build 36.1) were also shown. The 1-bp mismatch (marked by asterisk) between the breakpoint junction sequence of patient 3 and its distal reference sequence resulted from the SNP-rs6861831.

Figure 4.

Photographs of patients. Front and side views of patients 1 (A, B), 2 (C, D) and 5 (I, J); (E) patient 3 at 25 years of age; (F) patient 3 at 12 years of age; K: patient 6. Note the mild craniofacial defects. G and H: CT scan of skull of patient 3. Note the severe skull deformity. We have obtained the written consent from the patient or his/her legal guardian for publication of the images in print and online.

Table 1

Genomic regions duplicated in patients

Pt	chromosome	Duplication Size (Mb)	Genomic region *	Gene(s)	BACs detecting duplication in CMA analysis	CMA version
1	5	0.39	36845462-37231819	<i>NIPBL</i> , <i>FLJ13231</i> (exon 3-3'UT)	RP11-317I23, RP11-7M4, RP11-209M10	Version 5 BAC array
2	5	0.39	36845462-37231819	<i>NIPBL</i> , <i>FLJ13231</i> (exon 3-3'UT)	RP11-7M4, RP11-209M10	Version 6 oligo array
3	5	1.07	36487985-37556112	<i>SLC1A3</i> , <i>NIPBL</i> , <i>FLJ13231</i> , <i>NUP155</i> , <i>WDR70</i> (exon 1-9)	RP11-317I23, RP11-7M4, RP11-209M10	Version 6 oligo array
4	5	0.25	36870689-37120908	<i>NIPBL</i>	RP11-317I23, RP11-7M4	Version 6 oligo array
5	5	0.33	37084359-37417228	<i>NIPBL</i> (exon 39-47), <i>FLJ13231</i> , <i>NUP155</i> , <i>WDR70</i> (exon 1-2)	RP11-7M4, RP11-209M10	Version 6 oligo array
6	X	7.14	46824905-53961694	~174 genes including <i>WAS</i> , <i>CLCN5</i> , <i>SMC1A</i>	RP11-416B14, RP11-52N6, RP11-70P16	Version 6 BAC array

* based on the 244K aCGH data

Table 2
Clinical features of patients

Patient	1	2	3	4	5	6	CdLS*
Gender	F	F	F	M	F	F	
Age	18 years	6 years	30 years	8 months	5 years	6 years	
Birth height	NA	3%	3%	NA	NA	normal	low
Birth weight	NA	<5%	<5%	2.29 kg	50%	normal	low
Height	normal	normal	162.6 cm	<5%	<3%	normal	Short stature
Weight	> 95%	90%	59.2 Kg	<5%	50%	> 98%	low
Mental retardation	+	+	+	NA	+	+	+
Developmental delay	+	+	+	+	+	+	+ (global psychomotor delay)
Behavioral abnormalities	Sleep disturbance	Sleep disturbance	Suggested diagnosis of pervasive disorder	NA	No particular abnormalities	Sleep abnormalities	Self-injurious
	Self-injurious					Aggressive	Autistic-like features
	Repetitive stereotypic movements					Pain insensitivity	Hyperactivity
	Obsessive-compulsive behavior					Anxiety	Attention disorder
							Anxiety
Hypotonia	-	+	NA	+	-	+	Compulsive disorders
Seizures	-	staring spells	NA	no	+	+, staring spells	Sleep disturbance
Facial features and skull defects	Macrocephaly	Frontal bossing	Frontal bossing	Multiple suture cramosynostosis turricephaly brachiocephaly	Large flat forehead	Microcephaly	Microbrachycephaly
	Broad, tall forehead		Biparietal narrowing				
	Bitemporal narrowing						

Ocular anomalies	Short and upslanting palpebral fissures Exotropia	Astigmatism Strabismus	Hypotelorism Right epicanthic fold	Hypertelorism Proptosis	Short palpebral fissures	Long palpebral fissures Excess skin infraorbitally Esotropia	Palpebral ptosis Myopia Nystagmus
Nose	Bulbous nasal tip Broad nasal root	flat, wide base			Groove between nasal root and the face	Tubular nose with a pointed tip Prominent columella	Small nose Depressed nasal bridge Antverted nostrils
Mouth	Short philtrum Down-turned corners of mouth Prognathism Narrow palate	Short philtrum	Short philtrum Micrognathia Highly arched palate	High-arched palate	Poorly defined philtrum Several incisors missing	High palate	Micrognathia Crescent-shaped mouth Long philtrum Thin lips High arched palate
Ears	Low set	Low set, posteriorly rotated, cupped		Low set, small, and simplified in shape	Small pits anterior to the ear helixes Hearing disorder		Low set Posteriorly augulated Hearing disorder
Hairs, eyebrow			Low posterior hairline	Hair scant Absent eyebrows		Thick hair Heavy, long eyebrows	Low posterior hairline Synophrys Long eyelashes
Limb defects	Large hands, feet Very long fingers Right thumb: ulnar deviation Single palmer crease	Long fingers Right knee genu valgum	Normal	Single transverse palmar crease on the left hand	Short 5th fingers and a prominent hockey stick crease	Long gracile fingers 5 th finger clinodactyly Pes planus	Small hands and feet Short digits Phocomelia Oligodactyly 5 th finger clinodactyly Proximally placed thumbs Single palmer crease
Other defects	Scoliosis			Agenesis of the corpus callosum		Hirsutism Eczema Recurrent sinus and ear infections	Hirsutism Gastroesophageal reflux Heart defects

	Vesicoureteral reflux (VUR)	Vesicoureteral reflux (VUR)
--	-----------------------------	-----------------------------

F:female; M: male; NA: information not available
* common features for CdLS20,21,23,54