The I κ B kinase complex in NF- κ B regulation and beyond

Michael Hinz & Claus Scheidereit*

Abstract

The I κ B kinase (IKK) complex is the signal integration hub for NF- κ B activation. Composed of two serine-threonine kinases (IKK α and IKK β) and the regulatory subunit NEMO (also known as IKK γ), the IKK complex integrates signals from all NF- κ B activating stimuli to catalyze the phosphorylation of various I κ B and NF- κ B proteins, as well as of other substrates. Since the discovery of the IKK complex components about 15 years ago, tremendous progress has been made in the understanding of the IKK architecture and its integration into signaling networks. In addition to the control of NF- κ B, IKK subunits mediate the crosstalk with other pathways, thereby extending the complexity of their biological function. This review summarizes recent advances in IKK biology and focuses on emerging aspects of IKK structure, regulation and function.

Keywords cancer; development; differentiation; inflammation; immunity

DOI 10.1002/embr.201337983 | Received 11 September 2013 | Revised 31 October 2013 | Accepted 6 November 2013

See the Glossary for abbreviations used in this article.

Introduction

NF- κ B is an inducible transcription factor that coordinates specific gene expression programs to impact the regulation of multiple physiological functions. The most important and evolutionarily conserved role of NF- κ B is as mediator of the immune and inflammatory response. However, in addition, NF- κ B pathways contribute to cell adhesion, differentiation, proliferation, autophagy, senescence and protection against apoptosis. In line with its multi-layered physiological functions, deregulated NF- κ B activity is found in a number of disease states, including cancer, arthritis, chronic inflammation, asthma, neurodegenerative diseases, and heart disease [1,2].

The NF- κ B family consists of NFKB1 (p50/p105), NFKB2 (p52/p100), RelA (p65), c-Rel and RelB, which form various homo- and heterodimers. In resting cells, NF- κ B dimers are sequestered in the cytoplasm through interaction with I κ B proteins (I κ B α , I κ B β and I κ B ϵ) or the precursor proteins p100 and p105. Induction of NF- κ B depends on the phosphorylation of I κ Bs on critical serine residues.

As a consequence, I κ Bs are ubiquitinated by the E3 ubiquitin ligase SCF^{BT α CP} and degraded by the proteasome, which in turn allows the nuclear translocation of NF- κ B heterodimers (Fig 1; [3,4]). NF- κ B precursors are – constitutively or in a stimulus-dependent manner – processed by the proteasome to produce the mature transcription factors p50 and p52 [5]. As the small I κ Bs, p105 can undergo signal-dependent phosphorylation and degradation, which frees p50 homodimers, leading to the formation of Bcl-3:p50 complexes [6].

The discovery of the phosphorylation-dependent NF- κ B activation mechanism initiated the search for an I κ B kinase (IKK). An initial biochemical study described a high-molecular-weight kinase complex of approximately 700 kDa, which required non-degradative ubiquitination for its activity [7]. However, the identity of the kinase was not revealed. The serine/threonine kinases IKK α (85 kDa; also known as IKK1 or CHUK) and IKK β (87 kDa; also known as IKK2), were then identified as the two catalytic components of the IKK complex, which form dimers and are able to phosphorylate I κ Bs *in vitro* [8–13]. As a third component, NEMO (48-kDa; also known as IKK γ , IKKAP1 or Fip-3), was identified by complementation of an NF- κ B-unresponsive cell line and by sequencing of IKK-associated polypeptides [14,15]. NEMO is a regulatory non-enzymatic scaffold protein. Gene ablation studies of the IKK subunits then revealed that NF- κ B activation is achieved alternatively through the mechanistically different canonical and non-canonical pathways.

Inflammatory cytokines, radiation, stress signals and pathogenic assaults evoke the rapid canonical pathway and generally involve ubiquitin-mediated complex formation of signaling molecules, ultimately resulting in phosphorylation of two serine residues located in the IKK α and β activation loops, similar to the activation of many other kinases (Figs 1 and 2). Canonical signaling strictly depends on NEMO, while the catalytic subunits seem to be more redundant [16].

The non-canonical pathway is activated by a restricted group of stimuli, such as the TNF family members lymphotoxin- α/β , BAFF or CD40L, which trigger posttranslational stabilization of NF- κ B interacting kinase (NIK) (Fig 2). Crystal structural analysis showed that the kinase domain of NIK adopts an intrinsically active conformation, so that downstream targets can be phosphorylated without requiring an additional phosphorylation step [17]. In unstimulated cells, a TRAF-cIAP complex catalyzes K48-linked ubiquitination of NIK, leading to constitutive NIK degradation. Ligand binding induces the recruitment of the TRAF-cIAP complex to the receptor, whereupon TRAF2-mediated, K63-linked ubiquitination of cIAP1/2

Glossary

aa	amino acid
ABINs	A20-binding inhibitor of NF- κ B and NEMO
BAFF	Tumor necrosis factor (ligand) superfamily, member 13b
β-TrCP	β -transducin repeat containing E3 ubiquitin protein ligase
CBM	CARMA1 BCL10 MALT1
CBP	CREB-Binding-Protein
DARPin	Designed ankyrin repeat protein
DD	Death domains
Dok1	Docking protein 1
DUB	Deubiquitinase
CARD	Caspase activation and recruitment domains
CYLD	Cylindromatosis
EDA-ID	Ectodermal dysplasia with immunodeficiency
FLICE	FADD-like interleukin-1 beta-converting enzyme
FOXO3a	Forkhead box O3
HLH	Helix-loop-helix
HTLV1	Human T-lymphotropic virus 1
IκB	Inhibitor of nuclear factor- κ B
IL-1/IL-1R	Interleukin-1/interleukin-1 receptor
IRF	Interferon regulatory factor
IRS-1	Insulin receptor substrate 1
JNK	Jun N-terminal kinase
LUBAC	Linear ubiquitin chain assembly complex
LZ	Leucine zipper
MAPK	Mitogen-activated protein kinase
MAVS	Mitochondrial antiviral-signaling protein
MEKK3	MAPK/ERK kinase kinase 3
mTOR	Mammalian (also: mechanistic) target of rapamycin
NCOA3	Nuclear receptor coactivator 3
NBD	NEMO binding domain
NEMO	NF- κ B essential modifier
NF-κB	Nuclear factor- κ B
NLS	Nuclear localization sequence
OTU	Ovarian tumor domain
OTULIN	OTU DUB with linear linkage specificity
PARP-1	Poly-ADP-polymerase-1
PI3K	Phosphatidylinositol 3-kinases
PIAS1	Protein inhibitor of activated STAT, 1
RHIM	RIP homotypic interaction motifs (RHIMs)
S6K1	Ribosomal S6 protein kinase 1
SMRT	Silencing mediator for retinoid or thyroid-hormone receptors
SNAP23	Synaptosomal-associated protein, 23kDa
TAB1/2	TAK1 binding protein 1/2
TAK1	TGF- β -activated kinase 1
TBK1	TANK-binding kinase 1
TIR	Toll IL-1R (TIR) domain
TNFR	Tumor necrosis factor receptor
TNFRSF	Tumor necrosis factor receptor superfamily
TLR	Toll-like receptor
TRAF	TNF receptor associated factor
TSC1	Tuberous sclerosis 1
UBAN	Ubiquitin binding in ABINs
ULD	Ubiquitin-like domain

switches its K48 ubiquitin ligase activity from NIK to TRAF3. The ensuing TRAF3 degradation destabilizes the TRAF-cIAP complex and enables the accumulation of newly synthesized NIK [18,19]. NIK is thought to phosphorylate IKK α , which in turn phosphorylates and marks p100 for C-terminal processing to generate p52-containing complexes, primarily p52/RelB. The non-canonical pathway is characterized by a distinctive slower kinetics, requires neither IKK β

nor NEMO and plays a critical role in the development of lymphoid organs [20]. Attenuation of the non-canonical NF- κ B pathway is controlled through the DUB OTUD7B, which deubiquitinates TRAF3 in a K48-directed manner [21].

The substrate spectrum of the IKK complex is not restricted to I κ Bs and precursors, but includes a wide range of NF- κ B pathway inherent (such as RelA) and extraneous (such as p53 and TCS1) substrates (Table 1). Thus, the IKK complex is both a signaling hub for NF- κ B activation and an interface for crosstalk between NF- κ B activating pathways and other physiological processes [16,22].

Despite the steady increase in IKK-related publications, important questions still remain unanswered (Sidebar A). How a signaling cascade mechanistically induces the activation of the IKK complex, how IKK activity is turned off and how substrate specificity is achieved has to be resolved. Recent genetic, biochemical and structural data have shed new light on IKK structure, regulation and function.

IKK complex – structures and composition

I κ B kinase-mediated activation of NF- κ B is differentially regulated by IKK α and IKK β . Both kinases have high sequence homology (approximately 50% identity), and contain an N-terminal kinase domain, a dimerization domain and a C-terminal NEMO-binding domain (NBD) [16,22,23] (Fig 3A), although only IKK α has a predicted NLS [24]. Kinase activity critically depends on lysine 44 and phosphorylated T-loop serines – S176 and 180 for IKK α , S177 and 181 for IKK β [9–12,25]. IKK β contains a conserved ubiquitin-like domain (ULD), which is critical for its catalytic activation [26]. Much-anticipated 3D structural information was obtained in the last few years that allowed further elucidation of IKK structure-function relationships (Table 2).

IKK structures reveal complex intra- and inter-molecular interactions

As a major breakthrough, the long-awaited crystal structures of IKK β were recently presented, revealing the domain organization of this catalytic IKK subunit and shedding new light on protein-protein interactions, mechanisms of activation and the mechanism of substrate recognition [27–29]. IKK β has a trimodular structure, comprising the N-terminal kinase domain (KD), a central ULD and an elongated α -helical scaffold/dimerization domain (SDD) at the C-terminus (Fig 3B and Table 2). Neither of the previously predicted LZ and HLH motifs form as these residues are part of the SDD. Notably, the IKK-related kinase TBK1, which is essential for the induction of type I interferon (IFN), has the same trimodular structure [30–32]. Both IKK β and TBK1 form stable homodimers through the SDD domain. A comparison of the IKK β and TBK1 structures further revealed that the overall dimer conformation differs, depending on the activation status. Dimers of inactive kinases show a more compact geometry, resembling closed shears [29,31], while activated kinases adopt a more open conformation [27,28]. A detailed inspection of IKK β and TBK1 kinase domains revealed that the activation loop, when unphosphorylated, adopts conformations that are incompatible with protein substrate binding, explaining the essential role of T-loop phosphorylation at S177/S181 for IKK β and S172 for TBK1 kinase activity [27,31,33]. The previously predicted ULD

Figure 1. Prototypic canonical NF-κB pathway.

The activation of a cell surface receptor induces the recruitment and oligomerization of adapter proteins and E3 ligases, such as TRAF6 or LUBAC, which in turn synthesize non-degradative Ub chains (K11, K63, and M1) attached to various adaptor proteins, or to themselves. Subsequently, TAK1 and IKK complexes are recruited to upstream ubiquitin polymers through ubiquitin-binding domains. After IKK complex recruitment, NEMO is modified with poly- and mono-ubiquitin, and IKKs are phosphorylated in their T-loop either by TAK1 or through trans-autophosphorylation. IKK activation results in phosphorylation of IκB and p65. IκB phosphorylation leads to its ubiquitin-mediated proteasomal degradation, enabling NF-κB dimers translocation to the nucleus, where they bind to DNA and induce transcription. In addition, IKKs phosphorylate a variety of other substrates, resulting in crosstalk with other, NF-κB independent regulatory systems in the cytoplasm and the nucleus (Table 1).

is a critical element of IKKβ, required for kinase activity and, together with the SDD, involved in the exact positioning of the kinase substrate IκBα. Most likely, IκBα is recruited to IKKβ through NEMO, which ensures specific substrate recognition (see also below) [34]. The striking interdependence of the three domains is reflected by their extensive intramolecular interactions [28,29]. Intriguingly, the sequence and conformation of the SDD significantly differ between IKKβ and TBK1 [31,32]. The dimerization interface of TBK1 is almost twice as large as that of IKKβ, resulting in a stimulus-independent compact composition and a TBK1-specific overall domain organization. These structural differences might reflect specific activation requirements. IKKβ activation strictly depends on NEMO, whereas TBK1 activation requires K63-linked polyubiquitination of the conserved Lys30 and Lys401 residues [22,32]. In fact, structure-directed mutagenesis revealed that SDD-mediated kinase dimerization of IKKβ is required for NEMO binding and the kinase activation process, but not for enzymatic activity once its activation loop is phosphorylated [29]. Likewise, dimer-disrupting mutations on TBK1 had markedly decreased K63-linked polyubiquitination [32].

How signal transmission results in phosphorylation of the IκB kinase T-loop is still a matter of debate. The existence of IKK kinases (IKKKs) and/or IKK trans-autophosphorylation are both feasible scenarios [3,23]. In the closed dimer structure of inactive IKKβ, the active sites of the two neighboring KDs cannot interact with each other. However, activated IKKβ dimers seem to switch to a more open V-shaped conformation, allowing oligomerization and KD-KD interactions, with the potential to facilitate *trans*-autophosphorylation [28,29]. Of note, how the structure and relative position of the KDs in IKKα-IKKβ heterodimers, which are more relevant *in vivo*, may differ remains to be solved.

Structural insights into NEMO

Important structural information has been obtained for NEMO. On the basis of biochemical assays, NEMO had been predicted to be a mainly α-helical protein containing two coiled-coil domains (CC), a LZ and a C-terminal zinc-finger (ZF) region (Fig 3A; [22,23]). In addition, NEMO has a minimal oligomerization domain (MOD) and an N-terminal dimerization domain that enable the formation of higher order oligomers [35,36]. Finally, the kinase-binding domain (KBD) was determined by deletion analysis to be located in aa 44-111 of NEMO [37]. The ZF seems to be required for efficient IκBα binding and might direct IκBα to the ULD/SDD of IKKβ [34].

Although a crystal structure for full-length NEMO is still missing, the structures of several NEMO fragments have been resolved (Fig 3C and Table 2) [37–41]. The X-ray structure of the NEMO-IKKβ interface revealed an asymmetrical four-helix bundle, composed of a parallel NEMO dimer. Each NEMO molecule forms a crescent-shaped α-helix, which associates with mainly α-helical IKKβ peptide monomers via hydrogen bonding and hydrophobic interactions [37]. An X-ray structure of the central region, associated with a fragment of the viral IKK activating protein vFLIP, showed that the coil-coil domain encompasses aa 192–252 [38] (Fig 3C).

Nuclear factor-κB essential modulator has a central role in polyubiquitin-mediated IKK activation, as it specifically recognizes polyubiquitins through the CC2 – LZ region domain and becomes itself ubiquitinated. Although the type of ubiquitination is a matter of

Figure 2. Flow chart of the IKK signaling network.

Cell-surface receptors and stress response pathways trigger a dynamic regulatory network involving protein phosphorylation, non-degradative ubiquitination, adapter protein interactions and higher order oligomerization events to activate the IKK complex. Signaling is inhibited by deubiquitinases, phosphatases and IKK-mediated negative feedback regulation (coloured text). Enzymatic IKK activation, modulated by Hsp90-Cdc37 activity, may be achieved by trans-autophosphorylation and the action of IKK kinases (IKKs). Viral proteins, such as Kaposi's Sarcoma-Associated Herpesvirus vFLIP, have been shown to activate the IKK complex through direct interaction. The IKK complex preferentially phosphorylates IκBs and NF-κBs, but has also many other substrates. A subset of the TNF superfamily receptors (TNFSFR) activates the non-canonical pathway, which depends on NIK and selectively on IKKα and results in p100 processing. See text for details.

debate (discussed below) there is ample evidence that both events are important for IKK activation [42,43]. Mutations in the NEMO ubiquitin-binding region (UBAN, also known as NUB, NOA or CoZi), which impair ubiquitin binding and NF-κB activation, have been identified in patients suffering from ectodermal dysplasia with immunodeficiency (EDA-ID) [44]. The crystal structures of different NEMO fragments including the UBAN domain revealed a dimer, which contains two coil-coil domains representing CC2 and LZ, respectively [40,41,45]. These NEMO fragments were shown to bind to linear, M1-linked di-ubiquitin with higher affinity than K63- or K11-linked di-ubiquitins [41,45–47]. However, a larger NEMO fragment containing the UBAN domain together with the C-terminal ZF —another *bona fide* ubiquitin-binding domain— had an increased affinity for K63-linked polyubiquitin [39,48,49]. *In vitro* binding studies demonstrated a high preference of NEMO in solution for M1-linked ubiquitin oligomers, while immobilization enhances the affinity towards K63-linked ubiquitin [50]. Competition analyses indicated that NEMO functions as a high-affinity receptor for M1-linked ubiquitin chains and a low-affinity receptor for different types of long lysine-linked ubiquitin chains [51]. The possible existence of M1-K63-mixed-link-

age ubiquitin polymers may pose a new problem [52]. Taken together, these findings indicate that different types of polyubiquitin are able to bind to NEMO and contribute to IKK activation. Whether the differences in binding affinities determined in *in vitro* studies with NEMO fragments and di-ubiquitins are physiologically relevant and how they impact IKK activation remains to be determined.

Oligomeric composition of the IKK complex

The apparent molecular weight of the IKK complex in gel filtration chromatography is around 700 to 900 kDa. Although numerous proteins have been proposed to interact with IKK-components [23], co-immunoprecipitation studies with S³⁵ labeled proteins and size exclusion chromatography analysis with recombinant proteins clearly indicate a tripartite IKK composition [53–55]. However, the exact stoichiometry of IKKα, β and NEMO in the IKK complex remains an open question.

Crystallographic and quantitative analyses of the binding interactions between N-terminal NEMO and C-terminal IKK fragments suggest that IKKβ dimers would interact with NEMO dimers [37,56]. Such a model is supported by the recent crystal structures of IKKβ

Table 1. Substrates and function of IKK α and IKK β

Substrate	Affected molecular process	Biological function	References
IKKα-dependent			
Transcriptional regulation			
β -catenin	Interferes with ubiquitination mediated degradation and increases β -catenin-dependent transcription	Cell cycle regulation/cancer	[121–124]
CBP	Enhances NF- κ B-dependent transcription	NF- κ B dependent transcription $\uparrow$	[142]
c-Rel	Accelerates c-Rel turnover	NF- κ B dependent transcription $\downarrow$	[118]
ER α	Enhances estrogen receptor-mediated gene activation	Hormone response	[155]
IRF5	Inhibits TLR mediated interferon production	Inflammatory response	[156]
IRF7	Enhances TLR-mediated interferon production	Immune response	[157]
NCOA3	Enhances nuclear hormone receptor-mediated gene activation	Steroid hormone response	[155]
NF- κ B2/p100	Induces processing of p100 into p52	NF- κ B dependent transcription $\uparrow$	[57]
PIAS1	Represses NF- κ B-dependent transcription	Restriction of inflammation	[158]
RelA/p65	Accelerates RelA turnover	NF- κ B gene expression $\downarrow$	[118]
Unknown	Cofactor in TGF β -Smad2/3 signaling	Keratinocyte differentiation	[112]
Unknown	Required for efficient induction of mTOR	Autophagy/growth control	[133]
Chromatin regulation			
Histone H3	Enhances NF- κ B-dependent transcription	NF- κ B dependent transcription $\uparrow$	[159–161]
SMRT	De-represses NF- κ B target genes	NF- κ B dependent transcription $\uparrow$	[138,139]
Cell fate decision			
Cyclin D1	Triggers cyclin D1 degradation	Cell cycle regulation	[127]
p27/Kip1	Stimulates nuclear export of p27	Cell cycle regulation/cancer	[136]
Signaling			
Unknown	Suppress the serine protease inhibitor Maspin	Metastasis	[134]
IKKβ-dependent			
Transcriptional regulation			
FOXO3a	Promotes degradation of FOXO3a	Growth control/cancer	[147]
I κ Bs	Induces proteasomal degradation of I κ Bs	NF- κ B dependent transcription $\uparrow$	[23]
NCOA3	Enhances NCOA3 nuclear import	Hormone response	[162]
NF- κ B1/p105	Induces p105 proteolysis, resulting in Bcl-3-p50 Complex formation and TPL-2 activation	NF- κ B dependent transcription $\uparrow$ and inflammatory response	[6,163,164]
NF- κ B p65	Enhances transcriptional activity	NF- κ B dependent transcription $\uparrow$	[119]
Cell fate decision			
BAD	Primes BAD for inactivation	Cell survival	[128]
p53	Induces proteasomal degradation	Growth control/cancer	[145]
p85/PI3K	Interferes with Akt and mTOR inhibition after nutrient depletion	Autophagy/growth control	[132]
p85/S6K1	Activation of p85 S6K1 upon oxidative stress	Apoptosis	[129]
Signaling			
14-3-3 β	Dislocates TTP/14-3-3 β from AU-rich elements	mRNA stability	[165]
Aurora A	Induces proteasomal degradation	Genome integrity	[166]
Bcl10	Attenuates TCR signaling	IKK feedback regulation	[95,96]
β -catenin	Inhibits the β -catenin-dependent transcription	Cell cycle regulation	[123]
CARMA1	Facilitates CBM complex formation	IKK feedback regulation	[167]
CYLD	Inhibits deubiquitinase activity	IKK feedback regulation	[168]
DOK1	Inhibits ERK1 and ERK2 activation	Cell motility	[169]
IRS-1	Inhibits insulin signaling	Diabetes	[170]
NEMO	Modulates IKK structure	Feedback regulation	[93]
SNAP-23	Promotes exocytosis in mast cells	Immune response	[171]
TSC1	Activates the mTOR pathway and VEGF production	Growth control/cancer	[146]

(see above), and a similar situation is anticipated for IKK β :IKK α heterodimers. Chemical cross-linking and equilibrium sedimentation analyses of NEMO suggested a tetrameric oligomerization (dimers of dimers), depending on a C-terminal coiled-coil minimal oligomerization domain (MOD) and subsequent dimerization of the dimers with their N-terminal sequences [36]. Tetrameric NEMO could sequester four kinase molecules, yielding an IKK α_2 IKK β_2 NEMO $_4$ stoichiometry. Such a higher-order oligomerization could provide the basis for an IKK *trans*-autophosphorylation mechanism (as discussed before). Conformational changes in the scaffold induced by polyubiquitin binding to NEMO could bring the catalytic domains of two kinase dimers into proximity. Such a model is supported by the recent finding that IKK β dimers reversibly oligomerize in solution and the active kinase forms higher order oligomers in the crystal [28,29].

Taken together, tremendous progress has been made in understanding the structures of the IKK components, which provides the basis to finally understand the dynamic architecture of the IKK complex. Notably, although the tripartite IKK structure is perhaps the most abundant form, other IKK complexes might exist. In fact, co-expression experiments and *in vitro* studies have shown that NEMO can interact with IKK1 or IKK2 homodimers [54,55]. It is a tempting assumption that different complex compositions might be required for tissue-specific or stimulus-specific NF- κ B dependent and independent signaling events. Likewise, NIK-dependent activation of the non-canonical NF- κ B pathway was proposed to occur through phosphorylation of IKK α homodimers [57]. However,

whether specific IKK complexes with distinct oligomeric compositions and functions exist in cells remains to be demonstrated.

IKK activation and inhibition

I κ B kinases are activated by a plethora of agents and conditions, including extracellular ligands that bind membrane receptors, such as TNFR, TLR, or IL-1R, intracellular stress, such as DNA damage and reactive oxygen species, as well as the recognition of intracellular pathogens mediated by the NOD and RIG-I-like (NLR) family of proteins (Fig 2). The activated receptor structures nucleate dynamic regulatory networks, where protein phosphorylation, non-degradative ubiquitination, adapter protein interactions and most likely higher order oligomerization events all contribute to IKK activation (Figs 1 and 2). Moreover, canonical and non-canonical NF- κ B signaling pathways can be activated by human oncogenic viruses, including the human T-cell leukemia virus type 1, the Kaposi sarcoma-associated herpesvirus, and the Epstein-Bar virus [58]. Recent findings indicate that the virus-encoded oncoproteins either use components of the IKK upstream signaling network or directly act on the IKK complex to activate NF- κ B [59,60].

IKK phosphorylation

How signal transmission results in the phosphorylation of the I κ B kinase T-loop is still an important unsolved question. In analogy to

Table 2. Structural analysis of IKK subunits and their relatives

Protein	Origin	Details	Resolution	Structural features	References
IKK β (aa 4–675)	Xenopus laevis	Complex with inhibitors Cmpd1 or Cmpd2	3.6 Å	KD + ULD + SDD; dimer ('a pair of shears')	[29]
IKK β (aa 11–669)	Human	Constitutively active mutant (S177E/S181E)	4 Å	See above; open conformation; permits higher order oligomerization	[28]
His-IKK β (aa 1–664; HIS tagged)	Human	Partially phosphorylated; bound to the staurosporine analog K252a	2.8 Å	See above; comparison of active and inactive kinase domains	[27]
NEMO (aa 44–111)	Human	Hybrid complex containing NEMO 44–111 and IKK α / β or IKK β peptides	2.25/2.2 Å	Elongated and atypical parallel four-helix bundle	[37]
NEMO (aa 150–272)	Human	Hybrid complex containing NEMO 150–272 and vFLIP 1–178	3.2 Å	Coiled-coil dimer	[38]
NEMO (aa 251–337 His-tagged)	Mouse	Hybrid complex containing NEMO 251–337 and DARPin 1D5	2.95 Å	Coiled-coil dimer	[40]
NEMO (aa 254–337)	Human	NEMO 254–337 alone and in complex with di-ubiquitin	2.8/2.7 Å	Coiled-coil dimer	[41]
NEMO (ZF-WT and ZF-C417F)	Human	Solution structures determined by NMR	–	WT and mutant ZFs adopt a global $\beta\beta\alpha$ structure and bind zinc with comparable affinities; mutation causes instability	[49]
TBK1 (1–657; Δ CTD)	Human	A: Complex with inhibitors MRT67303 and BX795 B: D135N mutant (kinase dead)	2.4/2.5 Å and 3.3 Å	KD + ULD + SDD; Dimer; extensive interactions between different domains	[32]
TBK1 (1–657; Δ CTD)	Human	Complex of S172A mutant (inactive) or phosphorylated D135N mutant (active) with MRT67307/215 and BX795	2.6/4.0 Å	See above; Activation rearranges KD into an active conformation and maintains overall structure	[31]

Figure 3. The structure of IKKs.

(A) Domain organization. (B) Crystal structure of *Xenopus* IKK β . IKK β was found as a dimer of dimers, a conformation with the potential to facilitate trans-autophosphorylation [29]. (C) Crystal structures of NEMO fragments: N-terminal kinase-binding domain (HLX1) of NEMO in complex with IKK β NBD, HLX2 in complex with vFLIP, CC2-LZ in complex with linear di-Ub, ZF [37,38,40,41,49]. All crystal structures are from the database of the NCBI structure group (<http://www.ncbi.nlm.nih.gov/Structure/index.shtml>).

other signaling pathways, IKK kinases (IKKKs) have been suggested to mediate it. A prominent example is TAK1, which is also engaged in the JNK pathway. TAK1, together with the adaptor proteins TAB 1 and TAB 2, was found to act as TRAF6-regulated IKK activator in cell-free assays [61,62]. TAB 2 can recruit TAK1 to K63-linked polyubiquitin chains of upstream regulators which most likely cause induced proximity-driven IKK β phosphorylation. However, TAK1 is not an essential general IKKK, but rather a regulatory module with a stimulus and cell-type specific impact on IKK activation [16,42]. MEKK3 was proposed as another potential IKKK, as the kinase can phosphorylate IKK *in vitro* and NF- κ B activation is reduced in MEKK3 deficient cells in response to TNF, IL-1 or TLR stimulation [63,64]. IL-1-induced NF- κ B activation has been proposed to involve MEKK3 in addition to TAK1 [65,66]. However, IKK subunits could also be activated by trans-autophosphorylation, as previously proposed [67], instead of through an IKKK. This latter possibility is supported by recent structural and composition analyses, as discussed above. Trans-autophosphorylation and IKKK-dependent phosphorylation could even operate successively or in parallel to reach maximum kinase activation (Fig 1).

Ubiquitin-dependent signaling

The activation of the IKKs appears to depend on the induced proximity resulting from the dense organization of signaling complexes and on binding of adaptor proteins, such as NEMO or TAB proteins. Non-degradative polyubiquitination, a crucial prerequisite of IKK complex activation, triggers both processes (Fig 1) [42].

TRAF6 was identified as the first ubiquitin E3 ligase that could—together with Ubc13 and Uev1A—catalyze K63-linked auto-ubiquitination and subsequently trigger IKK activation [61]. TRAF6 is involved in a wide variety of NF- κ B-stimulating signaling pathways, including those triggered by IL-1R, TLR, TCR, RIG-I-like receptor and DNA double strand breaks [42,43]. In the case of IL-1 signaling, it has been demonstrated that TRAF6 enzymatic activity but not auto-ubiquitination, is required for NF- κ B activation [68]. In fact, TRAF6 has been shown not only to undergo self-ubiquitination, but also to mediate K63-linked polyubiquitination of several pathway components, such as IRAK1, MALT1 and TAK1 [42,43]. Furthermore, TRAF6 has been proposed to generate free, unanchored K63-linked polyubiquitin which would act as docking platform in the IKK activation process [69]. Additional K63-specific E3

ligases involved in specific NF- κ B signaling cascades have been identified, such as TRAF2/5, pellino proteins or TRIM25 in the TNFR, IL-1R/TLR and RIG-I pathways, respectively. Likewise, there is a growing list of proposed NF- κ B pathway regulators that are substrates of inducible K63 ubiquitination, such as Bcl10, NOD2, RIP1, RIP2 and ELKS [42,43].

In contrast to IL-1 β , TNF α signaling does not depend on K63-linked ubiquitination, indicating that alternative forms of non-degradative polyubiquitination are important in this pathway [70]. Accordingly, linear, M1-linked ubiquitination of NEMO and RIP1, which is catalyzed by the LUBAC complex, has been shown to be important for NF- κ B activation [71,72]. This E3 complex, consisting of HOIL1, HOIP and SHARPIN, is recruited to the TNFR1 signaling complex in a TRADD, TRAF2 and cIAP1/2 dependent manner. Furthermore, LUBAC mediated M1-linked ubiquitination contributes to IL-1R, CD40, TACI, parkin and DNA damage-mediated NF- κ B activation, but is dispensable for B-cell receptor-mediated signaling [43,73–75]. The physiological relevance of LUBAC was first demonstrated in mice with chronic proliferation dermatitis (Cpdm), which results from a spontaneous null mutation in the *Sharpin* gene. These mice partially mimic the phenotype of patients suffering X-linked hyper IGM syndrome and hypohydrotic ectodermal dysplasia, which is caused by NEMO mutations, and Cpdm-derived cells display attenuated TNF, CD40 ligand and IL-1 β signaling [71,76,77]. Biallelic inactivating mutations in *HOIL1* have been identified in patients suffering an inherited disorder with immunodeficiency, autoinflammation and amylopectinosis [78]. *HOIL1* deficiency, and subsequent LUBAC destabilization, resulted in impaired NF- κ B-mediated I-1 β responses, which differed depending on the tissue [78]. Thus, the ubiquitin linkage type seems not only to confer stimulus-specific, but also cell type-specific restraints in NF- κ B activation.

There is accumulating evidence that alternative and even hybrid ubiquitin linkages play a role in NF- κ B signaling, thereby increasing the complexity of ubiquitin-mediated processes [43]. *In vitro* assays have demonstrated that the E3 ligases cIAP1 and TRIM23 can catalyze K6-linked or K27-linked polyubiquitination of NEMO, respectively [79,80]. Modification of RIP1 seems to be a special case, because TNF α treatment induces modification with degradative K48 as well as with non-degradative M1-, K63-, and K11-linked ubiquitin chains [47,71,81–83]. The fact that only one ubiquitin acceptor site (K377) has been determined until now [81], raises the question of whether TNF α induces distinct populations of modified RIP1 or, alternatively, the modification of RIP1 with a poly-ubiquitin chain containing mixed linkages. Indeed, an analogous process has been recently reported for IRAK1, IRAK4 and MYD88, which are modified with K63/M1-linked hybrid ubiquitin chains in IL-1R or TLR-stimulated cells [52]. In addition to polyubiquitin, monoubiquitination of proteins has a functional impact [84]. In fact, a modified NEMO species 8 kDa larger, which most likely indicates a mono-ubiquitinated form, is generated in cells stimulated with various agents [68,72,85,86]. K285 could be determined as the acceptor site using mass spectrometry and mutational analysis [68,85]. The same site, in addition to K309, was determined as acceptor for M1-linked ubiquitination [72]. Site-specific ubiquitination of NEMO was shown to be crucial for NF- κ B activation in both cell culture and in a mouse model [68,72,85,87]. Thus, IKK activation coincides with two parallel or sequentially occurring modifications of NEMO. As an appealing

possibility, NEMO ubiquitination and NEMO-dependent ubiquitin binding might trigger and stabilize stimulus-induced, conformational changes to facilitate *trans*-autophosphorylation of the kinases. The Hsp90-Cdc37 chaperone complex, which transiently interacts with IKK, could have a supportive function in this process [53].

NF- κ B-inducing lymphogenic virus proteins, such as Kaposi's sarcoma associated herpes virus encoded vFLIP, have been recently shown to bind NEMO directly. This would probably induce conformational changes that would allow bypass of the ubiquitin-dependent signaling cascade [38,60]. In addition to ubiquitination, post translational modifications with ubiquitin-like proteins—such as genotoxic stress induced, PIASy-mediated SUMOylation of NEMO—are critically involved in signal transmission and NF- κ B activation [73]. Based on our current knowledge, modifications with mono-ubiquitin, non-degradative polyubiquitins and ubiquitin-like proteins can be concluded to have a crucial role in coordinating appropriate protein-protein interactions within specific NF- κ B signaling pathways. However, many details of the non-degradative actions of ubiquitin and ubiquitin-like proteins remain to be discovered.

Protein-protein interactions and higher order oligomerization

A typical early event in NF- κ B signaling is the receptor-mediated recruitment of adapter proteins that contain protein-protein interaction domains, such as DDs, CARDs, RHIMs and TIRs. These adapters have the potential to form higher order signaling platforms [3,88]. X-ray structure analyses showed that the TLR/IL-1R signaling molecules MyD88, IRAK4 and IRAK2 associate in helical assemblies, dependent on the DDs of the individual proteins [89]. Likewise, virus-induced RIG-I catalyzes the prion-like aggregation of MAVS that depends on CARD interaction. Oligomeric MAVS interacts with TRAF6 and TRAF2, resulting in IKK and TBK1 activation [90]. TRAF6 itself can also oligomerize to form a distinct network structure mediated by C-terminal trimerization and N-terminal dimerization with the ubiquitin-conjugating enzyme Ubc13 [91]. As a consequence, the increase in local concentration and proximity could promote TRAF6 auto-ubiquitination and downstream signaling. Lastly, activated IKKs have the potential to form higher-order oligomers, which might trigger rapid signal amplification by *trans*-autophosphorylation [28]. The weak interactions between IKK kinases observed *in vitro* could be further stabilized by the interaction with clustered upstream signal components and/or through polyubiquitin scaffolds.

Taken together, oligomerization appears to occur at all levels of canonical signaling cascades, in which multiple signaling oligomers combine to perform multiple reactions simultaneously and efficiently. The clustering of signaling molecules is an attractive mechanism that could provide temporal and spatial control of signal transmission and might help to increase the signal-to-noise ratio [92]. Therefore, a focus of future research should be to prove whether higher-order assemblies of signaling molecules are a general feature of canonical and even non-canonical NF- κ B signaling.

Control of transience and attenuation of IKK activity by feedback regulation

To maintain the transient nature of NF- κ B activity, signaling through the pathway is controlled by various levels of negative feed-

back mechanisms, including direct regulation of IKK as well as negative feedback mechanisms that affect upstream signaling components (Fig. 2). Intrinsic attenuation of IKK has been proposed to involve IKK β -mediated autophosphorylation of C-terminal serines, either close to the HLH or in and around the NEMO-binding domain, as well as phosphorylation of NEMO S68 [67,93,94]. Such events might disrupt the domain interactions of IKK subunits. Moreover, IKK β can terminate TCR signaling by phosphorylation of the pathway component BCL10 [95,96]. Dephosphorylation of serines in the activation loops of IKKs by phosphatases PP2A and PP2C has also been suggested as another direct inhibitory mechanism, as reviewed earlier [16,23]. An important step in IKK activation is the binding of the NEMO UBAN domain to polyubiquitin chains. This interaction can be disrupted by optineurin, a negative regulator of TNF α -induced NF- κ B activation, which contains a similar UBAN domain and competes with NEMO for binding to polyubiquitin [97]. Mutations of optineurin in amyotrophic lateral sclerosis interfere with the inhibitory effect towards NEMO, and thus exaggerate NF- κ B activation [98]. Another NEMO-dependent feedback mechanism has recently been described: p47 (also known as NSFL1C)—which is major adaptor of the cytosolic triple-A ATPase p97—binds to polyubiquitinated NEMO and induces its lysosomal degradation, resulting in reduced IKK activity [99].

Non-degradative polyubiquitination of signaling molecules by various E3 ligases plays a key role in IKK activation. A series of DUBs, which cleave specific linkage types, counteract these activities and terminate or attenuate the signaling process. The K63-directed DUBs A20 and CYLD have been shown to de-ubiquitinate RIP1, TRAF6, RIP2 and MALT1. USP21 and K11-specific Cezanne have been implicated in the removal of poly-ubiquitin from RIP1 (for review see [100–103]). A recently identified DUB known as OTULIN or Gumby, which is specific for M1-linked ubiquitin, has been shown to interact with the LUBAC component HOIP, decreasing M1-linked ubiquitination and attenuating NF- κ B activity. OTULIN-depleted cells spontaneously accumulate M1-linked ubiquitin chains on LUBAC components and, upon TNFR1 or NOD2 stimulation, on RIPK1 and RIPK2, respectively [104–106]. The non-canonical NF- κ B pathway is attenuated by the DUB OTUD7B, which deubiquitinates TRAF3, thereby counteracting its degradation and preventing NIK-mediated p100 processing [21].

NEMO SUMOylation is a critical step in the response to DNA double-strand breaks. Amongst the Sentrin/SUMO-specific proteases (SENPs), SENP2 has been shown to specifically deSUMOylate NEMO and attenuate NF- κ B activation. Since NF- κ B promotes SENP2 expression, a negative feedback loop is established [107].

IKK functions

Although the tripartite structure of the IKK complex suggests common physiological roles for IKK α , IKK β and NEMO, the phenotypes of the single knockouts suggest they have common, but also distinct functions. IKK β -deficient mice are embryonically lethal and die, like p65-deficient mice, at embryonic day 13, primarily due to TNF-induced liver apoptosis. These results underscore the importance of IKK β in canonical NF- κ B signaling [108,109]. However, subsequent studies with IKK β -deficient cells indicate that the absolute requirement for IKK β in canonical NF- κ B signaling depends on the stimu-

lus. IKK α can substitute IKK β function at least in the case of IL-1R signaling [110].

IKK α -deficient mice can survive for a month after birth, but suffer from striking morphological defects, such as markedly hyperplastic epidermis [111]. Follow-up studies determined that IKK α is a major cofactor in a TGF β -Smad2/3 signaling pathway that is required for cell cycle exit and induction of terminal differentiation of keratinocytes [112]. Accordingly, IKK α has been shown to be a critical suppressor of skin cancer in humans and mice [113,114]. In addition to its specific function in TGF- β -mediated keratinocyte differentiation, IKK α is a crucial regulator of non-canonical NF- κ B signaling, required for B-cell maturation and formation of secondary lymphoid organs [57]. It is also involved in the maturation of dendritic cells and pancreatic homeostasis [115,116].

Genetic depletion of the scaffold protein NEMO results in a complete loss of canonical signaling and mutant embryos die at E12.5–E13.0 from severe liver damage due to massive apoptosis [22]. NEMO exists either as a component of the IKK complex or as an unbound form that shuttles between cytoplasm and nucleus. In response to genotoxic stress, NEMO undergoes sequential post-translational modifications and has a central role in a dual, PARP-1/PIASy and ATM dependent signaling pathway that links the cellular DNA damage response to NF- κ B [117].

The key function of the IKK complex is to phosphorylate I κ Bs and the NF- κ B precursors p105 and p100 [23]. In addition, IKKs directly modulate the function of RelA and c-Rel. IKK β -dependent phosphorylation of p65 on Ser536 enhances its transactivation potential, while IKK α mediated C-terminal phosphorylation of p65 and c-Rel attenuates their activity [118,119]. However, IKK α has been shown to accumulate in the nucleus in a stimulus-dependent manner and to phosphorylate chromatin components (Table 1), implicating a more far-ranging spectrum of biological functions [16]. There is ample evidence that IKK activity is not restricted to NF- κ B-dependent pathways but can also mediate cross talk with other signaling cascades, such as mTOR and MAPK pathways [1,16,120]. Hence, numerous additional kinase substrates have been identified, which link IKK activity to a variety of biological functions including immune responses and transcriptional regulation and chromatin remodeling (Table 1).

I κ B kinase dependent but NF- κ B independent signaling events have been shown to influence various cell fate decision processes. The signal transducer for Wnt-dependent cell proliferation, β -catenin, was described as one of the first alternative IKK substrates. IKK β -mediated phosphorylation of β -catenin was proposed to induce its ubiquitin-dependent degradation, whereas phosphorylation by IKK α stabilizes β -catenin expression and induces β -catenin-dependent cyclin D1 transcription [121–123]. Indeed, IKK α knockdown in multiple myeloma cells did not inhibit NF- κ B activation, but correlated with impaired β -catenin expression and significant growth inhibition [124]. However, other studies showed IKK α -mediated cyclin D1 phosphorylation and degradation, as well as NF- κ B dependent transcriptional regulation of cyclin D1, indicating a complex IKK/NF- κ B dependent regulation of cyclin D1 [125–127]. IKKs not only affect cell proliferation, but also cell survival pathways, in an NF- κ B-independent manner. IKK β phosphorylates the BH3-only protein BAD at serine-26 (Ser26), which primes it for inactivation and suppresses TNF α -induced apoptosis [128]. In response to oxidative stress, however, IKK β mediates pro-apoptotic functions through the activation of

p85 S6K1 [129]. The IKK complex also has a direct role in the induction of autophagy [130,131]. Nutrient depletion induces IKK-dependent phosphorylation of the p85 subunit of PI3K, thereby blocking Akt and mTOR inhibition [132]. This crosstalk may contribute to the induction of autophagic genes, such as Beclin 1 [130]. However, studies in PTEN-inactive prostate cancer cells indicate that IKK α functions as a mediator of mTOR activation, which in turn suppresses autophagy [133]. Thus, it will be important to elucidate the physiological signaling events and outcomes associated with IKK-dependent regulation of autophagy in different settings.

Notably, both IKK α and IKK β have crucial NF- κ B-dependent and independent functions in various oncogenic scenarios, which are often correlated with inflammation-mediated tumorigenesis. Analysis of prostate cancer mouse models indicates that tumor-infiltrating T cells and macrophages express RANKL, which in turn induces nuclear IKK α activation and subsequent transcriptional inhibition of the tumor suppressor Maspin (also known as serpin B5). The amount of active nuclear IKK α in mouse and human prostate cancer correlates with reduced Maspin expression and with metastatic progression [134]. Nuclear IKK α activity, induced through a NIK-dependent pathway, is also involved in ErbB2-induced mammary tumorigenesis [135]. Activated IKK α phosphorylates the cyclin-dependent kinase inhibitor p27/Kip1 and stimulates its nuclear export or exclusion, which in turn correlates with the expansion of tumor-initiating cells. Notably, nuclear IKK α expression in human breast cancer is inversely correlated with nuclear p27 abundance and metastasis-free survival [136]. Colorectal tumors also have active IKK α in the nucleus, in this case concomitant with derepressed SMRT repressor (also known as NCoR2), which is aberrantly localized in the cytoplasm [137]. IKK α phosphorylates SMRT, which is part of a multisubunit repressor complex that includes histone deacetylases [138]. IKK α -mediated phosphorylation induces the exchange of corepressor for coactivator complexes on chromatin and potentiates the acetylation of RelA/p65 by p300, inducing full transcriptional activity [139]. A truncated p45-IKK α variant has been recently identified as the major IKK α form present in the nucleus of colorectal cancer cells. This truncated IKK α is generated by cathepsin activity, which is increased in these cells, and is constitutively active [140]. The role of IKK α in tumor development is dependent on the tissue, it can act as promoter of tumorigenesis in breast and prostate cancer, but is a tumor suppressor in lung carcinomas [134–136,141] and skin cancer [113,114].

Cellular stress responses, which might serve as anti-tumor barriers, activate two major pathways: NF- κ B and p53. The regulation of both pathways shares many similar features and many studies have suggested that crosstalk exists between them. Both transcription factors could compete for a limiting pool of the transcriptional co-activator CBP, as both require such interaction to maximize their activities [1]. IKK α has been shown to phosphorylate CBP at Ser1382 and Ser1386, thereby switching its binding preference from p53 to NF- κ B [142]. CBP phosphorylation status and IKK α activation are directly correlated in several tumor cell lines, suggesting that IKK α -mediated p53/NF- κ B cross-regulation may be a critical factor for cell proliferation and tumor growth [142]. The p53/NF- κ B crosstalk is not restricted to transcriptional regulation, but extends to cytoplasmic IKK function and regulation. IKK and subsequent NF- κ B activities are increased in p53-deficient cells, thereby promoting aerobic glycolysis, which cancer cells typically use as energy source

(Warburg effect). The catalytic activity of IKK β is boosted through an O-linked-N-acetyl-glucosamine modification, establishing a positive feedback regulation from increased glucose metabolism. p53 was also suggested to restrict IKK activation through the suppression of aerobic glycolysis [143,144], and IKK seems to regulate p53 protein levels through direct phosphorylation at Ser362 and Ser366, which leads to its β -TrCP-dependent degradation [145], further emphasizing the crosstalk between these two pathways.

Other tumor suppressors that can be regulated by IKK β are the TSC1/2 complex and the transcription factor FOXO3a. In breast cancer cells, TNF α -induced IKK β phosphorylates TSC1 at S487 and S511, resulting in the disruption of the TSC1/2 tumor suppressor complex and, consequently, in activation of the mTOR pathway. Tumor models expressing a TSC1 phosphomimetic mutant present enhanced angiogenesis and tumor growth. In agreement with this, activated IKK β is associated with TSC1 phosphorylation and VEGF production in different tumor types and correlates with poor clinical outcome of breast cancer patients [146]. The expression of the transcription factor FOXO3a is inversely correlated with that of IKK β in human breast tumor specimens and positively correlated with the survival rate in breast cancer. This is in line with the observation that IKK β can phosphorylate FOXO3a and trigger its ubiquitination-mediated proteasomal degradation [147]. Similarly, pharmacological inhibition of IKK in leukemic cells (AML; T-ALL) was shown to restore FOXO3a expression, correlating with impaired cell proliferation and induced apoptosis [148,149].

Taken together, the various IKK substrates identified to date underscore the enormous complexity of IKK function in (patho) biology (Table 1). In many cases, the detailed regulatory mechanisms are still elusive and there are questions regarding substrate specificity and the signaling outcome depending on the stimulus, tissue and environment. Moreover, cell type-specific variations in IKK expression levels and altered IKK complex composition might contribute to specific functions.

Concluding remarks and perspectives

Our understanding of the impact that IKK/NF- κ B signaling has in mammalian physiology and pathophysiology is continuously growing, as the number of diseases with an involvement of the IKK/NF- κ B system steadily increases. IKK/NF- κ B signaling has also been shown to be one of the key mediators in aging [150]. As an example, hypothalamic programming of systemic ageing in mice depends on IKK β and NF- κ B [151]. Intriguingly, IKK α and IKK β also trigger a wide variety of NF- κ B-independent signaling events, which control various physiological functions and impact disease states like cancer and diabetes (Table 1).

In parallel to the growing insight into its physiological relevance, enormous progress has been made in the mechanistic understanding of IKK/NF- κ B regulation. Signal transmission to IKK is controlled by a complex network of distinct regulatory modules, many of which are not essential, but most likely assure a tailored response to individual stimuli or cell type restrains, or modulate the strength and duration of signaling. Nevertheless, there are major open questions, ensuring many interesting years to come in the field (see Sidebar A).

The recently reported IKK β crystal structures mark an important step in the understanding of IKK function. Further insight can be

Sidebar A. In need of answers

- (i) The structures of activated and resting IKK holo-complexes will provide important insights into the activation mechanism. Another important question is if different cell types—for example in adult or embryonic tissues—contain different subpopulations of IKK complexes with distinct compositions.
- (ii) How the IKK activation loop phosphorylation is mechanistically achieved remains unclear. What are the functions of NEMO ubiquitination and ubiquitin binding in this process? How does the Hsp90/Cdc37 complex influence kinase activity?
- (iii) How are the specificities of IKK for its different substrates—of the NF- κ B pathway or others—determined? What is the role of IKK α in the tripartite complex? Why is IKK α required in the non-canonical pathway?
- (iv) How is IKK activity regulated by diverse types of ubiquitin chains?
- (v) How dynamic is the IKK complex? How is the cytoplasmic IKK complex activated by modified NEMO after its nuclear export? Why do NEMO and IKK α , but not IKK β , appear in the nucleus? How is IKK α recruited to specific gene loci?

anticipated from the complete structures of the remaining subunits and of the holo-complex, as well as from studies of the composition and stimulus-dependent dynamics of the latter.

Ubiquitin-dependent processes are undoubtedly the major driving force of IKK activation. An outstanding issue in this regard is the precise validation of putative ubiquitination targets, which can be addressed through mass spectrometry and genetic complementation experiments, as recently demonstrated for NEMO (see above). In addition to the qualitative determination of ubiquitin-dependent processes, also quantitative analyses may aid to understand the individual impact of the different regulatory modules on IKK regulation. So far, IKK research has been focused on K63-linked and M1-linked ubiquitin chains. However, analysis of *Saccharomyces cerevisiae* revealed that ubiquitin linkages at all seven lysine residues contribute to an unexpected diversity in polyubiquitin chain topology and function [152]. The biological relevance of these ‘unconventional’ ubiquitin chains is underscored by the fact that individual DUBs have evolved distinct ubiquitin linkage specificities [153]. *In vitro* studies have demonstrated that the proteasome can cleave polyubiquitin chains with diverse linkages with preferences following the order K48 > K6, K11, K33 > K27, K29, K63 [154]. Thus, unconventional polyubiquitins could provide properties that are important for the temporal control of NF- κ B signaling.

Although the NF- κ B/IKK pathway is the target of long-standing and intense research, its pharmacological intervention in human disease is still very limited. Given that IKK is involved in numerous physiological processes, it will be important to target the specific upstream modules that control IKK activation in response to particular signals or conditions. However, the design of such a pathway-tailored therapy requires a complete and detailed knowledge of the mechanisms that control IKK activation.

Acknowledgements

We apologize to all colleagues whose contributions to this field could not be cited due to space restrictions. This work was in part supported by grants from Deutsche Forschungsgemeinschaft (DFG) to M.H. and C.S. and the BMBF to C.S.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Baldwin AS (2012) Regulation of cell death and autophagy by IKK and NF-kappaB: critical mechanisms in immune function and cancer. *Immunol Rev* 246: 327–345
- Courtois G, Gilmore TD (2006) Mutations in the NF-kappaB signaling pathway: implications for human disease. *Oncogene* 25: 6831–6843
- Hayden MS, Ghosh S (2012) NF-kappaB, the first quarter-century: remarkable progress and outstanding questions. *Genes Dev* 26: 203–234
- Hinz M, Arslan SC, Scheidereit C (2012) It takes two to tango: IkappaBs, the multifunctional partners of NF-kappaB. *Immunol Rev* 246: 59–76
- Kanarek N, Ben-Neriah Y (2012) Regulation of NF-kappaB by ubiquitination and degradation of the IkappaBs. *Immunol Rev* 246: 77–94
- Heissmeyer V, Krappmann D, Wulczyn FG, Scheidereit C (1999) NF-kappaB p105 is a target of IkappaB kinases and controls signal induction of Bcl-3-p50 complexes. *EMBO J* 18: 4766–4778
- Chen ZJ, Parent L, Maniatis T (1996) Site-specific phosphorylation of I-kappaBalpha by a novel ubiquitination-dependent protein kinase activity. *Cell* 84: 853–862
- DiDonato JA, Hayakawa M, Rothwarf DM, Zandi E, Karin M (1997) A cytokine-responsive IkappaB kinase that activates the transcription factor NF-kappaB. *Nature* 388: 548–554
- Mercurio F, Zhu H, Murray BW, Shevchenko A, Bennett BL, Li J, Young DB, Barbosa M, Mann M, Manning A, Rao A (1997) IKK-1 and IKK-2: cytokine-activated IkappaB kinases essential for NF-kappaB activation. *Science* 278: 860–866
- Regnier CH, Song HY, Gao X, Goeddel DV, Cao Z, Rothe M (1997) Identification and characterization of an IkappaB kinase. *Cell* 90: 373–383
- Woronicz JD, Gao X, Cao Z, Rothe M, Goeddel DV (1997) IkappaB kinase-beta: NF-kappaB activation and complex formation with IkappaB kinase-alpha and NIK. *Science* 278: 866–869
- Zandi E, Rothwarf DM, Delhase M, Hayakawa M, Karin M (1997) The IkappaB kinase complex (IKK) contains two kinase subunits, IKKalpha and IKKbeta, necessary for IkappaB phosphorylation and NF-kappaB activation. *Cell* 91: 243–252
- Zandi E, Chen Y, Karin M (1998) Direct phosphorylation of IkappaB by IKKalpha and IKKbeta: discrimination between free and NF-kappaB-bound substrate. *Science* 281: 1360–1363
- Rothwarf DM, Zandi E, Natoli G, Karin M (1998) IKK-gamma is an essential regulatory subunit of the IkappaB kinase complex. *Nature* 395: 297–300
- Yamaoka S, Courtois G, Bessia C, Whiteside ST, Weil R, Agou F, Kirk HE, Kay RJ, Israel A (1998) Complementation cloning of NEMO, a component of the IkappaB kinase complex essential for NF-kappaB activation. *Cell* 93: 1231–1240
- Liu F, Xia Y, Parker AS, Verma IM (2012) IKK biology. *Immunol Rev* 246: 239–253

17. de Leon-Boenig G, Bowman KK, Feng JA, Crawford T, Everett C, Franke Y, Oh A, Stanley M, Staben ST, Starovasnik MA, Wallweber HJ, Wu J, Wu LC, Johnson AR, Hymowitz SG (2012) The crystal structure of the catalytic domain of the NF-kappaB inducing kinase reveals a narrow but flexible active site. *Structure* 20: 1704–1714
18. Vallabhapurapu S, Matsuzawa A, Zhang W, Tseng PH, Keats JJ, Wang H, Vignali DA, Bergsagel PL, Karin M (2008) Nonredundant and complementary functions of TRAF2 and TRAF3 in a ubiquitination cascade that activates NIK-dependent alternative NF-kappaB signaling. *Nat Immunol* 9: 1364–1370
19. Zarnegar BJ, Wang Y, Mahoney DJ, Dempsey PW, Cheung HH, He J, Shiba T, Yang X, Yeh WC, Mak TW, Korneluk RG, Cheng G (2008) Noncanonical NF-kappaB activation requires coordinated assembly of a regulatory complex of the adaptors cIAP1, cIAP2, TRAF2 and TRAF3 and the kinase NIK. *Nat Immunol* 9: 1371–1378
20. Sun SC (2012) The noncanonical NF-kappaB pathway. *Immunol Rev* 246: 125–140
21. Hu H, Brittain GC, Chang JH, Puebla-Orsorio N, Jin J, Zal A, Xiao Y, Cheng X, Chang M, Fu YX, Zal T, Zhu C, Sun SC (2013) OTUD7B controls non-canonical NF-kappaB activation through deubiquitination of TRAF3. *Nature* 494: 371–374
22. Israel A (2010) The IKK complex, a central regulator of NF-kappaB activation. *Cold Spring Harb Perspect Biol* 2: a000158
23. Scheidereit C (2006) IkappaB kinase complexes: gateways to NF-kappaB activation and transcription. *Oncogene* 25: 6685–6705
24. Sil AK, Maeda S, Sano Y, Roop DR, Karin M (2004) IkappaB kinase-alpha acts in the epidermis to control skeletal and craniofacial morphogenesis. *Nature* 428: 660–664
25. Ling L, Cao Z, Goeddel DV (1998) NF-kappaB-inducing kinase activates IKK-alpha by phosphorylation of Ser-176. *Proc Natl Acad Sci U S A* 95: 3792–3797
26. May MJ, Larsen SE, Shim JH, Madge LA, Ghosh S (2004) A novel ubiquitin-like domain in IkappaB kinase beta is required for functional activity of the kinase. *J Biol Chem* 279: 45528–45539
27. Liu S, Misquitta Y. R., Olland A., Johnson M. A., Kelleher K. S., Kriz R., Lin L. L., Stahl M., Mosyak L. (2013) Crystal Structure of A Human IkappaB Kinase beta Asymmetric Dimer. *J Biol Chem* 288: 22758–22767
28. Polley S, Huang DB, Hauenstein AV, Fusco AJ, Zhong X, Vu D, Schrofelbauer B, Kim Y, Hoffmann A, Verma IM, Ghosh G, Huxford T (2013) A structural basis for IkappaB kinase 2 activation via oligomerization-dependent trans auto-phosphorylation. *PLoS Biol* 11: e1001581
29. Xu G, Lo YC, Li Q, Napolitano G, Wu X, Jiang X, Dreano M, Karin M, Wu H (2011) Crystal structure of inhibitor of kappaB kinase beta. *Nature* 472: 325–330
30. Clement JF, Meloche S, Servant MJ (2008) The IKK-related kinases: from innate immunity to oncogenesis. *Cell Res* 18: 889–899
31. Larabi A, Devos JM, Ng SL, Nanao MH, Round A, Maniatis T, Panne D (2013) Crystal structure and mechanism of activation of TANK-binding kinase 1. *Cell Rep* 3: 734–746
32. Tu D, Zhu Z, Zhou AY, Yun CH, Lee KE, Toms AV, Li Y, Dunn GP, Chan E, Thai T, Yang S, Ficarro SB, Marto JA, Jeon H, Hahn WC, Barbie DA, Eck MJ (2013) Structure and ubiquitination-dependent activation of TANK-binding kinase 1. *Cell Rep* 3: 747–758
33. Ma X, Helgason E, Phung QT, Quan CL, Iyer RS, Lee MW, Bowman KK, Starovasnik MA, Dueber EC (2012) Molecular basis of Tank-binding kinase 1 activation by transautophosphorylation. *Proc Natl Acad Sci U S A* 109: 9378–9383
34. Schrofelbauer B, Polley S, Behar M, Ghosh G, Hoffmann A (2012) NEMO ensures signaling specificity of the pleiotropic IKKbeta by directing its kinase activity toward IkappaBalpha. *Mol Cell* 47: 111–121
35. Agou F, Traincard F, Vinolo E, Courtois G, Yamaoka S, Israel A, Veron M (2004) The trimerization domain of NEMO is composed of the interacting C-terminal CC2 and LZ coiled-coil subdomains. *J Biol Chem* 279: 27861–27869
36. Tegethoff S, Behlke J, Scheidereit C (2003) Tetrameric oligomerization of IkappaB kinase gamma (IKKgamma) is obligatory for IKK complex activity and NF-kappaB activation. *Mol Cell Biol* 23: 2029–2041
37. Rushe M, Silvian L, Bixler S, Chen LL, Cheung A, Bowes S, Cuervo H, Berkowitz S, Zheng T, Guckian K, Pellegrini M, Lugovskoy A (2008) Structure of a NEMO/IKK-associating domain reveals architecture of the interaction site. *Structure* 16: 798–808
38. Bagneris C, Ageichik AV, Cronin N, Wallace B, Collins M, Boshoff C, Waksman G, Barrett T (2008) Crystal structure of a vFlip-IKKgamma complex: insights into viral activation of the IKK signalosome. *Mol Cell* 30: 620–631
39. Cordier F, Grubisha O, Traincard F, Veron M, Delepiepierre M, Agou F (2009) The zinc finger of NEMO is a functional ubiquitin-binding domain. *J Biol Chem* 284: 2902–2907
40. Grubisha O, Kaminska M, Duquerroy S, Fontan E, Cordier F, Haouz A, Raynal B, Chiaravalli J, Delepiepierre M, Israel A, Veron M, Agou F (2010) DARPIn-assisted crystallography of the CC2-LZ domain of NEMO reveals a coupling between dimerization and ubiquitin binding. *J Mol Biol* 395: 89–104
41. Rahighi S, Ikeda F, Kawasaki M, Akutsu M, Suzuki N, Kato R, Kensche T, Uejima T, Bloor S, Komander D, Randow F, Wakatsuki S, Dikic I (2009) Specific recognition of linear ubiquitin chains by NEMO is important for NF-kappaB activation. *Cell* 136: 1098–1109
42. Chen ZJ (2012) Ubiquitination in signaling to and activation of IKK. *Immunol Rev* 246: 95–106
43. Schmukle AC, Walczak H (2012) No one can whistle a symphony alone – how different ubiquitin linkages cooperate to orchestrate NF-kappaB activity. *J Cell Sci* 125: 549–559
44. Hubeau M, Ngadjieu F, Puel A, Israel L, Feinberg J, Chrabieh M, Belani K, Bodemer C, Fabre I, Plebani A, Boisson-Dupuis S, Picard C, Fischer A, Israel A, Abel L, Veron M, Casanova JL, Agou F, Bustamante J (2011) New mechanism of X-linked anhidrotic ectodermal dysplasia with immunodeficiency: impairment of ubiquitin binding despite normal folding of NEMO protein. *Blood* 118: 926–935
45. Lo YC, Lin SC, Rospigliosi CC, Conze DB, Wu CJ, Ashwell JD, Eliezer D, Wu H (2009) Structural basis for recognition of diubiquitins by NEMO. *Mol Cell* 33: 602–615
46. Komander D, Reyes-Turcu F, Licchesi JD, Odenwaelde P, Wilkinson KD, Barford D (2009) Molecular discrimination of structurally equivalent Lys 63-linked and linear polyubiquitin chains. *EMBO Rep* 10: 466–473
47. Dynek JN, Goncharov T, Dueber EC, Fedorova AV, Izrael-Tomasevic A, Phu L, Helgason E, Fairbrother WJ, Deshayes K, Kirkpatrick DS, Vucic D (2010) c-IAP1 and UbcH5 promote K11-linked polyubiquitination of RIP1 in TNF signalling. *EMBO J* 29: 4198–4209
48. Laplantine E, Fontan E, Chiaravalli J, Lopez T, Lakisic G, Veron M, Agou F, Israel A (2009) NEMO specifically recognizes K63-linked poly-ubiquitin chains through a new bipartite ubiquitin-binding domain. *EMBO J* 28: 2885–2895
49. Cordier F, Vinolo E, Veron M, Delepiepierre M, Agou F (2008) Solution structure of NEMO zinc finger and impact of an anhidrotic ectodermal

- dysplasia with immunodeficiency-related point mutation. *J Mol Biol* 377: 1419–1432
50. Hadian K, Griesbach RA, Dornauer S, Wanger TM, Nagel D, Metlitzky M, Beisker W, Schmidt-Supprian M, Krappmann D (2011) NF-kappaB essential modulator (NEMO) interaction with linear and lys-63 ubiquitin chains contributes to NF-kappaB activation. *J Biol Chem* 286: 26107–26117
 51. Kensche T, Tokunaga F, Ikeda F, Goto E, Iwai K, Dikic I (2012) Analysis of nuclear factor-kappaB (NF-kappaB) essential modulator (NEMO) binding to linear and lysine-linked ubiquitin chains and its role in the activation of NF-kappaB. *J Biol Chem* 287: 23626–23634
 52. Emmerich C. H., Ordureau A., Strickson S., Arthur J. S., Pedrioli P. G., Komander D., Cohen P. (2013) Activation of the canonical IKK complex by K63/M1-linked hybrid ubiquitin chains. *Proc Natl Acad Sci U S A* 110: 15247–15252
 53. Hinz M, Broemer M, Arslan SC, Otto A, Mueller EC, Dettmer R, Scheidereit C (2007) Signal responsiveness of IkappaB kinases is determined by Cdc37-assisted transient interaction with Hsp90. *J Biol Chem* 282: 32311–32319
 54. Krappmann D, Hatada EN, Tegethoff S, Li J, Klippel A, Giese K, Baeuerle PA, Scheidereit C (2000) The I kappa B kinase (IKK) complex is tripartite and contains IKK gamma but not IKAP as a regular component. *J Biol Chem* 275: 29779–29787
 55. Miller BS, Zandi E (2001) Complete reconstitution of human IkappaB kinase (IKK) complex in yeast. Assessment of its stoichiometry and the role of IKKgammam in the complex activity in the absence of stimulation. *J Biol Chem* 276: 36320–36326
 56. Drew D, Shimada E, Huynh K, Bergqvist S, Talwar R, Karin M, Ghosh G (2007) Inhibitor kappaB kinase beta binding by inhibitor kappaB kinase gamma. *Biochemistry* 46: 12482–12490
 57. Senftleben U, Cao Y, Xiao G, Greten FR, Krahn G, Bonizzi G, Chen Y, Hu Y, Fong A, Sun SC, Karin M (2001) Activation by IKKalpha of a second, evolutionary conserved, NF-kappa B signaling pathway. *Science* 293: 1495–1499
 58. Vallabhapurapu S, Karin M (2009) Regulation and function of NF-kappaB transcription factors in the immune system. *Annu Rev Immunol* 27: 693–733
 59. Gewurz BE, Towfic F, Mar JC, Shinnars NP, Takasaki K, Zhao B, Cahir-McFarland ED, Quackenbush J, Xavier RJ, Kieff E (2012) Genome-wide siRNA screen for mediators of NF-kappaB activation. *Proc Natl Acad Sci U S A* 109: 2467–2472
 60. Matta H, Gopalakrishnan R, Graham C, Tolani B, Khanna A, Yi H, Suo Y, Chaudhary PM (2012) Kaposi's sarcoma associated herpesvirus encoded viral FLICE inhibitory protein K13 activates NF-kappaB pathway independent of TRAF6, TAK1 and LUBAC. *PLoS ONE* 7: e36601
 61. Deng L, Wang C, Spencer E, Yang L, Braun A, You J, Slaughter C, Pickart C, Chen ZJ (2000) Activation of the IkappaB kinase complex by TRAF6 requires a dimeric ubiquitin-conjugating enzyme complex and a unique polyubiquitin chain. *Cell* 103: 351–361
 62. Wang C, Deng L, Hong M, Akkaraju GR, Inoue J, Chen ZJ (2001) TAK1 is a ubiquitin-dependent kinase of MKK and IKK. *Nature* 412: 346–351
 63. Huang Q, Yang J, Lin Y, Walker C, Cheng J, Liu ZG, Su B (2004) Differential regulation of interleukin 1 receptor and Toll-like receptor signaling by MEKK3. *Nat Immunol* 5: 98–103
 64. Yang J, Lin Y, Guo Z, Cheng J, Huang J, Deng L, Liao W, Chen Z, Liu Z, Su B (2001) The essential role of MEKK3 in TNF-induced NF-kappaB activation. *Nat Immunol* 2: 620–624
 65. Qin J, Yao J, Cui G, Xiao H, Kim TW, Fraczek J, Wightman P, Sato S, Akira S, Puel A, Casanova JL, Su B, Li X (2006) TLR8-mediated NF-kappaB and JNK activation are TAK1-independent and MEKK3-dependent. *J Biol Chem* 281: 21013–21021
 66. Yao J, Kim TW, Qin J, Jiang Z, Qian Y, Xiao H, Lu Y, Qian W, Gulen MF, Sizemore N, DiDonato J, Sato S, Akira S, Su B, Li X (2007) Interleukin-1 (IL-1)-induced TAK1-dependent Versus MEKK3-dependent NFkappaB activation pathways bifurcate at IL-1 receptor-associated kinase modification. *J Biol Chem* 282: 6075–6089
 67. Delhase M, Hayakawa M, Chen Y, Karin M (1999) Positive and negative regulation of IkappaB kinase activity through IKKbeta subunit phosphorylation. *Science* 284: 309–313
 68. Walsh MC, Kim GK, Maurizio PL, Molnar EE, Choi Y (2008) TRAF6 auto-ubiquitination-independent activation of the NFkappaB and MAPK pathways in response to IL-1 and RANKL. *PLoS ONE* 3: e4064
 69. Xia ZP, Sun LJ, Chen X, Pineda G, Jiang XM, Adhikari A, Zeng WW, Chen ZJ (2009) Direct activation of protein kinases by unanchored polyubiquitin chains. *Nature* 461: 114–U125
 70. Xu M, Skaug B, Zeng W, Chen ZJ (2009) A ubiquitin replacement strategy in human cells reveals distinct mechanisms of IKK activation by TNFalpha and IL-1beta. *Mol Cell* 36: 302–314
 71. Gerlach B, Cordier SM, Schmukle AC, Emmerich CH, Rieser E, Haas TL, Webb AI, Rickard JA, Anderton H, Wong WW, Nachbur U, Gangoda L, Warnken U, Purcell AW, Silke J, Walczak H (2011) Linear ubiquitination prevents inflammation and regulates immune signalling. *Nature* 471: 591–596
 72. Tokunaga F, Sakata S, Saeki Y, Satomi Y, Kirisako T, Kamei K, Nakagawa T, Kato M, Murata S, Yamaoka S, Yamamoto M, Akira S, Takao T, Tanaka K, Iwai K (2009) Involvement of linear polyubiquitylation of NEMO in NF-kappaB activation. *Nat Cell Biol* 11: 123–132
 73. Iwai K (2012) Diverse ubiquitin signaling in NF-kappaB activation. *Trends Cell Biol* 22: 355–364
 74. Muller-Rischart AK, Pils A, Beaudette P, Patra M, Hadian K, Funke M, Peis R, Deinlein A, Schweimer C, Kuhn PH, Lichtenthaler SF, Motori E, Hrelia S, Wurst W, Trumbach D, Langer T, Krappmann D, Dittmar G, Tatzelt J, Winklhofer KF (2013) The E3 ligase parkin maintains mitochondrial integrity by increasing linear ubiquitination of NEMO. *Mol Cell* 49: 908–921
 75. Sasaki Y., Sano S., Nakahara M., Murata S., Kometani K., Aiba Y., Sakamoto S., Watanabe Y., Tanaka K., Kurosaki T., Iwai K. (2013) Defective immune responses in mice lacking LUBAC-mediated linear ubiquitination in B cells. *EMBO J* 32: 2463–2476
 76. Ikeda F, Deribe YL, Skanland SS, Stieglitz B, Grabbe C, Franz-Wachtel M, van Wijk SJ, Goswami P, Nagy V, Terzic J, Tokunaga F, Androulidaki A, Nakagawa T, Pasparakis M, Iwai K, Sundberg JP, Schaefer L, Rittinger K, Macek B, Dikic I (2011) SHARPIN forms a linear ubiquitin ligase complex regulating NF-kappaB activity and apoptosis. *Nature* 471: 637–641
 77. Tokunaga F, Nakagawa T, Nakahara M, Saeki Y, Taniguchi M, Sakata S, Tanaka K, Nakano H, Iwai K (2011) SHARPIN is a component of the NF-kappaB-activating linear ubiquitin chain assembly complex. *Nature* 471: 633–636
 78. Boisson B, Laplantine E, Prando C, Giliani S, Israelsson E, Xu Z, Abhyankar A, Israel L, Trevejo-Nunez G, Bogunovic D, Cepika AM, MacDuff D, Chrabieh M, Hubeau M, Bajolle F, Debre M, Mazzolari E, Vairo D, Agou F, Virgin HW, et al (2012) Immunodeficiency, autoinflammation and amylopectinosis in humans with inherited HOIL-1 and LUBAC deficiency. *Nat Immunol* 13: 1178–1186

79. Arimoto K, Funami K, Saeki Y, Tanaka K, Okawa K, Takeuchi O, Akira S, Murakami Y, Shimotohno K (2010) Polyubiquitin conjugation to NEMO by tripartite motif protein 23 (TRIM23) is critical in antiviral defense. *Proc Natl Acad Sci U S A* 107: 15856–15861
80. Tang ED, Wang CY, Xiong Y, Guan KL (2003) A role for NF-kappaB essential modifier/IkappaB kinase-gamma (NEMO/IKKgamma) ubiquitination in the activation of the IkappaB kinase complex by tumor necrosis factor-alpha. *J Biol Chem* 278: 37297–37305
81. Ea CK, Deng L, Xia ZP, Pineda G, Chen ZJ (2006) Activation of IKK by TNFalpha requires site-specific ubiquitination of RIP1 and polyubiquitin binding by NEMO. *Mol Cell* 22: 245–257
82. Newton K, Matsumoto ML, Wertz IE, Kirkpatrick DS, Lill JR, Tan J, Dugger D, Gordon N, Sidhu SS, Fellouse FA, Komuves L, French DM, Ferrando RE, Lam C, Compaan D, Yu C, Bosanac I, Hymowitz SG, Kelley RF, Dixit VM (2008) Ubiquitin chain editing revealed by polyubiquitin linkage-specific antibodies. *Cell* 134: 668–678
83. Wu CJ, Conze DB, Li T, Srinivasula SM, Ashwell JD (2006) Sensing of Lys 63-linked polyubiquitination by NEMO is a key event in NF-kappaB activation [corrected]. *Nat Cell Biol* 8: 398–406
84. Hoeller D, Crosetto N, Blagoev B, Raiborg C, Tikkanen R, Wagner S, Kowanetz K, Breitling R, Mann M, Stenmark H, Dikic I (2006) Regulation of ubiquitin-binding proteins by monoubiquitination. *Nat Cell Biol* 8: 163–169
85. Hinz M, Stilmann M, Arslan SC, Khanna KK, Dittmar G, Scheidereit C (2010) A cytoplasmic ATM-TRAF6-clAP1 module links nuclear DNA damage signaling to ubiquitin-mediated NF-kappaB activation. *Mol Cell* 40: 63–74
86. Huang TT, Wuerzberger-Davis SM, Wu ZH, Miyamoto S (2003) Sequential modification of NEMO/IKKgamma by SUMO-1 and ubiquitin mediates NF-kappaB activation by genotoxic stress. *Cell* 115: 565–576
87. Jun J. C., Kertesz S., Jones M. B., Marini J. M., Cobb B. A., Tigno-Aranjuez J. T., Abbott D. W. (2013) Innate Immune-Directed NF-kappaB Signaling Requires Site-Specific NEMO Ubiquitination. *Cell Rep* 4: 352–361
88. Napetschnig J, Wu H (2013) Molecular basis of NF-kappaB signaling. *Annu Rev Biophys* 42: 443–468
89. Lin SC, Lo YC, Wu H (2010) Helical assembly in the MyD88-IRAK4-IRAK2 complex in TLR/IL-1R signalling. *Nature* 465: 885–890
90. Hou F, Sun L, Zheng H, Skaug B, Jiang QX, Chen ZJ (2011) MAVS forms functional prion-like aggregates to activate and propagate antiviral innate immune response. *Cell* 146: 448–461
91. Yin Q, Lin SC, Lamothe B, Lu M, Lo YC, Hura G, Zheng L, Rich RL, Campos AD, Myszkowski DG, Lenardo MJ, Darnay BG, Wu H (2009) E2 interaction and dimerization in the crystal structure of TRAF6. *Nat Struct Mol Biol* 16: 658–666
92. Wu H (2013) Higher-order assemblies in a new paradigm of signal transduction. *Cell* 153: 287–292
93. Palkowitsch L, Leidner J, Ghosh S, Marienfeld RB (2008) Phosphorylation of serine 68 in the IkappaB kinase (IKK)-binding domain of NEMO interferes with the structure of the IKK complex and tumor necrosis factor-alpha-induced NF-kappaB activity. *J Biol Chem* 283: 76–86
94. Schomer-Miller B, Higashimoto T, Lee YK, Zandi E (2006) Regulation of IkappaB kinase (IKK) complex by IKKgamma-dependent phosphorylation of the T-loop and C terminus of IKKbeta. *J Biol Chem* 281: 15268–15276
95. Lobry C, Lopez T, Israel A, Weil R (2007) Negative feedback loop in T cell activation through IkappaB kinase-induced phosphorylation and degradation of Bcl10. *Proc Natl Acad Sci U S A* 104: 908–913
96. Wegener E, Oeckinghaus A, Papadopolou N, Lavitas L, Schmidt-Suppran M, Ferch U, Mak TW, Ruland J, Heissmeyer V, Krappmann D (2006) Essential role for IkappaB kinase beta in remodeling Carma1-Bcl10-Malt1 complexes upon T cell activation. *Mol Cell* 23: 13–23
97. Zhu G, Wu CJ, Zhao Y, Ashwell JD (2007) Optineurin negatively regulates TNFalpha-induced NF-kappaB activation by competing with NEMO for ubiquitinated RIP. *Curr Biol* 17: 1438–1443
98. Maruyama H, Morino H, Ito H, Izumi Y, Kato H, Watanabe Y, Kinoshita Y, Kamada M, Nodera H, Suzuki H, Komure O, Matsuura S, Kobatake K, Morimoto N, Abe K, Suzuki N, Aoki M, Kawata A, Hirai T, Kato T, et al (2010) Mutations of optineurin in amyotrophic lateral sclerosis. *Nature* 465: 223–226
99. Shibata Y, Oyama M, Kozuka-Hata H, Han X, Tanaka Y, Gohda J, Inoue J (2012) p47 negatively regulates IKK activation by inducing the lysosomal degradation of polyubiquitinated NEMO. *Nat Commun* 3: 1061
100. Harhaj EW, Dixit VM (2012) Regulation of NF-kappaB by deubiquitinases. *Immunol Rev* 246: 107–124
101. Bremm A, Freund SM, Komander D (2010) Lys11-linked ubiquitin chains adopt compact conformations and are preferentially hydrolyzed by the deubiquitinase Cezanne. *Nat Struct Mol Biol* 17: 939–947
102. Enesa K, Zakkar M, Chaudhury H, le Luong A, Rawlinson L, Mason JC, Haskard DO, Dean JL, Evans PC (2008) NF-kappaB suppression by the deubiquitinating enzyme Cezanne: a novel negative feedback loop in pro-inflammatory signaling. *J Biol Chem* 283: 7036–7045
103. Xu G, Tan X, Wang H, Sun W, Shi Y, Burlingame S, Gu X, Cao G, Zhang T, Qin J, Yang J (2010) Ubiquitin-specific peptidase 21 inhibits tumor necrosis factor alpha-induced nuclear factor kappaB activation via binding to and deubiquitinating receptor-interacting protein 1. *J Biol Chem* 285: 969–978
104. Fiil BK, Damgaard RB, Wagner SA, Keusekotten K, Fritsch M, Bekker-Jensen S, Mailand N, Choudhary C, Komander D, Gyrd-Hansen M (2013) OTULIN restricts Met1-linked ubiquitination to control innate immune signaling. *Mol Cell* 50: 818–830
105. Keusekotten K, Elliott PR, Glockner L, Fiil BK, Damgaard RB, Kulathu Y, Wauer T, Hospenthal MK, Gyrd-Hansen M, Krappmann D, Hofmann K, Komander D (2013) OTULIN antagonizes LUBAC signaling by specifically hydrolyzing Met1-linked polyubiquitin. *Cell* 153: 1312–1326
106. Rivkin E, Almeida SM, Ceccarelli DF, Juang YC, MacLean TA, Srikumar T, Huang H, Dunham WH, Fukumura R, Xie G, Gondo Y, Raught B, Gingras AC, Sicheri F, Cordes SP (2013) The linear ubiquitin-specific deubiquitinase gumby regulates angiogenesis. *Nature* 498: 318–324
107. Lee MH, Mabb AM, Gill GB, Yeh ET, Miyamoto S (2011) NF-kappaB induction of the SUMO protease SENP2: a negative feedback loop to attenuate cell survival response to genotoxic stress. *Mol Cell* 43: 180–191
108. Li Q, Van Antwerp D, Mercurio F, Lee KF, Verma IM (1999) Severe liver degeneration in mice lacking the IkappaB kinase 2 gene. *Science* 284: 321–325
109. Li ZW, Chu W, Hu Y, Delhase M, Deerinc T, Ellisman M, Johnson R, Karin M (1999) The IKKbeta subunit of IkappaB kinase (IKK) is essential for nuclear factor kappaB activation and prevention of apoptosis. *J Exp Med* 189: 1839–1845
110. Solt LA, Madge LA, Orange JS, May MJ (2007) Interleukin-1-induced NF-kappaB activation is NEMO-dependent but does not require IKKbeta. *J Biol Chem* 282: 8724–8733
111. Li Q, Lu Q, Hwang JY, Buscher D, Lee KF, Izpisua-Belmonte JC, Verma IM (1999) IKK1-deficient mice exhibit abnormal development of skin and skeleton. *Genes Dev* 13: 1322–1328

112. Descargues P, Sil AK, Karin M (2008) IKKalpha, a critical regulator of epidermal differentiation and a suppressor of skin cancer. *EMBO J* 27: 2639–2647
113. Liu B, Park E, Zhu F, Bustos T, Liu J, Shen J, Fischer SM, Hu Y (2006) A critical role for I kappaB kinase alpha in the development of human and mouse squamous cell carcinomas. *Proc Natl Acad Sci U S A* 103: 17202–17207
114. Liu B, Xia X, Zhu F, Park E, Carbajal S, Kiguchi K, DiGiovanni J, Fischer SM, Hu Y (2008) IKKalpha is required to maintain skin homeostasis and prevent skin cancer. *Cancer Cell* 14: 212–225
115. Li N, Wu X, Holzer RG, Lee JH, Todoric J, Park EJ, Ogata H, Gukovskaya AS, Gukovsky I, Pizzo DP, VandenBerg S, Tarin D, Atay C, Arkan MC, Deerinck TJ, Moscat J, Diaz-Meco M, Dawson D, Erkan M, Kleeff J, et al (2013) Loss of acinar cell IKKalpha triggers spontaneous pancreatitis in mice. *J Clin Invest* 123: 2231–2243
116. Mancino A, Habbeldine M, Johnson E, Luron L, Bebie M, Memet S, Fong C, Bajenoff M, Wu X, Karin M, Caamano J, Chi H, Seed M, Lawrence T (2013) I kappa B kinase alpha (IKKalpha) activity is required for functional maturation of dendritic cells and acquired immunity to infection. *EMBO J* 32: 816–828
117. McCool KW, Miyamoto S (2012) DNA damage-dependent NF-kappaB activation: NEMO turns nuclear signaling inside out. *Immunol Rev* 246: 311–326
118. Lawrence T, Bebie M, Liu GY, Nizet V, Karin M (2005) IKKalpha limits macrophage NF-kappaB activation and contributes to the resolution of inflammation. *Nature* 434: 1138–1143
119. Sakurai H, Chiba H, Miyoshi H, Sugita T, Toriumi W (1999) IkappaB kinases phosphorylate NF-kappaB p65 subunit on serine 536 in the trans-activation domain. *J Biol Chem* 274: 30353–30356
120. Chariot A (2009) The NF-kappaB-independent functions of IKK subunits in immunity and cancer. *Trends Cell Biol* 19: 404–413
121. Albanese C, Wu K, D'Amico M, Jarrett C, Joyce D, Hughes J, Hult J, Sakamaki T, Fu M, Ben-Ze'ev A, Bromberg JF, Lamberti C, Verma U, Gaynor RB, Byers SW, Pestell RG (2003) IKKalpha regulates mitogenic signaling through transcriptional induction of cyclin D1 via Tcf. *Mol Biol Cell* 14: 585–599
122. Carayol N, Wang CY (2006) IKKalpha stabilizes cytosolic beta-catenin by inhibiting both canonical and non-canonical degradation pathways. *Cell Signal* 18: 1941–1946
123. Lamberti C, Lin KM, Yamamoto Y, Verma U, Verma IM, Byers S, Gaynor RB (2001) Regulation of beta-catenin function by the IkappaB kinases. *J Biol Chem* 276: 42276–42286
124. Hideshima T, Chauhan D, Kiziltepe T, Ikeda H, Okawa Y, Podar K, Raju N, Protopopov A, Munshi NC, Richardson PG, Carrasco RD, Anderson KC (2009) Biologic sequelae of I{kappa}B kinase (IKK) inhibition in multiple myeloma: therapeutic implications. *Blood* 113: 5228–5236
125. Guttridge DC, Albanese C, Reuther JY, Pestell RG, Baldwin AS Jr (1999) NF-kappaB controls cell growth and differentiation through transcriptional regulation of cyclin D1. *Mol Cell Biol* 19: 5785–5799
126. Hinz M, Krappmann D, Eichten A, Heder A, Scheidereit C, Strauss M (1999) NF-kappaB function in growth control: regulation of cyclin D1 expression and G0/G1-to-S-phase transition. *Mol Cell Biol* 19: 2690–2698
127. Kwak YT, Li R, Becerra CR, Tripathy D, Frenkel EP, Verma UN (2005) IkappaB kinase alpha regulates subcellular distribution and turnover of cyclin D1 by phosphorylation. *J Biol Chem* 280: 33945–33952
128. Yan J, Xiang J, Lin Y, Ma J, Zhang J, Zhang H, Sun J, Danial NN, Liu J, Lin A (2013) Inactivation of BAD by IKK inhibits TNFalpha-induced apoptosis independently of NF-kappaB activation. *Cell* 152: 304–315
129. Jia CH, Li M, Liu J, Zhao L, Lin J, Lai PL, Zhou X, Zhang Y, Chen ZG, Li HY, Liu AL, Yang CL, Gao TM, Jiang Y, Bai XC (2013) IKK-beta mediates hydrogen peroxide induced cell death through p85 S6K1. *Cell Death Differ* 20: 248–258
130. Comb WC, Cogswell P, Sitcheran R, Baldwin AS (2011) IKK-dependent, NF-kappaB-independent control of autophagic gene expression. *Oncogene* 30: 1727–1732
131. Criollo A, Senovilla L, Authier H, Maiuri MC, Morselli E, Vitale I, Kepp O, Tasdemir E, Galluzzi L, Shen S, Tailler M, Delahaye N, Tesniere A, De Stefano D, Younes AB, Harper F, Pierron G, Lavandro S, Zitvogel L, Israel A, et al (2010) The IKK complex contributes to the induction of autophagy. *EMBO J* 29: 619–631
132. Comb WC, Hutt JE, Cogswell P, Cantley LC, Baldwin AS (2012) p85alpha SH2 domain phosphorylation by IKK promotes feedback inhibition of PI3K and Akt in response to cellular starvation. *Mol Cell* 45: 719–730
133. Dan HC, Adli M, Baldwin AS (2007) Regulation of mammalian target of rapamycin activity in PTEN-inactive prostate cancer cells by I kappa B kinase alpha. *Cancer Res* 67: 6263–6269
134. Luo JL, Tan W, Ricono JM, Korchynskyi O, Zhang M, Gonias SL, Cheresch DA, Karin M (2007) Nuclear cytokine-activated IKKalpha controls prostate cancer metastasis by repressing Maspin. *Nature* 446: 690–694
135. Cao Y, Luo JL, Karin M (2007) IkappaB kinase alpha kinase activity is required for self-renewal of ErbB2/Her2-transformed mammary tumor-initiating cells. *Proc Natl Acad Sci U S A* 104: 15852–15857
136. Zhang W, Tan W, Wu X, Poustovoitov M, Strasner A, Li W, Borcherdig N, Ghassemian M, Karin M (2013) A NIK-IKKalpha module expands ErbB2-induced tumor-initiating cells by stimulating nuclear export of p27/Kip1. *Cancer Cell* 23: 647–659
137. Fernandez-Majada V, Aguilera C, Villanueva A, Vilardell F, Robert-Moreno A, Aytes A, Real FX, Capella G, Mayo MW, Espinosa L, Bigas A (2007) Nuclear IKK activity leads to dysregulated notch-dependent gene expression in colorectal cancer. *Proc Natl Acad Sci U S A* 104: 276–281
138. Hoberg JE, Yeung F, Mayo MW (2004) SMRT derepression by the IkappaB kinase alpha: a prerequisite to NF-kappaB transcription and survival. *Mol Cell* 16: 245–255
139. Hoberg JE, Popko AE, Ramsey CS, Mayo MW (2006) IkappaB kinase alpha-mediated derepression of SMRT potentiates acetylation of RelA/p65 by p300. *Mol Cell Biol* 26: 457–471
140. Margalef P, Fernandez-Majada V, Villanueva A, Garcia-Carbonell R, Iglesias M, Lopez L, Martinez-Iniesta M, Villa-Freixa J, Mulero MC, Andreu M, Torres F, Mayo MW, Bigas A, Espinosa L (2012) A truncated form of IKKalpha is responsible for specific nuclear IKK activity in colorectal cancer. *Cell Rep* 2: 840–854
141. Xiao Z, Jiang Q, Willette-Brown J, Xi S, Zhu F, Burkett S, Back T, Song NY, Datla M, Sun Z, Goldszmid R, Lin F, Cohoon T, Pike K, Wu X, Schrumpp DS, Wong KK, Young HA, Trinchieri G, Wiltout RH, et al (2013) The pivotal role of IKKalpha in the development of spontaneous lung squamous cell carcinomas. *Cancer Cell* 23: 527–540
142. Huang WC, Ju TK, Hung MC, Chen CC (2007) Phosphorylation of CBP by IKKalpha promotes cell growth by switching the binding preference of CBP from p53 to NF-kappaB. *Mol Cell* 26: 75–87
143. Kawauchi K, Araki K, Tobiume K, Tanaka N (2008) p53 regulates glucose metabolism through an IKK-NF-kappaB pathway and inhibits cell transformation. *Nat Cell Biol* 10: 611–618

144. Kawauchi K, Araki K, Tobiume K, Tanaka N (2009) Loss of p53 enhances catalytic activity of IKKbeta through O-linked beta-N-acetyl glucosamine modification. *Proc Natl Acad Sci U S A* 106: 3431–3436
145. Xia Y, Padre RC, De Mendoza TH, Bottero V, Tergaonkar VB, Verma IM (2009) Phosphorylation of p53 by IkappaB kinase 2 promotes its degradation by beta-TrCP. *Proc Natl Acad Sci U S A* 106: 2629–2634
146. Lee DF, Kuo HP, Chen CT, Hsu JM, Chou CK, Wei Y, Sun HL, Li LY, Ping B, Huang WC, He X, Hung JY, Lai CC, Ding Q, Su JL, Yang JY, Sahin AA, Hortobagyi GN, Tsai FJ, Tsai CH, et al (2007) IKK beta suppression of TSC1 links inflammation and tumor angiogenesis via the mTOR pathway. *Cell* 130: 440–455
147. Hu MC, Lee DF, Xia W, Golfman LS, Ou-Yang F, Yang JY, Zou Y, Bao S, Hanada N, Saso H, Kobayashi R, Hung MC (2004) IkappaB kinase promotes tumorigenesis through inhibition of forkhead FOXO3a. *Cell* 117: 225–237
148. Buontempo F, Chiarini F, Bressanin D, Tabellini G, Melchionda F, Pesion A, Fini M, Neri LM, McCubrey JA, Martelli AM (2012) Activity of the selective IkappaB kinase inhibitor BMS-345541 against T-cell acute lymphoblastic leukemia: involvement of FOXO3a. *Cell Cycle* 11: 2467–2475
149. Chapuis N, Park S, Leotoing L, Tamburini J, Verdier F, Bardet V, Green AS, Willems L, Agou F, Ifrah N, Dreyfus F, Bismuth G, Baud V, Lacombe C, Mayeux P, Bouscary D (2010) IkappaB kinase overcomes PI3K/Akt and ERK/MAPK to control FOXO3a activity in acute myeloid leukemia. *Blood* 116: 4240–4250
150. Tilstra JS, Clauson CL, Niedernhofer LJ, Robbins PD (2011) NF-kappaB in Aging and Disease. *Aging Dis* 2: 449–465
151. Zhang G, Li J, Purkayastha S, Tang Y, Zhang H, Yin Y, Li B, Liu G, Cai D (2013) Hypothalamic programming of systemic ageing involving IKK-beta, NF-kappaB and GnRH. *Nature* 497: 211–216
152. Peng J, Schwartz D, Elias JE, Thoreen CC, Cheng D, Marsischky G, Roelofs J, Finley D, Gygi SP (2003) A proteomics approach to understanding protein ubiquitination. *Nat Biotechnol* 21: 921–926
153. Mevissen TE, Hospenthal MK, Geurink PP, Elliott PR, Akutsu M, Arnaudo N, Ekkebus R, Kulathu Y, Wauer T, El Oualid F, Freund SM, Ovaa H, Komander D (2013) OTU deubiquitinases reveal mechanisms of linkage specificity and enable ubiquitin chain restriction analysis. *Cell* 154: 169–184
154. Xu P, Duong DM, Seyfried NT, Cheng D, Xie Y, Robert J, Rush J, Hochstrasser M, Finley D, Peng J (2009) Quantitative proteomics reveals the function of unconventional ubiquitin chains in proteasomal degradation. *Cell* 137: 133–145
155. Park KJ, Krishnan V, O'Malley BW, Yamamoto Y, Gaynor RB (2005) Formation of an IKKalpha-dependent transcription complex is required for estrogen receptor-mediated gene activation. *Mol Cell* 18: 71–82
156. Balkhi MY, Fitzgerald KA, Pitha PM (2010) IKKalpha negatively regulates IRF-5 function in a MyD88-TRAF6 pathway. *Cell Signal* 22: 117–127
157. Hoshino K, Sugiyama T, Matsumoto M, Tanaka T, Saito M, Hemmi H, Ohara O, Akira S, Kaisho T (2006) IkappaB kinase-alpha is critical for interferon-alpha production induced by Toll-like receptors 7 and 9. *Nature* 440: 949–953
158. Liu B, Yang Y, Chernishof V, Loo RR, Jang H, Tahk S, Yang R, Mink S, Shultz D, Bellone CJ, Loo JA, Shuai K (2007) Proinflammatory stimuli induce IKKalpha-mediated phosphorylation of PIAS1 to restrict inflammation and immunity. *Cell* 129: 903–914
159. Anest V, Cogswell PC, Baldwin AS Jr (2004) IkappaB kinase alpha and p65/RelA contribute to optimal epidermal growth factor-induced c-fos gene expression independent of IkappaBalpha degradation. *J Biol Chem* 279: 31183–31189
160. Anest V, Hanson JL, Cogswell PC, Steinbrecher KA, Strahl BD, Baldwin AS (2003) A nucleosomal function for IkappaB kinase-alpha in NF-kappaB-dependent gene expression. *Nature* 423: 659–663
161. Yamamoto Y, Verma UN, Prajapati S, Kwak YT, Gaynor RB (2003) Histone H3 phosphorylation by IKK-alpha is critical for cytokine-induced gene expression. *Nature* 423: 655–659
162. Wu RC, Qin J, Hashimoto Y, Wong J, Xu J, Tsai SY, Tsai MJ, O'Malley BW (2002) Regulation of SRC-3 (pCIP/ACTR/AIB-1/RAC-3/TRAM-1) Coactivator activity by I kappa B kinase. *Mol Cell Biol* 22: 3549–3561
163. Beinke S, Robinson MJ, Hugunin M, Ley SC (2004) Lipopolysaccharide activation of the TPL-2/MEK/extracellular signal-regulated kinase mitogen-activated protein kinase cascade is regulated by IkappaB kinase-induced proteolysis of NF-kappaB1 p105. *Mol Cell Biol* 24: 9658–9667
164. Waterfield M, Jin W, Reiley W, Zhang M, Sun SC (2004) IkappaB kinase is an essential component of the Tpl2 signaling pathway. *Mol Cell Biol* 24: 6040–6048
165. Gringhuis SI, Garcia-Vallejo JJ, van Het Hof B, van Dijk W (2005) Convergent actions of I kappa B kinase beta and protein kinase C delta modulate mRNA stability through phosphorylation of 14-3-3 beta complexed with tristetraprolin. *Mol Cell Biol* 25: 6454–6463
166. Irelan JT, Murphy TJ, DeJesus PD, Teo H, Xu D, Gomez-Ferreria MA, Zhou Y, Miraglia LJ, Rines DR, Verma IM, Sharp DJ, Tergaonkar V, Chanda SK (2007) A role for IkappaB kinase 2 in bipolar spindle assembly. *Proc Natl Acad Sci U S A* 104: 16940–16945
167. Shinohara H, Maeda S, Watarai H, Kurosaki T (2007) IkappaB kinase beta-induced phosphorylation of CARMA1 contributes to CARMA1 Bcl10 MALT1 complex formation in B cells. *J Exp Med* 204: 3285–3293
168. Reiley W, Zhang M, Wu X, Granger E, Sun SC (2005) Regulation of the deubiquitinating enzyme CYLD by IkappaB kinase gamma-dependent phosphorylation. *Mol Cell Biol* 25: 3886–3895
169. Lee S, Andrieu C, Saltel F, Destaing O, Auclair J, Pouchkine V, Michelon J, Salaun B, Kobayashi R, Jurdic P, Kieff ED, Sylla BS (2004) IkappaB kinase beta phosphorylates Dok1 serines in response to TNF, IL-1, or gamma radiation. *Proc Natl Acad Sci U S A* 101: 17416–17421
170. Gao Z, Hwang D, Bataille F, Lefevre M, York D, Quon MJ, Ye J (2002) Serine phosphorylation of insulin receptor substrate 1 by inhibitor kappa B kinase complex. *J Biol Chem* 277: 48115–48121
171. Suzuki K, Verma IM (2008) Phosphorylation of SNAP-23 by IkappaB kinase 2 regulates mast cell degranulation. *Cell* 134: 485–495