CASE REPORT

Spontaneous passage of long, sharp gastrointestinal foreign body in a child

Vilvapathy Senguttuvan Karthikeyan,¹ Mohammed Gaffoor Ansari,²
Ramasamy Suresh,³ Bettaiyagowder Easwaran⁴

¹Department of Urology, Jawaharlal Institute of Postgraduate Medical Education and Research, Pondicherry, India

²Department of Radiodiagnosis, Sri Venkateshwarra Medical College Hospital and Research Centre, Pondicherry, India

³Department of Pediatrics, Sri Venkateshwarra Medical College Hospital and Research Centre, Pondicherry, India

⁴Department of General Surgery, Sri Venkateshwarra Medical College Hospital and Research Centre, Pondicherry, India

Correspondence to

Dr Vilvapathy Senguttuvan Karthikeyan, sengkarthik@yahoo.co.in

Accepted 30 December 2014

SUMMARY

Foreign body (FB) ingestion is a common problem in children. Up to 90% of these FBs pass spontaneously. FBs reaching the stomach usually pass out spontaneously. Exceptions to this spontaneous passage include a long FB that cannot cross the pylorus, duodenum or ileocaecal junction. We present a case of a 9-year boy who accidentally ingested a long paper pin, which spontaneously passed in 26 h. This case is being reported to highlight the successful spontaneous passage of a 4 cm long sharp foreign body in a child.

BACKGROUND

Ingestion of foreign bodies (FB) is a common problem in the paediatric population. Normally, 80–90% of these FBs pass spontaneously and 10–20% require endoscopic removal.¹ Surgical intervention is indicated in less than 1% of children.² FBs in the stomach usually pass out. Exceptions to spontaneous passage include a long FB that cannot cross the pylorus, duodenal loop or ileocaecal valve.³ This case is being reported to highlight successful spontaneous passage of a long sharp foreign body in a child.

CASE PRESENTATION

A 9-year-old boy presented to the emergency ward with history of having swallowed a paper pin at his school 10 h earlier. He was asymptomatic and clinical examination was unremarkable.

INVESTIGATIONS

Plain abdominal radiograph showed a long paper pin in the region of the fourth part of the duodenum/proximal jejunum (figure 1) and it was confirmed on fluoroscopy (figure 2). Plain chest radiograph showed no pneumoperitoneum or pneumomediastinum. Neck radiograph revealed no subcutaneous or prevertebral emphysema.

TREATMENT

As the boy did not show evidence of impaction or perforation and as the swallowed object was a sharp foreign body, he was admitted and serial radiographs were taken.

OUTCOME AND FOLLOW-UP

Twelve hours later, the pin had moved to the ileum on abdominal radiograph (figure 3) and fluoroscopy and 16 h later (26 h from ingestion) he passed the 4 cm long paper pin per rectum (figure 4). Post

Figure 1 Plain radiograph of the abdomen showing a long linear radiopaque foreign body in the abdomen, in the region of the fourth part of the duodenum/proximal jejunum (10 h after ingestion).

To cite: Karthikeyan VS, Ansari MG, Suresh R, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-206542

Figure 2 Fluoroscopy image confirming the position of the foreign body in the fourth part of the duodenum/proximal jejunum.

passage X-ray confirmed complete expulsion. He had no complications and was discharged the next morning.

DISCUSSION

FB ingestion is a common problem in the paediatric age group with 75% cases seen in children under 4 years of age.⁴ Most ingested FBs pass through the gastrointestinal tract (GIT) spontaneously⁵ and without complications. Impaction is most commonly seen at anatomic constrictions in the oesophagus⁶ and in over 75% patients, the commonest site is upper oesophagus at the cricopharyngeus.⁶ Beyond the gastro-oesophageal junction, FBs usually pass through the GIT without complications⁷ and impaction in the intestines is seen in less than 10% children.¹ FB impaction leads to mucosal abrasions, bleeding, GI

Figure 4 Photograph showing a 4 cm long paper pin passed spontaneously by the child (26 h after ingestion).

perforation, secondary mediastinitis and peritonitis,¹ and hence impaction is a strong indication for removal. The clinical spectrum of FBs in GIT can range from being asymptomatic to having GI symptoms such as vomiting, drooling, dysphagia, odynophagia, globus sensation or respiratory symptoms such as coughing, stridor and choking. In a review of paediatric foreign body ingestion, it was noted that spontaneous passage rates were described in just over a third of the studies¹ and that the spontaneous passage rate was higher in stomach and intestine FBs as compared with oesophageal FBs. If stomach and intestinal FBs are managed conservatively, patients can be safely discharged home. FBs impacted in the upper oesophagus often require medical intervention.^{1 8}

Children commonly ingest sharp objects such as fish and chicken bones, pins, needles, tooth picks, nails and even sharp toys. A single straight pin usually passes as it normally travels blunt end first as it traverses the GIT, following Jackson's axiom that, 'advancing points perforate and trailing points do not.'⁹ Removal of sharp FBs even if they have reached the stomach¹ is

Figure 3 Plain radiograph of the abdomen showing the foreign body in the region of distal ileum (12 h after ingestion).

advocated due to the sharp increase in complication rates from 1% to 15–35%.¹⁰ Ingested sharp FBs get impacted in the intestinal mucosa, which can erode into the peritoneal cavity or directly perforate the intestinal wall.¹ Sharp objects that are not amenable to endoscopic removal need to be followed up with daily radiographs, and surgical removal should be performed if the sharp FB fails to progress in 3 days.⁵

Sharp FBs reaching the stomach can usually pass spontaneously down the bowel. Exceptions to spontaneous passage of such sharp FBs are when the large FB cannot negotiate the pylorus, duodenal C-loop or ileocaecal valve.³ Normally, 80–90% of FBs pass spontaneously through the GIT but 10–20% will require endoscopic removal and less than 1% will need surgical intervention.²

Learning points

- ▶ In children, long sharp foreign bodies have the potential to lead to complications.
- ▶ Rapid passage through the gastrointestinal tract in an asymptomatic child is a case for watchful waiting.
- ▶ This approach avoids the potential morbidity of endoscopic or surgical intervention.

Contributors VSK and MGA conceptualised and designed the report, drafted the initial manuscript and approved the final manuscript as submitted. RS drafted the

initial manuscript and approved the final manuscript as submitted. BE performed the literature search, reviewed and revised the manuscript, and approved the final manuscript as submitted.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Jayachandra S, Eslick GD. A systematic review of paediatric foreign body ingestion: presentation, complications, and management. *Int J Pediatr Otorhinolaryngol* 2013;77:311–17.
- 2 Kay M, Wyllie R. Pediatric foreign bodies and their management. *Curr Gastroenterol Rep* 2005;7:212–18.
- 3 Jones A. Caustic ingestion and foreign bodies: damage to the upper gastrointestinal tract. In: Wyllie R, Hyams JS, eds. *Pediatric gastrointestinal and liver disease*. 3rd edn. Saunders, 2006:269.
- 4 McNeill MB, Sperry SL, Crockett SD, et al. Epidemiology and management of oesophageal coin impaction in children. *Dig Liver Dis* 2012;44:482–6.
- 5 Rodríguez H, Passali GC, Gregori D, et al. Management of foreign bodies in the airway and oesophagus. *Int J Pediatr Otorhinolaryngol* 2012;76(Suppl 1):S84–91.
- 6 Louie JP, Alpern ER, Windreich RM. Witnessed and unwitnessed esophageal foreign bodies in children. *Pediatr Emerg Care* 2005;21:582–5.
- 7 Hesham A-Kader H. Foreign body ingestion: children like to put objects in their mouth. *World J Pediatr* 2010;6:301–10.
- 8 Paul SP, Wilkinson R. Foreign body ingestion in children. *Nurs Times* 2012;108:25.
- 9 Spitz L. Management of ingested foreign bodies in childhood. *Br Med J* 1971;4:469–72.
- 10 Lee JH, Lee JS, Kim MJ, et al. Initial location determines spontaneous passage of foreign bodies from the gastrointestinal tract in children. *Pediatr Emerg Care* 2011;27:284–9.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>. BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow