Published in final edited form as:

Psychol Med. 2013 July ; 43(7): 1475–1486. doi:10.1017/S0033291712002383.

Prefrontal Dysfunction during Emotion Regulation in Generalized Anxiety and Panic Disorder

Tali Manber Ball, MA^{1,2}, Holly J. Ramsawh, PhD¹, Laura Campbell-Sills, PhD¹, Martin P. Paulus, MD^{1,3}, and Murray B. Stein, MD, MPH^{1,3,4}

¹Department of Psychiatry, University of California San Diego, La Jolla, CA, USA

²San Diego State University / University of California San Diego Joint Doctoral Program in Clinical Psychology, San Diego, CA, USA

³Psychiatry Service, Veterans Affairs San Diego Healthcare System, San Diego, CA, USA

⁴Department of Family and Preventive Medicine, University of California San Diego, La Jolla, CA, USA

Abstract

Background—The mechanisms that contribute to emotion dysregulation in anxiety disorders are not well understood. Two common disorders, Generalized Anxiety Disorder (GAD) and Panic Disorder (PD), were examined to test the hypothesis that both disorders are characterized by hypo-activation in prefrontal cortex (PFC) during emotion regulation. A competing hypothesis that GAD in particular is characterized by PFC *hyper*-activation during emotion regulation (reflecting overactive top-down control) also was evaluated.

Methods—Twenty-two medication-free Healthy Control (HC), 23 GAD, and 18 PD participants underwent functional magnetic resonance imaging (fMRI) during a task that required them to reappraise (i.e., reduce) or maintain emotional responses to negative images.

Results—GAD participants reported the least reappraisal use in daily life, and reappraisal use was inversely associated with anxiety severity and functional impairment in these participants. During fMRI, HC demonstrated greater activation during both reappraisal and maintenance than either GAD or PD (who did not differ) in brain areas important for emotion regulation (e.g., dorsolateral and dorsomedial PFC). Furthermore, across all anxious participants, activation during reappraisal in dorsolateral and dorsomedial PFC was inversely associated with anxiety severity and functional impairment.

Conclusions—Emotion dysregulation in GAD and PD may be the consequence of PFC hypo-activation during emotion regulation, consistent with insufficient top-down control. The relationship between PFC hypo-activation and functional impairment suggests that the failure to engage PFC during emotion regulation may be part of the critical transition from dispositionally high anxiety to an anxiety disorder.

Address correspondence to: Tali Manber Ball, MA, 8939 Villa La Jolla Dr, Suite 200, La Jolla, CA 92037, Tel: (858) 246-0628, Fax: (858) 534-6460, tmball@ucsd.edu.

Financial Disclosures

All other authors report no biomedical financial interests or potential conflicts of interest.

Introduction

Anxiety disorders are common (12 month prevalence 18%; Kessler *et al.*, 2005), and cause reduced quality of life (Mendlowicz & Stein, 2000; Lochner *et al.*, 2003; Barrera & Norton, 2009) and loss of productivity (DuPont *et al.*, 1996). Many investigations have sought to understand how pathological anxiety is maintained. Conceptual models have highlighted the role of anxious arousal and negative affectivity as predisposing factors (Clark & Watson, 1991), and avoidance as a key maintaining factor (Reddy *et al.*, 2006; Aupperle & Paulus, 2010). In addition, the role of emotion regulation is increasingly receiving attention (Mennin, 2006; Campbell-Sills & Barlow, 2007; Amstadter, 2008; Aldao *et al.*, 2010; Cisler *et al.*, 2010).

Emotion regulation has been defined as “cognitive and behavioral processes that influence the occurrence, intensity, duration, and expression of emotion” (Campbell-Sills & Barlow, 2007). One of the most studied forms of emotion regulation is reappraisal, or “changing how we think about a situation in order to decrease its emotional impact” (Gross, 2002). Reappraisal is associated with positive affect, interpersonal functioning, and well-being (Gross & John, 2003). Individuals with anxiety disorders are thought to have poorer emotion regulation skills, including reappraisal (Campbell-Sills & Barlow, 2007; Amstadter, 2008; Cisler *et al.*, 2010), however the evidence supporting this hypothesis is mixed (Decker *et al.*, 2008; Aldao *et al.*, 2010; Werner & Gross, 2010).

The neural substrates of reappraisal have been studied in both healthy adults and individuals with anxiety disorders. Studies of healthy adults have consistently found activation in dorsomedial, dorsolateral, and ventrolateral prefrontal cortex (PFC) during reappraisal (Ochsner & Gross, 2005; Lieberman, 2007). However, whether and how these substrates differ in individuals with anxiety disorders remains unclear. Studies of social anxiety disorder and posttraumatic stress disorder (PTSD) found less PFC activation in patients than healthy controls during reappraisal (Goldin *et al.*, 2009a; Goldin *et al.*, 2009b; New *et al.*, 2009). However, a study of young adults with high trait anxiety found the opposite: greater PFC activation during reappraisal in those with higher anxiety (Campbell-Sills *et al.*, 2011).

In the current study, our aim was to examine the common and distinct neural bases of emotion regulation in healthy and clinically anxious participants, while also comparing two common anxiety disorders, Generalized Anxiety Disorder (GAD) and Panic Disorder (PD). Emotion dysregulation has been hypothesized to play a key role in GAD: patients with GAD may not be as skilled at adaptively regulating negative emotions and may instead suppress emotions (Mennin *et al.*, 2002; Mennin *et al.*, 2005). Although to our knowledge the neural bases of reappraisal in GAD have not been examined, other neuroimaging studies of GAD have suggested a tendency to over-engage prefrontal top-down control (Mathew *et al.*, 2004; Etkin *et al.*, 2009).

The role of emotion dysregulation in PD has been even less well-studied than in GAD. One study found that reliance on suppression as an emotion regulation strategy maintains anxiety and avoidance in patients with PD (Levitt *et al.*, 2004). Furthermore, inflexible emotion regulation may moderate the relationship between anxiety sensitivity and PD (Cisler *et al.*,

2010). Biological models have pointed to insufficient top-down control as a factor in the generation of panic (Kent & Rauch, 2003). However, to our knowledge, there have been no neuroimaging studies of reappraisal in PD.

We examined two competing hypotheses. First, based on the conceptualization that GAD involves an overactive top-down control system, one hypothesis posits that individuals with GAD exhibit PFC hyper-activation during reappraisal. This would be consistent with findings in non-treatment-seeking, high trait anxiety participants, including many with GAD symptoms (Campbell-Sills *et al.*, 2011). A second hypothesis, based on the notion that emotion dysregulation and its neural bases are consistent across anxiety disorders, posits that both GAD and PD participants show *attenuated* prefrontal responding (reflecting inadequate top-down control) during reappraisal. This would be consistent with findings in other anxiety disorders.

Methods and Materials

Participants

This study was approved by the University of California San Diego Human Research Protections Program. After providing written informed consent, 139 participants underwent a semi-structured diagnostic interview (Sheehan *et al.*, 1998). Of these, 24 healthy control (HC) adults were eligible and didn't meet DSM-IV criteria (APA, 2000) for lifetime mood, anxiety, psychotic, or substance dependence disorders. In addition, 45 patients met DSM-IV criteria for clinically predominant PD ($n = 24$: 20 with and four without agoraphobia), or clinically predominant GAD ($n = 28$), and didn't meet criteria for lifetime psychosis, past-year substance dependence, or past-month substance abuse. Due to the high rate of comorbidity in anxiety disorders (Kessler *et al.*, 2005), comorbid mood and other anxiety disorders were allowed, however individuals with co-occurring PD and GAD ($n = 7$) were excluded. Two participants from each group were removed from the analysis due to poor fMRI data quality. The final dataset included 22 HC, 18 PD, and 23 GAD participants. In the GAD group, two participants had comorbid major depression, two had obsessive-compulsive disorder, and eight had social anxiety disorder. In the PD group, two participants had comorbid major depression, one had obsessive-compulsive disorder, and two had social anxiety disorder. All participants were free of psychotropic medications for six weeks prior to study enrollment (two weeks for benzodiazepines).

All participants met safety and eligibility criteria for fMRI scanning: no neurological conditions, no implanted ferrous metal, and no history of loss of consciousness greater than five minutes. Demographic characteristics of the 63 participants are presented in Table 1. There were group differences on age ($p < .07$) and gender ($p < .05$), therefore all group effects analyses were also run with age and gender as covariates.

Procedure

Eligible participants underwent a medical examination, including medical history, laboratory evaluation, EKG, and drug and pregnancy screen. Participants also completed questionnaires including the Overall Anxiety Severity and Impairment Scale (OASIS; Norman *et al.*, 2006),

the Penn State Worry Questionnaire (PSWQ; Meyer *et al.*, 1990), the Anxiety Sensitivity Index (ASI; Reiss *et al.*, 1986), and the Quick Inventory of Depressive Symptomatology (QIDS; Rush *et al.*, 2003). Fifty participants (78%) participated in an optional adjunct study, and completed the Emotion Regulation Questionnaire (ERQ; Gross & John, 2003). All participants completed one fMRI scan before receiving 10 sessions of weekly individual cognitive behavioral therapy. Relationship of fMRI findings and therapy outcomes will be considered in a separate paper.

Task

Two processes were examined with the emotion regulation task: in each trial, individuals either maintained or reappraised their emotional responses to negative images. Each trial was 24 seconds long, with a scrambled image presented for the first 12 seconds and a negative image for the last 12 seconds (Supplemental Figure 1). After the first 1 – 3 seconds (jittered) of scrambled image presentation the words “Rate Emotion (1–4)” appeared for 3 seconds, cuing participants to provide a baseline rating of their emotional experience from one (not at all negative) to four (very negative). 1 – 3 seconds later, participants received an instruction to either “Keep Up Emotion” (Maintain) or “Reduce Emotion” (Reappraise) to the negative image they were about to see. The instruction remained on the screen for 3 seconds, followed by continued presentation of the scrambled image (1 – 3 seconds). Participants then viewed the target negative image for 4 – 6 seconds, implementing the instruction they had just received. They then received the cue to rate their negative emotion (3 seconds), followed by continued presentation of the negative image (3 – 5 seconds). The period of interest was the 4 – 6 seconds when the participant was maintaining or reappraising their emotional reactions.

The images used in the task were twelve negatively valenced images for each condition, selected from the International Affective Picture System (Lang *et al.*, 2008), with a pixel-wise scrambled version of each image presented as a baseline at the start of each trial. More information about the images and the task are given in (Campbell-Sills *et al.*, 2011), though there is no overlap in subjects between those reported here and in (Campbell-Sills *et al.*, 2011).

Prior to fMRI scanning, participants were trained on the task. For the Maintain condition participants were instructed to “maintain your emotional reaction until the picture disappears.” For the Reappraise condition, participants were asked to “change the way that you think about the picture in order to decrease your negative emotions.” Suggested reappraisal strategies included generating a positive interpretation, or taking a more detached perspective. Participants practiced out loud on sample images, to confirm understanding.

Image Acquisition

One 9 minute 40 second BOLD fMRI run was acquired, using a Signa EXCITE 3.0 Tesla-GE scanner (T2*-weighted echo planar imaging, TR = 2000 ms, TE = 32 ms, FOV = 240 × 240 mm³, 64 × 64 matrix, thirty 2.6 mm axial slices with a 1.4 mm gap, 290 scans). For anatomical reference, a high resolution T1-weighted image (SPGR, TI = 450, TR = 8 ms,

TE = 3 ms, FOV = $250 \times 250 \text{ mm}^3$, flip angle = 12° , 172 sagittally acquired slices with 1 mm thickness) was obtained during the same session.

Image Processing

All structural and functional image processing was done with the Analysis of Functional NeuroImages (AFNI) software package (Cox, 1996). The AFNI function *3dToutcount* was used to identify the number of voxels classified as outliers at each time point: any time point with greater than two standard deviations more outlier voxels than the subject's mean were excluded from analysis. Voxel time series were interpolated to correct for non-simultaneous slice acquisition and corrected for three-dimensional motion. Anatomical and echo planar volumes were co-registered using an algorithm that minimizes the amount of image translation and rotation (Saad *et al.*, 2009).

Individual participant time series data were analyzed with AFNI's *3dDeconvolve* program. The orthogonal regressors of interest were those indicating the Maintain, Reappraise, and Baseline conditions (see Supplemental Figure 1). Additional regressors of non-interest modeled the emotion rating periods and post-rating viewing period at the end of each trial, as well as linear and quadratic trends in the time series and motion regressors based on the time series alignments in the roll, pitch, and yaw directions. Regressors were convolved with a modified gamma variate function to account for the hemodynamic response (Boynton *et al.*, 1996) using the AFNI program *waver*. Following deconvolution, data were converted to percent signal change, by dividing the coefficient by the zero-order regressor within each voxel. Data were normalized to Talairach coordinates (Talairach & Tournoux, 1988) and subjected to a 4mm Gaussian blur for spatial smoothing.

To protect for multiple comparisons, Monte Carlo simulations ($n=10000$) using AFNI's *3dClustSim* program demonstrated that minimum cluster volumes of 640 μL for a per-voxel threshold of $p < .01$ (one-sided) for whole brain analysis with 4 mm spatial smoothing would result in a cluster-wise $p < .05$. Therefore, only clusters of 640 μL or greater are reported.

Statistical Analysis

Analyses of self-report and behavioral data were performed with PASW Statistics (Version 18.0.0, Chicago IL). Single subject analysis of functional images was performed using the AFNI software package (Cox, 1996), and group level statistical analyses were completed using the R statistical package (<http://cran.r-project.org>). Group analysis was conducted at each voxel using the R library *fmri* to read in and write out full AFNI data files for voxel-wise analysis. Specifically, using the R library *nlme*, a linear mixed effects model was computed with group (HC, PD, GAD) as a fixed factor and condition (Reappraise, Maintain; each contrasted with Baseline) as a grouped factor within subjects. Subjects were modeled as random factors.

First, to determine whether there were condition-specific group differences, we examined the group by condition interaction. Next, because examining the main effect of group by combining both conditions would be difficult to interpret, we examined the main effect of

group in each condition separately, and also identified regions of overlap that showed significant group effects in both conditions. These analyses were subjected to a volume-adjusted threshold such that each voxel met a statistical threshold ($p < .01$, one-sided) and was part of a cluster of 640 μL , corresponding to a whole-brain adjusted $p < .05$. All group effects were also examined with age and gender as covariates. Finally, planned contrasts between HC and anxious participants, and between GAD and PD participants, were obtained from cluster averages to describe the characteristics of the group and interaction effects. Cohen's d was computed using pooled standard deviation.

The main effect of task (Reappraise versus Maintain) across all participants was then examined, however using the statistical and volume thresholds described above yielded clusters so large they were difficult to interpret and report. Therefore a per-voxel threshold of $p < .001$ with a minimum cluster size of 256 μL was used. Monte Carlo simulations ($n=10000$) using AFNI's *3dClustSim* program demonstrated that this statistical and volume combination also yields a whole-brain cluster-wise $p < .05$.

Results

Self-Report Measures

Table 1 summarizes the self-report questionnaires. One-way ANOVA analyses revealed a significant main effect of group on overall anxiety severity and impairment, worry, anxiety sensitivity, and depression ($p < .001$). Follow-up t -tests comparing patients and controls showed higher scores in patients on all measures ($p < .001$). Follow-up t -tests comparing GAD and PD revealed differences in anxiety sensitivity (PD higher, $p < .05$) and overall anxiety severity and impairment (GAD higher, $p < .05$) only.

The groups also differed on emotion regulation use in daily life (reappraisal $p < .005$; suppression $p = .05$). Follow-up t -tests showed greater use of suppression in patients ($p < .05$) with no difference between GAD and PD. Patients and controls did not differ on reappraisal use, but GAD reported less reappraisal use than PD ($p < .005$). Reappraisal use was inversely associated with overall anxiety severity and impairment in GAD ($r = -0.61$, $p < .05$) but not PD ($r = 0.14$, ns). When controlling for age, gender, and depression severity, all self-report differences remained significant ($p < .05$) except group differences in suppression use ($p = .08$).

Behavioral Task Effects

Figure 1A illustrates the results of a 3 (group: HC, PD, GAD) by 3 (condition: Baseline, Maintain, Reappraise) repeated measures ANOVA on the negative emotion ratings collected during the task. Emotion ratings differed by condition, with most negative emotion during the Maintain condition, and least during Baseline ($F(2,110) = 94.0$, $p < .001$). Paired t -tests confirmed that emotion ratings significantly differed across all conditions in each of the three groups ($p < .05$). Ratings also differed by group, with HC reporting less intense emotions than either PD or GAD ($F(2,55) = 7.3$, $p < .005$). There was also a group by condition interaction ($F(4,110) = 2.4$, $p = .05$): HC reported significantly less negative emotion than GAD or PD during Baseline and Reappraisal, and less negative emotion than

GAD during the Maintain condition ($p < .05$). Emotion ratings in the GAD and PD groups did not differ from each other in any condition ($p > .09$).

In order to control for differences in negative emotion at baseline, differential emotion ratings (i.e., Reappraise minus Baseline and Maintain minus Baseline) were also examined. No group differences were found for either condition ($p > .1$). All behavioral results were unchanged when controlling for age and gender.

fMRI Task Effects

Across all participants, greater activation to Maintain than Reappraise was observed in many brain areas, including medial supplemental motor area, bilateral inferior parietal lobule, posterior cingulate, bilateral posterior insula, and bilateral middle temporal gyrus. Greater activation to Reappraise than Maintain was observed across all participants in bilateral middle occipital gyrus, right fusiform gyrus, and right precuneus (Supplemental Table 1).

fMRI Group by Task Interaction Effects

Voxel-wise whole-brain analysis of group (HC, GAD, PD) by task (Reappraise, Maintain) interactions revealed a cluster in the left post-central gyrus extending to the inferior parietal lobule. The interaction was due to significantly greater activation in HC than either GAD or PD in the Maintain condition, but no group differences during the Reappraise condition (Table 2).

fMRI Group Effects – Maintain

During the Maintain condition, significant group differences were observed in several prefrontal regions: dorsomedial PFC, bilateral dorsolateral and ventrolateral PFC, and dorsal anterior cingulate. These were all characterized by significantly greater activation in HC than anxious participants ($d = 1.05 - 1.77$), with only right ventrolateral PFC and dorsal anterior cingulate demonstrating significant differences between the two anxiety groups (PD > GAD, $d = 0.90$ and 0.64 respectively). Of these prefrontal regions, all but right ventrolateral PFC and dorsal anterior cingulate continued to show significant group effects when controlling for age and gender.

HC also demonstrated significantly greater activation than anxiety patients in other regions, including left anterior insula, bilateral caudate and thalamus, left superior parietal lobule, left fusiform gyrus, bilateral middle and superior temporal gyrus, posterior cingulate, and left middle occipital gyrus ($d = 0.61 - 1.45$). GAD and PD differed in right middle and superior temporal gyrus and left superior parietal lobule (PD > GAD, $d = 1.01 - 1.14$), as well as left middle occipital gyrus (GAD > PD, $d = 0.83$). All regions except posterior cingulate continued to show significant group effects when controlling for age and gender (Table 3A, Figures 1 and 2).

fMRI Group Effects – Reappraisal

During reappraisal, HCs demonstrated significantly greater activation than anxiety patients in several regions, including dorsomedial and bilateral dorsolateral PFC, left fusiform gyrus and right middle occipital gyrus ($d = 0.82 - 1.37$). GAD and PD differed in right middle

occipital gyrus only (PD > GAD, $d = 0.77$). Dorsomedial PFC, left dorsolateral PFC, and left fusiform gyrus had at least four voxels of overlap with regions that also showed significant group effects during the Maintain condition. When controlling for age and gender, only bilateral dorsolateral PFC continued to show significant group effects (Table 3B, Figures 1 and 2).

Brain-Behavior Relationships

Relationships between self-report measures and average brain activation were examined for the two prefrontal regions (dorsomedial and left dorsolateral PFC) that differed between anxious and control participants during both reappraisal and maintenance. Because activation in these regions did not differ between GAD and PD, correlations were conducted on the combined group of anxious participants.

Among anxious participants, those with greater anxiety severity and associated impairment showed relatively less activation in PFC during both reappraisal and maintenance. Specifically, individuals reporting greater anxiety severity and functional impairment had less activation in left dorsolateral PFC during reappraisal ($r = -.43$, $p = .005$) and in dorsomedial PFC during both reappraisal ($r = -.38$, $p < .05$) and maintenance ($r = -.33$, $p < .05$) (Figures 1 and 2). The relationships between activation during reappraisal and anxiety severity remained significant when controlling for age and gender, and marginally significant ($p < .07$) when controlling for depression severity (in dorsolateral PFC; dorsomedial PFC, *ns*).

Similarly, across anxious participants, those who worried more had less activation in dorsomedial PFC during negative emotion maintenance ($r = -.34$, $p < .05$) (Figure 2). This relationship remained significant when controlling for age, gender, and depression severity. There were no associations between PFC activation and worry severity during reappraisal, or between anxiety sensitivity, regular reappraisal use, or negative emotion ratings and PFC activation during either condition.

Discussion

This study examined two alternative hypotheses regarding the relationship of PFC activation and emotion regulation in PD and GAD. First, based on the conceptualization that GAD involves an overactive top-down control system, we hypothesized that GAD would exhibit *hyper*-activation in PFC during regulation. Second, based on the emotion regulation deficits common across anxiety disorders, a competing hypothesis was that both GAD and PD involve inadequate top-down control and therefore would show *attenuated* prefrontal responding during emotion regulation. The main neuroimaging finding showed that HC participants activated dorsomedial and bilateral dorsolateral PFC more during reappraisal than those with GAD or PD. Therefore, the results of this study are consistent with the second hypothesis and support potential cross-disorder impairment in regulation of negative emotions.

Behaviorally, the groups did not differ in self-reported negative emotion during regulation when adjusting for baseline negative emotion ratings. This is consistent with previous

reports that anxious individuals can successfully regulate emotions when explicitly trained and cued to do so (Goldin *et al.*, 2009b; Campbell-Sills *et al.*, 2011; Aldao & Mennin, 2012). Furthermore, the lack of difference in emotion ratings aids in interpretation of the fMRI results, as they cannot be readily explained by differential changes in internal emotional experience.

Unlike previous investigations (Campbell-Sills *et al.*, 2011), no task effect or group by task interactions were found in PFC: group differences in prefrontal activation were present in both conditions. Both reappraisal and maintenance are active processes, and both involve conscious attempts to alter ongoing emotional experience. The use of such an active comparison condition may have contributed to the lack of differential effect in PFC, leading to the hypothesis that PFC recruitment may be broadly impaired in GAD and PD. For example, hypo-activations in PFC during a task requiring attentional control has been associated with trait anxiety (Bishop, 2009) consistent with concentration difficulties common in anxiety disorders (APA, 2000). Therefore, a decreased ability to recruit PFC may cut across multiple contexts relevant for anxiety disorders.

One region that did show a group by task interaction was the left inferior parietal lobule, where HC but not anxious individuals showed greater activation during maintenance than reappraisal. The inferior parietal lobule together with the dorsolateral PFC is part of the central executive network, which is critical for top-down control during decision making, goal implementation, and other executive functions (Menon, 2011). The attenuated activation of this system during emotion regulation is consistent with the hypothesis that anxious participants exert insufficient goal-directed top-down control over brain systems that process emotional information.

In a previous study using this task (Campbell-Sills *et al.*, 2011) PFC activation was greater in anxious participants during reappraisal, whereas here dorsolateral and dorsomedial PFC activation was greatest in HC during both reappraisal and maintenance. However, the previous study used a non-clinical (i.e., not treatment-seeking) sample. Studies of treatment-seeking adults with social anxiety disorder have found less PFC activation during reappraisal in anxious compared to healthy individuals (Goldin *et al.*, 2009a; Goldin *et al.*, 2009b), in line with the present results. This suggests that heightened PFC activation in anxious, non-clinical adults could be a compensatory mechanism in individuals with dispositionally high anxiety. This notion is consistent with previous investigations suggesting that dorsolateral PFC activation may serve a compensatory mechanism in less severe patients with GAD (Etkin *et al.*, 2009) or major depression (Etkin & Schatzberg, 2011).

In contrast, anxiety associated with clinically significant impairment (i.e., disordered anxiety) could be associated with an inability to sufficiently engage PFC in the service of emotion regulation. We have previously suggested that inadequate top-down control in anxiety disorders could be the consequence of amplified bottom-up signals that make identifying salience more difficult (Paulus & Stein, 2010). In other words, chronic over-responsiveness of limbic circuitry in anxiety disorders (Etkin & Wager, 2007) may fatigue the top-down system, rendering it unable to effectively exert control when needed. Such

emotion regulation deficits could perpetuate patients' belief that negative emotions are aversive and uncontrollable, continuing the cycle of amplified limbic inputs and fatigued top-down control and maintaining pathological anxiety. The inverse correlation between functional impairment and prefrontal activation supports the hypothesis that PFC hypo-activation could be a component of the transition from dispositionally high anxiety to disordered anxiety.

Although differences in neural activation between anxious and healthy participants were pronounced, there were few differences between GAD and PD. This supports the notion that decreased PFC engagement during emotion regulation is a common feature of the two disorders. However, activation in dorsomedial PFC during maintenance was inversely associated with worry severity. Worry, often conceptualized as an internal avoidance strategy, is a hallmark feature of GAD (Borkovec & Inz, 1990). Individuals with more severe worry may be less comfortable with affective arousal and more likely to avoid emotional stimuli, thereby showing diminished PFC activation during maintenance of negative emotions. On self-report measures, patients with GAD reported the least use of reappraisal in their daily lives. Furthermore, reappraisal use was associated with less anxiety severity and functional impairment in GAD, but not PD. These results support the theory (Mennin *et al.*, 2002) that emotion dysregulation is especially important in understanding GAD.

In conclusion, patients with GAD and PD demonstrated less PFC activation than healthy controls during cognitive modulation of emotion, and those with the least PFC activation reported the greatest anxiety severity and impairment. When considered in conjunction with similar results in social anxiety disorder and PTSD (Goldin *et al.*, 2009a; Goldin *et al.*, 2009b; New *et al.*, 2009), these results suggest a potential common neural basis of emotion dysregulation in anxiety disorders. Furthermore, these results highlight that emotion dysregulation may be related to functional impairment and treatment-seeking rather than a specific type of anxiety symptoms. However, given that patients with GAD reported least regular use of reappraisal, as well as the inverse association between worry and prefrontal activation, emotion dysregulation may be especially important in the etiology and maintenance of GAD (Mennin *et al.*, 2002).

This study has several limitations. One limitation is the lack of a passive viewing condition, which may have contributed to the lack of task effect in PFC. Although a previous study using the same task (Campbell-Sills *et al.*, 2011) did find such an effect, it was not replicated in this sample. The results should therefore be interpreted with caution. The absence of a passive viewing condition may also have contributed to the lack of group by task interaction in PFC. Without such an interaction, the alternative hypothesis that top-down control is broadly impaired in GAD and PD cannot be ruled out. Future studies should utilize a variety of comparison conditions to more clearly differentiate the processes that are impaired in anxiety disorders.

Another limitation is that patients with GAD and PD differed on overall anxiety severity and impairment (greater in GAD) and habitual use of reappraisal (less frequent in GAD). In addition, the measure of habitual emotion regulation was only available for some

participants, decreasing the power to detect effects related to this variable. However, there were few differences in neural activation between GAD and PD. Therefore it is unlikely that group differences on anxiety severity or emotion regulation use account for the results. As is common in anxiety disorders (Kessler *et al.*, 2005), 10% of patients in this sample had comorbid major depression, and an additional 24% reported past depressive episodes. However, associations between neural activation and anxiety changed only marginally when controlling for depressive symptoms, suggesting that depression severity cannot explain the findings.

The present study included only individuals with GAD or PD, but the results support the hypothesis that emotion dysregulation may be consistent across anxiety diagnoses. Future work should examine the neural bases of emotion regulation in other disorders. For example, emotion dysregulation in PTSD has been frequently noted (Roemer *et al.*, 2001; Eftekhari *et al.*, 2009; New *et al.*, 2009; Bonn-Miller *et al.*, 2011), suggesting that a better understanding of the neural bases of reappraisal in PTSD could be especially fruitful.

Lastly, successful emotion regulation can be accomplished in many ways, though here we only examined reappraisal and maintenance. Future work should examine subtypes of reappraisal (e.g., detachment, focusing on positive aspects) along with other emotion regulation strategies to understand how strategy selection impacts regulation success. A better grasp of the processes employed is important in moving past simply examining differences in activation location and intensity, and towards a more nuanced and mechanistic understanding.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Supported by NIMH Grants MH65413 and MH64122 to Dr. Stein. The authors are grateful to Sarah Sullivan and Taru Flagan for assistance with subject scanning and data management, to Shadha Cissell for assistance with subject evaluation, and to Ariel Lang for clinical supervision.

Dr. Stein reports that he is paid for his editorial work at UpToDate, Inc. and Depression and Anxiety (Wiley).

References

- Aldao A, Mennin D. Paradoxical cardiovascular effects of implementing adaptive emotion regulation strategies in generalized anxiety disorder. *Behaviour Research and Therapy*. 2012; 50:122–130. [PubMed: 22218164]
- Aldao A, Nolen-Hoeksema S, Schweizer S. Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clinical Psychology Review*. 2010; 30:217–237. [PubMed: 20015584]
- Amstadter A. Emotion regulation and anxiety disorders. *Journal of Anxiety Disorders*. 2008; 22:211–221. [PubMed: 17349775]
- APA. *Diagnostic and Statistical Manual of Mental Disorders DSM-IV-TR*. Washington DC: American Psychiatric Press; 2000.
- Aupperle RL, Paulus MP. Neural systems underlying approach and avoidance in anxiety disorders. *Dialogues in Clinical Neuroscience*. 2010; 12:305–319. [PubMed: 20954427]

- Barrera TL, Norton PJ. Quality of life impairment in generalized anxiety disorder, social phobia, and panic disorder. *Journal of Anxiety Disorders*. 2009; 23:1086–1090. [PubMed: 19640675]
- Bishop S. Trait anxiety and impoverished prefrontal control of attention. *Nature Neuroscience*. 2009; 12:92–98.
- Bonn-Miller MO, Vujanovic AA, Boden MT, Gross JJ. Posttraumatic stress, difficulties in emotion regulation, and coping-oriented marijuana use. *Cognitive Behaviour Therapy*. 2011; 40:34–44. [PubMed: 21337213]
- Borkovec TD, Inz J. The nature of worry in generalized anxiety disorder: A predominance of thought activity. *Behaviour Research and Therapy*. 1990; 28:153–158. [PubMed: 2183759]
- Boynton GM, Engel SA, Glover GH, Heeger DJ. Linear systems analysis of functional magnetic resonance imaging in human V1. *The Journal of Neuroscience*. 1996; 16:4207–4221. [PubMed: 8753882]
- Campbell-Sills, L.; Barlow, DH. Incorporating emotion regulation into conceptualizations and treatments of anxiety and mood disorders. In: Gross, JJ., editor. *Handbook of Emotion Regulation*. New York: Guilford Press; 2007. p. 542-559.
- Campbell-Sills L, Simmons AN, Lovero KL, Rochlin AA, Paulus MP, Stein MB. Functioning of neural systems supporting emotion regulation in anxiety-prone individuals. *NeuroImage*. 2011; 54:689–696. [PubMed: 20673804]
- Cisler JM, Olatunji BO, Feldner MT, Forsyth JP. Emotion regulation and the anxiety disorders: An integrative review. *Journal of Psychopathology and Behavioral Assessment*. 2010; 32:68–82. [PubMed: 20622981]
- Clark LA, Watson D. Tripartite model of anxiety and depression: Psychometric evidence and taxonomic implications. *Journal of Abnormal Psychology*. 1991; 100:316–336. [PubMed: 1918611]
- Cox R. AFNI: Software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical Research*. 1996; 29:162–173. [PubMed: 8812068]
- Decker ML, Turk CL, Hess B, Murray CE. Emotion regulation among individuals classified with and without generalized anxiety disorder. *Journal of Anxiety Disorders*. 2008; 22:485–494. [PubMed: 17512697]
- DuPont RL, Rice DP, Miller LS, Shiraki SS, Rowland CR, Harwood HJ. Economic costs of anxiety disorders. *Anxiety*. 1996; 2:167–172. [PubMed: 9160618]
- Eftekhari A, Zoellner LA, Vigil SA. Patterns of emotion regulation and psychopathology. *Anxiety, Stress & Coping*. 2009; 22:571–586.
- Etkin A, Prater KE, Schatzberg AF, Menon V, Greicius MD. Disrupted amygdalar subregion functional connectivity and evidence of a compensatory network in generalized anxiety disorder. *Archives of General Psychiatry*. 2009; 66:1361–1372. [PubMed: 19996041]
- Etkin A, Schatzberg AF. Common abnormalities and disorder-specific compensation during implicit regulation of emotional processing in generalized anxiety and major depressive disorders. *The American Journal of Psychiatry*. 2011; 168:968–978. [PubMed: 21632648]
- Etkin A, Wager TD. Functional neuroimaging of anxiety: A meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *The American Journal of Psychiatry*. 2007; 164:1476–1488. [PubMed: 17898336]
- Goldin PR, Manber-Ball T, Werner K, Heimberg R, Gross JJ. Neural mechanisms of cognitive reappraisal of negative self-beliefs in social anxiety disorder. *Biological Psychiatry*. 2009a; 66:1091–1099. [PubMed: 19717138]
- Goldin PR, Manber T, Hakimi S, Canli T, Gross JJ. Neural bases of social anxiety disorder: Emotional reactivity and cognitive regulation during social and physical threat. *Archives of General Psychiatry*. 2009b; 66:170–180. [PubMed: 19188539]
- Gross JJ. Emotion regulation: Affective, cognitive, and social consequences. *Psychophysiology*. 2002; 39:281–291. [PubMed: 12212647]
- Gross JJ, John OP. Individual differences in two emotion regulation processes: Implications for affect, relationships, and well-being. *Journal of Personality and Social Psychology*. 2003; 85:348–362. [PubMed: 12916575]

- Kent JM, Rauch SL. Neurocircuitry of anxiety disorders. *Current Psychiatry Reports*. 2003; 5:266–273. [PubMed: 12857529]
- Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*. 2005; 62:617–627. [PubMed: 15939839]
- Lang, PJ.; Bradley, MM.; Cuthbert, BN. International Affective Picture System (IAPS): Affective Ratings of Pictures and Instruction Manual, Technical Report A-8. Gainesville, FL: University of Florida; 2008.
- Levitt J, Brown T, Orsillo S, Barlow D. The effects of acceptance versus suppression of emotion on subjective and psychophysiological response to carbon dioxide challenge in patients with panic disorder. *Behavior Therapy*. 2004; 35:747–766.
- Lieberman MD. Social cognitive neuroscience: A review of core processes. *Annual Review of Psychology*. 2007; 58:259–289.
- Lochner C, Mogotsi M, du Toit PL, Kaminer D, Niehaus DJ, Stein DJ. Quality of life in anxiety disorders: A comparison of obsessive-compulsive disorder, social anxiety disorder, and panic disorder. *Psychopathology*. 2003; 36:255–262. [PubMed: 14571055]
- Mathew SJ, Mao X, Coplan JD, Smith ELP, Sackeim HA, Gorman JM, Shungu DC. Dorsolateral prefrontal cortical pathology in generalized anxiety disorder: A proton magnetic resonance spectroscopic imaging study. *American Journal of Psychiatry*. 2004; 161:1119–1121. [PubMed: 15169704]
- Mendlowicz MV, Stein MB. Quality of life in individuals with anxiety disorders. *American Journal of Psychiatry*. 2000; 157:669–682. [PubMed: 10784456]
- Mennin DS. Emotion Regulation Therapy: An integrative approach to treatment-resistant anxiety disorders. *Journal of Contemporary Psychotherapy*. 2006; 36:95–105.
- Mennin DS, Heimberg RG, Turk CL, Fresco DM. Applying an emotion regulation framework to integrative approaches to generalized anxiety disorder. *Clinical Psychology: Science and Practice*. 2002; 9:85–90.
- Mennin DS, Heimberg RG, Turk CL, Fresco DM. Preliminary evidence for an emotion dysregulation model of generalized anxiety disorder. *Behaviour Research and Therapy*. 2005; 43:1281–1310. [PubMed: 16086981]
- Menon V. Large-scale brain networks and psychopathology: A unifying triple network model. *Trends in Cognitive Sciences*. 2011; 15:506–483.
- Meyer TJ, Miller ML, Metzger RL, Borkovec TD. Development and validation of the Penn State Worry Questionnaire. *Behaviour Research and Therapy*. 1990; 28:487–495. [PubMed: 2076086]
- New AS, Fan J, Murrough JW, Liu X, Liebman RE, Guise KG, Tang CY, Charney DS. A functional magnetic resonance imaging study of deliberate emotion regulation in resilience and posttraumatic stress disorder. *Biological Psychiatry*. 2009; 66:656–664. [PubMed: 19589502]
- Norman SB, Cissell SH, Means-Christensen AJ, Stein MB. Development and validation of an Overall Anxiety Severity And Impairment Scale (OASIS). *Depression and Anxiety*. 2006; 23:245–249. [PubMed: 16688739]
- Ochsner KN, Gross JJ. The cognitive control of emotion. *Trends in Cognitive Sciences*. 2005; 9:242–249. [PubMed: 15866151]
- Paulus MP, Stein MB. Interoception in anxiety and depression. *Brain Structure and Function*. 2010; 214:451–463. [PubMed: 20490545]
- Reddy MK, Pickett SM, Orcutt HK. Experiential avoidance as a mediator in the relationship between childhood psychological abuse and current mental health symptoms in college students. *Journal of Emotional Abuse*. 2006; 6:67–85.
- Reiss S, Peterson RA, Gursky DM, McNally RJ. Anxiety sensitivity, anxiety frequency and the prediction of fearfulness. *Behaviour Research and Therapy*. 1986; 24:1–8. [PubMed: 3947307]
- Roemer L, Litz BT, Orsillo SM, Wagner AW. A preliminary investigation of the role of strategic withholding of emotions in PTSD. *Journal of Traumatic Stress*. 2001; 14:149–156.
- Rush AJ, Trivedi MH, Ibrahim HM, Carmody TJ, Arnow B, Klein DN, Markowitz JC, Ninan PT, Kornstein S, Manber R, Thase ME, Kocsis JH, Keller MB. The 16-item Quick Inventory of Depressive Symptomatology (QIDS), clinician rating (QIDS-C), and self-report (QIDS-SR): a

- psychometric evaluation in patients with chronic major depression. *Biological Psychiatry*. 2003; 54:573–583. [PubMed: 12946886]
- Saad ZS, Glen DR, Chen G, Beauchamp MS, Desai R, Cox RW. A new method for improving functional-to-structural MRI alignment using local Pearson correlation. *NeuroImage*. 2009; 44:839–848. [PubMed: 18976717]
- Sheehan D, Lecrubier Y, Sheehan K, Amorim P, Janavs J, Weiller E, Hergueta T, Baker R, Dunbar GC. The Mini-International Neuropsychiatric Interview (M.I.N.I.): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *The Journal of Clinical Psychiatry*. 1998; 59:22–33. [PubMed: 9881538]
- Talairach, J.; Tournoux, P. Co-planar stereotaxic atlas of the human brain: 3-dimensional proportional system: an approach to cerebral imaging. Thieme; 1988.
- Werner, K.; Gross, JJ. Emotion regulation and psychopathology: A conceptual framework. In: Kring, AM.; Sloan, DM., editors. *Emotion Regulation and Psychopathology: A Transdiagnostic Approach to Etiology and Treatment*. New York: Guilford Press; 2010. p. 13-37.

Figure 1.

(A) Negative emotion ratings during the task. HC = healthy control; GAD = Generalized Anxiety Disorder; PD = Panic Disorder. * indicates significant group differences within a condition, $p < .05$. (B and C) Attenuated left dorsolateral PFC activation during reappraisal. GAD and PD is correlated with anxiety severity. Activation extracted from the region showing activation in both the Reappraise and Maintain conditions. (B) Left dorsolateral PFC activation during reappraisal and maintenance (* $p < .05$; ** $p < .005$). (C) Left dorsolateral PFC activation during reappraisal associated with anxiety severity and impairment in patients with anxiety disorders. HC = healthy control; GAD = generalized anxiety disorder; PD = panic disorder

Figure 2.

Attenuated dorsomedial PFC activation in GAD and PD is correlated with anxiety severity during reappraisal and correlated with worry severity during maintenance. Activation extracted from the region showing activation in both the Reappraise and Maintain conditions. (A) Dorsomedial PFC activation during each condition (* $p < .05$; ** $p < .005$). (B) Dorsomedial PFC activation during reappraisal associated with anxiety severity and impairment in patients with anxiety disorders. (C) Dorsomedial PFC activation during maintenance associated with worry severity in patients with anxiety disorders. HC = healthy control; GAD = generalized anxiety disorder; PD = panic disorder

Table 1

Demographics and clinical features of participants

	HC	GAD	PD	ANX vs HC	GAD vs PD
N	22	23	18		
Age (Years, SD)	27 (9)	35 (11)	29 (7)	$t(61) = 2.0^{\dagger}$	$t(39) = 2.0^{\dagger}$
Gender (% Female)	50%	74%	83%	$\chi^2 = 5.2^*$	$\chi^2 = 0.5$
Education (Years, SD)	15 (2)	16 (2)	14 (2)	$t(61) = -0.6$	$t(39) = 1.9^{\dagger}$
Ethnicity (% Caucasian)	59%	74%	53%	$\chi^2 = 0.3$	$\chi^2 = 1.5$
Anxiety Severity (SD)	0.8 (1.2)	9.9 (3.2)	7.3 (3.4)	$t(61) = 10.2^*$	$t(39) = 2.5^*$
Worry (SD)	12.6 (1.1)	17.8 (2.7)	15.9 (3.7)	$t(61) = 6.1^*$	$t(39) = 2.0^{\dagger}$
Anxiety Sensitivity (SD)	7.5 (5.6)	28.1 (11.3)	35.3 (10.4)	$t(61) = 9.2^*$	$t(39) = -2.1^*$
Depression (SD)	2.1 (1.2)	8.8 (4.4)	6.2 (4.1)	$t(61) = 5.8^*$	$t(39) = 2.0^{\dagger}$
Reappraisal Use (SD)	31.0 (6.1)	24.8 (8.0)	32.7 (5.2)	$t(48) = -1.2$	$t(27) = -3.1^*$
Suppression Use (SD)	12.0 (2.8)	15.3 (5.5)	15.3 (4.9)	$t(48) = 2.5^*$	$t(27) = 0.02$
% with current MDD (past)	0% (0%)	9% (26%)	11% (22%)	$\chi^2 = 9.8^*$	$\chi^2 = 0.1$

Note. HC = Healthy Controls, ANX = GAD and PD groups combined, GAD = Generalized Anxiety Disorder, PD = Panic Disorder, SD = Standard Deviation, MDD = Major Depressive Disorder.

* = $p < .05$,

† = $p < .07$

Table 2
Regions demonstrating a significant differential effect of condition (Reappraise versus Maintain) across groups.

Region	BA	Peak			Vol
		x	y	z	
<i>Parietal Lobe</i>					
L postcentral gyrus / inferior parietal lobule	3, 40	-42	-29	48	7.53 2,944

Note. BA = Brodmann area, Vol = volume in μL , L = left, R = right, PFC = prefrontal cortex. Peak x y z coordinates in LPI orientation.

Table 3

A. Regions demonstrating a significant group effect in the Maintain versus Baseline contrast.
 B. Regions demonstrating a significant group effect in the Reappraise versus Baseline contrast.

Region	BA	Peak			Vol	Effect
		x	y	z		
A. Maintain versus Baseline contrast						
Frontal Lobe						
Dorsomedial PFC	6, 8	-10	31	56	14.03	19,712 HC > PD = GAD
Dorsal ACC [†]	32	10	15	36	8.12	640 HC > PD = GAD
Dorsal ACC [†]	32, 24	-6	31	20	8.90	1,536 HC > PD > GAD
L dorsolateral PFC	6, 9, 8	-42	-1	52	15.89	11,456 HC > PD = GAD
L dorsolateral PFC	9, 10	-30	19	28	12.16	5,696 HC > PD = GAD
R dorsolateral PFC	6, 8	42	-5	60	10.93	4,096 HC > PD = GAD
L ventrolateral PFC	45, 46	-42	19	20	8.19	1,472 HC > PD = GAD
R ventrolateral PFC [†]	44, 45	62	11	12	7.60	1,088 HC > PD = GAD
R ventrolateral PFC [†]	47, 45	42	39	-8	10.22	1,024 HC > PD > GAD
Temporal Lobe						
R middle and superior temporal gyrus	22, 21	62	-45	4	10.90	4,736 HC = PD > GAD
R temporal pole	38	46	3	-16	7.16	704 HC > PD = GAD
L middle temporal gyrus	21	-54	-45	0	11.03	1,088 HC > PD = GAD
L superior temporal gyrus	22	-50	-45	16	7.98	960 HC = PD > GAD
Parietal Lobe						
Posterior cingulate [†]	23, 31	-6	-45	20	7.25	1,344 HC > PD = GAD
L superior and inferior parietal lobule	7, 40	-50	-57	52	7.08	896 HC = PD > GAD
R supramarginal gyrus	40	38	-37	36	6.09	704 HC > PD > GAD
Occipital Lobe						
L fusiform gyrus	19, 18	-30	-61	-12	9.25	1,216 HC > PD = GAD
L middle occipital gyrus	37	-46	-61	-4	6.64	640 HC > GAD > PD
R cuneus	17	22	-85	0	7.63	1,024 HC > PD > GAD
Subcortical						

Region	BA	Peak			Vol	Effect
		x	y	z		
Bilateral thalamus/caudate, L anterior insula	13	-30	23	0	11.85	20,736 HC > PD = GAD
L cerebellum / L fusiform gyrus	37	-34	-41	-20	9.33	1,344 HC > PD = GAD
L cerebellum		-26	-69	-28	8.55	2,176 HC > PD > GAD
R cerebellum		34	-57	-24	10.61	1,600 HC > PD = GAD
R cerebellum		18	-57	-28	9.24	640 HC > PD = GAD
B. Reappraise versus Baseline contrast						
<i>Frontal Lobe</i>						
L dorsolateral PFC / premotor cortex	6, 9, 8	-38	3	32	11.49	4,544 HC > PD = GAD
R dorsolateral PFC [†]	9	58	15	28	6.17	960 HC > PD = GAD
R dorsolateral PFC	9	34	7	28	6.32	640 HC > PD = GAD
Dorsomedial PFC [†]	6	-10	27	56	5.23	640 HC > PD = GAD
<i>Temporal Lobe</i>						
L fusiform gyrus / cerebellum [†]	37	-34	-49	-20	7.95	1,216 HC > PD = GAD
<i>Occipital Lobe</i>						
R middle occipital gyrus [†]	17	30	-85	12	9.08	2,560 HC > PD > GAD
<i>Subcortical</i>						
L cerebellum [†]		-22	-73	-24	9.37	704 HC > PD > GAD

Note. BA = Brodmann area, Vol = volume in μL , L = left, R = right, PFC = prefrontal cortex, ACC = anterior cingulate cortex, HC = healthy control, PD = panic disorder, GAD = generalized anxiety disorder. Peak x y z coordinates in LPI orientation.

[†] = region no longer significant when controlling for age and gender.