Published in final edited form as:

J Hosp Med. 2015 February ; 10(2): 90–96. doi:10.1002/jhm.2280.

Characteristics associated with clinician diagnosis of aspiration pneumonia; a descriptive study of afflicted patients and their outcomes

Michael J. Lanspa, MD, MS^{1,2}, Paula Peyrani, MD³, Timothy Wiemken, PhD, MPH³, Emily Wilson, MStat¹, Julio A. Ramirez, MD³, and Nathan C. Dean, MD^{1,2}

Paula Peyrani: p0peyr01@louisville.edu; Timothy Wiemken: tim.wiemken@louisville.edu; Emily Wilson: emily.wilson@imail.org; Julio A. Ramirez: j.ramirez@louisville.edu; Nathan C. Dean: nathan.dean@imail.org

¹Division of Pulmonary and Critical Care Medicine, Intermountain Medical Center, Salt Lake City, UT

²Division of Pulmonary and Critical Care Medicine, University of Utah School of Medicine, Salt Lake City, UT

³Division of Infectious Diseases, University of Louisville Medical School, Louisville, KY

Abstract

Background—Aspiration pneumonia is a common disease, although less well characterized than other pneumonia syndromes.

Objective—We analyzed patient-level covariates associated with clinician-defined aspiration pneumonia.

Methods—We used the Community-Acquired Pneumonia Organization (CAPO) database, a multicenter, international population of patients with community-acquired pneumonia, using data from 2001 to 2012. Aspiration pneumonia was determined by the treating clinician. We analyzed covariates associated with clinician-defined diagnosis of aspiration pneumonia using logistic regression. We compared aspiration pneumonia patients to propensity-matched cases with non-aspiration pneumonia.

Results—We studied 5185 patients. 451 of these patients had aspiration pneumonia. Patients with aspiration pneumonia were older, had greater disease severity, and more comorbidities than patients with non-aspiration pneumonia. They were more likely cared for in the intensive care unit (19% vs. 13%, $p=0.002$), had longer unadjusted hospital length of stay (9 vs. 7 days, $p<0.001$), and took longer to achieve clinical stability (unadjusted 8 vs. 4 days, $p<0.001$). Confusion, nursing home residence, and cerebrovascular disease were most associated with clinician diagnosis of

Corresponding Author: Michael Lanspa, MD, Intermountain Medical Center, Shock-Trauma Intensive Care Unit, 5121 S. Cottonwood Street, Murray, UT 84107, USA, Tel: (801) 507-6450, Fax: (801) 507-4699, michael.lanspa@imail.org.

Disclosures: All authors report no relevant conflicts of interest.

Contributions: ML contributed to study design, data analysis, statistical analysis, writing manuscript. PP contributed to study design, revision of manuscript for important intellectual content. TW and EW contributed to study design, statistical analysis, and revision of manuscript for important intellectual content. JR and ND contributed to study design and revision of manuscript for important intellectual content. All authors read and approved final manuscript. ML takes responsibility for the integrity of the work as a whole, from inception to published article.

aspiration pneumonia (OR 4.4, 2.9, 2.3 respectively). Unadjusted inpatient mortality was higher (23% vs. 9%, $p < 0.001$). Aspiration pneumonia conferred a 2.3 odds ratio for inpatient mortality, after adjusting for age, disease severity, and comorbidities.

Conclusions—Among pneumonia patients, confusion, nursing home residence, and cerebrovascular disease are associated with a clinician diagnosis of aspiration. Aspiration pneumonia is associated with greater mortality among patients with community-acquired pneumonia, which is not explained by older age, measured indices of severity, or comorbidities.

Keywords

Aspiration; Pneumonia

Introduction

Pneumonia is a common clinical syndrome with well-described epidemiology and microbiology. Aspiration pneumonia comprises 5-15% of patients with pneumonia acquired outside of the hospital¹, but is less well-characterized despite being a major syndrome of pneumonia in the elderly.^{2,3} Difficulties in studying aspiration pneumonia include the lack of a sensitive and specific marker for aspiration as well as the potential overlap between aspiration pneumonia and other forms of pneumonia.⁴⁻⁶ Additionally, clinicians have difficulty distinguishing between aspiration pneumonia, which develops after the aspiration of oropharyngeal contents, and aspiration pneumonitis, wherein inhalation of gastric contents causes inflammation without the subsequent development of bacterial infection.^{7,8} Central to the study of aspiration pneumonia is whether it should exist as its own entity, or if aspiration is really a designation used for pneumonia in an older patient with greater comorbidities. The ability to clearly understand how a clinician diagnoses aspiration pneumonia, and whether that method has face validity with expert definitions may allow for improved future research, improved generalizability of current or past research, and possibly better clinical care.

Several validated mortality prediction models exist for community-acquired pneumonia (CAP), using a variety of clinical predictors, but their performance in patients with aspiration pneumonia is less well characterized. Most studies validating pneumonia severity scoring systems excluded aspiration pneumonia from their study population.⁹⁻¹¹ Severity scoring systems for CAP may not accurately predict disease severity in patients with aspiration pneumonia. The CURB-65⁹ and the eCURB¹² scoring systems are poor predictors of mortality in patients with aspiration pneumonia, perhaps because they do not account for patient comorbidities.¹³ The pneumonia severity index (PSI)¹⁰ might predict mortality better than CURB-65 in the aspiration population, due to inclusion of comorbidities.

Previous studies have demonstrated that patients with aspiration pneumonia are older, have greater disease severity, and more comorbidities.¹³⁻¹⁵ These single-center studies also demonstrated greater mortality, more frequent admission to an intensive care unit, and longer hospital lengths of stay in patients with aspiration pneumonia. These studies identified aspiration pneumonia by the presence of a risk factor for aspiration¹⁵, or by physician billing codes.¹³ In practice, however, the bedside clinician diagnoses a patient as

having aspiration pneumonia, but the logic is likely vague and inconsistent. Despite the potential for variability with individual judgment, an aggregate estimation from independent judgments may perform better than individual judgments.¹⁶ Because there is no gold standard for defining aspiration pneumonia, all previous research has been limited to definitions created by investigators. This multicenter study seeks to determine what clinical characteristics lead physicians to diagnose a patient as having aspiration pneumonia, and whether or not the clinician-derived diagnosis is distinct and clinically useful.

Our objectives were to: (1) Identify covariates associated with bedside clinicians diagnosing a pneumonia patient as having aspiration pneumonia; (2) compare aspiration pneumonia and non-aspiration pneumonia in regard to disease severity, patient demographics, comorbidities, and clinical outcomes; and (3) measure the performance of the PSI in aspiration pneumonia versus non-aspiration pneumonia.

Patients and Methods

Study Design and Setting

We performed a secondary analysis of the Community-Acquired Pneumonia Organization (CAPO) database, which contains retrospectively collected data from 71 hospitals in 16 countries between June 2001 and December 2012. In each participating center, primary investigators selected nonconsecutive, adult hospitalized patients diagnosed with CAP. To decrease systematic selection biases, the selection of patients with CAP for enrollment in the trial was based on the date of hospital admission. Each investigator completed a case report form that was transferred via the internet to the CAPO study center at the University of Louisville (Louisville, KY). A sample of the data collection form is available at the study web site (www.caposite.com). Validation of data quality was performed at the study center before the case was entered into the CAPO database. Local institutional review board approval was obtained for each study site.

Inclusion and Exclusion Criteria

Patients 18 years of age and satisfying criteria for CAP were included in this study. A diagnosis of CAP required a new pulmonary infiltrate at time of hospitalization, and at least one of the following: New or increased cough, leukocytosis, leukopenia, or left shift pattern on white blood cell count, temperature $> 37.8^{\circ}\text{C}$, or $< 35.6^{\circ}\text{C}$. We excluded patients with pneumonia attributed to mycobacterial or fungal infection, and patients infected with Human Immunodeficiency Virus (HIV), as we believed these types of pneumonia differ fundamentally from typical CAP.

Patient variables

Patient variables included presence of aspiration pneumonia, laboratory data, comorbidities, and measures of disease severity, including the PSI. The clinician made a clinical diagnosis of presence or absence of aspiration for each patient by marking a box on the case report form. Outcomes included in-hospital mortality, hospital length of stay up to 14 days, and time to clinical stability up to 8 days. All variables were obtained directly from the case report form. Clinical stability was defined according to previously published definitions; the

day the following criteria were all met: improved clinical signs (improved cough and shortness of breath), lack of fever for > 8 hours, improving leukocytosis (decreased at least 10% from the previous day), and tolerating oral intake.^{17,18}

Statistical analysis

Baseline characteristics of patients with aspiration and non-aspiration CAP were compared using Chi-squared or Fisher's exact tests for categorical variables and the Mann-Whitney U-test for continuous variables.

To determine which patient variables were important in the physician diagnosis of aspiration pneumonia, we performed logistic regression with initial covariates comprising of the demographic, comorbidity, and disease severity measurements listed in Table 1. We included interactions between cerebrovascular disease and age, nursing home status, and confusion, to improve model fit. We centered all variables (including binary indicators) according to the method outlined by Kraemer and Blasey to improve interpretation of the main effects.¹⁹

To determine if aspiration pneumonia had worse clinical outcomes compared to non-aspiration pneumonia, multiple methods were used. To compare the differences between the two groups with respect to time to clinical stability and length of hospital stay, we constructed Kaplan-Meier survival curves and Cox proportional hazards regression models. The Log-Rank test was used to determine statistical differences between the KM survival curves. To compare the impact of aspiration on mortality in patients with CAP, we conducted a propensity score matched analysis. We chose propensity score matching over traditional logistic regression to balance variables among groups and to avoid the potential for overfit and multicollinearity. We considered a variable balanced after matching if its standardized difference was < 10. All variables in the propensity score matched analysis were balanced.

Although our dataset contained minimal missing data, we imputed any missing values in order to maintain the full study population in the creation of the propensity score. Missing data were imputed using the `aregImpute` function of the `hmisc` package of R.^{20,21} We built the propensity score model using a variable selection algorithm described by Bursac et al.²² Our model included variables for region (USA/Canada, Europe, Asia/Africa, or Latin America) and the variables listed in Table 1, with the exception of the pneumonia severity index and the four clinical outcomes. Given that previous analyses accounting for clustering by physician did not substantially affect our results²³, our model did not include physician-level variables and did not account for the clustering effects of physicians. Using the propensity scores generated from this model, we matched a case of aspiration CAP with a case of non-aspiration CAP.²⁴ We then constructed a general linear model using the matched dataset to obtain the magnitude of effect of aspiration on mortality.

We used receiver-operating characteristic (ROC) curves to define the diagnostic accuracy of the pneumonia severity index for the prediction of mortality among patients with aspiration pneumonia and those with non-aspiration pneumonia. SAS v9.3 (SAS Institute, Cary, North

Carolina) and R v2.15.3 (R Development Core Team, Vienna, Austria), were used for all analyses. P-values of ≤ 0.05 were considered statistically significant in all analyses.

Results

Our initial query, after exclusion criteria, yielded a study population of 5185 patients (Figure 1). We compared 451 patients diagnosed with aspiration pneumonia to 4734 with CAP (Figure 1). Patient characteristics are summarized in Table 1. Patients with aspiration pneumonia were older, more likely to live in a nursing home, had greater disease severity, and more likely to be admitted to an intensive care unit (ICU). Patients with aspiration pneumonia had longer adjusted hospital lengths-of-stay and took more days to achieve clinical stability than patients with non-aspiration pneumonia (Figure 2). After adjusting for all variables in Table 1, the Cox proportional hazards models demonstrated that aspiration pneumonia was associated with ongoing hospitalization (HR for discharge 0.77, 95% confidence interval 0.65-0.91, $p = 0.002$) and clinical instability (HR for attaining clinical stability 0.72, 95% confidence interval 0.61-0.84, $p < 0.001$). Patients with aspiration pneumonia presented with greater disease severity than those with non-aspiration pneumonia. While there was no difference between groups in regard to temperature, respiratory rate, hyponatremia, or presence of pleural effusions or cavitory lesions, all other measured indices of disease severity were worse in patients with aspiration pneumonia. Patients with aspiration pneumonia were more likely to have cerebrovascular disease than those with non-aspiration pneumonia. Aspiration pneumonia patients also had increased prevalence of congestive heart failure. There was no appreciable difference between groups among other measured comorbidities.

The patient characteristics most associated with a physician diagnosis of aspiration pneumonia, identified using logistic regression, were confusion, residence in nursing home, and presence of cerebrovascular disease, (odds ratios of 4.4, 2.9, and 2.3 respectively), while renal disease was associated with decreased physician diagnosis of aspiration pneumonia over non-aspiration pneumonia (OR 0.58, Table 2).

Observed in-patient mortality of aspiration pneumonia was 23%. This mortality was considerably higher than a mean PSI score of 123 would predict (class-IV risk group, with expected 30-day mortality of 8-9%²⁵). The PSI score's ability to predict inpatient mortality in patients with aspiration pneumonia was moderate, with an area under the curve (AUC) of 0.71. This was similar to its performance in patients with non-aspiration pneumonia (AUC 0.75, Figure 3). These values are lower than the AUC of 0.81 for the PSI in predicting mortality, derived from a meta-analysis of 31 other studies.²⁶

Our regression model after propensity score matching, demonstrated that aspiration pneumonia independently confers a 2.3-fold increased odds for inpatient mortality (95% CI 1.56-3.45, $p < 0.001$).

Discussion

Pneumonia patients with confusion, nursing home residence, or cerebrovascular disease are more likely to be diagnosed with aspiration pneumonia by clinicians. While this is

unsurprising, it is notable that these patients are more than twice as likely to die in the inpatient setting, even after accounting for age, comorbidities, and disease severity. These findings are similar to three previously published studies comparing aspiration and non-aspiration pneumonia at single institutions, albeit using different aspiration pneumonia definitions.¹³⁻¹⁵ This study is the first large, multicenter, multinational study to demonstrate these findings.

Central to the interpretation of our results is the method of diagnosing aspiration vs. non-aspiration. A “bottom-up” method that relies on a clinician to check a box for “aspiration” may appear poorly reproducible. Because there is no diagnostic gold standard, clinicians may use different criteria to diagnose aspiration, creating potential for idiosyncratic noise. The strength of the “wisdom of the crowd” method used in this study is that an aggregate estimation from independent judgments may reduce the noise from individual judgments.¹⁶ Although clinicians may vary in why they diagnose a particular patient as having aspiration pneumonia, it appears that the overwhelming reason for diagnosing a patient as having aspiration pneumonia is the presence of confusion, followed by previous nursing home residence or cerebrovascular disease. This finding has some face validity when compared with studies using an investigator definition, as altered mental status, chronic debility, and cerebrovascular disease are either prominent features of the definition of aspiration pneumonia,⁸ or frequently observed in patients with aspiration pneumonia.^{13,15} The distribution of cerebrovascular disease among our study's aspiration and non-aspiration pneumonia patients was similar to studies which used formal criteria in their definitions.^{13,15} Although nursing home residence was more likely in aspiration pneumonia patients, the majority of aspiration pneumonia patients were residing in the community, suggesting that aspiration is not simply a surrogate for healthcare-associated pneumonia (HCAP). While patients with aspiration pneumonia are typically older than their non-aspiration counterparts, it appears that age is not a key determinant in the diagnosis of aspiration. With aspiration pneumonia, confusion, nursing home residence, and the presence of cerebrovascular disease are the greatest contributors in the clinical diagnosis, more than age.

Our data demonstrate that aspiration pneumonia confers increased odds for mortality, even after adjustment for age, disease severity, and comorbidities. These data suggest that aspiration pneumonia is a distinct entity from non-aspiration pneumonia, and that this disease is worse than non-aspiration CAP. If aspiration pneumonia is distinct from non-aspiration pneumonia, some unrecognized host factor other than age, disease severity, or the captured comorbidities decreases survival in aspiration pneumonia patients. However, it is also possible that aspiration pneumonia is merely a clinical designation for one end of the pneumonia spectrum, and we and others have failed to completely account for all measures of disease severity or all measures of comorbidities. Examples of unmeasured comorbidities would include presence of oropharyngeal dysphagia, which is not assessed in the database, but could have a significant effect on clinical diagnosis. Unmeasured covariates can include measures beyond that of disease severity or comorbidity, such as the presence of a Do Not Resuscitate (DNR) order, which could have a significant confounding effect on the observed association. A previous, single-center study demonstrated that increased 30-day mortality in aspiration pneumonia was mostly attributable to greater disease severity and comorbidities, although aspiration pneumonia independently conferred greater risk for adverse long-term

outcomes.¹⁵ We propose that aspiration pneumonia represents a clinically distinct entity from non-aspiration pneumonia. Patients with chronic aspiration are often chronically malnourished and may have different oral flora than patients without chronic aspiration.^{27,28} Chronic aspiration has been associated with granulomatous reaction, organizing pneumonia, diffuse alveolar damage, and chronic bronchiolitis.²⁹ Chronic aspiration may elicit changes in the host physiology, and may render the host more susceptible to the development of secondary bacterial infection, with morbid consequences.

The ability of the PSI to predict inpatient mortality was moderate (AUC only 0.7), with no significant additional discrimination between the aspiration and non-aspiration pneumonia groups. While the PSI had moderate ability to predict inpatient mortality, the observed mortality was considerably higher than predicted. It is possible that the PSI incompletely captures clinically relevant comorbidities (e.g., malnutrition). Further study to improve mortality prediction of aspiration pneumonia patients could employ sensitivity analysis to determine optimal thresholds and weighting of the PSI components.

Patients with aspiration pneumonia had longer hospital lengths-of-stay and took longer to achieve clinical stability than their non-aspiration counterparts. Time to clinical stability has been associated with increased post-hospitalization mortality and is associated with time to switch from intravenous to oral antibiotics.¹⁷ While some component of hospital length-of-stay is subject to local practice patterns, time to clinical stability has explicit criteria for clinical improvement and failure, and therefore is less likely to be affected by local practice patterns.

We noted a relatively high (16-21%) incidence of prior antibiotic use among patients in this database. Analysis of antibiotic prescription patterns was limited, given the several different countries from which the database draws its cases. Although we used accepted criteria to define community-acquired pneumonia cases, it is possible that this population may have a higher rate of resistant or uncommon pathogens than other studies of community-acquired pneumonia that have populations with lower incidence of prior antibiotic use. Although not assessed, we suspect a significant component of the prior antibiotic use represented outpatient pneumonia treatment during the few days prior to visiting the hospital.

This study has several limitations, of which the most important may be that we used clinical determination for defining presence of aspiration pneumonia. This method is susceptible to the subjective perceptions of the treating clinician. We did not account for the effect of individual physicians in our model, although we did adjust for regional differences. The retrospective identification of patients allows for the possibility of selection bias, and therefore we have not attempted to make inferences regarding the relative incidence of pneumonia, nor did we adjust for temporal trends in diagnosis. The ratio of aspiration pneumonia patients to non-aspiration pneumonia patients may not necessarily reflect that observed in reality. Microbiologic and antibiotic data was unavailable for analysis. This study cannot inform on non-hospitalized patients with aspiration pneumonia, as only hospitalized patients were enrolled. The database identified cases of pneumonia, so it is possible for a patient to enter into the database more than once. Detection of mortality was limited to the inpatient setting rather than a set interval of 30 days. Inpatient mortality

depends on length-of-stay patterns that may bias the mortality endpoint.³⁰ Also not assessed was the presence of a DNR order. It is possible that an older patient with greater comorbidities and disease severity may have care intentionally limited or withdrawn early by the family or clinicians.

Strengths of the study include its size and its multicenter, multinational population. The CAPO database is a large and well-described population of patients with CAP.^{17,31} These attributes, as well as the clinician-determined diagnosis, increase the generalizability of the study, compared to a single-center, single-country study that employs investigator-defined criteria.

Conclusion

Pneumonia patients with confusion, nursing home residence, and cerebrovascular disease are more likely to be diagnosed with aspiration pneumonia by clinicians. Our clinician-diagnosed cohort appears similar to those derived using an investigator definition. Patients with aspiration pneumonia are older, have greater disease severity, and more comorbidities than patients with non-aspiration pneumonia. They have greater mortality than their PSI score class would predict. Even after accounting for age, disease severity, and comorbidities, the presence of aspiration pneumonia independently conferred a greater than twofold increase in inpatient mortality. These findings together suggest that aspiration pneumonia should be considered a distinct entity from typical pneumonia, and that additional research should be done in this field.

Acknowledgments

This investigation was partly supported with funding from the National Center for Research Resources and the National Center for Advancing Translational Sciences, National Institutes of Health [Grant 8UL1TR000105 (formerly UL1RR025764)].

References

1. Torres A, Serra-Batlles J, Ferrer A, et al. Severe community-acquired pneumonia. Epidemiology and prognostic factors. *Am Rev Respir Dis*. 1991; 144(2):312–318. [PubMed: 1859053]
2. Koivula I, Sten M, Makela PH. Risk factors for pneumonia in the elderly. *Am J Med*. 1994; 96(4): 313–320. [PubMed: 8166149]
3. Marik PE, Kaplan D. Aspiration pneumonia and dysphagia in the elderly. *Chest*. 2003; 124(1):328–336. [PubMed: 12853541]
4. Mylotte JM, Goodnough S, Naughton BJ. Pneumonia versus aspiration pneumonitis in nursing home residents: diagnosis and management. *J Am Geriatr Soc*. 2003; 51(1):17–23. [PubMed: 12534840]
5. Marik PE. Aspiration pneumonia: mixing apples with oranges and tangerines. *Crit Care Med*. 2004; 32(5):1236. author reply 1236–1237. [PubMed: 15190985]
6. Kozlow JH, Berenholtz SM, Garrett E, Dorman T, Pronovost PJ. Epidemiology and impact of aspiration pneumonia in patients undergoing surgery in Maryland, 1999–2000. *Crit Care Med*. 2003; 31(7):1930–1937. [PubMed: 12847385]
7. Marik PE. Aspiration syndromes: aspiration pneumonia and pneumonitis. *Hosp Pract (Minneap)*. 2010; 38(1):35–42.
8. Marik PE. Aspiration pneumonitis and aspiration pneumonia. *N Engl J Med*. 2001; 344(9):665–671. [PubMed: 11228282]

9. Lim WS, van der Eerden MM, Laing R, et al. Defining community acquired pneumonia severity on presentation to hospital: an international derivation and validation study. *Thorax*. 2003; 58(5):377–382. [PubMed: 12728155]
10. Fine MJ, Hanusa BH, Lave JR, et al. Comparison of a disease-specific and a generic severity of illness measure for patients with community-acquired pneumonia. *Journal of general internal medicine*. 1995; 10(7):359–368. [PubMed: 7472683]
11. Espana PP, Capelastegui A, Gorordo I, et al. Development and validation of a clinical prediction rule for severe community-acquired pneumonia. *Am J Respir Crit Care Med*. 2006; 174(11):1249–1256. [PubMed: 16973986]
12. Jones BE, Jones J, Bewick T, et al. CURB-65 Pneumonia Severity Assessment Adapted for Electronic Decision Support. *Chest*. 2011; 140(1):156–163. [PubMed: 21163875]
13. Lanspa MJ, Jones BE, Brown SM, Dean NC. Mortality, morbidity, and disease severity of patients with aspiration pneumonia. *Journal of hospital medicine : an official publication of the Society of Hospital Medicine*. 2013; 8(2):83–90. [PubMed: 23184866]
14. Heppner HJ, Sehlhoff B, Niklaus D, Pientka L, Thiem U. Pneumonia Severity Index (PSI), CURB-65, and mortality in hospitalized elderly patients with aspiration pneumonia. *Z Gerontol Geriatr*. 2011; 44(4):229–234. [PubMed: 21769515]
15. Taylor JK, Fleming GB, Singanayagam A, Hill AT, Chalmers JD. Risk factors for aspiration in community-acquired pneumonia: analysis of a hospitalized UK cohort. *The American journal of medicine*. 2013; 126(11):995–1001. [PubMed: 24054176]
16. Yi SK, Steyvers M, Lee MD, Dry MJ. The wisdom of the crowd in combinatorial problems. *Cognitive science*. 2012; 36(3):452–470. [PubMed: 22268680]
17. Aliberti S, Peyrani P, Filardo G, et al. Association between time to clinical stability and outcomes after discharge in hospitalized patients with community-acquired pneumonia. *Chest*. 2011; 140(2):482–488. [PubMed: 21330383]
18. Ramirez JA. Clinical stability and switch therapy in hospitalised patients with community-acquired pneumonia: are we there yet? *The European respiratory journal : official journal of the European Society for Clinical Respiratory Physiology*. 2013; 41(1):5–6.
19. Kraemer HC, Blasey CM. Centring in regression analyses: a strategy to prevent errors in statistical inference. *International journal of methods in psychiatric research*. 2004; 13(3):141–151. [PubMed: 15297898]
20. Harrell F. Hmisc: Harrell miscellaneous. 2012
21. Heitjan DF, Little RJA. Multiple Imputation for the Fatal Accident Reporting System. *Journal of the Royal Statistical Society (Applied Statistics)*. 1991; 40(1):13–29.
22. Bursac Z, Gauss CH, Williams DK, Hosmer DW. Purposeful selection of variables in logistic regression. *Source code for biology and medicine*. 2008; 3:17. [PubMed: 19087314]
23. Arnold FW, Wiemken TL, Peyrani P, Ramirez JA, Brock GN, authors C. Mortality differences among hospitalized patients with community-acquired pneumonia in three world regions: results from the Community-Acquired Pneumonia Organization (CAPO) International Cohort Study. *Respiratory medicine*. 2013; 107(7):1101–1111. [PubMed: 23660396]
24. Parsons L. Reducing bias in a propensity score matched-pair sample using greedy matching techniques. *SAS SUGI*. 2001:214–226.
25. Fine MJ, Auble TE, Yealy DM, et al. A prediction rule to identify low-risk patients with community-acquired pneumonia. *N Engl J Med*. 1997; 336(4):243–250. [PubMed: 8995086]
26. Chalmers JD, Singanayagam A, Akram AR, et al. Severity assessment tools for predicting mortality in hospitalised patients with community-acquired pneumonia. *Systematic review and meta-analysis*. *Thorax*. 2010; 65(10):878–883. [PubMed: 20729231]
27. Cabre M, Serra-Prat M, Palomera E, Almirall J, Pallares R, Clave P. Prevalence and prognostic implications of dysphagia in elderly patients with pneumonia. *Age and ageing*. 2010; 39(1):39–45. [PubMed: 19561160]
28. Pace CC, McCullough GH. The association between oral microorganisms and aspiration pneumonia in the institutionalized elderly: review and recommendations. *Dysphagia*. 2010; 25(4):307–322. [PubMed: 20824288]

29. Yousem SA, Faber C. Histopathology of aspiration pneumonia not associated with food or other particulate matter: a clinicopathologic study of 10 cases diagnosed on biopsy. *The American journal of surgical pathology*. 2011; 35(3):426–431. [PubMed: 21317714]
30. Jencks SF, Daley J, Draper D, Thomas N, Lenhart G, Walker J. Interpreting hospital mortality data. The role of clinical risk adjustment. *JAMA*. 1988; 260(24):3611–3616. [PubMed: 3057250]
31. Arnold FW, Ramirez JA, McDonald LC, Xia EL. Hospitalization for community-acquired pneumonia: the pneumonia severity index vs clinical judgment. *Chest*. 2003; 124(1):121–124. [PubMed: 12853513]

Figure 1.

Patient selection from June 2001 –December 2012. HIV = Human Immunodeficiency Virus.

Figure 2. Kaplan Meier graph of hospital length of stay (A, above), and time to clinical stability (B, below)

Figure 3.
Receiver-operating characteristic curve of pneumonia severity index score and inpatient mortality.

Table 1

Patient characteristics of the CAPO database, stratified by aspiration pneumonia. All continuous data are median values followed by interquartile ranges, unless otherwise specified. Significance testing between groups was assessed using Chi-square or Mann-Whitney U, where appropriate. BUN: blood urea nitrogen.

	Aspiration Pneumonia (N=451)	Non-Aspiration pneumonia (N=4734)	p-value
Demographics			
Age (years)	79 (22)	69 (27)	<0.001
% male	59%	60%	0.58
Nursing Home Residence	25%	5%	<0.001
Recent (30 day) antibiotic use	21%	16%	0.017
Comorbidities			
Cerebrovascular disease	35%	14%	<0.001
Chronic Obstructive Pulmonary Disease	25%	27%	0.62
Congestive Heart Failure	23%	19%	0.027
Diabetes	18%	18%	0.85
Cancer	12%	10%	0.12
Renal disease	10%	11%	0.53
Liver disease	6%	5%	0.29
Disease Severity			
Pneumonia Severity Index	123 (54)	92 (49)	<0.001
Confusion	49%	12%	<0.001
PaO ₂ < 60 mmHg	43%	33%	<0.001
BUN > 30 g/dL	42%	23%	<0.001
Multilobar pneumonia	34%	28%	0.003
Pleural effusion	25%	21%	0.07
Respiratory Rate > 30 breaths/min	21%	20%	0.95
pH < 7.35	13%	5%	<0.001
Hematocrit < 30%	11%	6%	0.001
Temperature > 37.8° C or < 35.6° C	9%	7%	0.30
Systolic Blood Pressure < 90 mmHg	8%	9%	0.003
Sodium < 130 mEq/L	8%	6%	0.08
Heart rate > 125 beats/min	8%	5%	0.71
Glucose > 250 mg/dL	6%	7%	0.06
Cavitary lesion	0%	0%	0.67
Clinical Outcomes			
In-hospital mortality	23%	9%	<0.001
Intensive care unit admission	19%	13%	0.002
Hospital Length of stay (days)	9 (10)	7 (8)	<0.001
Time to clinical stability (days)	8 (4)	4 (5)	<0.001

Table 2

Final logistic regression model for physician diagnosis of aspiration pneumonia. Initial model included all demographic, comorbidity, and disease severity measurements from Table 1. Parameter estimates are for mean-centered variables. Renal disease is defined as having a clinical diagnosis in the medical record. Although other interaction terms were used in the initial model, they were eliminated from the final model. We centered all variables (including binary indicators) according to the method described by Kraemer and Blasey.¹⁹ The area under the curve of the final model is 0.79

Covariate	Odds ratio	95% confidence intervals	P value
Demographics			
Age (years)	1.00	0.99-1.01	0.948
Male	1.20	0.94 – 1.54	0.148
Nursing Home Residence	2.93	2.13-4.00	<0.001
Comorbidities			
Cerebrovascular disease	2.26	1.53-3.32	<0.001
Renal disease	0.58	0.39 – 0.85	0.006
Disease Severity			
Confusion	4.41	3.40-5.72	<0.001
Hematocrit < 30%	1.59	1.06 – 2.33	0.020
pH < 7.35	1.67	1.10 – 2.47	0.013
Temperature > 37.8° C or < 35.6° C	1.60	1.07 – 2.35	0.019
Multilobar pneumonia	1.29	1.00 – 1.65	0.047
Interaction terms			
Age * Cerebrovascular disease	0.98	0.96 – 0.99	0.011
Nursing home * Cerebrovascular disease	0.51	0.27 – 0.96	0.037
Confusion * Cerebrovascular disease	0.70	0.42 – 1.17	0.175