Published in final edited form as:

Bioorg Med Chem Lett. 2015 February 1; 25(3): 685–689. doi:10.1016/j.bmcl.2014.11.078.

Diversity Oriented Synthesis and IKK inhibition of Aminobenzimidazole Tethered quinazoline-2,4-diones, thioxoquinazolin-4-ones, benzodiazepine-2,3,5-triones, isoxazoles and isoxazolines

Sureshbabu Dadiboyena^a, Aïcha Arfaoui^b, Hassen Amri^b, F. Javier Piedrafita^c, and Adel Nefzi^{a,*}

^aTorrey Pines Institute for Molecular Studies, 11350 SW Village Parkway, Port St. Lucie, FL 34987 USA

^bLaboratory of Selective Organic Chemistry & Biological Activity, Faculty of Science of Tunis, University El Manar, 2092, Tunisia

^cTorrey Pines Institute for Molecular Studies, General Atomics Court, San Diego, CA 92121 USA

Abstract

The derivatization of resin-bound aminobenzimidazole toward The parallel solid-phase synthesis of aminobenzimidazole tethered pharmacologically important heterocycles such as quinazoline-2,4-diones, thioxoquinazolin-4-ones, benzodiazepine-2,3,5-triones, isoxazoles and isoxazolines is reported. All the compounds were tested for IKK inhibition. Only one compound elicited significant inhibition of IKK ϵ , TBK-1 and IKK2.

Keywords

Solid-phase synthesis; aminobenzimidazoles; quinazolinediones; thioxoquinazolinones; benzodiazepinetriones; Intramolecular cyclization; Isoxazoles; Isoxazolines; 1,3-Dipolar cycloaddition; heterocycles

One of the central objectives of organic and medicinal chemistry is the design, synthesis, and production of molecules having value as human therapeutic agents. Historically, the identification of such compounds has been carried out using compounds from plant and animal tissue extracts, microbial broth extracts, as well as individual compound collections resulting from fifty years of effort by synthetic chemists in academic and pharmaceutical organizations. Diversity-Oriented Synthesis (DOS), is a process by which multiple compounds are generated simultaneously, in a predictable fashion using techniques that

© 2014 Elsevier Ltd. All rights reserved

Corresponding author: adeln@tpims.org Phone: 1-772-345-4739, Fax: 1-772-345-3649.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

involve parallel chemical transformations.¹⁻⁴ It allows chemists to achieve more structural complexity than in the early days of combinatorial chemistry.⁴⁻⁷ Benzimidazoles are an important class of heterocycles displaying a wide array of biological properties,⁸⁻¹⁰ and represent a key structural motif in angiotensin-II-antagonists, NMDA antagonists, anticoagulants, and gastric protonpump inhibitors.¹¹⁻¹⁴ We previously reported the use of resin-bound aminobenzimidazoles for the synthesis of a variety of fused and/or tethered heterocyclic compounds.¹⁵⁻¹⁹

Continuing with our interest with the diversification of aminobenzimidazoles, we describe herein a multistep approach for the parallel synthesis of structurally diverse aminobenzimidazole tethered pharmacologically known heterocycles such as quinazoline-2,4-diones, thioxoquinazolin-4-ones, benzodiazepine-2,3,5-triones, isoxazoles and isoxazolines. Substituted quinazolinone, thioxoquinazolinone and benzodiazepinetrione are found in natural products,²⁰ and in various drug based p38 kinase inhibitors,²¹ aldose reductase inhibitors,²² 5-HT_{2C} agonist,²³ and CB₂ agonists.²⁴ Recently, quinazolinones have been developed as typical anti-psychotic agents for treating Schizophrenia and Alzheimer's diseases.²⁵ Likewise, benzodiazepines are found in several drugs including serotonin and dopamine receptors,²⁶ 5-HT_{2C} receptors,²⁷ and glycogen synthase kinase-3 inhibitors.²⁸

The parallel solid-phase synthesis (tea-bag technology)²⁹ of aminobenzimidazole tethered quinazoline-2,4-diones, thioxoquinazolin-4-ones and benzodiazepine-2,3,5-triones is outlined in Scheme 1. Resin-bound aminobenzimidazole template **1**³⁰ was coupled to 2-nitrobenzoic acid in the presence of PyBOP to provide an essential N-benzimidazolyl nitrobenzamide precursor **2** which, following tin (II) chloride reduction generated an amine **3**. Treatment of the amine **3** with 1,1'-carbonyldiimidazole generated an isocyanate intermediate which, upon intramolecular cyclization furnished the resin-bound quinazoline-2,4-diones **4**. Similarly, the separate treatment of compound **3** with 1,1'-thiocarbonyldiimidazole and oxalyldiimidazole afforded following thioxoquinazolin-4-ones **5** and benzodiazepine-2,3,5-triones **6**.

The resin was cleaved with HF/anisole and the desired quinazoline-2,4-diones **4**, thioxoquinazolin-4-ones **5**, and benzodiazepine-2,3,5-triones **6** were isolated in moderate yields (30-55%) (Table 1).

The application of this solid-phase methodology was explored to a further extent and a series of aminobenzimidazole tethered isoxazoles and isoxazolines were synthesized. We envisioned that aminobenzimidazole coupled alkyne or alkene template would serve as a convenient partner for 1,3-dipolar cycloaddition studies. Recently, we documented the synthesis of an array of isoxazoles and isoxazolines via 1,3-dipolar cycloaddition using resin-bound alkenes and alkynes.³¹ In order to build upon this premise, we decided to study the application of aminobenzimidazole based alkyne or alkene template to access a variety of tethered cycloaddition products. Aminobenzimidazole tethered isoxazoles and isoxazolines were obtained following on resin-bound 1,3-dipolar cycloaddition of aminobenzimidazole acylated with carboxylic acids bearing alkyne or alkene with nitrile oxides. Isoxazoles and isoxazolines are very important class of active compounds. They

display antiviral,³² antitubulin,³³ as well as anti-inflammatory activities.^{34,35} Isoxazoline core is a prevalent feature for several spiroisoxazoline natural products³⁶⁻⁴¹ and isoxazole motif is found in pharmaceutical drugs such as bextra® and parecoxib.⁴²⁻⁴⁶ Due to aforementioned applications, the syntheses of these isoxazole (isoxazoline) based structural units have received greater attention and a few examples representing isoxazoline (**7-12**) and isoxazole (**13,14**) structural motifs are presented in Figure 1.

Our approach toward the synthesis of aminobenzimidazole tethered isoxazoles and isoxazolines is outlined in Scheme 2. Following the coupling of phenylpropionic acid or 4-vinylbenzoic acid to resin-bound aminobenzimidazole **1**, the generated alkyne **15** or alkene precursors **16** were treated with freshly prepared hydroximoyl chlorides in the presence of diisopropyldiethylamine (DIEA). The *in situ* formed nitrile oxides reacted with alkenes or alkynes in a 1,3-dipolar fashion^{31,47-52} to furnish the corresponding resin-bound isoxazoles **18** or isoxazolines **19**. The cycloaddition reactions on solid-phase occurred with complete regiochemical integrity and the products were isolated in good yields (Table 2). The resin was cleaved with HF/anisole and the desired aminobenzimidazole tethered isoxazoles **20** and racemic isoxazolines **21** were obtained following purification in reasonable yields (12-41%).

We evaluated the effect of all the compounds on the kinase activity of the atypical Inhibitor of kappaB kinases (IKKs), IKK epsilon (IKK ϵ) and TANK-Binding Kinase-1 (TBK-1),⁵⁴ which play an essential role in carcinogenesis, inflammation and immunity. In particular, IKKs was identified as a breast protooncogene and is overexpressed in many breast cancer cell lines and primary samples. All compounds were first tested at a concentration of 10 μ M, and only the Benzimidazole tethered thioxoquinazolin-4-one **5a** elicited significant inhibition (> 50%) of both kinases *in vitro*. The rest of the compounds showed activity less than 10%. A follow up dose response for **5a** was carried out and we included another member of the IKK family, IKK2, as counter screen. We found that compound **5a** was actually a better inhibitor of IKK2 (Figure 2) with an IC₅₀ of 2.643 μ M. The IC₅₀s for IKKs and TBK-1 were 3.774 and 6.224 μ M, respectively.

In conclusion, we have developed different multi-step solid-phase strategies for the construction of aminobenzimidazole separately tethered with a variety of biologically important heterocycles, such as quinazoline-2,4-diones, thioxoquinazolin-4-ones, benzodiazepine-2,3,5-triones, isoxazoles and isoxazolines.^{52, 53} Following the screening of all the compounds for IKK inhibition, compound **5a** showed significant inhibition of IKK1, IKK ϵ and IKK2.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was funded in part through the Florida Drug Discovery Acceleration Program by the State of Florida, Department of Health. NIH grant 1R01AI105836-01A1(Piedrafita/Nefzi)

References and Notes

1. Schreiber SL. *Science*. 2000; 287:1964. [PubMed: 10720315]
2. Schreiber SL, Nicolaou KC, Davies K. *Chem Biol*. 2002; 9:1. [PubMed: 11841933]
3. Stockwell BR. *Nature*. 2004; 432:846. [PubMed: 15602550]
4. Thompson LA, Ellman JA. *Chem. Rev.* 1996; 96:555. [PubMed: 11848765]
5. Nefzi A, Ostresh JM, Houghten RA. *Chem. Rev.* 1997; 97:449. [PubMed: 11848878]
6. Dolle RE. *J Comb. Chem.* 2001; 4:369. [PubMed: 12217012]
7. Nicolaou KC, Pfefferkorn JA, Barluenga S, Mitchell HJ, Roecker AJ, Cao G-Q. *J Am Chem Soc.* 2000; 122:9968.
8. Brown HD, Matzuk AR, Ilves IR, Peterson LH, Harris SA, Sarett LH, Egerton JR, Yakstis JJ, Campbell WC, Cuckler AC. *J. Am. Chem. Soc.* 1961; 83:1764.
9. Craig WA, LeSueur BW, Skibo EB. *J. Med. Chem.* 1999; 42:3324. [PubMed: 10464019]
10. Wienen W, Huel N, van Meel JC, Narr B, Ries U, Entzeroth M. *Br J. Pharmacol.* 1993; 110:245. [PubMed: 8220885]
11. McCanley JA, Munson PM, Thompson W, Claremon DA. 2000 WO0132634.
12. McCanley JA, Munson PM, Thompson W, Claremon DA. 2000 WO0132634.
13. Rogers SA, Huigens RW III, Melander CJ. *Am. Chem. Soc.* 2009; 131:9868.
14. Seth PA, Jefferson EA, Risen LM, Osgood SA. *Bioorg. Med. Chem. Lett.* 2003; 13:1669. [PubMed: 12729638]
15. Hoesl CE, Nefzi A, Houghten RA. *J. Comb. Chem.* 2004; 6:220. [PubMed: 15002970]
16. Nefzi A, Giulianotti M, Houghten RA. *Tetrahedron Lett.* 2000; 41:2283.
17. Dadiboyena S, Nefzi A. *Tetrahedron Lett.* 2011; 52:7030. [PubMed: 22199408]
18. Dadiboyena S, Nefzi A. *Synthesis*. 2012; 44:215.
19. Dadiboyena S, Nefzi A. *Tetrahedron Lett.* 2012; 53:6897. [PubMed: 23209332]
20. Niwa K, Yoshida Y, Yamada KJ. *Nat. Prod.* 1988; 51:343.
21. Schlapbach A, Heng R, Di Padova F. *Bioorg. Med. Chem. Lett.* 2004; 14:357. [PubMed: 14698158]
22. Malamas MS, Millen J. *J. Med. Chem.* 1991; 34:1492. [PubMed: 1901912]
23. Ahmad S, Ngu K, Miller KJ, Wu G, Hung C, Malmstrom S, Zhang G, O'Tanyi E, Keim WJ, Cullen MJ, Rohrbach KW, Thomas M, Ung T, Qu Q, Gan J, Narayanan R, Pelleymounter MA, Robl JA. *Bioorg. Med. Chem. Lett.* 2010; 20:1128. [PubMed: 20022752]
24. Omura H, Kawai M, Shima A, Iwata Y, Ito F, Masuda T, Ohta A, Makita N, Omoto K, Sugimoto H, Kikuchi A, Iwata H, Ando K. *Bioorg. Med. Chem. Lett.* 2008; 18:3310. [PubMed: 18440813]
25. Bhat L, Mohapatra PP, Dadiboyena S, Adiey K. 2010 WO 099503A1.
26. Guzzo PR, Molino BF, Cui W, Liu S, Olson RE. *PTC Int. Appl.* 2008 2008141081.
27. Sabb AL, Vogel RL, Welmaker GS, Sabalski JE, Coupet J, Dunlop J, Resenzweig-Lipson S, Harrison B. *Bioorg. Med. Chem. Lett.* 2004; 14:2603. [PubMed: 15109661]
28. Magnus NA, Ley CP, Pollock PM, Wepsiec JP. *Org. Lett.* 2010; 12:3700. [PubMed: 20704418]
29. Houghten RA. *Proc. Natl. Acad. Sci. U.S.A.* 1985; 82:5131. [PubMed: 2410914]
30. Dadiboyena S, Nefzi A. *Tetrahedron Lett.* 2012; 53:2096. [PubMed: 22715296]
31. Lee Y-S, Kim BH. *Bioorg. Med. Chem. Lett.* 2002; 12:1395. [PubMed: 11992785]
32. Srivastava S, Bajpai LK, Batra S, Bhaduri AP, Maikhuri JP, Gupta G, Dhar JD. *Bioorg. Med. Chem.* 1999; 7:2607. [PubMed: 10632071]
33. Simoni D, Grisolia G, Giannini G, Roberti M, Rondanin R, Piccagli L, Baruchello R, Rossi M, Romagnoli R, Invidiata FP, Grimaudo S, Jung MK, Hamel E, Gebbia N, Crosta L, Abbadessa V, Di Cristina A, Dusonchet L, Meli M, Tolomeo M. *J. Med. Chem.* 2005; 48:723. [PubMed: 15689156]
34. Kaffy J, Pontikis R, Carrez D, Croisy A, Monneret C, Florent J-C. *Bioorg. Med. Chem.* 2006; 14:4067. [PubMed: 16510288]

35. Berquist, PR.; Wells, RJ. Chemotaxonomy of the Porifera: The Development and Current Status of the Field. In: Scheuer, PJ., editor. Marine Natural Products: Chemical and Biological Perspectives. Vol. 5. Academic Press; New York: 1993. p. 1-50.
36. Faulkner DJ. Nat. Prod. Rep. 1998;113. [PubMed: 9586223]
37. Faulkner DJ. Nat. Prod. Rep. 1997;259.
38. Berquist, PR.; Wells, RJ. Marine Natural Products. Scheuer, PJ., editor. Vol. V. Academic Press; New York: 1983. Chapter 1
39. Encarnacion RD, Sandoval E, Malmstrom J, Christophersen CJ. Nat. Prod. 2000; 63:874.
40. Dadiboyena S, Hamme AT II. Tetrahedron Lett. 2011; 52:2536. [PubMed: 21686095]
41. Talley JJ, Brown DL, Carter JS, Graneto MJ, Koboldt CM, Masferrer JL, Perkins WE, Rogers RS, Shaffer AF, Zhang YY, Zweifel BS, Seibert KJ. Med. Chem. 2000; 43:775.
42. Talley JJ, Bertenshaw SR, Brown DL, Carter JS, Graneto MJ, Kellogg MS, Koboldt CM, Yuan J, Zhang YY, Seibert KJ. Med. Chem. 2000; 43:1661.
43. Toyokuni T, Dileep Kumar JS, Walsh JC, Shapiro A, Talley JJ, Phelps ME, Herschman HR, Barrio JR, Satyamurthy N. Bioorg. Med. Chem. Lett. 2005; 15:4699. [PubMed: 16153836]
44. Penning TD, Talley JJ, Bertenshaw SR, Carter JS, Collins PW, Docter S, Graneto MJ, Lee LF, Malecha JW, Miyashiro JM, Rogers RS, Rogier DJ, Yu SS, Anderson GD, Burton EG, Cogburn JN, Gregory SA, Koboldt CM, Perkins WE, Seibert K, Veenhuizen AW, Zhang YY, Isakson PC. J. Med. Chem. 1997; 40:1347. [PubMed: 9135032]
45. Dadiboyena S, Nefzi A. Eur. J. Med. Chem. 2010; 45:4697. [PubMed: 20724040]
46. Dadiboyena S, Xu J, Hamme AT II. Tetrahedron Lett. 2007; 48:1295. [PubMed: 18265873]
47. Sandanayaka VP, Youjun Y. Org. Lett. 2000; 2:3087. [PubMed: 11009352]
48. Huisgen R. Angew. Chem. Intl. Ed. 1963; 2:565.
49. Huisgen R. Angew. Chem. Intl. Ed. 1963; 2:633.
50. Croce PD, La Rosa C, Zecchi GJ. Chem. Soc., Perkin Trans. 1985:2621. 1.
51. Crossley JA, Browne DL. J. Org. Chem. 2010; 75:5414. [PubMed: 20608670]
52. General procedure for the synthesis of amino-benzimidazole tethered quinazoline-2,4-dione, thioxoquinazolin-4-one, and benzodiazepine-2,3,5-triones: p-Methylbenzhydrylamine (MBHA) resin (100 mg, 1.10 meq/g, 100-200 mesh) was sealed inside a polypropylene mesh packet. Polypropylene bottles were used for all of the reactions. Resin bound aminobenzimidazoles **1** were synthesized according to a previous literature.^{15,16} 2-Nitrobenzoic acid (8 equiv.) was coupled to resin bound aminobenzimidazole for 8h using PyBOP (8 equiv.).^{17,18} The excess solution was decanted and the resin was washed with DMF (3 times) and DCM (3 times). Reduction of the nitro group was achieved with tin(II) chloride (10 eq., 1.0M in DMF) and the resin was washed with DMF (10 times).¹⁵ Intramolecular cyclizations were performed in a 100 mL glass vial fitted with a screw cap. The N-terminal free amine **3** was treated with 1,1'-carbonyldiimidazole (or) 1,1'-thiocarbonyldiimidazole (or) 1,1'-oxalyldiimidazole (0.5M in anhydrous DMF) and allowed to stir at 80 °C for 8h.¹⁷ The excess solution was decanted, and the resulting resin-bound aminobenzimidazolyl heterocycles were washed with DMF (2 times) and DCM (2 times). The resin was cleaved with HF/anisole (95:5) for 90 min at 0 °C and the desired quinazoline-2,4-diones (or) thioxodihydroquinazolin-4-ones (or) benzodiazepine-2,3,5-triones were obtained following extraction with 95% AcOH in H₂O, lyophilized as a colorless powder, and purified by preparative reverse-phase HPLC. **4a**: ¹H NMR (DMSO-d₆): δ 1.77-1.81 (m, 2H), 2.07-2.10 (m, 5H), 2.29-2.34 (m, 2H), 5.33 (t, J = 8Hz, 1H), 7.39 (t, J = 8 Hz, 2H), 7.62-7.67 (m, 2H), 7.81 (t, J = 8Hz, 1H), 8.05 (d, J = 8Hz, 1H), 8.13 (br s, 1H), 8.28 (d, J = 8Hz, 1H), 9.05 (s, 1H); LC-MS m/z data calcd. for C₂₁H₁₉N₅O₃ (M⁺): 389.40; found: (MH⁺) 390.5; **4b**: ¹H NMR (DMSO-d₆): δ 0.93 (t, J = 8 Hz, 4H), 1.33-1.42 (m, 2H), 1.68-1.76 (m, 2H), 3.98 (t, J = 8Hz, 2H), 7.33 (br s, 1H), 7.46 (d, J = 8Hz, 1H), 7.52 (t, J = 8Hz, 1H), 7.76 (d, J = 8Hz, 1H), 7.83-7.88 (m, 2H), 8.05 (br s, 1H), 8.15 (d, J = 8Hz, 1H), 8.81(br s, 1H), 12.43 (br s, 1H); LC-MS m/z data calcd. for C₂₀H₁₉N₅O₃ (M⁺): 377.40; found (MNa⁺): 400.0; **4c**: ¹H NMR (DMSO-d₆): δ 0.89-1.07 (m, 6H), 1.47-1.59 (m, 5H), 1.77-1.82 (m, 1H), 3.94 (d, J = 8Hz, 2H), 7.24-7.29 (m, 3H), 7.72-7.78 (m, 2H), 7.90 (d, J = 8Hz, 2H), 7.98 (d, J = 8Hz, 1H), 8.02 (br s, 1H), 8.25 (br s, 1H); LC-MS m/z data calcd. for C₂₃H₂₃N₅O₃ (M⁺): 417.46; found (MH⁺): 418.5; **4d**: ¹H NMR (DMSO-d₆): δ 0.83 (t, J = 8Hz, 7H), 2.09-2.15 (m, 1H), 3.95 (d, J = 8Hz, 2H), 7.29-7.33 (m, 3H), 7.74-7.82 (m, 2H), 7.91 (d, J =

8Hz, 1H), 8.00 (d, $J = 8$ Hz, 1H), 8.03 (br s, 1H), 8.27 (s, 1H); LC-MS m/z data calcd. for $C_{20}H_{19}N_5O_3$ (M^+): 377.40; found (MH⁺): 378.6; **4e**: 1H NMR (DMSO- d_6): δ 5.33 (s, 2H), 7.34 (br s, 1H), 7.40 (d, $J = 8$ Hz, 1H), 7.55 (t, $J = 8$ Hz, 1H), 7.61 (t, $J = 8$ Hz, 1H), 7.68-7.73 (m, 2H), 7.77-7.83 (m, 2H), 7.87-7.91 (m, 3H), 7.99 (br s, 1H), 8.17 (d, $J = 8$ Hz, 1H), 8.82 (s, 1H); LC-MS m/z data calcd. for $C_{24}H_{16}F_3N_5O_3$ (M^+): 479.41; found (MH⁺): 480.6; **5a**: 1H NMR (DMSO- d_6): δ 1.59-1.66 (m, 2H), 1.91 (m, 3H), 2.03-2.09 (m, 2H), 2.11-2.17 (m, 2H), 6.50 (s, 1H), 7.27-7.34 (m, 2H), 7.42 (m, 1H), 7.67 (d, $J = 8$ Hz, 1H), 7.78-7.82 (m, 1H), 7.86 (d, $J = 8$ Hz, 1H), 7.95 (d, $J = 8$ Hz, 1H), 8.01 (br s, 1H), 8.25 (s, 1H); LC-MS m/z data calcd. for $C_{21}H_{19}N_5O_2S$ (M^+): 405.47; found (MH⁺): 407.0; **5b**: 1H NMR (DMSO- d_6): δ 0.83 (t, $J = 8$ Hz, 4H), 1.28-1.33 (m, 2H), 1.71-1.76 (m, 2H), 4.12 (t, $J = 8$ Hz, 2H), 7.29 (br s, 1H), 7.42 (t, $J = 8$ Hz, 1H), 7.50 (d, $J = 8$ Hz, 1H), 7.72 (d, $J = 8$ Hz, 1H), 7.85-7.92 (m, 2H), 8.03 (d, $J = 8$ Hz, 2H), 8.25 (br s, 1H); LC-MS m/z data calcd. for $C_{21}H_{19}N_5O_2S$ (M^+): 393.46; found (MH⁺): 394.6; **5c**: 1H NMR (DMSO- d_6): δ 0.88-1.08 (m, 6H), 1.48-1.61 (m, 5H), 1.87-1.92 (m, 1H), 3.94-3.97 (m, 2H), 7.29 (br s, 1H), 7.43 (t, $J = 8$ Hz, 1H), 7.51 (d, $J = 8$ Hz, 1H), 7.74 (d, $J = 8$ Hz, 1H), 7.86-7.91 (m, 2H), 8.03-8.05 (m, 2H), 8.25 (s, 1H); LC-MS m/z data calcd. for $C_{23}H_{23}N_5O_2S$ (M^+): 433.52; found (MH⁺): 434.6; **5d**: 1H NMR (DMSO- d_6): δ 0.86 (dd, $J = 8$ Hz, 28Hz, 7H), 2.15-2.22 (m, 1H), 3.89-4.00 (m, 2H), 7.29 (br s, 1H), 7.43 (t, $J = 8$ Hz, 1H), 7.51 (d, $J = 8$ Hz, 1H), 7.76 (d, $J = 8$ Hz, 1H), 7.86-7.91 (m, 2H), 8.03 (d, $J = 8$ Hz, 2H), 8.26 (br s, 1H); LC-MS m/z data calcd. for $C_{20}H_{19}N_5O_2S$ (M^+): 393.46; found (MH⁺): 394.6; **5e**: 1H NMR (DMSO- d_6): δ 5.56 (s, 2H), 6.50 (s, 1H), 7.29-7.36 (m, 3H), 7.43 (d, $J = 8$ Hz, 1H), 7.51 (t, $J = 8$ Hz, 1H), 7.58 (d, $J = 8$ Hz, 1H), 7.62 (d, $J = 8$ Hz, 1H), 7.71 (s, 1H), 7.79-7.81 (m, 2H), 7.91 (d, $J = 8$ Hz, 1H), 8.00 (br s, 1H), 8.28 (s, 1H); LC-MS m/z data calcd. for $C_{24}H_{16}F_3N_5O_2S$ (M^+): 495.48; found (MH⁺): 496.4.

53. General procedure for the 1,3-dipolar cycloaddition reaction: p-Methylbenzhydrylamine (MBHA) resin (100 mg, 1.10 meq/g, 100-200 mesh) was sealed inside a polypropylene mesh packet. Polypropylene bottles were used for all of the reactions. Resin bound amino-benzimidazoles were synthesized according to a previous literature.^{15,16} Phenylpropionic acid (4-vinylbenzoic acid) (8 eq., 0.2M in anhyd. DMF) was coupled to MBHA resin bound amino-benzimidazole for 8 h at room temperature using PyBOP (8 eq.), HOBt (8 eq.) and DIEA (8 eq.) coupling conditions. The excess solution was decanted and the resin was washed with DMF (3 times) and DCM (3 times). The resin-bound amino-benzimidazole tethered alkyne or alkene was treated with a solution of the hydroximoyl chloride (10 eq., 0.2M) in 10 mL of dry DCM and the reaction mixture was stirred overnight at 35°C. The excess solution was decanted, and the resin was washed with DCM (3 times). The resin was cleaved with HF/anisole (95:5) for 90 min at 0°C, and the desired isoxazole (isoxazoline) was obtained following extraction with 95% AcOH in H₂O and lyophilization as a colorless powder. The isoxazole (isoxazoline) was purified by preparative reverse-phase HPLC. **21a**: 1H NMR (DMSO- d_6): δ 1.75-1.80 (m, 2H), 2.04 (m, 5H), 2.25-2.33 (m, 2H), 3.45 (dd, $J = 8$ Hz, 20Hz, 1H), 3.95 (dd, $J = 8$ Hz, 20Hz, 1H), 5.35 (m, 1H), 5.80-5.85 (m, 1H), 7.30 (br s, 1H), 7.46-7.55 (m, 7H), 7.71-7.79 (m, 3H), 7.95 (br s, 1H), 8.03 (s, 1H), 8.23 (d, $J = 8$ Hz, 2H), 12.88 (br s, 1H); LC-MS m/z data calcd. for $C_{29}H_{27}N_5O_3$ (M^+): 493.55; found (MH⁺): 495.0; **21b**: 1H NMR (DMSO- d_6): δ 0.93 (t, $J = 8$ Hz, 4H), 1.32-1.38 (m, 2H), 1.76-1.83 (m, 2H), 3.45 (dd, $J = 8$ Hz, 20Hz, 1H), 3.95 (dd, $J = 8$ Hz, 20 Hz, 1H), 4.28 (t, $J = 8$ Hz, 2H), 5.80-5.85 (m, 1H), 7.29 (br s, 1H), 7.46-7.51 (m, 4H), 7.55 (d, $J = 8$ Hz, 1H), 7.71-7.76 (m, 2H), 7.79 (d, $J = 8$ Hz, 1H), 7.95 (br s, 1H), 8.01 (s, 1H), 8.25 (d, $J = 8$ Hz, 2H), 12.83 (br s, 1H); LC-MS m/z data calcd. for $C_{28}H_{27}N_5O_3$ (M^+): 481.55; found (MH⁺): 483.0; **21c**: 1H NMR (DMSO- d_6): δ 1.12-1.16 (m, 5H), 1.58-1.66 (m, 5H), 1.98 (m, 1H), 3.47 (dd, $J = 8$ Hz, 20 Hz, 1H), 3.94 (dd, $J = 8$ Hz, 20Hz, 1H), 4.22 (d, $J = 8$ Hz, 2H), 5.80-5.85 (m, 1H), 7.29 (br s, 1H), 7.46-7.56 (m, 6H), 7.72-7.77 (m, 2H), 7.78 (d, $J = 8$ Hz, 1H), 7.95 (br s, 1H), 8.01 (s, 1H), 8.25 (d, $J = 8$ Hz, 2H), 12.81 (br s, 1H); LC-MS m/z data calcd. for $C_{31}H_{31}N_5O_3$ (M^+): 521.61; found (MH⁺): 523.0; **21d**: 1H NMR (DMSO- d_6): δ 0.95 (d, $J = 8$ Hz, 7H), 2.28-2.35 (m, 1H), 3.45 (dd, $J = 8$ Hz, 20Hz, 1H), 3.95 (dd, $J = 8$ Hz, 20Hz, 1H), 4.09 (d, $J = 8$ Hz, 2H), 5.80-5.85 (m, 1H), 7.29 (br s, 1H), 7.46-7.51 (m, 4H), 7.56 (d, $J = 8$ Hz, 1H), 7.71-7.78 (m, 2H), 7.79 (d, $J = 8$ Hz, 1H), 7.96 (br s, 1H), 8.02 (s, 1H), 8.25 (d, $J = 8$ Hz, 2H), 12.81 (br s, 1H); LC-MS m/z data calcd. for $C_{28}H_{27}N_5O_3$ (M^+): 481.55; found (MH⁺): 483.0; **21e**: 1H NMR (DMSO- d_6): δ 3.44 (dd, $J = 8$ Hz, 20Hz, 1H), 3.94 (dd, $J = 8$ Hz, 20Hz, 1H), 5.60 (s, 2H), 5.80-5.85 (m, 1H), 7.30 (br s, 1H), 7.46-7.49 (m, 5H), 7.56-7.61 (m, 2H), 7.64-7.78 (m, 5H), 7.94 (br s, 1H), 7.96 (s, 1H), 8.02 (s, 1H), 8.25 (d, $J = 8$ Hz, 2H), 12.88 (br s, 1H); LC-MS m/z data calcd. for $C_{32}H_{24}F_3N_5O_3$ (M^+): 583.56; found (MH⁺): 585.0; **21f**: 1H NMR (DMSO- d_6): δ 1.75-1.77 (m, 2H), 2.05 (m, 4H), 2.27-2.33 (m, 2H), 3.45 (dd, $J = 8$ Hz,

20Hz, 1H), 3.98 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 5.33-5.35 (m, 1H), 5.83-5.87 (m, 1H), 7.29 (br s, 1H), 7.41 (t, $J = 8\text{Hz}$, 1H), 7.48-7.55 (m, 5H), 7.72-7.84 (m, 7H), 7.95 (br s, 1H), 8.03 (s, 1H), 8.23 (d, $J = 8\text{Hz}$, 2H), 12.90 (br s, 1H); LC-MS m/z data calcd. for $\text{C}_{35}\text{H}_{31}\text{N}_5\text{O}_3$ (MH^+): 569.65; found (MH^+): 571.0; **21g**: ^1H NMR ($\text{DMSO}-d_6$): δ 0.93 (t, $J = 8\text{Hz}$, 4H), 1.30-1.40 (m, 2H), 1.76-1.83 (m, 2H), 3.50 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 3.98 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 4.29 (t, $J = 8\text{Hz}$, 2H), 5.82-5.87 (m, 1H), 7.29 (br s, 1H), 7.38-7.43 (m, 1H), 7.48-7.56 (m, 4H), 7.73 (d, $J = 8\text{Hz}$, 2H), 7.77 (m, 5H), 7.95 (br s, 1H), 8.02 (s, 1H), 8.26 (d, $J = 8\text{Hz}$, 2H), 12.81 (br s, 1H); LC-MS m/z data calcd. for $\text{C}_{34}\text{H}_{31}\text{N}_5\text{O}_3$ (M^+): 557.65; found (MH^+): 559.0; **21h**: ^1H NMR ($\text{DMSO}-d_6$): δ 1.13-1.16 (m, 5H), 1.58-1.66 (m, 5H), 1.98 (m, 1H), 3.51 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 3.98 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 4.13 (d, $J = 8\text{Hz}$, 2H), 5.83-5.87 (m, 1H), 7.29 (br s, 1H), 7.38-7.43 (m, 1H), 7.48-7.53 (m, 5H), 7.73 (d, $J = 8\text{Hz}$, 2H), 7.77-7.84 (m, 5H), 7.95 (br s, 1H), 8.01 (s, 1H), 8.26 (d, $J = 8\text{Hz}$, 2H), 12.81 (br s, 1H); LC-MS m/z data calcd. for $\text{C}_{37}\text{H}_{35}\text{N}_5\text{O}_3$ (M^+): 597.71; found (MH^+): 599.0; **21i**: ^1H NMR ($\text{DMSO}-d_6$): δ 0.92 (d, $J = 8\text{Hz}$, 7H), 2.29-2.35 (m, 1H), 3.50 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 3.98 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 4.10 (d, $J = 8\text{Hz}$, 2H), 5.83-5.87 (m, 1H), 7.30 (br s, 1H), 7.38-7.43 (m, 1H), 7.48-7.53 (m, 3H), 7.57 (d, $J = 8\text{Hz}$, 1H), 7.73 (d, $J = 8\text{Hz}$, 2H), 7.77-7.84 (m, 5H), 7.95 (s, 1H), 8.02 (s, 1H), 8.26 (d, $J = 8\text{Hz}$, 2H), 12.83 (br s, 1H); LC-MS m/z data calcd. for $\text{C}_{31}\text{H}_{31}\text{N}_5\text{O}_3$ (M^+): 557.64; found (MH^+): 559.0; **21j**: ^1H NMR ($\text{DMSO}-d_6$): δ 3.48 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 3.98 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 5.62 (s, 2H), 5.83-5.87 (m, 1H), 7.31 (br s, 1H), 7.38-7.43 (m, 1H), 7.48-7.51 (m, 4H), 7.55-7.61 (m, 2H), 7.64-7.74 (m, 5H), 7.76-7.83 (m, 4H), 7.94 (br s, 1H), 7.96 (s, 1H), 8.02 (s, 1H), 8.25 (d, $J = 8\text{Hz}$, 2H), 12.90 (br s, 1H); LC-MS m/z data calcd. for $\text{C}_{38}\text{H}_{28}\text{F}_3\text{N}_5\text{O}_3$ (M^+): 659.66; found (MH^+): 661.0; **21k**: ^1H NMR ($\text{DMSO}-d_6$): δ 1.75-1.78 (m, 2H), 2.05 (m, 4H), 2.25-2.30 (m, 2H), 3.45 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 3.87-3.94 (m, 5H), 5.35 (m, 1H), 5.78-5.83 (m, 1H), 7.25 (d, $J = 8\text{Hz}$, 1H), 7.30 (br s, 1H), 7.49 (d, $J = 8\text{Hz}$, 2H), 7.54 (d, $J = 8\text{Hz}$, 1H), 7.68 (d, $J = 8\text{Hz}$, 1H), 7.76-7.79 (m, 2H), 7.96 (br s, 1H), 8.04 (s, 1H), 8.22 (d, $J = 8\text{Hz}$, 2H), 12.86 (br s, 1H); LC-MS m/z data calcd. for $\text{C}_{30}\text{H}_{29}\text{N}_5\text{O}_4$ (M^+): 523.58; found (MH^+): 525.0; **21l**: ^1H NMR ($\text{DMSO}-d_6$): δ 0.93 (t, $J = 8\text{Hz}$, 4H), 1.30-1.40 (m, 2H), 1.76-1.83 (m, 2H), 3.45 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 3.87-3.94 (m, 4H), 4.28 (t, $J = 8\text{Hz}$, 2H), 5.78-5.83 (m, 1H), 7.25 (d, $J = 8\text{Hz}$, 1H), 7.29 (br s, 1H), 7.49 (d, $J = 8\text{Hz}$, 2H), 7.56 (d, $J = 8\text{Hz}$, 1H), 7.68 (d, $J = 8\text{Hz}$, 1H), 7.77-7.81 (m, 2H), 7.95 (br s, 1H), 8.01 (s, 1H), 8.25 (d, $J = 8\text{Hz}$, 2H), 12.79 (br s, 1H); LC-MS m/z data calcd. for $\text{C}_{29}\text{H}_{29}\text{N}_5\text{O}_4$ (M^+): 511.57; found (MH^+): 513.0; **21n**: ^1H NMR ($\text{DMSO}-d_6$): δ 0.96 (d, $J = 4\text{Hz}$, 7H), 2.28-2.35 (m, 1H), 3.45 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 3.87-3.94 (m, 4H), 4.09 (d, $J = 8\text{Hz}$, 2H), 5.78-5.83 (m, 1H), 7.25 (d, $J = 8\text{Hz}$, 1H), 7.29 (br s, 1H), 7.49 (d, $J = 8\text{Hz}$, 2H), 7.57 (d, $J = 8\text{Hz}$, 1H), 7.68 (d, $J = 8\text{Hz}$, 1H), 7.77-7.80 (m, 2H), 7.95 (br s, 1H), 8.01 (s, 1H), 8.24 (d, $J = 8\text{Hz}$, 2H), 12.86 (br s, 1H); LC-MS m/z data calcd. for $\text{C}_{29}\text{H}_{29}\text{N}_5\text{O}_4$ (M^+): 511.57; found (MH^+): 513.0; **21o**: ^1H NMR ($\text{DMSO}-d_6$): δ 3.45 (dd, $J = 8\text{Hz}$, 20Hz, 1H), 3.87-3.94 (m, 3H), 5.61 (s, 2H), 5.78-5.83 (m, 1H), 7.25 (d, $J = 8\text{Hz}$, 1H), 7.30 (br s, 1H), 7.47 (d, $J = 8\text{Hz}$, 2H), 7.54-7.60 (m, 2H), 7.64-7.70 (m, 4H), 7.75-7.78 (m, 3H), 7.93 (br s, 1H), 7.96 (s, 1H), 8.02 (s, 1H), 8.25 (d, $J = 8\text{Hz}$, 2H), 12.90 (br s, 1H); LC-MS m/z data calcd. for $\text{C}_{33}\text{H}_{26}\text{F}_3\text{N}_5\text{O}_4$ (M^+): 613.58; found (MH^+): 615.0.

54. **Kinase assays:** Kinase activity was measured using a LANCE Ultra time-resolved fluorescence resonance energy transfer (TR-FRET) assay and purified recombinant IKKs (Carna Biosciences). Kinases were diluted in kinase buffer (50 mM Hepes pH 7.4, 10 mM MgCl_2 , 1 mM EGTA, 2 mM DTT, and 0.01% Tween-20) to a final concentration of 2 nM (IKK ϵ), 4 nM (TBK-1), or 1 nM (IKK2). 50 nM Ulight-rpS6 and Ulight-I κ B α (Perkin Elmer) were used as peptide substrates for IKK ϵ /TBK-1 and IKK2, respectively. All assays were performed with an ATP concentration close to the apparent K_m for each enzyme (5 μM for IKK ϵ , 10 μM for TBK-1 and 1.25 μM for IKK2). After 1 h (IKK2) or 2 h incubation (IKK ϵ /TBK-1) at room temperature, the reaction was stopped by addition of 20 mM EDTA in LANCE detection buffer, containing 2 nM Europium-labelled phospho-specific antibody (Perkin Elmer). Two hours later, the TR-FRET signals at 620 and 665 nm were measured in a CLARIOstar (BMGLabtech) multilabel reader. 665/620 ratios and delta F values were calculated with MARS data analysis program and the IC_{50} values for active compounds were determined using a 7 point titration experiment with GraphPrism.

Scheme 1.

Synthesis of quinazoline-2,4-dione, thioxoquinazolin-4-one, and benzodiazepine-2,3,5-trione derivatives: (a) 2-nitrobenzoic acid (8 eq., in anhyd. DMF), PyBOP (8 eq.), DIEA (8 eq.), 8h; (b) $\text{SnCh} \cdot 2\text{H}_2\text{O}$ (10 eq., 1.0M DMF), 24 h; (c) 1, 1'-carbonyldiimidazole (or) 1,1'-thiocarbonyldiimidazole (10 eq., 0.5 M in anhyd. DMF), 80 °C, 8h; (d) Oxalyl diimidazole (10 eq., 0.5M in anhydrous DMF), 80 °C, 8h; (e) HF/anisole (99:1), 90 min, 0 °C

Figure 1.
Biologically active isoxazoles and isoxazolines of synthetic and natural origins.

Scheme 2.

Synthesis of aminobenzimidazole tethered isoxazoles and isoxazolines: (a) Carboxylic acid (8 eq., 0.2M in anhydrous DMF), PyBOP (8 eq.), HOBT (8 eq.), DIEA (8 eq.); (b) hydroxirnoyl chloride (10 eq., in anhydrous DCM), DIEA (15 eq.) 35 °C; (c) HF/anisole (95:5), 90 min, 0 °C.

Figure 2.
IKK inhibition of Benzimidazole tethered thioxoquinazolin-4-one **5a**

Table 1

Benzimidazole tethered Quinazoline-2,4-diones, thioxoquinazolin-4-ones and benzodiazepine-2,3,5-triones.

Entry	R	Yield ^a (%)
4a	Cyclopentyl	37
4b	<i>n</i> -Butyl	40
4c	Cyclohexanemethyl	44
4d	<i>i</i> -Butyl	35
4e	3-(trifluoromethyl)benzyl	38
5a	Cyclopentyl	46
5b	<i>n</i> -Butyl	42
5c	Cyclohexanemethyl	54
5d	<i>i</i> -Butyl	40
5e	3-(trifluoromethyl)benzyl	34
6a	Cyclopentyl	51
6b	<i>n</i> -Butyl	48
6c	Cyclohexanemethyl	55
6d	<i>i</i> -Butyl	44
6e	3-(trifluoromethyl)benzyl	30

Isolated yields of aminobenzimidazole tethered quinazoline-2,4-diones, thioxoquinazolin-4-ones and benzodiazepine-2,3,5-triones: The products were run on a Vydac column, gradients 5–95% formic acid in ACN in 7 min.

^aThe yields are based on the weight of purified products and are relative to the initial loading of the resin.

Table 2

Aminobenzimidazole tethered isoxazoles and isoxazolines

Entry	R ¹	R ²	Yield ^a (%)
20a	Cyclopentyl	H	37
20b	<i>n</i> -Butyl	H	22
20c	Cyclohexanemethyl	H	33
20d	<i>i</i> -Butyl	H	24
20e	3-(trifluoromethyl)benzyl	H	36
20f	Cyclopentyl	Ph	41
20g	<i>n</i> -Butyl	Ph	33
20h	Cyclohexanemethyl	Ph	31
20i	<i>i</i> -Butyl	Ph	14
20j	3-(trifluoromethyl)benzyl	Ph	28
20k	Cyclopentyl	OMe	25
20l	<i>n</i> -Butyl	OMe	12
20m	Cyclohexanemethyl	OMe	40
20n	<i>i</i> -Butyl	OMe	15
20o	3-(trifluoromethyl)benzyl	OMe	29
21a	Cyclopentyl	H	41
21b	<i>n</i> -Butyl	H	18
21c	Cyclohexanemethyl	H	39
21d	<i>i</i> -Butyl	H	14
21e	3-(trifluoromethyl)benzyl	H	35
21f	Cyclopentyl	Ph	32
21g	<i>n</i> -Butyl	Ph	30
21h	Cyclohexanemethyl	Ph	41
21i	<i>i</i> -Butyl	Ph	17
21j	3-(trifluoromethyl)benzyl	Ph	23
21k	Cyclopentyl	OMe	24
21l	<i>n</i> -Butyl	OMe	32
21m	Cyclohexanemethyl	OMe	41
21n	<i>i</i> -Butyl	OMe	35
21o	3-(trifluoromethyl)benzyl	OMe	27

Isolated yields of aminobenzimidazole tethered isoxazoles and isoxazolines: The products were run on a Vydac column, gradients 5–95% formic acid in ACN in 7 min.

^aThe yields are based on the weight of purified products and are relative to the initial loading of the resin.