Published in final edited form as:

Biochem J. 2014 December 1; 464(2): 241–249. doi:10.1042/BJ20140787.

Cytochrome P450 System Proteins Reside in Different Regions of the Endoplasmic Reticulum

Ji Won Park, James R. Reed, Lauren M. Brignac-Huber, and Wayne L. Backes

Department of Pharmacology and Experimental Therapeutics, Stanley S. Scott Cancer Center, Louisiana State University Health Science Center, 533 Bolivar St., New Orleans, LA 70112

Abstract

Cytochrome P450 function is dependent on the ability of these enzymes to successfully interact with their redox partners, NADPH-cytochrome P450 reductase (CPR) and cytochrome b_5 , in the endoplasmic reticulum (ER). Because the ER is heterogeneous in lipid composition, membrane microdomains with different characteristics are formed. Ordered microdomains are more tightly packed, and enriched in saturated fatty acids, sphingomyelin and cholesterol, whereas disordered regions contain higher levels of unsaturated fatty acids. The goal of this study was to determine if the P450 system proteins localize to different regions of the ER. The localization of CYP1A2, CYP2B4, and CYP2E1 within the ER was determined by partial membrane solubilization with Brij 98, centrifugation on a discontinuous sucrose gradient, and immune blotting of the gradient fractions to identify ordered and disordered microdomains. CYP1A2 resided almost entirely in the ordered regions of the ER with CPR also localized predominantly to this region. CYP2B4 was equally distributed between the ordered and disordered domains. In contrast, CYP2E1 localized to the disordered membrane regions. Removal of cholesterol (an important constituent of ordered domains) led to the relocation of CYP1A2, CYP2B4 and CPR to the disordered regions. Interestingly, CYP1A1 and CYP1A2 localized to different membrane microdomains, despite their high degree of sequence similarity. These data demonstrate that P450 system enzymes are organized in specific membrane regions, and their localization can be affected by depletion of membrane cholesterol. The differential localization of different P450s in specific membrane regions may provide a novel mechanism for modulating P450 function.

Keywords

Cytochrome P450; endoplasmic reticulum; lipid microdomains; Brij 98

Corresponding author: Wayne L Backes, PhD, LSU Health Sciences Center, 533 Bolivar St., New Orleans, LA 70112, Phone-(504) 568- 6557, FAX- (504) - 568-2361, wbacke@lsuhsc.edu.

Author Contribution

Participated in Research Design: Park, Reed, Backes

Conducted Experiments: Park, Brignac-Huber

Performed Data Analysis: Park, Reed, Brignac-Huber, Backes

Wrote or contributed to the writing of the manuscript: Park, Reed, Backes

Introduction

Cytochromes P450 (P450) comprise a superfamily of heme-containing proteins, found in most organisms and catalyze oxidative reactions of both endogenous and exogenous compounds, with P450s enzymes from families CYP1, CYP2 and CYP3 being primarily involved in the metabolism of drugs [1]. Unlike bacterial P450s that are generally expressed as soluble enzymes, the mammalian forms responsible for drug and foreign compound metabolism are membrane-bound proteins that are widely expressed in the endoplasmic reticulum (ER) [2]. Even though the N-terminal region of P450 is known to be responsible for its insertion into the membrane, recent evidence suggests that amino acids in the catalytic region are also involved in membrane localization and orientation of the protein [3,4].

For functional reactions to occur, P450 enzymes must interact with other integral membrane proteins, particularly NADPH-cytochrome P450 reductase (CPR) and cytochrome b₅, which serve as electron donors [5,6]. Besides the interaction of P450 with its redox partners, many studies have shown that P450s can form complexes with other P450 enzymes in the membrane. In many cases, these complexes alter P450 function. Interactions between CYP1A2-CYP2B4, CYP1A2-CYP2E1, CYP2C9-CYP2D6, CYP2C9-CYP3A4, CYP3A4-CYP3A4 and CYP1A2-CYP1A2, have been reported [7–12]. Although a few studies have shown that these interactions occur in microsomes [13] and living cells [14–16], most of these studies were done using lipid reconstituted systems (RCS), which contain a single class of lipids that generate homogenous lipid membranes. However, biological membranes are composed of numerous types of lipids and the diversity of lipid species provides heterogeneity to the lipid bilayer. These phospholipids are not randomly distributed in the membrane, but associate in a manner to generate membrane regions with different characteristics. Some regions contain phospholipids that are enriched in unsaturated fatty acids and are referred to as liquid-disordered domains. Other regions, known as liquid-ordered domains have higher concentrations of saturated fatty acids, are enriched in cholesterol and sphingomyelin, and are more tightly packed [17]. These ordered domains are commonly referred to as lipid rafts [18], and because they are not readily solubilized by non-ionic detergent, they are also referred to as detergent-resistant membranes (DRM).

Phospholipids are pivotal components in the formation of the lipid bilayer, and are also essential for normal P450 function [19]. According to studies using mixtures of different phospholipids, alterations in membrane lipid composition were shown to affect P450 behavior. For example, several groups showed that addition of other phospholipids such as, phosphatidic acid (PA), phosphatidylinositol (PI) and phosphatidylserine (PS) into phosphatidylcholine (PC) vesicles stimulated P450 catalytic activity. These effects were attributed to the negative charges on PA, PI and PS. Although the mechanism for modulation of catalysis is not known, it has been demonstrated that the negatively charged phospholipids increase the P450 insertion rate into model membranes, and stability of the enzymes by causing conformational changes of P450 [20–23]. These effects may be mediated by a direct interaction of fatty acid chains of the phospholipids with the N-terminus or other membrane-binding regions of the P450 molecule. As ordered domains contain different phospholipid compositions, such as lower levels of PC and higher concentrations of PS, when compared to disordered regions [24], localization of P450 could

also be a significant factor affecting P450 function and their interaction with other proteins. The heterogeneity of the membranes not only might affect P450 activities directly, but also might affect their activities indirectly by either segregating or concentrating the P450s and their electron-donating partners in specific membrane regions.

Previously, our lab has examined the potential for domain formation in the ER. These experiments demonstrated that 1) ordered domains in the ER have similar lipid compositions to those found in the plasma membrane and 2) CPR and CYP1A2 selectively localize in the ordered membranes, instead of being randomly distributed throughout the ER [25]. In a follow-up study, both proteins were also shown to distribute into ordered regions when reconstituted into membranes with compositions similar to those found in the ER [26]. These studies raised a question related to whether other P450s are similarly located in the ordered regions of the ER. Therefore, we compared the localization of CYP2B4 and CYP2E1 in the ER membrane to that of CYP1A2 and CPR. In this study, we demonstrate that different P450 enzymes reside in different membrane regions, with CYP1A2 congregating predominantly in the ordered regions [25], CYP2B4 localizing in both ordered and disordered regions, and CYP2E1 segregating into the disordered membranes. The data indicate that components of the P450 system are not randomly distributed throughout the ER, but are organized in specific microdomains.

EXPERIMENTAL

Materials

Primary antibodies against CPR were obtained from Stressgen (Ann Arbor, MI). Anti-CYP2B4 and anti-CYP2E1 were purchased from Oxford Biomedical Research (Oxford, MI). The antibody for CYP1A2 was provided by Dr. Kristopher Krausz from NIH. Amplex Red cholesterol assay kit was purchased from Invitrogen (Eugene, OR). Bicinchoninic acid (BCA) protein assay kit and phenobarbital were purchased from Fisher Scientific (Rockford, IL). Methyl- β -cyclodextrin, Brij 98, pyrazole and all other reagents were obtained from Sigma-Aldrich (St. Louis, MO).

Induction of P450 and isolation of microsomes

Male New Zealand white rabbits were treated with either 80 mg/kg phenobarbital or 200 mg/kg pyrazole for induction of CYP2B4 and CYP2E1, respectively. Phenobarbital was administered by daily intraperitoneal injections for 3 days, and pyrazole was administered as a single intraperitoneal injection. Rabbits were euthanized 24 h after the final injection, the livers removed and placed into ice-cold 0.25 M sucrose. Microsomes were then prepared by differential centrifugation [27] and the final pellets were resuspended in 10 mM potassium phosphate buffer (pH 7.25). Protein concentration was determined using the BCA assay and the samples were stored as aliquots at -80°C .

Microsomal membrane solubilization by Brij 98

Detergent solubilization of microsomes was performed as described previously [25]. Briefly, Brij 98 was diluted to a 10% stock solution using membrane solubilization buffer (50 mM HEPES, pH 7.25, containing 150 mM NaCl and 5 mM EDTA and a protease inhibitor

cocktail (Roche)). The Brij 98 solution was added to the microsomal samples (2 mg/ml) to a final concentration of 1% Brij 98. Membrane solubilization buffer was added to each sample bringing the total volume to 1 ml. Samples were then incubated in a water bath at 37°C for 5 min prior to either direct centrifugation or sucrose gradient centrifugation as described.

Isolation of detergent-resistant membranes by discontinuous sucrose gradient centrifugation

Solubilized samples were mixed with 1 ml of 80% sucrose solution. A 6 ml layer of 38% sucrose was carefully placed on top followed by a 3 ml layer of 5% sucrose. The samples were centrifuged for 19 h in a SW41 rotor at $210,000 \times g$ at 4°C. After centrifugation, 1 ml aliquots were taken from the top and placed in eleven separate tubes. One ml of solubilization buffer was added to the empty tube to re-homogenize pellets. Samples were then analyzed by immune blotting. The DRM were designated as fractions 2–5 at the 5%/38% sucrose interface as described previously [25].

Isolation of detergent-resistant membranes alternatively by pelleting

For a more rapid separation of soluble proteins from those in detergent-resistant membranes, microsomes were treated, as described above, with 1% Brij at 37 °C for 5 min. Subsequently, membrane solubilization buffer was added to obtain a final volume of 3 ml. The samples were then centrifuged using a SW55Ti rotor at $290,000 \times g$ for 15 min. The supernatants were saved and pellets were resuspended in 3 ml of membrane solubilization buffer. The supernatants and pellets were then analyzed by immune blotting.

Immune blot analysis

Samples were subjected to 10% SDS-PAGE and transferred to nitrocellulose membranes (Bio-Rad). Membranes were probed using mouse anti- CYP1A1/1A2, mouse anti-rat CYP2B1, mouse anti-rat CYP2E1 or rabbit anti- rat CPR antibodies. Horseradish peroxidase-conjugated anti-rabbit IgG or anti-mouse secondary antibodies were used at 1:4000. Treated blots were visualized using Supersignal West Pico chemiluminescent substrate (Thermo Fisher Scientific). Densitometry was performed with Quantity One (Bio-Rad) or Image Lab software (Bio-Rad).

Microsomal cholesterol depletion with methyl- β -cyclodextrin (M β CD) treatment

Cholesterol depletion was performed as described previously with some modifications [25]. A 100 mM stock solution of M β CD was prepared in buffer (50 mM HEPES and 150 mM NaCl, pH 7.4). Microsomes (2 mg/ml) were incubated with or without 50 mM M β CD at 37°C for 30 min. Then, samples were centrifuged and the resulting cholesterol-depleted microsomal pellets were resuspended in 1 ml membrane solubilization buffer as described previously [25]. To confirm cholesterol depletion, 20 μ l of the resuspended pellet was diluted 10-fold and the cholesterol content of each sample was measured using the Amplex Red Cholesterol Assay Kit (Invitrogen). The samples were then treated with 1% Brij 98 and subjected to discontinuous sucrose gradient centrifugation.

Results

Microdomain localization of CYP2B4 in the ER of phenobarbital-treated rabbits

As noted from our previous studies, CPR and CYP1A2 were found to localize in “ordered” regions of the ER that contain higher concentrations of cholesterol and sphingomyelin. This raised the question of whether all P450 proteins similarly localize into these microdomains, and led to the current experiments examining the localization of CYP2B4 and CYP2E1. To examine microdomain localization of CYP2B4 proteins, liver microsomes isolated from phenobarbital-treated rabbits were subjected to detergent treatment with 1% Brij 98, followed by discontinuous sucrose gradient centrifugation. After centrifugation, 1 ml fractions were collected from the top of the tube, and each fraction was analyzed for cholesterol and protein content. Because of their high lipid/protein ratio, membranes containing ordered domains float within the low-density sucrose fractions [28]. Fractions between 2 to 5 possessed the highest cholesterol/protein ratios (supplementary Fig. 1A) as reported previously for ER from untreated rabbits [25]. These fractions correspond to the detergent-resistant (ordered) membrane regions. To analyze protein localization, each fraction was subjected to SDS-PAGE and visualized by immune blotting. The majority of CYP1A2 and CPR proteins were found in the low density sucrose fractions, which is consistent with our previous results from microsomes isolated from untreated rabbits showing their localization in the ordered DRM membranes [25]. However, unlike the distribution of CYP1A2 and CPR, CYP2B4 appeared to reside in both the ordered and disordered microdomains (Fig. 1). These differences appeared to be a characteristic of the proteins, as the relative composition of lipid classes (PC, phosphatidylethanolamine, PI, PS, PA, sphingomyelin and cholesterol) were not affected by PB treatment (unpublished and [25]). Although it has been reported that amino acid motifs and posttranslational modifications, such as glycosylphosphatidylinositol (GPI) anchor and palmitoylation are involved in ordered domain localization [29,30], it has not been examined with P450s. Therefore, further study is required to identify the factors responsible for localization of P450s in ordered domains.

Distribution and differential detergent solubility of microsomal proteins from pyrazole-treated rabbits

Next, CYP2E1 localization was examined. Since the CYP2E1 expression level is low in untreated and phenobarbital-treated rabbits, pyrazole was used to enhance CYP2E1 levels. The increased expression level of CYP2E1 was confirmed by immune blot (data not shown). As described for phenobarbital-treated rabbits, ordered domains were isolated by detergent treatment of pyrazole microsomes followed by sucrose gradient centrifugation. Pyrazole microsomes were similar in the cholesterol/protein ratio to PB microsomes (supplementary Fig. 1B). CYP2E1 composition from each fraction was analyzed by immune blotting and a dramatic difference was observed in the distribution of this P450. CYP2E1 was exclusively found in high-density sucrose fractions indicative of their presence in the detergent soluble “disordered” domains whereas CYP1A2 and CPR were mainly localized in the ordered domains (fractions 2–5, Fig. 2).

Because disordered domains are less tightly packed than ordered domains, we hypothesized that proteins segregating primarily in disordered domains would be more easily released from the microsomal membrane with lower concentrations of Brij 98. Therefore, the solubilities of CYP1A2, CPR and CYP2E1 were compared with varying concentrations of Brij 98 (0, 0.125, 0.25 and 1%) in pyrazole-treated microsomes. As expected, CYP2E1 was partially solubilized by 0.125% Brij 98 and was completely solubilized at 1% Brij (Fig. 3). In contrast, both CYP1A2 and CPR showed solubility profiles that differed from CYP2E1 – in the Brij 98 concentration range from 0.25 to 1% there was little solubilization of these proteins from their membrane environment. These data demonstrate that different P450 system enzymes localize in different membrane microdomains.

Confirmation of different localization of P450s by pelleting

While high-speed centrifugation with sucrose gradients make insoluble fractions float to the interface between the 5 and 38% sucrose layers, pelleting (high-speed centrifugation without sucrose) pulls down insoluble membrane fractions to the bottom of centrifuge tubes [28]. Using this method, the supernatant contains the solubilized proteins. This method has the advantage of more rapidly separating the ordered and disordered regions (15 min versus 19 hr). In order to validate this methodology, 1% Brij 98-treated samples were separated by ultracentrifugation and the results compared to the sucrose gradient method. After the ultracentrifugation of microsomal samples, Brij 98 treatment caused CYP2B4 and CPR to distribute into both pellet and supernatant, whereas most of CYP1A2 remained in the pellet (Fig. 4). In contrast, CYP2E1 was completely solubilized by Brij 98 and was found only in the supernatant of the samples (Fig. 4). These results are consistent with the previous experiments utilizing the discontinuous sucrose gradient, and confirmed that different P450s localize into distinct microdomains.

We obtained a particularly interesting result when examining CYP1A protein distribution (Fig. 4). Both CYP1A1 and CYP1A2 are expressed in rabbit liver [31], and can be seen in the figure with the upper and lower bands representing CYP1A1 and CYP1A2, respectively. Despite their high level of sequence similarity, these proteins showed contrasting results after microsomal treatment with Brij 98. Whereas CYP1A2 proteins localized in the ordered membrane regions (found in the pellet after centrifugation), a significant portion of CYP1A1 was found in the disordered regions (i.e. the supernatant) (Fig. 4), suggesting that subtle differences in the protein sequence can influence P450 localization.

CYP2B4 localization was altered by cholesterol depletion

Cholesterol and sphingomyelin are known as important constituents leading to the formation of ordered microdomains. Therefore, having demonstrated that CYP2B4 can reside in these ordered regions, we tested whether depletion of cholesterol would disturb the distribution of DRM-localized proteins in a manner similar to that observed with CYP1A2 and CPR [25]. For cholesterol depletion, PB microsomes were treated with methyl- β -cyclodextrin (M β CD) that selectively extracts cholesterol from membranes via a binding pocket on the molecule [32]. After incubation at 37°C for 30 min, M β CD-treated microsomes were centrifuged to remove extracted cholesterol. Cholesterol contents of resuspended microsomes after M β CD treatment and centrifugation were decreased by almost 80% (Fig. 5A). Next, the microsomes

were solubilized by 1 % Brij 98 and applied to a sucrose gradient and centrifuged. Protein compositions of each fraction were then analyzed by immune blot. The results showed that proteins in DRM fractions were shifted to fractions of higher sucrose density after M β CD treatment (Fig. 5B and 5C), consistent with disruption of the ordered domains that contain CYP2B4, CPR and CYP1A2. Interestingly, the susceptibility to detergent solubilization after M β CD pre-treatment differed among the proteins. In Table 1, averages of relative distribution (%) of proteins with or without cholesterol depletion were calculated from sucrose gradient fractions 2 to 5. For example, with microsomes from phenobarbital-treated animals, disruption of the DRM caused the release of more than 60% of the CPR; however, only about 45% of the DRM-localized CYP1A2, and 40% of CYP2B4 were released by M β CD/detergent treatment (Table 1). These results suggest that *at least some* of the P450s located in the ordered domains may not be co-localized in the same ordered regions of the membrane.

Non-DRM localization of CYP2E1 is independent of cholesterol depletion

As reported in figure 4, the distribution of CPR, CYP1A2, and CYP2B4 enzymes in ordered fractions was affected by cholesterol depletion from the ER membrane. Conversely, because CYP2E1 appeared to reside in the less ordered membrane regions, its distribution should not be altered by M β CD pretreatment. Therefore, pyrazole-treated microsomes were subjected to M β CD/detergent and the localization of CPR, CYP1A2 and CYP2E1 were determined after sucrose gradient centrifugation. M β CD treatment decreased cholesterol content by more than 80% (Fig. 6A) and shifted DRM localized proteins (CYP1A2 and CPR) to later higher-density fractions (Fig. 6B and 6C). In contrast, most of CYP2E1 was found in high density sucrose fractions without M β CD treatment, and this distribution was not affected by the removal of cholesterol by M β CD (Fig. 6B and 6C). These results clearly demonstrate that CYP2E1 localization was not affected by removal of cholesterol, which is consistent with it residing in the disordered membrane regions.

Consistent with the results shown with PB microsomes, differences in the release of the proteins that reside in the ordered microdomains (CPR and CYP1A2) were observed after M β CD treatment with microsomes from pyrazole-treated rabbits. Whereas greater than 75% of the CPR was released from the ordered membranes, M β CD only led to the release of 40% of CYP1A2 (Table 1). Again, the results are consistent with the idea that even though multiple proteins can exist within the ordered microdomains, the microdomains are heterogeneous with regard to protein composition.

Discussion

It has been established that the biological lipid bilayer is composed of numerous types of lipid that are heterogeneously distributed to form ordered and disordered microdomains. The ordered domains in cellular membranes have been reported to play important roles in cellular signaling and transport of lipids and proteins [33–35]. Because of the complexity of cellular membranes and technical limitations related to their isolation, it is difficult to directly visualize ordered domains in biological membranes. However, the organization of microdomains in model membranes has been well characterized and visualized [36,37].

Furthermore, the lipid domains isolated by detergent treatment in living cells showed similar properties to liquid ordered domains from model membranes [38]. Microdomains from cell membranes have mostly been characterized from the plasma membrane where the cholesterol and sphingomyelin compositions range from 20–30% [24]. In contrast, the cholesterol/phospholipid ratios in other organelles range from 5–20% [39]. The cholesterol and sphingomyelin compositions of the ER are particularly low – with both being in the 4–5% range [25,40]. The low concentration of cholesterol in the ER raised the question as to whether this organelle contains cholesterol-rich lipid microdomains. However, several groups have shown the potential for the formation of ordered domains within the ER membrane, as well as in other organelles such as golgi, mitochondria and late endosomes [25,41–44]. Our laboratory did show that DRM from the ER are enriched in cholesterol [25], a result that also is borne out from the current study (Supplemental Fig. 1).

Even though cholesterol is a key component responsible for the formation/stability of ordered domains found in the plasma membrane [45], some raft marker proteins including glycosylphosphatidylinositol (GPI)-anchored proteins, galectin-4 and flotillins are not affected by cholesterol depletion while other proteins in ordered domains are released from the membrane [46–48]. These findings suggest that there is heterogeneity within the population of ordered domains and the requirement of cholesterol in the formation/stability of ordered domains is different among microdomains [49]. This may explain why CPR distribution was most disturbed by M β CD treatment among proteins in ordered domains in Table 1, and reflect the existence of different types of DRM. In other words, even though some proteins can be found in the same fractions of the discontinuous sucrose gradient, it does not guarantee that those proteins co-localize in the same microdomains.

Despite evidence that the ER membrane contains microdomains [28,41] and that membrane lipid composition affects P450 function [20,22,23], little is known about whether P450 system components reside in different regions of the ER membrane and if this localization influences P450 function. Previously, our laboratory isolated ordered domains in rabbit microsomes by Brij 98 solubilization and sucrose gradient centrifugation and showed higher concentrations of cholesterol and sphingomyelin in the ordered domains [25]. The study showed that CYP1A2 and CPR proteins were predominantly localized to ordered regions of the microsomal membrane. In addition, when purified CYP1A2 and CPR were incorporated into reconstituted systems, vesicles containing higher levels of cholesterol and sphingomyelin were associated with an increase in CYP1A2-mediated substrate metabolism and a decrease in apparent K_D for CPR in comparison to PC vesicles. The data suggested that increased CYP1A2 activity in the vesicles containing higher levels of cholesterol and sphingomyelin was mediated by increasing the efficiency of interaction of CYP1A2 and CPR [25]. It is currently unknown how much of this change in apparent K_D^{CPR} is due to concentration of the proteins into a smaller surface area (i.e. the ordered microdomains), and how much is a direct effect of the anionic phospholipids on protein activity.

These data led us to investigate the localization of other P450s, CYP2B4 and CYP2E1 in the ER. Unlike CYP1A2 and CPR, CYP2B4 was equally distributed between the ordered and disordered domains and CYP2E1 was almost exclusively distributed in disordered domains (Figs. 1 and 2). With CPR residing predominantly in the ordered regions, interactions with

CYP1A2 and the fraction of CYP2B4 localized in the ordered regions would be facilitated. However, interactions with CYP2E1 might be limited by its segregation from CPR into different domains. CYP2E1 is known to generate significant amounts of hydrogen peroxide due to its high degree of uncoupling. Therefore, segregation of CPR and CYP2E1 in different microdomains might serve the physiological function of limiting the generation of this damaging reactive oxygen species.

The current study is the first to show the differential localization of P450 system components into different membrane regions. It is well established that some signal transduction pathways, such as immune receptor signaling, work efficiently through specific microdomains [29,50]. These microdomains can bring proteins together to enhance their ability to interact, or segregate the proteins into different regions to limit their interaction. In like manner, the presence of microdomains in the ER could affect P450 function by influencing the ability of particular P450 enzymes to interact with CPR. Although the ratio of CPR to P450 is variable in different tissues, in most tissues such as the liver the CPR:P450 ratio ranges from 1:5 to 1:30 [12]. The co-localization of CPR and a P450 (e.g. CPR and CYP1A2) would be expected to increase their relative concentrations within the microdomain and enhance their ability to interact. In contrast, segregation of proteins into different microdomains, as shown with CPR and CYP2E1, would be expected to diminish activities. Consequently, for CYP2E1 to function in this system, a mechanism for complex formation is required. Likely possibilities involve (1) the translocation of CYP2E1 to the ordered lipid domains due to some stimulus such as the presence of substrate, (2) the transfer of CPR to the disordered domains, or (3) the interaction of CYP2E1 with the small amount of CPR that already resides in the disordered region, a condition that would limit its activity. Therefore, further study is required to determine if stimuli such as substrate or NADPH induce translocation of P450s and the reductase. Protein localization will be expected not only to affect formation of the functional CPR•P450 complex, but also to influence interactions with other protein partners such as P450s and cytochrome b₅. Several studies have reported that P450 enzymes are capable of both homomeric [11,14,16] and heteromeric interactions [7–10,12], which result in altered P450 activities. However, many of these studies were done in reconstituted systems containing a single lipid type. The results from the current study showing localization of P450s into different microdomains in the ER can either limit or stimulate these potential interactions. As an example, our lab has previously reported that CYP1A2 and CYP2E1 interact in reconstituted systems [7]. The segregation of CYP2E1 and CYP1A2 into disordered and ordered domains, respectively, has the potential to significantly limit the ability of these proteins to form CYP1A2•CYP2E1 complexes.

The sensitivity for localization of a protein into ordered or disordered domains is exemplified by the comparison of CYP1A1 and CYP1A2. Even though these proteins showed about 80% amino acid sequence similarity, they produced contrasting results regarding their regional localization, with CYP1A1 residing primarily in the disordered regions and CYP1A2 being found in the ordered microdomains (Fig. 4). When comparing the amino acid sequences of these two proteins, most variations were found in early N-terminus and catalytic regions. Unlike conserved subcellular targeting motifs, such as KDEL for ER localization, very little is known about microdomain targeting motifs within the

subcellular compartments. These data suggest that specific sequence motifs may be responsible for targeting CYP1A proteins into ordered or disordered microdomains. A similar explanation could be provided for the pan-domain distribution of CYP2B4. In this case, a subtle change in the sequence of the putative domain-targeting motif could lead to incomplete localization of the protein into ordered microdomains. Studies to identify the protein sequences responsible for their selective incorporation into membrane microdomains are currently ongoing.

In conclusion, the data shown here demonstrate that different forms of P450s are organized in specific membrane regions of the ER. The differential localization of different P450s may provide a novel mechanism for modulating P450 function. Future study will be required to determine if there are changes in dynamic movement of the proteins with stimuli and under particular physiological or pathological conditions modulating membrane lipid composition.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding

These studies were supported by Grants ES004344 and ES013648 from the National Institute of Environmental Health Sciences.

Abbreviations

P450	Cytochrome P450
DRM	detergent-resistant membrane
ER	endoplasmic reticulum
MβCD	methyl- β -cyclodextrin
CPR	NADPH-cytochrome P450 reductase
PC	phosphatidylcholine
PS	phosphatidylserine
PI	phosphatidylinositol
PA	phosphatidic acid
RCS	reconstituted systems

References

1. Guengerich FP. Cytochromes P450, drugs, and diseases. *Mol Interv.* 2003; 3:194–204. [PubMed: 14993447]
2. Werck-Reichhart D, Feyereisen R. Cytochromes P450: a success story. *Genome Biol.* 2000; 1:Reviews3003.1–3003.9. [PubMed: 11178272]

3. Hu G, Johnson EF, Kemper B. CYP2C8 exists as a dimer in natural membranes. *Drug Metab Dispos.* 2010; 38:1976–1983. [PubMed: 20699412]
4. Cojocaru V, Balali-Mood K, Sansom MS, Wade RC. Structure and dynamics of the membrane-bound cytochrome P450 2C9. *PLoS Comput Biol.* 2011; 7:e1002152. [PubMed: 21852944]
5. Gigon PL, Gram TE, Gillette JR. Studies on the rate of reduction of hepatic microsomal cytochrome P-450 by reduced nicotinamide adenine dinucleotide phosphate: effect of drug substrates. *Mol Pharmacol.* 1969; 5:109–122. [PubMed: 4389181]
6. Porter TD. The roles of cytochrome b5 in cytochrome P450 reactions. *J Biochem Mol Toxicol.* 2002; 16:311–316. [PubMed: 12481306]
7. Kelley RW, Cheng D, Backes WL. Heteromeric Complex Formation between CYP2E1 and CYP1A2: Evidence for the Involvement of Electrostatic Interactions. *Biochem.* 2006; 45:15807–15816. [PubMed: 17176103]
8. Reed JR, Eyer M, Backes WL. Functional interactions between cytochromes P450 1A2 and 2B4 require both enzymes to reside in the same phospholipid vesicle: evidence for physical complex formation. *J Biol Chem.* 2010; 285:8942–8952. [PubMed: 20071338]
9. Subramanian M, Low M, Locuson CW, Tracy TS. CYP2D6-CYP2C9 protein-protein interactions and isoform-selective effects on substrate binding and catalysis. *Drug Metab Dispos.* 2009; 37:1682–1689. [PubMed: 19448135]
10. Subramanian M, Zhang H, Tracy TS. CYP2C9-CYP3A4 protein-protein interactions in a reconstituted expressed enzyme system. *Drug Metab Dispos.* 2010; 38:1003–1009. [PubMed: 20215413]
11. Davydov DR, Sineva EV, Sistla S, Davydova NY, Frank DJ, Sligar SG, Halpert JR. Electron transfer in the complex of membrane-bound human cytochrome P450 3A4 with the flavin domain of P450BM-3: the effect of oligomerization of the heme protein and intermittent modulation of the spin equilibrium. *Biochim Biophys Acta.* 2010; 1797:378–390. [PubMed: 20026040]
12. Reed JR, Backes WL. Formation of P450. P450 complexes and their effect on P450 function. *Pharmacol Ther.* 2012; 133:299–310. [PubMed: 22155419]
13. Cawley GF, Zhang S, Kelley RW, Backes WL. Evidence supporting the interaction of CYP2B4 and CYP1A2 in microsomal preparations. *Drug Metab Dispos.* 2001; 29:1529–1534. [PubMed: 11717170]
14. Reed JR, Connick JP, Cheng D, Cawley GF, Backes WL. Effect of Homomeric P450•P450 Complexes on P450 Function. *Biochem J.* 2012; 446:489–497. [PubMed: 22738171]
15. Ozalp C, Szczesna-Skorupa E, Kemper B. Bimolecular fluorescence complementation analysis of cytochrome p450 2c2, 2e1, and NADPH-cytochrome p450 reductase molecular interactions in living cells. *Drug Metab Dispos.* 2005; 33:1382–1390. [PubMed: 15980100]
16. Szczesna-Skorupa E, Mallah B, Kemper B. Fluorescence Resonance Energy Transfer Analysis of Cytochromes P450 2C2 and 2E1 Molecular Interactions in Living Cells. *J Biol Chem.* 2003; 278:31269–31276. [PubMed: 12766165]
17. London E, Brown DA. Insolubility of lipids in triton X-100: physical origin and relationship to sphingolipid/cholesterol membrane domains (rafts). *Biochim Biophys Acta.* 2000; 1508:182–195. [PubMed: 11090825]
18. Rajendran L, Simons K. Lipid rafts and membrane dynamics. *J Cell Sci.* 2005; 118:1099–1102. [PubMed: 15764592]
19. Strobel HW, Lu AYH, Heidema J, Coon MJ. Phosphatidylcholine requirement in the enzymatic reduction of hemoprotein P-450 and in fatty acid, hydrocarbon, and drug hydroxylation. *J Biol Chem.* 1970; 245:4851–4854. [PubMed: 4393962]
20. Ingelman-Sundberg M, Hagbjörk AL, Ueng YF, Yamazaki H, Guengerich FP. High rates of substrate hydroxylation by human cytochrome p450 3A4 in reconstituted membranous vesicles: Influence of membrane charge. *Biochem Biophys Res Commun.* 1996; 221:318–322. [PubMed: 8619853]
21. Ahn T, Guengerich FP, Yun CH. Membrane insertion of cytochrome P450 1A2 promoted by anionic phospholipids. *Biochem.* 1998; 37:12860–12866. [PubMed: 9737864]

22. Cho EY, Yun CH, Chae HZ, Chae HJ, Ahn T. Anionic phospholipid-induced regulation of reactive oxygen species production by human cytochrome P450 2E1. *FEBS Lett.* 2008; 582:1771–1776. [PubMed: 18472009]
23. Jang HH, Kim DH, Ahn T, Yun CH. Functional and conformational modulation of human cytochrome P450 1B1 by anionic phospholipids. *Arch Biochem Biophys.* 2010; 493:143–150. [PubMed: 19857456]
24. Pike LJ, Han X, Chung KN, Gross RW. Lipid rafts are enriched in arachidonic acid and plasmalogen phospholipids and their composition is independent of caveolin-1 expression: a quantitative electrospray ionization/mass spectrometric analysis. *Biochem.* 2002; 41:2075–2088. [PubMed: 11827555]
25. Brignac-Huber LM, Reed JR, Backes WL. Organization of NADPH-cytochrome P450 reductase and CYP1A2 in the endoplasmic reticulum – Microdomain localization affects monooxygenase function. *Mol Pharmacol.* 2011; 79:549–557. [PubMed: 21156755]
26. Brignac-Huber LM, Reed JR, Eyer MK, Backes WL. Relationship between CYP1A2 localization and lipid microdomain formation as a function of lipid composition. *Drug Metab Dispos.* 2013; 41:1896–1905. [PubMed: 23963955]
27. Sequeira DJ, Cawley GF, Eyer CS, Backes WL. Temporal changes in P-450 2E1 expression with continued ethylbenzene exposure. *Biochim Biophys Acta.* 1994; 1207:179–186. [PubMed: 8075152]
28. Pielsticker LK, Mann KJ, Lin WL, Sevlever D. Raft-like membrane domains contain enzymatic activities involved in the synthesis of mammalian glycosylphosphatidylinositol anchor intermediates. *Biochem Biophys Res Commun.* 2005; 330:163–171. [PubMed: 15781246]
29. Webb Y, Hermida-Matsumoto L, Resh MD. Inhibition of protein palmitoylation, raft localization, and T cell signaling by 2-bromopalmitate and polyunsaturated fatty acids. *J Biol Chem.* 2000; 275:261–270. [PubMed: 10617614]
30. Legler DF, Doucey MA, Schneider P, Chapatte L, Bender FC, Bron C. Differential insertion of GPI-anchored GFPs into lipid rafts of live cells. *FASEB J.* 2005; 19:73–75. [PubMed: 15516372]
31. Pineau T, Daujat M, Pichard L, Girard F, Angevain J, Bonfils C, Maurel P. Developmental expression of rabbit cytochrome P450 CYP1A1, CYP1A2 and CYP3A6 genes. Effect of weaning and rifampicin. *Eur J Biochem.* 1991; 197:145–153. [PubMed: 2015817]
32. Taylor CM, Coetzee T, Pfeiffer SE. Detergent-insoluble glycosphingolipid/cholesterol microdomains of the myelin membrane. *J Neurochem.* 2002; 81:993–1004. [PubMed: 12065611]
33. Fessler MB, Parks JS. Intracellular lipid flux and membrane microdomains as organizing principles in inflammatory cell signaling. *J Immunol.* 2011; 187:1529–1535. [PubMed: 21810617]
34. Colin D, Limagne E, Jeanningros S, Jacquel A, Lizard G, Athias A, Gambert P, Hichami A, Latruffe N, Solary E, Delmas D. Endocytosis of resveratrol via lipid rafts and activation of downstream signaling pathways in cancer cells. *Cancer Prev Res(Phila).* 2011; 4:1095–1106. [PubMed: 21467134]
35. Brown DA, London E. Functions of lipid rafts in biological membranes. *Annu Rev Cell Dev Biol.* 1998; 14:111–136. [PubMed: 9891780]
36. Chiantia S, Kahya N, Ries J, Schwille P. Effects of ceramide on liquid-ordered domains investigated by simultaneous AFM and FCS. *Biophys J.* 2006; 90:4500–4508. [PubMed: 16565041]
37. Veatch W, Stryer L. The dimeric nature of the gramicidin A transmembrane channel: conductance and fluorescence energy transfer studies of hybrid channels. *J Mol Biol.* 1977; 113:89–102. [PubMed: 69713]
38. Brown DA, London E. Structure and function of sphingolipid- and cholesterol-rich membrane rafts. *J Biol Chem.* 2000; 275:17221–17224. [PubMed: 10770957]
39. van Meer G, Voelker DR, Feigenson GW. Membrane lipids: where they are and how they behave. *Nat Rev Mol Cell Biol.* 2008; 9:112–124. [PubMed: 18216768]
40. Glaumann H, Dallner G. Lipid composition and turnover of rough and smooth microsomal membranes in rat liver. *J Lipid Res.* 1968; 9:720–729. [PubMed: 5685264]

41. Browman DT, Resek ME, Zajchowski LD, Robbins SM. Erlin-1 and erlin-2 are novel members of the prohibitin family of proteins that define lipid-raft-like domains of the ER. *J Cell Sci.* 2006; 119:3149–3160. [PubMed: 16835267]
42. Babiychuk EB, Draeger A. Biochemical characterization of detergent-resistant membranes: a systematic approach. *Biochem J.* 2006; 397:407–416. [PubMed: 16608442]
43. Sobo K, Chevallier J, Parton RG, Gruenberg J, van der Goot FG. Diversity of raft-like domains in late endosomes. *PLoS One.* 2007; 2:e391. [PubMed: 17460758]
44. Poston CN, Duong E, Cao Y, Bazemore-Walker CR. Proteomic analysis of lipid raft-enriched membranes isolated from internal organelles. *Biochem Biophys Res Commun.* 2011; 415:355–360. [PubMed: 22037461]
45. Silvius JR. Role of cholesterol in lipid raft formation: lessons from lipid model systems. *Biochim Biophys Acta.* 2003; 1610:174–183. [PubMed: 12648772]
46. Schroeder RJ, Ahmed SN, Zhu Y, London E, Brown DA. Cholesterol and sphingolipid enhance the Triton X-100 insolubility of glycosylphosphatidylinositol-anchored proteins by promoting the formation of detergent-insoluble ordered membrane domains. *J Biol Chem.* 1998; 273:1150–1157. [PubMed: 9422781]
47. Rajendran L, Masilamani M, Solomon S, Tikkanen R, Stuermer CA, Plattner H, Illges H. Asymmetric localization of flotillins/reggies in preassembled platforms confers inherent polarity to hematopoietic cells. *Proc Natl Acad Sci USA.* 2003; 100:8241–8246. [PubMed: 12826615]
48. Hansen GH, Immerdal L, Thorsen E, Niels-Christiansen LL, Nystrom BT, Demant EJ, Danielsen EM. Lipid rafts exist as stable cholesterol-independent microdomains in the brush border membrane of enterocytes. *J Biol Chem.* 2001; 276:32338–32344. [PubMed: 11389144]
49. Pike LJ. Lipid rafts: heterogeneity on the high seas. *Biochem J.* 2004; 378:281–292. [PubMed: 14662007]
50. Dykstra M, Cherukuri A, Pierce SK. Rafts and synapses in the spatial organization of immune cell signaling receptors. *J Leukoc Biol.* 2001; 70:699–707. [PubMed: 11698488]

Summary Statement

The endoplasmic reticulum (ER) is heterogeneous in nature. It is comprised of disordered regions containing unsaturated phospholipids, and ordered microdomains containing saturated phospholipids, cholesterol and sphingomyelin. P450 enzymes are not randomly distributed, but reside in specific regions of the ER.

Figure 1. Distribution of CYP2B4 between ordered and disordered membrane regions

Microsomes from phenobarbital-treated rabbit liver (2 mg/ml) were incubated with 1% Brij 98 at 37°C for 5 min, followed by sucrose gradient centrifugation. Eleven fractions (1 ml each) were collected from the top, and samples from each fraction were subjected to SDS-PAGE. The localization of CYP2B4, CYP1A2 and CPR was shown by immune blotting (A). Relative distribution (%) of proteins was measured by densitometry (B–D). The results are representative of at least three independent experiments.

Figure 2. Distribution of CYP2E1 between ordered and disordered membrane regions

Microsomes from pyrazole-treated rabbit liver (2 mg/ml) were separated by sucrose gradient centrifugation after being solubilized by 1% Brij 98 at 37°C for 5 min. Protein distribution of collected fractions was analyzed by immune blotting (A). Relative distribution (%) of proteins was measured by densitometry (B–D).

Figure 3. Differential solubility of DRM- and non-DRM- localized proteins by varying concentrations of Brij 98

Pyrazole-induced microsomes (2 mg/ml) were solubilized with different concentrations (0, 0.125, 0.25, and 1%) of Brij 98 for 5 min and DRMs were isolated via sucrose gradient centrifugation. Samples from each fraction were analyzed by immune blotting, probing for CYP2E1 (A), CYP1A2 (B), and CPR (C).

Figure 4. Isolation of DRM- and non-DRM- proteins by high-speed centrifugation

Phenobarbital- or pyrazole-induced liver microsomes were solubilized by 1% Brij 98 and then diluted 3-fold with 50 mM HEPES, 150 mM NaCl and 5 mM EDTA with protease inhibitor cocktail (Roche), pH 7.4. Diluted microsomes were separated by centrifugation using an SW55Ti rotor at $290,000 \times g$ for 15 min into supernatant (S) and pellet (P). The supernatant was saved and the precipitate was resuspended in same volume as the supernatant. P450s and CPR were visualized by immune blotting.

Figure 5. Effect of cholesterol depletion on P450 localization in phenobarbital-treated microsomes

Microsomes (2 mg/ml) were incubated in the presence or absence of 50 mM methyl- β -cyclodextrin (M β CD) at 37°C for 30 min. Microsomal samples were centrifuged at $20,000 \times g$ for 20 min and the pellet was resuspended in solubilization buffer. (A) Cholesterol contents were assayed by using Amplex Red Cholesterol assay kit. Resuspended microsomes were treated with 1% Brij 98 at 37°C for 5 min and subjected to sucrose gradient centrifugation. Protein composition of each fraction was analyzed by immune blotting (B) and percentages of protein composition were estimated by densitometric analysis (C).

Figure 6. Effect of cholesterol depletion on P450 distribution in pyrazole-treated microsomes
 Microsomes were treated with or without 50 mM MβCD at 37°C for 30 min. Resuspended microsomes, following centrifugation at $20,000 \times g$ for 20 min were used to determine cholesterol contents (A) and subjected to sucrose gradient centrifugation after 1% Brij 98 treatment at 37°C for 5 min. Protein concentrations in each fraction were analyzed by immune blotting (B) and densitometry (C).

Table 1
Differential detergent solubility of P450s and CPR after cholesterol depletion

Rabbit liver microsomes were treated with M β CD to remove cholesterol from the membranes, or with buffer as a control. After treatment, the samples were treated with 1% Brij 98 to solubilize the membranes. The amount of protein remaining in the more ordered, detergent-resistant membranes was identified as those proteins remaining in fractions 2–5 after discontinuous sucrose gradient centrifugation as described in Materials and Methods.

	% remaining in ordered membrane regions	
	Phenobarbital-pretreated microsomes	Pyrazole-pretreated microsomes
	No treatment	50 mM M β CD
CYP2B4	32.3 $\pm$ 1.8	19.6 $\pm$ 1.3 [*]
CYP1A2	84.9 $\pm$ 9.2	44.7 $\pm$ 3.7 [*]
CPR	66.0 $\pm$ 4.3	23.7 $\pm$ 1.9 ^{**}
	No treatment	50 mM M β CD
CYP2E1	14.0 $\pm$ 1.5	10.6 $\pm$ 0.8
CYP1A2	85.6 $\pm$ 5.4	50.0 $\pm$ 6.7 ^{**}
CPR	75.1 $\pm$ 2.7	17.6 $\pm$ 4.1 ^{**}

These data represent the mean $\pm$ S.E.M. for three experiments using a paired t-test –

* p < 0.05;

** p < 0.01.