Off-Label Drug Uses

Rifaximin: Irritable Bowel Syndrome

Joyce A. Generali, RPh, MS, FASHP (Editor),* and Dennis J. Cada, PharmD, FASHP, FASCP†

This *Hospital Pharmacy* feature is extracted from *Off-Label Drug Facts*, a publication available from Wolters Kluwer Health. *Off-Label Drug Facts* is a practitioner-oriented resource for information about specific drug uses that are unapproved by the US Food and Drug Administration. This new guide to the literature enables the health care professional or clinician to quickly identify published studies on off-label uses and determine if a specific use is rational in a patient care scenario. References direct the reader to the full literature for more comprehensive information before patient care decisions are made. Direct questions or comments regarding *Off-Label Drug Uses* to jgeneral@ku.edu.

BACKGROUND

Irritable bowel syndrome (IBS) is a chronic gastrointestinal (GI) disorder associated with altered bowel habits and abdominal pain or discomfort in the absence of organic disease. Subtypes include IBS with constipation (IBS-C), IBS with diarrhea (IBS-D), mixed IBS (IBS-M), and unclassified IBS (IBS-U).¹ Although symptoms vary among subtypes and patients, most efforts in the management of IBS are directed at symptom control and have included dietary modification, physical activity, and adjunctive pharmacological therapy, which is typically reserved for patients with moderate to severe symptoms that impair quality of life. Use of antibiotics, including rifaximin, is not routinely recommended in the management of IBS and is typically reserved for patients with moderate to severe IBS without constipation who have failed to respond to other therapies. Recommendations for antibiotic use are based on the proposed relationship between small intestine microbial overgrowth and malabsorption or excessive gas production.^{2–4}

PATIENT POPULATION

Adults with signs and symptoms of IBS.

DOSAGE AND DURATION

550 mg 3 times daily for 2 weeks.⁵

RESULTS

Rifaximin for the management of IBS has been studied in several placebo-controlled trials (enrolling more than 1,500 treated patients) demonstrating modest effect on global IBS symptoms and benefit in bloating in IBS patients without constipation.^{5,6} Most trials included a short-term follow-up (less than 3 months) after therapy completion. An American College of Gastroenterology (ACG) monograph on the management of IBS provides a weak recommendation for rifaximin use, based on moderate evidence indicating modest but consistent efficacy in IBS without constipation.

Position Statement

An ACG monograph on the management of IBS and chronic idiopathic constipation states that therapies supported by high levels of evidence for improvement of IBS symptoms include linaclotide for IBS-C and antidepressants for IBS. Therapies supported by moderate levels of evidence include lubiprostone for IBS-C; alosetron in women with IBS-D; peppermint oil, psyllium, and fiber for overall symptom relief in IBS; and rifaximin for total IBS symptoms and bloating in IBS-D. Therapies supported by low levels of evidence include probiotics for global symptoms, bloating, and flatulence and certain antispasmodics for short-term relief of symptoms. Specialized diets are supported by

*Editor-in-Chief, *Hospital Pharmacy*, and Clinical Professor, Emeritus, Department of Pharmacy Practice, University of Kansas, School of Pharmacy, Kansas City/Lawrence, Kansas, e-mail: jgeneral@ku.edu; †Founder and Contributing Editor, *The Formulary*, and Editor, *Off-Label Drug Facts*, e-mail: Dennis.Cada@wolterskluwer.com.

very low levels of evidence. According to the statement, there is insufficient evidence to recommend prebiotics, synbiotics, loperamide, mixed 5-HT₄ agonists/5-HT₃ antagonists, or polyethylene glycol.

Although moderate evidence supports rifaximin for IBS, the recommendation for its use is classified as weak, based on results from 5 trials in 1,805 IBS patients. Rifaximin has demonstrated modest but consistent efficacy in the management of IBS without constipation. The monograph notes that there was a statistically significant benefit in favor of rifaximin (relative risk [RR], 0.84; 95% CI, 0.78 to 0.9), with a number needed to treat (NNT) of 9 (95% CI, 6 to 12.5). There was no difference between antibiotic therapy and placebo for risk of adverse events (RR, 0.7; 95% CI, 0.42 to 1.16).¹

Controlled Trials

In 2 identically designed, double-blind, placebo-controlled trials (TARGET 1 and TARGET 2) in 1,258 adult patients with IBS without constipation, patients were randomized to receive rifaximin (550 mg 3 times daily) or placebo for 2 weeks and were monitored for an additional 10-week period after therapy completion. The primary outcome was the percentage of patients achieving adequate relief of global IBS symptoms, as measured by patient response, for at least 2 of the first 4 weeks after treatment. The secondary outcome was patient-reported relief of IBS-related symptoms, including bloating and abdominal pain and discomfort, according to a 7-point scale. Combined results from both studies revealed that significantly more patients receiving rifaximin reported relief of global IBS symptoms compared with patients receiving placebo (40.7% vs 31.7%; $P < .001$). Similar results were also reported for IBS-related bloating (40.2% vs 30.3%; $P < .001$) and abdominal pain and discomfort (44.3% vs 36.3% in TARGET 1 [$P = .03$]; 42.9% vs 34.4% in TARGET 2 [$P = .02$]). Continued relief of symptoms was maintained in the rifaximin group during all 3 months in both studies.⁵

Noncontrolled Trial

A retrospective review of patients with IBS without constipation who had previously received at least 1 course of rifaximin treatment was conducted to identify the number of rifaximin retreatments, the success of additional rifaximin treatments, and the duration of benefit with retreatment. Of 522 patient charts reviewed, 187 subjects had IBS without constipation; of these subjects, 169 had received rifaximin and 148

were evaluated at follow-up visits. Rifaximin produced clinical improvement in most patients ($n = 111$; 75%). Seventy-one patients experienced symptomatic relapse with documented rifaximin retreatment; the majority of this group was female ($n = 44$), with a mean age of 47 years. Forty-eight patients received retreatment with a second course of rifaximin, 22 received a third course, 9 received a fourth course, and 4 received a fifth course. More than 75% of patients who initially responded to rifaximin experienced clinical benefit with additional treatment. Although relapse rates varied for all retreatments, median duration of benefit was at least 4 months. The authors concluded that in general, patients who respond to 1 rifaximin treatment course also respond to subsequent courses.⁷

Meta-analysis

A meta-analysis evaluated 5 double-blind, randomized, placebo-controlled trials ($N = 1,803$) that compared the efficacy of rifaximin versus placebo for the management of IBS, as defined by accepted symptom criteria. The primary outcome was global improvement of IBS symptoms. Rifaximin was associated with improvement of global symptoms compared with placebo (summary odds ratio [OR], 1.57; 95% CI, 1.22 to 2.01; $P < .001$). Significantly more patients in the rifaximin group reported global improvement in symptoms compared with placebo (42.2% vs 32.4%), with an NNT of 10.2 for therapeutic gain (9.8%). Mild heterogeneity was identified ($I^2 = 26\%$; $P = .25$). When patient and study characteristics were evaluated to determine impact on response rates, age was considered a significant predictor of symptom improvement (OR, 1.16; 95% CI, 1.07 to 1.25). Secondary outcomes included bloating, abdominal pain, and stool consistency. Rifaximin was associated with improvement in bloating compared with placebo (OR, 1.55; 95% CI, 1.23 to 1.96; $P < .001$). Therapeutic gain was 9.9%, with an NNT of 10.1. Raw data regarding abdominal pain were not available, and data addressing stool consistency were conflicting in the 3 studies for which data were available. The authors concluded that rifaximin is more effective than placebo for relief of global symptoms and bloating in IBS and that therapeutic gain is similar to that demonstrated by other currently available IBS therapies.⁶

SAFETY

This is a limited safety profile. Refer to package labeling for complete prescribing information (eg, Warnings/Precautions, Adverse Reactions, Drug Interactions).

In large placebo-controlled trials, rates of serious adverse events were similar for rifaximin and placebo (1.6% vs 2.4%, respectively).⁵

In a meta-analysis, adverse events were available for 4 of the 5 studies that met inclusion criteria. Overall, the incidence of adverse events and discontinuation rates due to adverse events were similar between the placebo and rifaximin groups. The most frequently cited adverse events (up to 6%) included headache, upper respiratory tract infection, nausea, nasopharyngitis, diarrhea, and abdominal pain.⁶

A pooled analysis of study data from 3 double-blind, placebo-controlled trials enrolling 1,940 patients evaluated the safety and tolerability of rifaximin for treatment of IBS without constipation. Mean age of patients was 46 years, and the majority (73%) was female. Most rifaximin dosage regimens were administered for 2 weeks and included 275 mg, 550 mg, or 1,100 mg twice daily, or 550 mg 3 times daily; 1 study included 550 mg twice daily for 4 weeks. The overall safety profile indicated that the majority of adverse events were mild to moderate in nature, with a similar overall incidence rate between rifaximin and placebo (12.1% vs 10.7%). Most reactions included GI-associated complaints or infection.⁸

No cases of *Clostridium difficile*-associated diarrhea or death have been reported in reviewed trials or meta-analyses.^{5,6,8}

Concerns have been raised regarding long-term or repeated use of antibiotics, including rifaximin, suggesting the drug should be limited to single, acute, short-term treatment.⁹

THERAPY CONSIDERATIONS

Rifaximin for the management of IBS has been studied in several placebo-controlled trials demonstrating modest effects on global IBS symptoms and benefit in bloating in IBS patients without constipa-

tion. Most trials included a short-term follow-up (less than 3 months) after therapy completion. An ACG monograph on the management of IBS provides a weak recommendation for rifaximin use, based on moderate evidence indicating modest but consistent efficacy in IBS without constipation.

REFERENCES

1. Ford AC, Moayyedi P, Lacy BE, et al. American College of Gastroenterology monograph on the management of irritable bowel syndrome and chronic idiopathic constipation. *Am J Gastroenterol*. 2014;109(Suppl 1):S2-S26.
2. Saadi M, McCallum RW. Rifaximin in irritable bowel syndrome: Rationale, evidence and clinical use. *Ther Adv Chronic Dis*. 2013;4(2):71-75.
3. Saha L. Irritable bowel syndrome: Pathogenesis, diagnosis, treatment, and evidence-based medicine. *World J Gastroenterol*. 2014;20(22):6759-6773.
4. Wall GC, Bryant GA, Bottenberg MM, Maki ED, Miesner AR. Irritable bowel syndrome: A concise review of current treatment concepts. *World J Gastroenterol*. 2014;20(27):8796-8806.
5. Pimentel M, Lembo A, Chey WD, et al; TARGET Study Group. Rifaximin therapy for patients with irritable bowel syndrome without constipation. *N Engl J Med*. 2011;364(1):22-32.
6. Menees SB, Maneerattannaporn M, Kim HM, Chey WD. The efficacy and safety of rifaximin for the irritable bowel syndrome: A systematic review and meta-analysis. *Am J Gastroenterol*. 2012;107(1):28-35.
7. Pimentel M, Morales W, Chua K, et al. Effects of rifaximin treatment and retreatment in nonconstipated IBS subjects. *Dig Dis Sci*. 2011;56(7):2067-2072.
8. Schoenfeld P, Pimentel M, Chang L, et al. Safety and tolerability of rifaximin for the treatment of irritable bowel syndrome without constipation: A pooled analysis of randomised, double-blind, placebo-controlled trials. *Aliment Pharmacol Ther*. 2014;39(10):1161-1168.
9. Farrell DJ. Rifaximin in the treatment of irritable bowel syndrome: Is there a high risk for development of antimicrobial resistance? *J Clin Gastroenterol*. 2013;47(3):205-211. ■