PSB27: A thylakoid protein enabling *Arabidopsis* to adapt to changing light intensity

Xin Hou, Aigen Fu¹, Veder J. Garcia², Bob B. Buchanan³, and Sheng Luan³

Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720

Contributed by Bob B. Buchanan, December 16, 2014 (sent for review October 9, 2014; reviewed by Roberto Bassi and Marc M. Nowaczyk)

In earlier studies we have identified FKBP20-2 and CYP38 as soluble proteins of the chloroplast thylakoid lumen that are required for the formation of photosystem II supercomplexes (PSII SCs). Subsequent work has identified another potential candidate functional in SC formation (PSB27). We have followed up on this possibility and isolated mutants defective in the PSB27 gene. In addition to lack of PSII SCs, mutant plants were severely stunted when cultivated with light of variable intensity. The stunted growth was associated with lower PSII efficiency and defective starch accumulation. In response to high light exposure, the mutant plants also displayed enhanced ROS production, leading to decreased biosynthesis of anthocyanin. Unexpectedly, we detected a second defect in the mutant, namely in CP26, an antenna protein known to be required for the formation of PSII SCs that has been linked to state transitions. Lack of PSII SCs was found to be independent of PSB27, but was due to a mutation in the previously described *cp26* gene that we found had no effect on light adaptation. The present results suggest that PSII SCs, despite being required for state transitions, are not associated with acclimation to changing light intensity. Our results are consistent with the conclusion that PSB27 plays an essential role in enabling plants to adapt to fluctuating light intensity through a mechanism distinct from photosystem II supercomplexes and state transitions.

photosystem | PSII supercomplex | photosynthesis | anthocyanin biosynthesis | reactive oxygen species

Photosynthetic light reactions entail the coordinated function of several large membrane complexes: photosystem I (PSI), photosystem II (PSII), cytochrome *b6f* complex, and CF₀-CF₁ complex. PSII catalyzes the initial step of photosynthesis, the light-dependent oxidation of water that yields molecular oxygen and reduced plastoquinone. The native form of PSII residing in the thylakoid membrane is believed to be organized into several types of supercomplexes (SCs), including the PSII core and the peripheral light harvesting complex II (LHCII), that play a primary role in the harvesting of light, transfer of excitation energy to the reaction center and regulation of light utilization. Several monomeric antenna proteins including CP24, CP26, and CP29 regulate the interaction of the PSII core with LHCII trimers (1–4). Regulation is achieved through photoprotective mechanisms that dissipate absorbed excess energy as heat in response to stress conditions such as high light intensity (5, 6).

In natural settings, both the intensity and the spectral quality of light vary extensively, sometimes within very short periods. The changes in light intensity result in imbalanced excitation of PSI and PSII, and with that may lower the efficiency of the photosynthetic light reactions. In acclimating to the changing conditions, plants modify their thylakoid proteins and reorganize their photosynthetic machinery (7, 8). In a rapid response, designated state transitions (9), LHCII associates reversibly with PSII or PSI. Under high light intensity, excessive activation of PSII increases the reduced plastoquinone pool and thereby activates protein kinase STN7 which, in turn, phosphorylates LHCII and prompts the migration of LHCII from PSII (state 1) to PSI (state 2) (10, 11). Oxidation of the plastoquinone pool by the higher activity of PSI then activates protein phosphatase

PPH1 that dephosphorylates LHCII and promotes a return to the original association of LHCII with PSII (state 1) (12, 13). In this context, the formation of LHCII-PSII SCs is expected to be a prerequisite for state transitions and, therefore, essential for adaptation to changing light intensity. However, it remains to be rigorously tested if state transitions play a crucial role in plant adaptation to changing light intensity.

In the present study, we have identified an *Arabidopsis* mutant that lacked PSII supercomplexes and grew poorly under changing light intensity. The mutant harbored a T-DNA insertion in the gene encoding the thylakoid lumen protein, PSB27, implicating its function in both the assembly of PSII SCs and adaptation to changing light. A detailed comparative genetic and biochemical analysis confirmed the requirement for PSB27 in enabling plants to adapt to changing light, but, surprisingly, revealed that this adaptation is independent of PSII SC assembly. A second, previously unrecognized defect was localized in CP26, a protein unrelated to PSB27, that is linked to PSII SC assembly. These results prompt the conclusion that PSB27 plays a fundamental role in enabling plants to adapt to changes in light intensity independently of the formation of PSII SCs.

Results and Discussion

Identification of a Mutant with a Defect in the Assembly of PS II Supercomplexes. Our earlier work suggests that FKBP20-2 and CYP38 chaperones function in the assembly of PSII SCs (14, 15). Specifically, these proteins appear to be localized to the chloroplast thylakoid lumen where a major portion of the PSII complex resides. These findings raise the possibility that additional luminal proteins may be required for PSII assembly. To gain information on this point, we have taken a systematic reverse genetic approach and isolated 58 *Arabidopsis* T-DNA insertion mutants in genes

Significance

Our results provide evidence that PSB27, a soluble protein of the chloroplast thylakoid lumen, is essential for enabling plants to adapt to changes in light intensity. Adaptation takes place independently of photosystem II supercomplexes and, based on work of others, of state transitions for the redistribution of light harvesting proteins. This finding opens the door to pursue the question of how photosynthesis adjusts to fluctuating sunlight and enables plants to grow under ever-changing environmental conditions.

Author contributions: X.H., A.F., and S.L. designed research; X.H., A.F., and V.J.G. performed research; X.H., A.F., B.B.B., and S.L. analyzed data; and X.H., B.B.B., and S.L. wrote the paper.

Reviewers: R.B., Università di Verona; and M.M.N., Ruhr-Universität Bochum.

The authors declare no conflict of interest.

¹Present address: College of Life Sciences, Northwest University, Xi'an, Shanxi 710069, China.

²Present address: Department of Plant Biology, Carnegie Institution for Science, Stanford, CA 94305.

³To whom correspondence may be addressed. Email: sluan@berkeley.edu or view@berkeley.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1424040112/-DCSupplemental.

encoding 30 luminal proteins (16, 17). The mutants were screened for defects in PSII complex organization by blue native gel electrophoresis. The mutant line with the most marked defect in PSII organization (SALK_004769) contained a T-DNA insertion in the 5' UTR of the *PSB27* gene (At1g03600, Fig. S1A). As reported earlier (18), this mutant was devoid of *PSB27* mRNA determined by RT-PCR [Fig. S1B, compare wild type (WT) and mutant line (ML)] as well as the corresponding protein probed by SDS/PAGE and Western blot (Fig. S1C). Protein fractionation of subchloroplast compartments confirmed that *PSB27* is located in the thylakoid lumen (Fig. S1D).

In an effort to localize the defect in PSII, we next conducted more quantitative blue native gel analysis with WT and mutant thylakoid membrane proteins and found pronounced differences in the abundance of individual PSII SC bands: notably, the mutant plants were almost completely devoid of PSII SCs that were readily detected in the WT (Fig. 1A). By contrast, the band representing the free forms of the corresponding LHC trimers (LHCII-T) showed a relative increase in abundance. When subjecting the blue native gel slices to the second electrophoresis dimension by SDS/PAGE, we observed a dramatic decrease in abundance of protein subunits associated with the PSII SCs (Fig. 1B, see red boxes).

To further visualize changes in the PSII complexes, we used immunoblots to examine the distribution and abundance of subunits of the four major thylakoid membrane complexes. To this end, the individual proteins resolved by 2D-SDS PAGE were transferred to a nitrocellulose membrane and probed with antibody against subunits of each of the four thylakoid membrane complexes: PSII: D1, D2, CP43, CP47, PsbO, and LHCII; PSI: PsaA/B,

PsaD, and PsaF; cytochrome *b₆f* complex: CytF, *b₆*, and PetC; and CF₀-CF₁ complex: ATP α . The results confirmed that the mutant (ML) lacked the protein subunits in the PSII SCs that were observed in the WT (Fig. 1C). The subunit distribution pattern of other complexes including PSI, Cyt *b₆f*, and CF₀-CF₁ remained unchanged (Fig. 1C).

It was difficult to compare the abundance of each individual protein between WT and mutant using 2D Western blots because the gels were processed separately. To circumvent this difficulty, we quantified these proteins by loading WT and mutant samples on the same SDS/PAGE gel, followed by Western blot. The results showed no significant difference in the abundance of PSII subunits between the WT and mutant (ML) cultivated under growth chamber (Fig. 1D) or greenhouse conditions (Fig. 1E). Taken together, these results suggested although the total amount of individual PSII subunits was not affected, the mutant plants were incapable of assembling the individual subunits to form PSII SCs.

We noted that the mutant showed two new weak bands in the PSII SCs region of the blue native gel (Fig. 1A): one was located between 1,250 and 1,100 kDa and the other was just below the 880 kDa band of WT PSII SC. To determine whether these bands represented altered forms of the PSII SC or were unrelated to this photosystem, we performed blue native gel analysis of both WT and mutant (ML) with a larger amount of total protein (Fig. S2A). The major bands above the PSII dimer in both WT and mutant samples in Fig. S2A were isolated and subjected to SDS PAGE analysis. The results showed that the top band, which was identified as NDH (labeled WT-1 and ML-1 in Fig. S2A), and the bottom band, which was identified as PSI SC (19, 20) (labeled WT-6 and ML-4), displayed identical patterns of subunit composition in the WT and ML (Fig. S2B). The four major PSII SC bands of WT (WT-2–WT-5) all contained typical PSII subunits (red lettered a–d bands in Fig. S2B), whereas the two unique bands (ML-2 and ML-3) in mutant lacked these subunits. We further showed by mass spectrometry that these proteins (a–d bands in the WT sample in Fig. S2B) are true PSII subunits (Fig. S2C). Interestingly, both the 880-kDa band of the mutant (ML-3) and the upper band (ML-2), neither of which is obvious in the WT gel, displayed a subunit pattern similar to PSI SC (WT-6) instead of PSII SC (Fig. S2A and B). This result again supports our conclusion that all detectable forms of PSII SCs are eliminated in the mutant.

Mutant Line Is Deficient in Adaptation to Changing Light Intensity.

When cultivated in a growth chamber under normal growth conditions (Fig. 2A), the ML failed to show a noticeable phenotypic difference from WT in either soil (Fig. 2E) or MS medium (Fig. 2I) despite lacking PSII SCs. Similar results were reported previously (18). By contrast, when grown in the greenhouse (Fig. 2B), leaves of ML plants were much smaller and less green compared with WT (Fig. 2F and J). Clearly, environmental differences (growth chamber vs. greenhouse) caused a dramatic change in mutant phenotype. To identify the factor responsible for such a difference, we cultivated the plants under several growth conditions—i.e., temperature, day length (8 h light/16 h dark for short day, 16 h light/8 h dark for long day), and light intensity (20 $\mu\text{mol m}^{-2} \text{s}^{-1}$ to 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$). The results indicated that none of these factors was responsible for the observed difference in phenotype. We then realized that the growth chamber supplies light at constant intensity during the light cycle, whereas the greenhouse experiences light at fluctuating intensity during the day akin to a natural setting.

To pursue the possibility that change in light intensity led to altered growth of the mutant, we designed growth chamber conditions such that plants experienced a gradient of light intensity during the light cycle (Fig. 2C), mimicking greenhouse conditions (Fig. 2B). Indeed, the ML plants subjected to this light intensity regime clearly differed from WT, similar to those cultivated in the greenhouse (Fig. 2G and K). When the light intensity was changed every hour from 50 to 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ during the light cycle (Fig. 2D), growth of ML plants was also slow relative to WT

Fig. 1. Analysis of thylakoid membrane protein complexes. (A) Blue native gel analysis of thylakoid proteins of WT and ML plants (15 μg chlorophyll as loading standard). (B) Thylakoid proteins separated by BN gel in A were further subjected to SDS PAGE and silver-stained. The red frames indicate the individual protein components of the PSII SCs. Protein identification was according to Fu et al. (15). (C) 2D-SDS PAGE immunoblots of thylakoid proteins. Antibodies were applied as indicated. (D) 1D-SDS PAGE Western blots of thylakoid proteins (2.5 μg chlorophyll) from WT and ML plants grown for 4 wk in a growth chamber under constant light intensity. (E) 1D-SDS PAGE immunoblots of thylakoid proteins (2.5 μg chlorophyll) from WT and ML grown in a greenhouse for 4 wk. Antibodies applied and plants grown as indicated.

Fig. 2. Mutant line is compromised in light adaptation. (A–D) Light intensity regimes for plant growth. The light/dark cycle was 16 h light/8 h dark for all regimes. (A) Growth chamber conditions with constant light intensity ($100 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$) during the light periods. (B) Greenhouse conditions with light intensity changes during the day time. (C) Growth chamber conditions with light intensity changes from $75 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ to $250 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ during the light periods. (D) Growth chamber conditions with light intensity switched between $50 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ and $150 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ every hour during the light periods. (E–H) WT and ML were grown in soil for 4 wk under the various light regimes in A–D. (I–L) WT and ML were grown on 1/2 MS medium for 10 d under light regimes in A–D. Growth conditions depicted in A–D were used, respectively, for growth in E–H and I–L.

(Fig. 2H). A similar phenotypic difference was observed with plants grown on agar plates containing 1/2 MS medium (Fig. 2L). These results suggested the ML plants have defects in adaptation to changing light intensity.

Mutant Line Is Defective in PSII Function and Starch Biosynthesis. The stunted growth of ML plants observed under changing light intensity suggests that the mutant had lower photosynthetic capacity under these conditions. To test this idea, photosynthetic parameters were determined by measuring chlorophyll fluorescence in leaves of whole plants. The ratio of variable fluorescence to maximum fluorescence (F_v/F_m), which represents the maximum photochemical efficiency of PSII (21), was lower in the ML than in the WT even with plants cultivated in the growth chamber which showed no visible phenotypic difference (Fig. 3A, Upper). The decrease in the F_v/F_m ratio became more pronounced with plants grown in the greenhouse (Fig. 3B, Upper). ΦPSII , which measures the proportion of PSII-associated light harvesting, was also lower in the ML (Fig. 3A, Lower). Like the F_v/F_m ratio, ΦPSII was further decreased when plants were grown in the greenhouse (Fig. 3B, Lower). Taken together, the chlorophyll fluorescence assays suggested that ML plants are compromised in PSII function.

The fluorescence results prompted us to test the possibility that PSII SC formation is needed for thylakoid membrane organization. If so, lack of PSII SCs could lead to defects in chloroplast morphology in mutant plants. To pursue this idea, we examined leaf sections of both WT and ML under the electron microscope. Under growth chamber conditions, the chloroplasts

from ML and WT showed similar size, grana, and starch accumulation (Fig. 3C). By contrast, electron microscopy revealed a striking difference between WT and ML specimens grown in the greenhouse: The chloroplasts in wild type accumulated copious amounts of starch, whereas, consistent with its stunted growth, the ML was almost completely devoid of starch grains (Fig. 3D), indicating a much lower efficiency of starch biosynthesis under changing light conditions. This inefficiency could result, in part, from an impairment in the regulation of enzymes of starch biosynthesis due to defective PSII activity and the attendant reduction of thioredoxin (22).

Mutant Line Accumulates Less Anthocyanin and Produces More ROS Under High Light Stress. Because PSII deficiency and photo-inhibition can lead to damage under high light, we investigated whether the mutant line is defective in adaptation to this condition. To this end, WT and ML were grown on 1/2 MS medium for

Fig. 3. Mutant line is deficient in utilization of light. (A) Chlorophyll fluorescence analyses of WT and ML plants grown under constant light intensity conditions in the growth chamber ($80 \mu\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ for 16 h and dark for 8 h) for 4 wk. (B) Chlorophyll fluorescence analyses of WT and ML plants grown under greenhouse conditions for 4 wk. F_v/F_m , the maximum efficiency of PSII photochemistry; ΦPSII , efficiency of PSII electron transport. Data represent mean $\pm$ SEM ($n = 4$). (C) Electron micrographs of chloroplasts from 4 wk-old WT and ML grown under constant light intensity conditions. (D) Electron micrographs of chloroplasts from WT and ML grown in greenhouse for 4 wk. Transmission Electron Microscope (FEI) was used to examine leaf cross sections as described (42).

contribute to light adaptation at least under the conditions used in this study.

Concluding Remarks. The present experiments have uncovered a protein of the chloroplast thylakoid lumen, PSB27, that is required for adaptation to changing light intensity. Using an *Arabidopsis* mutant lacking PSB27, we found that, although not essential under conditions of constant light intensity, this protein was required for normal growth under changing light regimes in either a natural setting or a growth chamber. The stunted growth observed under changing light was associated with lower PSII efficiency and defective starch accumulation. In response to high light exposure, mutant plants also displayed enhanced ROS production and reduced synthesis of the protectant anthocyanin although such changes did not affect plant growth.

We observed that the monomeric antenna protein CP26, a component of PSII SCs, is required for the assembly of the photosynthetic apparatus. An *Arabidopsis* mutant lacking CP26 was found to be essentially devoid of PSII SCs. Although it was reported that the LHCII S trimer could associate with the PSII core

Fig. 5. Genetic linkage of *cp26* and *psb27* mutant alleles with plant growth defects. (A) The original T-DNA ML containing both *cp26* and *psb27* mutant alleles was compared with the WT and with segregated single mutants of *cp26* and *psb27* under constant light intensity in a growth chamber (Upper) or under varying light intensity in a greenhouse (Lower). (B) The plants were grown on 1/2 MS medium in growth chamber (Left) or greenhouse (Center) for 10 d. The same plants grown under constant low light for 10 d were then transferred to high light ($350 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 2 d (Right). (C) Blue native gel analysis showing PSII SCs of thylakoid proteins of WT, ML, *psb27*, *cp26*, ML complemented with CP26 or/and PSB27. (D) Tris-Tricine SDS PAGE analyses showing CP26 proteins of thylakoids of the same plants in C.

Fig. 6. Comparison of *cp26*, *psb27*, *stn7*, *stn8*, and *stn7 stn8* double mutant in their adaptation to light conditions. (A) WT, *cp26*, *psb27*, *stn7*, *stn8*, and *stn7 stn8* double mutant were grown in soil in growth chamber (Upper) and greenhouse (Lower) for 4 wk (the growth conditions were as in Fig. 2 A and B). (B) The same plants were grown on 1/2 MS medium for 10 d under light regimes in Fig. 2A (Growth Chamber) and Fig. 2B (Greenhouse). (C) phosphorylation of thylakoid proteins. Total proteins of 2-wk-old seedlings were separated by 12% SDS PAGE and immunoblotted with anti-phosphothreonine antibodies.

in the absence of CP26 (28), our results suggest that this association is not stable, consistent with the sucrose gradient results (1).

The impact of state transitions on energy balance between PSI and PSII has been established for the green alga *Chlamydomonas* (29). However, state transitions play a less prominent role in land plants than in green algae. Growth is only marginally affected in *Arabidopsis* mutants impaired in state transitions (27, 30), even in fluctuating light (10). Our results have shown that state transition mutants also display minor, if any, phenotypic change under conditions that severely inhibit growth of *psb27* mutant plants. Our results suggest that land plants adapt to changing light intensity through a PSB27-dependent mechanism that is independent of both state transitions and PSII SCs. Thus, based on our results it seems possible that the recently reported effects of PSB27 on PSII SC formation and state transitions (31) may have been the result of the *cp26* point mutation that was present together with the *psb27* T-DNA insert in the mutant line that was used (SALK_004769).

A final point, PSB27 is present in cyanobacteria as well as land plants (16, 32). In cyanobacteria PSB27 is tightly associated with the thylakoid membrane via an N-terminal lipid modification that is absent in PSB27 from land plants (33). Therefore, PSB27 of *Arabidopsis* is present exclusively in the soluble phase of the thylakoid lumen (Fig. S1D). This difference in location may, in part, be responsible for functional differences of PSB27 in the two groups of organisms. In cyanobacteria, the membrane-bound PSB27 facilitates the assembly of the manganese cluster of PSII (34–38) by interacting tightly with the CP43 core protein in the assembly of intermediates. In land plants, on the other hand, PSB27 functions in acclimation to changing light. Elucidation of further details of PSB27 function may well add insight into the evolution of oxygenic photosynthesis.

Materials and Methods

Plant Materials and Growth Conditions. *Arabidopsis thaliana*, ecotype Columbia-0, was used in this study. The T-DNA insertion mutant line (SALK_004769, locus At1g03600) was obtained from the *Arabidopsis* Resource Center (Columbus, OH). For soil-grown plants, sown seeds were cold-treated for 2 d, and then transferred to the indicated growth conditions. For plate-grown plants, surface-sterilized seeds were planted on 1/2 MS medium with 0.8% phytigel, cold-treated for 2 d, and grown under the same conditions as soil-grown plants.

Blue Native PAGE and 2D-SDS PAGE. Blue native gel electrophoresis was performed as described in ref. 14. For 2D-SDS PAGE, lanes of blue native gel were excised with a razor blade and incubated in 2× SDS sample buffer containing 2.5% (vol/vol) β-mercaptoethanol for 15 min at 25 °C and then for 15 min at 70 °C. Each lane with denatured proteins was placed on top of a 12% SDS PAGE gel and subjected to second dimension separation (14).

Immunoblot Analysis. Protein samples corresponding to equal amounts of chlorophyll were separated on 12% SDS PAGE gels and transferred to nitrocellulose membranes, followed by Western blot analysis (39). After blocking nonspecific binding with 5% milk, the blot was subsequently incubated with antibodies generated against the indicated proteins and detected using SuperSignal West Pico Chemiluminescent Substrate kit (Pierce).

Chlorophyll Fluorescence Measurements. Chlorophyll fluorescence was measured with a FMS2 fluorometer (Hansatech) attached to intact leaves that had been dark adapted overnight. The F_0 , F_m , and F_s were measured and calculated as described (14). F_v/F_m was calculated as $(F_m - F_0)/F_m$, and $\Phi PSII$ was defined as $(F'_m - F_s)/F'_m$.

RNA Isolation and Quantitative Reverse Transcription-PCR. Total RNA was isolated from leaves using an RNeasy Plant Mini Kit (Qiagen). cDNA was synthesized using SuperScript II Reverse Transcriptase kit (Invitrogen). Real-time quantitative RT-PCR was performed with a DNA Engine Opticon System (MJ Research) using DyNAmo HS SYBR Green qPCR Kit (New England BioLabs). *Actin2* gene was used as the endogenous control. Relative expression levels were determined as described (39).

Anthocyanin and Hydrogen Peroxide Measurements. Anthocyanin were extracted in water from 0.2 g leaves by grinding in liquid N₂. After centrifugation at 16,000g, the supernatant was diluted with 0.4 M sodium acetate (pH 4.5) and 25 mM potassium chloride (pH 1.0). Total monomeric anthocyanin was measured by recording the A_{520} as outlined in Giusti and Wrolstad (40).

For hydrogen peroxide staining, plants were vacuum-infiltrated with 0.1 mg/mL 3,3'-diaminobenzidine in 50 mM Tris-acetate buffer, pH 5.0. Samples were incubated for 24 h at room temperature in the dark. To remove chlorophylls, the stained samples were transferred to 80% ethanol.

Mutant Complementation. The coding region of *PSB27* gene was amplified by RT-PCR from total RNA and cloned in the vector pCB2004H to produce a construct that expresses PSB27 protein under the control of CaMV 35S promoter (39). For complementation by genomic DNA, a 2.2-kb genomic DNA fragment containing the coding region, 1.4-kb upstream, and 220-bp downstream sequences was amplified by PCR and cloned in the vector pCB2004. For complementation of *cp26* mutation, a 2.5-kb genomic DNA fragment containing the *CP26* gene and the 1.1-kb promoter was cloned in the vector pCB2004. The primers are shown in Table S1. These constructs were transferred into mutant plants using the floral dip method (41).

ACKNOWLEDGMENTS. We thank Dr. Kent McDonald (Electron Microscope Laboratory at the University of California, Berkeley) for preparing the electron micrographs. We thank Drs. A. Melis, S. Park, R. Malkin (University of California, Berkeley) and S. Theg (University of California, Davis) for providing antibodies and K. Niyogi (University of California, Berkeley) for advice on chlorophyll fluorescence measurements. This work was supported by grants from U.S. Department of Energy (to S.L.) and the National Science Foundation (to B.B.B. and S.L.).

- Caffarri S, Kouril R, Kereiche S, Boekema EJ, Croce R (2009) Functional architecture of higher plant photosystem II supercomplexes. *EMBO J* 28(19):3052–3063.
- Boekema EJ, Van Roon H, Van Breemen JF, Dekker JP (1999) Supramolecular organization of photosystem II and its light-harvesting antenna in partially solubilized photosystem II membranes. *Eur J Biochem* 266(2):444–452.
- de Bianchi S, et al. (2011) *Arabidopsis* mutants deleted in the light-harvesting protein Lhcb4 have a disrupted photosystem II macrostructure and are defective in photoprotection. *Plant Cell* 23(7):2659–2679.
- de Bianchi S, Dall'Osto L, Tognon G, Morosinotto T, Bassi R (2008) Minor antenna proteins CP24 and CP26 affect the interactions between photosystem II subunits and the electron transport rate in grana membranes of *Arabidopsis*. *Plant Cell* 20(4):1012–1028.
- Schmid VH (2008) Light-harvesting complexes of vascular plants. *Cell Mol Life Sci* 65(22):3619–3639.
- Nixon PJ, Michoux F, Yu J, Boehm M, Komenda J (2010) Recent advances in understanding the assembly and repair of photosystem II. *Ann Bot (Lond)* 106(1):1–16.
- Kanervo E, Suorsa M, Aro EM (2005) Functional flexibility and acclimation of the thylakoid membrane. *Photochem Photobiol Sci* 4(12):1072–1080.
- Walters RG (2005) Towards an understanding of photosynthetic acclimation. *J Exp Bot* 56(411):435–447.
- Bonaventura C, Myers J (1969) Fluorescence and oxygen evolution from *Chlorella pyrenoidosa*. *Biochim Biophys Acta* 189(3):366–383.
- Bellafiore S, Barneche F, Peltier G, Rochaix JD (2005) State transitions and light adaptation require chloroplast thylakoid protein kinase STN7. *Nature* 433(7028):892–895.
- Allen JF (1992) Protein phosphorylation in regulation of photosynthesis. *Biochim Biophys Acta* 1098(3):275–335.
- Shapiguzov A, et al. (2010) The PPH1 phosphatase is specifically involved in LHClI dephosphorylation and state transitions in *Arabidopsis*. *Proc Natl Acad Sci USA* 107(10):4782–4787.
- Pribil M, Pesaresi P, Hertle A, Barbato R, Leister D (2010) Role of plastid protein phosphatase TAP38 in LHClI dephosphorylation and thylakoid electron flow. *PLoS Biol* 8(1):e1000288.
- Lima A, et al. (2006) A redox-active FKBP-type immunophilin functions in accumulation of the photosystem II supercomplex in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 103(33):12631–12636.
- Fu A, et al. (2007) A chloroplast cyclophilin functions in the assembly and maintenance of photosystem II in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 104(40):15947–15952.
- Peltier JB, et al. (2002) Central functions of the luminal and peripheral thylakoid proteome of *Arabidopsis* determined by experimentation and genome-wide prediction. *Plant Cell* 14(1):211–236.
- Schubert M, et al. (2002) Proteome map of the chloroplast lumen of *Arabidopsis thaliana*. *J Biol Chem* 277(10):8354–8365.
- Chen H, et al. (2006) A Psb27 homologue in *Arabidopsis thaliana* is required for efficient repair of photodamaged photosystem II. *Plant Mol Biol* 61(4–5):567–575.
- Nelson N, Yocum CF (2006) Structure and function of photosystems I and II. *Annu Rev Plant Biol* 57:521–565.
- Sárvári E, et al. (2014) Comparison of thylakoid structure and organization in sun and shade *Haberlea rhodopensis* populations under desiccation and rehydration. *J Plant Physiol* 171(17):1591–1600.
- Maxwell K, Johnson GN (2000) Chlorophyll fluorescence—a practical guide. *J Exp Bot* 51(345):659–668.
- Nikkanen L, Rintamäki E (2014) Thioredoxin-dependent regulatory networks in chloroplasts under fluctuating light conditions. *Philos Trans R Soc Lond B Biol Sci* 369(1640):20130224.
- Steyn WJ, Wand SJE, Holcroft DM, Jacobs G (2002) Anthocyanins in vegetative tissues: A proposed unified function in photoprotection. *New Phytol* 155(3):349–361.
- Vanderauwera S, et al. (2005) Genome-wide analysis of hydrogen peroxide-regulated gene expression in *Arabidopsis* reveals a high light-induced transcriptional cluster involved in anthocyanin biosynthesis. *Plant Physiol* 139(2):806–821.
- Gadjev I, et al. (2006) Transcriptomic footprints disclose specificity of reactive oxygen species signaling in *Arabidopsis*. *Plant Physiol* 141(2):436–445.
- Vainonen JP, Hansson M, Vener AV (2005) STN8 protein kinase in *Arabidopsis thaliana* is specific in phosphorylation of photosystem II core proteins. *J Biol Chem* 280(39):33679–33686.
- Bonardi V, et al. (2005) Photosystem II core phosphorylation and photosynthetic acclimation require two different protein kinases. *Nature* 437(7062):1179–1182.
- Yakushevskaya AE, et al. (2003) The structure of photosystem II in *Arabidopsis*: Localization of the CP26 and CP29 antenna complexes. *Biochemistry* 42(3):608–613.
- Eberhard S, Finazzi G, Wollman FA (2008) The dynamics of photosynthesis. *Annu Rev Genet* 42:463–515.
- Lunde C, Jensen PE, Haldrup A, Knoetzel J, Scheller HV (2000) The PSI-H subunit of photosystem I is essential for state transitions in plant photosynthesis. *Nature* 408(6812):613–615.
- Dietzel L, et al. (2011) Photosystem II supercomplex remodeling serves as an entry mechanism for state transitions in *Arabidopsis*. *Plant Cell* 23(8):2964–2977.
- Rajalahti T, et al. (2007) Proteins in different *Synechocystis* compartments have distinguishing N-terminal features: A combined proteomics and multivariate sequence analysis. *J Proteome Res* 6(7):2420–2434.
- Nowaczyk MM, et al. (2006) Psb27, a cyanobacterial lipoprotein, is involved in the repair cycle of photosystem II. *Plant Cell* 18(11):3121–3131.
- Roose JL, Pakrasi HB (2008) The Psb27 protein facilitates manganese cluster assembly in photosystem II. *J Biol Chem* 283(7):4044–4050.
- Cormann KU, et al. (2009) Structure of Psb27 in solution: Implications for transient binding to photosystem II during biogenesis and repair. *Biochemistry* 48(37):8768–8770.
- Mabbitt PD, et al. (2009) Solution structure of Psb27 from cyanobacterial photosystem II. *Biochemistry* 48(37):8771–8773.
- Liu H, Roose JL, Cameron JC, Pakrasi HB (2011) A genetically tagged Psb27 protein allows purification of two consecutive photosystem II (PSII) assembly intermediates in *Synechocystis* 6803, a cyanobacterium. *J Biol Chem* 286(28):24865–24871.
- Grasse N, et al. (2011) Role of novel dimeric Photosystem II (PSII)-Psb27 protein complex in PSII repair. *J Biol Chem* 286(34):29548–29555.
- Hou X, Xie K, Yao J, Qi Z, Xiong L (2009) A homolog of human ski-interacting protein in rice positively regulates cell viability and stress tolerance. *Proc Natl Acad Sci USA* 106(15):6410–6415.
- Giusti MM, Wrolstad RE (2001) Characterization and Measurement of Anthocyanins by UV-Visible Spectroscopy. *Current Protocols in Food Analytical Chemistry*, eds Wrolstad RE, et al. (John Wiley & Sons, New York).
- Clough SJ, Bent AF (1998) Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* 16(6):735–743.
- Bauer J, et al. (2000) The major protein import receptor of plastids is essential for chloroplast biogenesis. *Nature* 403(6766):203–207.