Potential Risks of Excess Iodine Ingestion and Exposure: Statement by the American Thyroid Association Public Health Committee

Angela M. Leung,^{1,*} Anca M. Avram,² Alina V. Brenner,³ Leonidas H. Duntas,⁴ Joel Ehrenkranz,⁵
James V. Hennessey,⁶ Stephanie L. Lee,⁷ Elizabeth N. Pearce,⁷ Sanziana A. Roman,⁸
Alex Stagnaro-Green,⁹ Erich M. Sturgis,¹⁰ Krishnamurthi Sundaram,¹¹
Michael J. Thomas,¹² and Jason A. Wexler¹³

for the American Thyroid Association Public Health Committee

Dear Editor:

Iodine is a micronutrient required for normal thyroid function. In the United States, recommended daily allowances (RDA) for iodine intake are 150 μg in adults, 220–250 μg in pregnant women, and 250–290 μg in breastfeeding women (1,2). The U.S. diet generally contains enough iodine to meet these needs, with common sources being iodized salt, dairy products, some breads, and seafood. During pregnancy and lactation, women require higher amounts of iodine for the developing fetus and infant. The American Thyroid Association (ATA) recommends that women take a multivitamin containing 150 μg of iodine daily in the form of potassium iodide (KI) (3) during preconception, pregnancy, and lactation to meet these needs (4).

Ingestion of greater than 1100 μg of iodine per day (tolerable upper limits for iodine) (1) is not recommended and may cause thyroid dysfunction. During pregnancy and lactation, when the risk of excess iodine are primarily related to the fetus and newborn infant, the recommendations for the upper limit vary and range from 500–1100 μg of iodine daily (2). In particular, infants, the elderly, pregnant and lactating women, and individuals with preexisting thyroid disease (such as autoimmune Hashimoto's disease, Graves' disease, nontoxic thyroid nodules, history of partial thyroidectomy, and other conditions) are susceptible to adverse effects of excess iodine intake and exposure

(5). The public is advised that many iodine, potassium iodide, and kelp supplements contain iodine in amounts that are up to a hundred times higher than the daily tolerable upper limits for iodine. The ATA advises against the ingestion of iodine and kelp supplements containing in excess of 500 μg iodine daily for children and adults and during pregnancy and lactation. Long-term iodine intake in amounts greater than the tolerable upper limits should be closely monitored by a physician. There are only equivocal data supporting the benefit of iodine at higher doses than these, including a possible benefit for patients with fibrocystic breast disease (6). There is no known thyroid benefit of routine daily iodine doses in excess of the U.S. RDA.

There are a limited number of medical conditions in which the short-term use of high amounts of iodine is indicated. Exceptions for the recommendations to not exceed the tolerable upper limits include closely monitored patients prescribed Lugol's solution or saturated solution of potassium iodide (SSKI) in their treatment of severe hyperthyroidism, such as thyroid storm and prior to surgery in patients with Graves' disease, and individuals in the vicinity of a nuclear power plant who are recommended to take KI in the event of a nuclear accident. SSKI is not indicated nor recommended in individuals with thyroid nodules. Finally, patients receiving the large amounts of iodine in iodinated contrast dyes, as required for radiologic studies, should be

¹Division of Endocrinology, UCLA David Geffen School of Medicine, Los Angeles, California.

²Division of Nuclear Medicine/Radiology, University of Michigan Medical Center, Ann Arbor, Michigan.

³Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Department of Health and Human Services, Bethesda, Maryland.

⁴Unit of Endocrinology, Metabolism, and Diabetes, Evgenidion Hospital, University of Athens, Athens, Greece.

⁵Department of Medicine, Intermountain Healthcare, Salt Lake City, Utah.

⁶Division of Endocrinology, Department of Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts.

⁷Section of Endocrinology, Diabetes, and Nutrition, Boston University School of Medicine, Boston, Massachusetts.

⁸Department of Surgery, Duke University School of Medicine, Durham, North Carolina.

⁹University of Illinois College of Medicine at Rockford, Rockford, Illinois.

¹⁰Department of Head and Neck Surgery, UT MD Anderson Cancer Center, Houston, Texas.

¹¹Department of Otolaryngology, SUNY Downstate Medical Center, Brooklyn, New York.

¹²Carolina Endocrine, P.A., Raleigh, North Carolina.

¹³Division of Endocrinology and Metabolism, Georgetown University School of Medicine, Washington, District of Columbia.

*Chair of American Thyroid Association Public Health Committee.

monitored for iodine-induced thyroid dysfunction if risk factors are present.

Key points include:

- Adequate iodine intake is required for normal thyroid function.
- The recommended iodine intake in nonpregnant adults is 150 μg daily.
- Pregnant and breastfeeding women should take a prenatal vitamin that contains 150 μg of potassium iodine daily.
- Given a tolerable upper limit of 1100 μg iodine daily, ingestion of an iodine or kelp supplement containing in excess of 500 μg iodine daily should not be done.
- Certain exceptions to these recommendations include those for specific medical conditions, which usually require only a limited number of doses for a short-term duration; such individuals should be closely monitored for thyroid dysfunction.

REFERENCES

1. Food and Nutrition Board, Institute of Medicine 2006 Dietary Reference Intakes. National Academy Press, Washington, D.C.
2. De Groot L, Abalovich M, Alexander EK, Amino N, Barbour L, Cobin RH, Eastman CJ, Lazarus JH, Luton D, Mandel SJ, Mestman J, Rovet J, Sullivan S 2012 Management of thyroid dysfunction during pregnancy and postpartum: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* **97**:2543–2565.
3. Leung AM, Pearce EN, Braverman LE 2009 Iodine content of prenatal multivitamins in the United States. *N Engl J Med* **360**:939–940.
4. Stagnaro-Green, A, Abalovich, M, Alexander, E, Azizi, F, Mestman, J, Negro, R, Nixon, A, Pearce, EN, Soldin, OP, Sullivan, S, Wiersinga, W, American Thyroid Association Taskforce on Thyroid Disease During Pregnancy and Postpartum 2011 Guidelines of the American Thyroid Association for the diagnosis and management of thyroid disease during pregnancy and postpartum. *Thyroid* **21**:1081–1125.
5. Ghent WR, Eskin BA, Low DA, Hill LP 1993 Iodine replacement in fibrocystic disease of the breast. *Can J Surg* **36**:453–460.
6. Office of Dietary Supplements, National Institutes of Health. Dietary Supplement Fact Sheet: Iodine. Available at: <http://ods.od.nih.gov/factsheets/Iodine-HealthProfessional> (Last accessed May 24, 2014).

Address correspondence to:

*Angela M. Leung, MD, MSc
Division of Endocrinology (111D)
UCLA David Geffen School of Medicine
11301 Wilshire Boulevard
Los Angeles, CA 90073*

E-mail: amleung@mednet.ucla.edu