CASE REPORT

Cardiac asystoles misdiagnosed as epileptic seizures

Valentina Chiesa,¹ Aglaia Vignoli,² Maria Paola Canevini²¹Epilepsy Centre, San Paolo Hospital, Milan, Italy²Epilepsy Centre, San Paolo Hospital, Department of Health Sciences, University of Milan, Italy

Correspondence to

Dr Aglaia Vignoli;
aglaia.vignoli@ao-sanpaolo.it

Accepted 9 January 2015

SUMMARY

We report a case of a 78-year-old man who presented several episodes of transient loss of consciousness preceded by sensation of ascending heat from the feet, with increasing frequency and duration within a few weeks. One month later he was admitted because he started to fall during the episodes, which became daily. Brain MRI detected a gliotic right frontal area, and EEG showed slow activity in the same region. Carbamazepine was started without clinical response despite high plasmatic level. Video-EEG polygraphic monitoring was performed and the patient showed an episode of left hemifacial clonic jerks followed by loss of consciousness, starting 17 s after the beginning of asystole. Sinus rhythm reappeared 32 s later and the patient regained consciousness in few seconds. A pacemaker was implanted and carbamazepine was withdrawn. No further episodes occurred.

BACKGROUND

In clinical practice, the assessment of a patient with a transient loss-of-consciousness may be difficult. Differential diagnosis between epileptic seizure and syncope due to cerebral hypoperfusion might be challenging because clinical manifestations often overlap: both conditions may present a paroxysmal onset with loss of consciousness and loss of postural tone. Patients might manifest eye deviations, fragmentary myoclonus and tonic posturing and vocalisation.¹ Since the management of the two conditions is extremely different, a prompt diagnosis is crucial.

CASE PRESENTATION

A 78-year-old man with a case history of hypertension, diabetes mellitus, paroxysmal atrial fibrillation, severe left carotid stenosis and chronic renal failure started to show brief episodes characterised by a sensation of ascending heat starting from the feet, followed by a feeling of being estranged from the environment. Initially sporadic, the episodes occurred daily and for longer durations in the following months. Additionally, the patient started to lose consciousness immediately after the aforementioned sensations, showing left hemifacial clonic jerks and having occasional falls; consciousness was regained in a few seconds, followed by slight confusion. A month later, one of these episodes led to hospitalisation. The patient had a family history of epilepsy. His blood pressure, pulse, ECG and clinical neurological examination were normal. Brain MRI detected a right frontal gliotic area (figure 1) and EEG showed slow activity in the same region. A diagnosis of probable epilepsy with focal seizures was made, and antiepileptic treatment with

Figure 1 Brain MRI detected a right frontal gliotic area.

carbamazepine was started, without clinical response despite high plasmatic level. Owing to the persistence of daily episodes, Video-EEG polygraphic monitoring was performed.

INVESTIGATIONS

During the Video-EEG polygraphic monitoring, the patient, asleep, showed an episode of left hemifacial clonic jerks followed by loss of consciousness (video 1), starting 17 s after the beginning of asystole (figure 2A). During the episode, the EEG showed a progressive slowing of background activity and voltage reduction, followed by electrocerebral silence (figure 2B). No epileptic discharges were recognisable. Sinus rhythm reappeared 32 s later and the patient regained consciousness in a few seconds (figure 2C).

DIFFERENTIAL DIAGNOSIS

- ▶ Focal seizures in symptomatic epilepsy
- ▶ Syncope
- ▶ Other causes of loss of consciousness

Video 1 Left hemifacial clonic jerks started 17 s after the beginning of asystole (see figure 2). The patient regained consciousness in a few seconds after the sinus rhythm reappearance.

CrossMark

To cite: Chiesa V, Vignoli A, Canevini MP. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2014-206969

Figure 2 Asystole started during Video-EEG monitoring (A) 17 s later the beginning of the complete atrioventricular block, EEG activity showed diffuse slowing evolving into electrocerebral silence (B); prompt reappearance of background activity followed the sinus rhythm reappearance (C).

TREATMENT

A pacemaker was implanted and antiepileptic drug withdrawn.

OUTCOME AND FOLLOW-UP

No further episodes occurred after pacemaker implantation (follow-up 24 months).

DISCUSSION

The incidence of epilepsy in the elderly has increased steadily over the past few decades, and stroke is the most frequent aetiology.² Usually, focal epilepsy symptomatic of ischaemic cerebrovascular disease in elderly people has a good outcome using low dosages of antiepileptic drugs (AEDs), such as carbamazepine or lamotrigine.² The persistence of seizures not responsive to AEDs ought to suggest a different aetiology. Moreover, sodium channel-blocking drugs such as carbamazepine or phenytoin might have negative chronotropic and dromotropic effects in a patient with a sick heart.³ Therefore caution is required when starting treatment for episodes of uncertain aetiology.

Differential diagnosis with ictal asystole should be considered: this phenomenon consists of bradycardia or asystole caused by epileptic activity involving temporal lobe structures. In patients with temporal lobe epilepsy and ictal asystole, the loss of tone, due to cerebral hypoperfusion, occurs relatively late during the event and is often preceded by semiological features seen during typical dyscognitive seizures.^{4 5}

Focal neurological signs were the first clinical feature of the episode in our patient, starting more than 15 s after the beginning of asystole, without evidence of any epileptic activity on the EEG. As already reported,¹ focal myoclonus involving facial muscles can often occur during syncopal episodes, presumably related to disinhibition of subcortical structures.

However, the cerebrovascular condition of this patient might have a role in developing focal clinical signs: the severe left carotid stenosis and the right gliotic frontal lesion, together with the decrease of blood perfusion due to asystolia, could hypothetically lead to seizure onset. However, the Video-EEG recorded episode did not support this hypothesis since focal epileptic activity was absent. The EEG, instead, showed the typical features of syncope.⁶

Since cardiac asystoles might resemble focal epileptic seizures with vegetative prodromes and focal neurological signs, Video-EEG monitoring with simultaneous ECG is mandatory when the diagnosis is unreliable.

Learning points

- ▶ The persistence of paroxysmal episodes not responsive to antiepileptic drugs in patients who are believed to have post-stroke focal epilepsy ought to suggest a different nature of episodes.
- ▶ Since cardiogenic syncope might resemble focal epileptic seizures with vegetative prodromes and focal neurological signs, Video-EEG monitoring with simultaneous ECG is mandatory when the diagnosis is unreliable.
- ▶ Cooperation between neurologist and cardiologist can improve the management of patients with transient loss of consciousness.

Contributors VC followed the patient and wrote the manuscript. AV wrote and revised the manuscript. MPC revised the manuscript for intellectual content.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Lempert T, Bauer M, Schmidt D. Syncope: a videometric analysis of 56 episodes of transient cerebral hypoxia. *Ann Neurol* 1994;36:233–6.
- 2 Stefan H. Epilepsy in the elderly: facts and challenges. *Acta Neurol Scand* 2011;124:223–37.
- 3 Bergfeldt L. Differential diagnosis of cardiogenic syncope and seizure disorders. *Heart* 2003;89:353–8.
- 4 Shuele SU, Bermeo AC, Alexopoulos AV, et al. Video-electrographic and clinical features in patients with ictal asystole. *Neurology* 2007;69:434–41.
- 5 Kukla P, Jastrzębski M, Czamara M, et al. [Epileptic asystole—a case report]. *Kardiologia Pol* 2012;70:64–5. Polish.
- 6 Gastaut H, Fischer-Williams M. Electro-encephalographic study of syncope. Its differentiation from epilepsy. *Lancet* 1957;2:1018–25.

Copyright 2015 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit

<http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow