Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2015 February ; 24(2): 379–385. doi:
10.1158/1055-9965.EPI-14-0841.

Efficacy of a couple-based randomized controlled trial to help Latino fathers quit smoking during pregnancy and postpartum: The Parejas Trial

Kathryn I. Pollak, PhD^{1,2}, Pauline Lyna, MPH¹, Alicia K. Bilheimer, MPH, Kristina C. Gordon, PhD³, Bercedis L. Peterson, PhD⁴, Xiaomei Gao, MA¹, Geeta K. Swamy, MD⁵, Susan Denman, PhD⁶, Alicia Gonzalez, MSN⁷, Pilar Rocha, MS⁸, and Laura J. Fish, PhD²

¹Cancer Control and Population Science, Duke Cancer Institute

²Department of Community and Family Medicine, Duke University School of Medicine

³Department of Psychology, University of Tennessee at Knoxville

⁴Department of Biostatistics and Bioinformatics, Duke University School of Medicine

⁵Department of Obstetrics and Gynecology, Duke University School of Medicine

⁶School of Nursing, Duke University

⁷Duke AHEC Program, Duke University School of Medicine

⁸El Centro Hispano, Durham, NC

Abstract

Background—Although many Latinos in the US smoke, they receive assistance to quit less often than non-Latinos. To address this disparity, we recruited Latino couples into a randomized controlled trial and provided a smoking cessation program during a teachable moment, when men's partners were pregnant.

Methods—We compared two interventions: 1) written materials plus nicotine replacement therapy (NRT) to 2) materials, NRT, and couple-based counseling that addressed smoking cessation and couples communication. We recruited 348 expectant fathers who smoked via their pregnant partners from county health departments. Our primary outcome was 7-day point prevalence smoking abstinence and was collected from November 2010 through April 2013 and analyzed it in February 2014.

Results—We found high rates of cessation but no arm differences in smoking rates at the end of pregnancy (0.31 vs. 0.30, materials only vs. counseling, respectively) and 12 months after randomization (postpartum: 0.39 vs. 0.38). We found high quit rates among non-daily smokers but no arm differences (0.43 vs. 0.46 in pregnancy and 0.52 vs. 0.48 postpartum). Among daily smokers, we found lower quit rates with no arm differences but effects favoring the intervention arm (0.13 vs. 0.16 in pregnancy and 0.17 vs. 0.24 postpartum).

Corresponding author: Kathryn I. Pollak, Ph.D., 2424 Erwin Road, Suite 602, Durham, NC 27705, Tel: 919-681-4757 Fax: 919-681-4785, kathryn.pollak@duke.edu.

This trial is registered with ClinicalTrials.gov: NCT01040949.

Conclusions—A less intensive intervention promoted cessation equal to more intensive counseling. Postpartum might be a more powerful time to promote cessation among Latino men.

Impact—Less intensive interventions when delivered during teachable moments for Latino men could result in a high smoking cessation rate and could reduce disparities.

The smoking rate among Latino men living in the US is comparable to non-Hispanic White men;¹ Yet Latinos are at higher risk for cardiovascular disease than Whites.² Also, given Latinos are less likely to receive cancer screening and present with later stage cancer³, preventing cancer is imperative in this population. Despite these disparities, Latinos are underrepresented in smoking cessation studies^{4,5-7}. A limited number of trials have tested culturally sensitive cessation interventions among Latino smokers, and fewer have been tested using experimental designs⁸. Interventions have included self-help written materials, nicotine replacement therapy (NRT), group, individual, and telephone counseling^{5, 9-13}. Results from the few trials testing cessation interventions for Latinos indicate that NRT and telephone counseling may help Latinos quit, but biochemically-validated rates are low (~20%), and none has examined or found significant effects 12 months after the intervention. One study indicated that acculturation might play a role in cessation¹⁴; however, this trial did not include random assignment. Clearly, more randomized controlled trials with long follow-up are needed to assess whether an effective intervention can promote smoking cessation among Latinos.

An important factor to consider when developing interventions for Latinos is that most are non-daily smokers or light smokers (i.e., smoke 10 or fewer cigarettes per day¹⁵). Because they differ from traditional heavy smokers, it might be that less intensive approaches are enough to promote cessation. One less intensive and easily disseminable approach is using written materials; it is unknown whether intensive interventions are needed to help Latino smokers quit smoking.

Further, existing interventions can be strengthened by intervening at a time when smokers feel compelled to quit, namely a teachable moment. A teachable moment is a naturally occurring event or situation that increases motivation for behavior change through changes in risk perceptions, emotions, and self-image¹⁶. For instance, when their partner becomes pregnant, Latino men might feel differently about the risk of smoking around her, might have positive or negative emotions related to smoking that they did not have before she was pregnant, and might view themselves differently especially when thinking about their smoking (e.g., a good role model for my children is someone who doesn't smoke). Pregnancy has been shown to be a teachable moment for women as many quit smoking upon learning of their pregnancy¹⁶; whether it is a teachable moment for partners, and particularly Latino partners, needs further exploration.

Our pilot work indicates that Latino expectant fathers have a strong desire to quit smoking¹⁷; thus, targeting smoking cessation during pregnancy might have increased efficacy over other times. What also is unknown is whether motivation to quit increases after the birth of the baby as pregnancy might be a relatively abstract event for men (not women) until the baby is born.

Further, the cultural value of *familismo* emphasizes trust between family members, loyalty to the family, and a general orientation to the family.¹⁸ This central aspect of Latino culture suggests that intervening with individuals, particularly those who are less acculturated, may not be the best approach for promoting smoking cessation among Latino men. Thus, couple-based interventions may be powerful, sustainable, and culturally appropriate as it capitalizes on *familismo*, the influence of the male in the family (*machismo*), and the fathers' motivation for protecting the fetus and new baby. It also is important to examine whether an intervention during pregnancy is more or less effective than promoting cessation postpartum. Further, the effectiveness of an intensive couples-based intervention may vary for daily smokers versus non-daily smokers, as daily smoking likely might require a more intensive intervention. To answer these many questions about how best to promote cessation among Latino men, the aim of this randomized controlled trial was to compare a culturally tailored couples-based intervention plus written materials (self-help smoking cessation guide or *Guia*) and free nicotine replacement therapy (NRT) to a minimal intervention involving written materials plus NRT in promoting smoking cessation among expectant Latino fathers during pregnancy and postpartum. The main aim was to determine whether a more intensive intervention outperformed a less intensive intervention during these potential teachable moments.

Method

Sample

This study was approved by the Duke University School of Medicine Institutional Review Board. We recruited pregnant Latinas and their partners who smoked from 10 urban and rural county health departments in North Carolina. Eligible women had to be at least 16 years of age, between 13 and 29 weeks gestation, and not smoking. Eligible men had to have smoked at least 100 cigarettes in their lifetime, have smoked in the past 30 days (to include non-daily smokers), and be at least 16 years old. For women and men who were 16-18 years old, they were required to be married and thus emancipated (does not require parental consent).

At all of the clinics, nurses introduced the “*Parejas*” (couples) study during women's initial prenatal appointment. Nurses asked women to sign a form agreeing to be contacted by our research staff about the study. *Parejas* study staff called only women who signed the contact form and assessed eligibility. Staff asked women if their partner had ever smoked (some women were unaware of men's current smoking but knew he had smoked at some point). If women were eligible and willing to participate, we asked permission to contact her partner. Staff called men and assessed eligibility. If the man was eligible and willing to participate, staff went to the couples' home to obtain written consent and conduct a baseline survey with the couple. After the baseline survey, staff randomized the couple to one of two study arms from a preset randomization list that was stratified on whether men were daily or non-daily smokers and first time fathers or not. Men randomized to the written materials only arm received the booklet, *No lo deje para mañana deje de fumar hoy: Guía para Dejar de Fumar* and the option of up to 6 weeks of free NRT (nicotine gum or patch). Men randomized to the intensive intervention arm received the booklet, an option of up to 6 weeks of NRT, and also

were told they would receive three counseling sessions during pregnancy (one face-to-face and two via phone) and three postpartum (one face-to-face and two via phone) to help them identify and achieve their goals for quitting smoking and improve their communication with their partner. In both arms, staff briefly discussed elements of the written materials as some of the participants could not read. Staff went to couples' homes again to administer a follow-up survey at the end of pregnancy (~34 weeks gestation) and 12 months post-randomization. We paid all participants \$10 for each survey and an extra \$10 for any saliva samples provided to biochemically validate 7-day point-prevalence abstinence. We collected data from November 2010 through April 2013 and analyzed it in February 2014.

Intervention

This study was a community-based partnership from the inception of the project idea throughout the entire project. In the first year of this study, we worked with our community partner, the local Latino community center, El Centro Hispano, to culturally adapt the intervention. We also convened a Community Advisory Panel that met monthly to review the intervention components, recruitment procedures, and surveys. In addition, we conducted focus groups with expectant Latino couples to obtain their input on integral intervention components.

One thing we learned during our formative work was that for men to be receptive to working on their smoking, we needed to also ask women to make behavior changes. If both members worked on changing a behavior, men reported that they would not "lose face" and feel stigmatized by our intervention. Based on our formative work, we identified written materials and revised the counseling sessions to also promote healthy nutrition and physical activity for the pregnant partners. In addition, the couple-based as well as individual counseling sessions focused on teaching couple's communication skills, namely speaking/listening skills and problem solving. It was hypothesized that when couples communicate better, they will feel less stress and more effective support for quitting and staying quit, which will lead to higher cessation rates.

The first session was with both couple members together in the couple's home and lasted approximately two hours. A male and female counselor presented didactic information on smoking and quitting including a narrated Powerpoint presentation. The first part of the two-part presentation was designed to specifically evoke emotions related to family such as love, pride, protectiveness, and responsibility by including pictures of fathers with infants and children. Information presented in the second part included the health risks of smoking for the smokers and health risks from environmental tobacco smoke for the family. The male counselor talked individually with man to build motivation to quit, identify barriers to quitting, and to set a goal to quit or move towards quitting during the pregnancy. The female counselor talked individually with the woman to identify whether she wanted to work on nutrition or physical activity, build motivation for change, identify barriers to change, and to set realistic goal to work towards during pregnancy. In each sessions, the counselors also taught and asked couples to practice effective communication skills to understand each other's barriers to changing their health behaviors and problem-solve on how to support each other in making these changes. The two subsequent counseling sessions were

conducted over the phone (~15 minutes) with men only and focused on reviewing progress towards quitting and setting new goals, if necessary, to quit smoking and practicing couples communication skills. Postpartum, the male counselor conducted a phone counseling session with men around 4 weeks postpartum, a face-to-face counseling session with both couple members together at 8-10 weeks, and one last counseling call with men at 4 months postpartum.

The counseling protocol was based on Social Cognitive Theory¹⁹ and the teachable moment model.¹⁶ The teachable moment model asserts that for a moment to motivate people to change, it should affect their risk perceptions, emotions, and self-concept. The elements of the face-to-face sessions attempted to address each of these components (e.g., smoking around pregnant women/newborns puts babies at risk, children want their dads to be around for important events, and the role of the dad is to protect his children). The counseling also was informed by Motivational Interviewing (MI).²⁰ Counselors attended a 40-hour training to learn the counseling protocol and MI. Finally, the dyadic portion of the session used communication skills protocols from Cognitive-Behavioral Couple Therapy.²¹ Counselors audio recorded all of their sessions with participants. The clinical supervisors listened to the first three cases for each counselor and then randomly chose 10% of cases to review and provide feedback throughout the study to minimize drift.

Measures

We assessed 7-day point prevalence abstinence and 30-day point-prevalence abstinence at baseline, at the end of pregnancy (~ 34 weeks gestation), and 12 months post-randomization. We also assessed continuous abstinence at both follow-up time-points. Our primary outcome was 7-day point prevalence abstinence; however, given many of the smokers were not daily smokers, we also analyzed secondary effects on 30-day point prevalence abstinence. We biochemically validated men's reports of 7-day point prevalence abstinence. We asked men to provide a 1mL saliva sample. Saliva samples were analyzed for the presence of cotinine using radio-immunoassay (RIA). The presence of cotinine was tested based on an antiserum produced by injecting trans-4-carbozycotinine bound to albumin. Both the inter- and intra-assay variations are lower than 6%, and results are comparable to those produced by gas-liquid chromatography. The sensitivity of this procedure is 0.5mg/dl. Cotinine is a highly sensitive measure for identifying smokers. A cut point of 16 ng/ml for saliva cotinine was used to discriminate abstainers and smokers.

Statistical Methods

The primary objectives of the trial were to test for arm differences in 7-day point prevalence abstinence rates at end of pregnancy and at 12 months post-randomization. Each test was calculated in a logistic regression model with a one-sided alpha of 0.0125; thus the overall alpha level of the trial was controlled at 0.025. As planned at the time the trial was designed, baseline covariates previously shown to be related to the outcome were used as covariates: education (9th grade vs. >9 grade, wantedness of pregnancy (yes/no), and number of cigarettes per day (5 vs. > 5). Secondary objectives were to estimate arm differences in 30-day point-prevalence abstinence and continuous abstinence at end of pregnancy and at 12 months post-randomization. Logistic regression models were used to estimate these two arm

effects, using the three covariates listed above; p-values for these outcomes were not assessed for significance. In all analyses, men who were lost to follow-up were imputed to be smokers, as were men whose cotinine levels indicated that they were active smokers. Given we expected larger intervention effects among non-daily smokers, we analyzed arm differences in non-daily and daily smokers separately.

Results

Sample characteristics

Figure 1 shows our recruitment numbers. Table 1 shows the demographic characteristics of the sample. Follow-up rates for end of pregnancy and 12 months post-randomization were 0.89 and 0.81, respectively. More couples were lost in the intervention arm at the end of pregnancy survey than the control arm (Tx: 14% and Control: 7%). There were no lost-to-follow-up differences by arm at 12 months post-randomization (Tx: 21% and Control: 17%). We examined differential attrition by baseline characteristics. At the end of pregnancy, men who reported monthly income \$1001 and Spanish as the only language spoken at home were lost to follow-up more than those with less monthly income (14% vs. 7%, $p=0.03$) and Spanish and English spoken at home (14% vs. 5%, $p=0.01$). NRT was given about equally to men in the intervention and control arms (Tx: 52% vs. 45%). However, more men in the intervention arm than the control arm reported using it (Tx: 41% vs. Control: 28%).

Cessation

Table 2 shows cessation rates by intervention arm. At the end of pregnancy and 12 months post-randomization, we found high cessation rates in both arms with no added benefit of the more intensive couples-based counseling for 7-day (primary) and 30-day (secondary) cessation. Because we expected differential responses from daily and non-daily smokers, we examined arm differences within these two subgroups at both follow-up time-points (Table 2). Cessation rates were higher among non-daily smokers than daily smokers; in both subgroups, we found no clinically meaningful arm differences. However, unlike among non-daily smokers, arm differences among daily smokers were in the expected direction, favoring the intervention arm for both 7-day and 30-day abstinence but were not statistically significant.

Among both non-daily smokers and daily smokers, cessation rates were higher postpartum than they were in pregnancy. For continuous abstinence (secondary), we found no arm difference in cessation rates with the same pattern for non-daily and daily smokers' cessation rates (Table 2). Sensitivity analyses were run to assess the impact of lost-to-follow-up on the primary and secondary effects of interest. The findings were similar to the intent-to-treat analyses (data not shown).

Discussion

There are three important findings from this trial. First, providing culturally adapted written materials and NRT to Latino expectant and postpartum fathers seems to be as powerful an intervention as providing one-on-one and couple-based counseling, particularly among non-

daily smokers. Second, daily smokers might benefit from a more intensive intervention. Third, quit rates among Latino men are high postpartum.

It was surprising to find no arm differences given the intensity of the intervention. This might be because most of the smokers did not smoke daily, and non-daily smokers did an intensive intervention when their motivation was naturally increased. It also could be that we offered NRT, and more used it than we anticipated as others have reported low rates of NRT use among Latino smokers.²² This cessation effect is promising for future interventions and is higher than has been found in previous trials⁸. If written materials and NRT alone plus home visits for assessment can produce close to 50% cessation rates during pregnancy and postpartum, then less intensive approaches should be used to promote cessation among Latinos, especially among those who are non-daily smokers. Written materials can only be effective, however, with men who know how to read. Future trials should assess the efficacy of written materials that are distributed without having a research assistant explain the highlights as this could serve as an intervention and without home assessment visits.

The arm differences for daily smokers were small and not statistically significant, but the trend favored the more intensive intervention. In the postpartum period, the 30-day cessation rates were almost double in the intervention arm and were almost the exact rates upon which we powered the trial. We might have been underpowered to detect this difference with only 134 daily smokers, less than a third of our sample. However, results show that the combination of couples-based and individual cessation counseling can help daily smokers quit more than materials.

Finally, cessation rates both among daily and non-daily smokers were highest postpartum. This is the opposite effect seen among pregnant smokers, where up to 50% quit smoking when learning they are pregnant and up to 90% return to smoking postpartum. It could be that even though *familismo* is a strong component of Latino culture, pregnancy was not as strong a trigger to motivate cessation as seeing an actual baby. Having the baby in his arms and seeing how environmental tobacco smoke and third hand smoke can affect that child might be the impetus a Latino father needs to quit smoking. They might still believe that when they smoke away from their pregnant wife, that they are protecting the baby, but they also might feel more compelled to quit once the baby has arrived. This effect should be replicated.

This study has limitations that affect interpretation of the results. First, we limited our study to Spanish speakers, and many were from Mexico; thus, our findings might not generalize to English-speaking Latinos or those from other countries. Second, although our attrition rate was low, we were unable to reach some couples at follow up, which also might affect generalizability. While our attrition rates were significantly lower than those in other recent RCT conducted with Latino smokers⁵, we did find reasons other than wanting to participate in our study that contributed to our high no contact rate (e.g., fear, work schedules, etc.). We were unable to biochemically validate the 30-day abstinence rates, which for non-daily smokers is a better measure than 7-day point prevalence abstinence. It is possible that our rates are inflated by social desirability, which is more common among Latinos than

Whites.²³ We have confidence in our 30-day rates, however, given they are considerably lower than our 7-day rates, which we were able to validate. Finally, we recruited smokers via their partners rather than directly; this might have biased our sample.

This was the first study to try to promote cessation among expectant Latinos throughout pregnancy and postpartum. We included light and non-daily smokers as this is representative of Latino smokers in the US and worldwide. We found that for non-daily smokers, providing written materials and NRT during pregnancy promotes cessation. Daily smokers might need a more intensive intervention even in the teachable moment of pregnancy and postpartum. Future interventions might focus only on postpartum and, to be more cost-effective, offer only written materials to non-daily smokers and more intensive counseling only to daily smokers. Interventions also might target postpartum rather than pregnancy as a teachable moment to motivate cessation.

Acknowledgments

This work was supported by R01CA127307.

References

1. Prevention CfDCA. Health disparities experienced by Hispanics - United States (Reprinted from MMWR, vol 53, pg 935-937, 2004). Morbidity and Mortality Weekly Report. 2004;535-7. [PubMed: 15215743]
2. Mensah GA, Dietz WH, Harris VB, Henson R, Labarthe DR, Vinicor F, et al. Prevention and control of coronary heart disease and stroke--nomenclature for prevention approaches in public health: a statement for public health practice from the Centers for Disease Control and Prevention. American journal of preventive medicine. 2005; 29:152-7. [PubMed: 16389142]
3. Society AC. Cancer Facts and Figures. American Cancer Society; Atlanta, GA: 2014.
4. Cox LS, Okuyemi K, Choi WS, Ahluwalia JS. A review of tobacco use treatments in U.S. ethnic minority populations. American journal of health promotion : AJHP. 2011; 25:S11-30. [PubMed: 21510783]
5. Cabriaes JA, Cooper TV, Salgado-Garcia F, Naylor N, Gonzalez E. A randomized trial of a brief smoking cessation intervention in a light and intermittent Hispanic sample. Experimental and clinical psychopharmacology. 2012; 20:410-9. [PubMed: 22731733]
6. Hayes RB, Borrelli B. Differences between Latino daily light and heavier smokers in smoking attitudes, risk perceptions, and smoking cessation outcome. Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco. 2013; 15:103-11. [PubMed: 22589424]
7. Webb MS, Rodriguez-Esquivel D, Baker EA. Smoking cessation interventions among Hispanics in the United States: A systematic review and mini meta-analysis. American journal of health promotion : AJHP. 2010; 25:109-18. [PubMed: 21039291]
8. Castro Y, Basen-Engquist K, Fernandez ME, Strong LL, Eakin EG, Resnicow K, et al. Design of a randomized controlled trial for multiple cancer risk behaviors among Spanish-speaking Mexican-origin smokers. BMC public health. 2013; 13:237. [PubMed: 23506397]
9. Wetter DW, Mazas C, Daza P, Nguyen L, Fouladi RT, Li Y, et al. Reaching and treating Spanish-speaking smokers through the National Cancer Institute's Cancer Information Service. A randomized controlled trial. Cancer. 2007; 109:406-13. [PubMed: 17149758]
10. Woodruff SI, Edwards CC, Conway TL. Enhancing response rates to a smoking survey for enlisted U.S. Navy women. Evaluation review. 1998; 22:780-91. [PubMed: 10345198]
11. Munoz RF, Marin BV, Posner SF, Perez-Stable EJ. Mood management mail intervention increases abstinence rates for Spanish-speaking Latino smokers. American journal of community psychology. 1997; 25:325-43. [PubMed: 9332966]

12. Nevid JS, Javier RA. Preliminary investigation of a culturally specific smoking cessation intervention for Hispanic smokers. *American journal of health promotion : AJHP*. 1997; 11:198–207. [PubMed: 10165099]
13. Leischow SJ, Hill A, Cook G. The effects of transdermal nicotine for the treatment of Hispanic smokers. *Am J Health Behav*. 1996; 20:304–11.
14. Bock BC, Niaura RS, Neighbors CJ, Carmona-Barros R, Azam M. Differences between Latino and non-Latino White smokers in cognitive and behavioral characteristics relevant to smoking cessation. *Addict Behav*. 2005; 30:711–24. [PubMed: 15833576]
15. Trinidad, Perez-Stable EJ, Emery SL, White MM, Grana RA, Messer KS. Intermittent and light daily smoking across racial/ethnic groups in the United States. *Nicotine Tob Res*. 2009; 11:203–10. [PubMed: 19246433]
16. McBride CM, Emmons KM, Lipkus IM. Understanding the potential of teachable moments: the case of smoking cessation. *Health Educ Res*. 2003; 18:156–70. [PubMed: 12729175]
17. Pollak KI, Denman S, Gordon KC, Lyna P, Rocha P, Brouwer RN, et al. Is pregnancy a teachable moment for smoking cessation among US Latino expectant fathers? A pilot study. *Ethnicity & health*. 2010; 15:47–59. [PubMed: 20013439]
18. Lorenzo-Blanco EI, Unger JB, Ritt-Olson A, Soto D, Baezconde-Garbanati L. A longitudinal analysis of Hispanic youth acculturation and cigarette smoking: the roles of gender, culture, family, and discrimination. *Nicotine Tob Res*. 2013; 15:957–68. [PubMed: 23109671]
19. Bandura, A. Self-efficacy mechanism in physiological activation and health-promoting behavior. In: Madden, JM., editor. *Adaptation, Learning, and Affect*. Raven Press; New York: 1986.
20. Miller, WR.; Rollnick, S. *Motivational interviewing : preparing people for change*. 2nd ed.. Guilford Press; New York: 2002.
21. N, E.; DH, B. *Enhanced cognitive behavioral therapy for couples: A contextual approach*. American Psychological Association; Washington, DC: 2002.
22. Levinson AH, Perez-Stable EJ, Espinoza P, Flores ET, Byers TE. Latinos report less use of pharmaceutical aids when trying to quit smoking. *American journal of preventive medicine*. 2004; 26:105–11. [PubMed: 14751320]
23. Davis RE, Resnicow K, Couper MP. Survey Response Styles, Acculturation, and Culture Among a Sample of Mexican American Adults. *J Cross Cult Psychol*. 2011; 42:1219–36. [PubMed: 21927503]

Figure 1. Flow of Participants in Parejas Trial

Table 1

Baseline characteristics of participants

	<i>Totals</i>	<i>Materials alone N(175)</i>	<i>Materials + Counseling N(173)</i>
	M (SD)/ % (N)	M (SD)/ % (N)	M (SD)/ % (N)
Age M (SD)	30 (6)	29 (5)	30 (6)
Education (%)			
9 years	66 (229)	69 (120)	63 (109)
Employment (%)			
Full time	66 (226)	67 (116)	64 (110)
Part time	28 (96)	27 (46)	29 (50)
Not currently employed	6 (23)	6 (11)	7 (12)
Country of Origin (%)			
Mexico	78 (269)	79 (139)	76 (132)
Length of Time in US (%)			
>3 years	94 (326)	94 (165)	93 (161)
Language spoken at home (%)			
Spanish only	68 (237)	70 (133)	66 (114)
Length of relationship (%)			
> 3 years	67(231)	65 (113)	69 (118)
Number of children (%)			
1 or more	77 (267)	76 (133)	77 (134)
Wanted baby now (%)	65 (225)	69 (120)	61 (105)
Difficulty paying bills (%)	38 (131)	36 (62)	40 (69)
Smoking (%)			
Every day (n=134)	39 (134)	37 (64)	41 (70)
Some days (n=214)	61(214)	63 (111)	60 (103)
Cigarettes per day M(Range)			
Daily (n= 134)	8 (5)	8 (5)	8 (6)
Some days (n=214)	3 (3)	3 (2)	4(3)
Smoking inside home (%)	3 (11)	3 (6)	3 (5)

Table 2

Adjusted cessation rates and Odds Ratios (OR) with 95% Confidence Interval (95% CI) for men at the End of Pregnancy (EPS) and 12 month post-randomization (postpartum)

	Less intensive (N=175)		More intensive (N=173)		
	N	Rate	N	Rate	Adj. OR(95%CI)
Primary Outcome					
7-day point prevalence					
End of pregnancy	59	0.31	55	0.30	0.96 (0.60-1.55)
12 months post-randomization	69	0.39	66	0.38	-1.02 (0.65-1.60)
Secondary Outcomes					
30-day abstinence					
End of pregnancy	30	0.14	36	0.19	1.42 (0.81-2.49)
12 months post-randomization	51	0.30	52	0.30	1.14 (0.78-1.84)
Continuous abstinence					
End of pregnancy	19	0.09	27	0.15	1.63 (0.86-3.10)
12 months post-randomization	16	0.07	24	0.12	1.79 (0.90-3.60)
Non-daily smokers (n=214)					
7-day point prevalence					
End of pregnancy	51	0.46	44	0.43	0.92 (0.53-1.60)
12 months post-randomization	58	0.52	49	0.48	0.87 (0.50-1.50)
30-day point prevalence					
End of pregnancy	29	0.26	30	0.29	1.26 (0.68-2.33)
12 months post-randomization	45	0.41	41	0.40	1.00 (0.58-1.75)
Continuous					
End of pregnancy	19	0.15	23	0.19	1.4 (0.71-2.80)
12 months post-randomization	16	0.14	21	0.20	1.5 (0.74-3.18)
Daily Smokers (N=134)					
7-day point prevalence					
End of pregnancy	8	0.13	11	0.16	1.02 90.34-3.10)
12 months post-randomization	11	0.17	17	0.24	1.59 (0.64-3.92)
30-day abstinence					
End of pregnancy	1	0.02	6	0.09	----- [†]
12 months post-randomization	6	0.09	11	0.16	2.12 (0.68-6.63)
Continuous					
End of pregnancy	0	0.00	4	1.00	----
12 months post-randomization	1	0.002	4	0.04	----

[†] Effects are controlled for education, number of cigs smoked per day and wantedness of pregnancy.

[‡] Unable to model due to sample size.