Published in final edited form as:

Hum Psychopharmacol. 2009 April ; 24(3): 225–232. doi:10.1002/hup.1012.

Ziprasidone as an adjuvant for clozapine- or olanzapine-associated medical morbidity in chronic schizophrenia

David C. Henderson, M.D.^{1,2}, Xiaoduo Fan, M.D., M.S.^{1,2}, Paul M. Copeland, M.D.², Bikash Sharma, MBBS¹, Christina P. Borba, M.P.H.¹, Sharon I. Forstbauer¹, Kate Miley¹, Ryan Boxill, Ph.D.¹, Oliver Freudenreich, M.D.^{1,2}, Corey Cather, Ph.D.^{1,2}, A. Eden Evins, M.D.^{1,2}, and Donald C. Goff, M.D.^{1,2}

¹Schizophrenia Program, Massachusetts General Hospital, Boston, MA

²Harvard Medical School, Boston MA

Abstract

Objective—This study sought to examine the effect of ziprasidone on olanzapine or clozapine associated medical morbidity such as insulin resistance, diabetes mellitus and impaired fasting glucose, obesity and hyperlipidemia in patients with schizophrenia or schizoaffective disorder.

Method—This was a six-week, open label trial of ziprasidone 160 mg/day added to a stable dose of olanzapine or clozapine in twenty-one schizophrenia or schizoaffective patients with diabetes mellitus, impaired fasting glucose, or insulin resistance.

Results—Ten olanzapine-treated subjects and eleven clozapine-treated subjects were enrolled in the study. There were no significant differences between the two groups at baseline for age, gender, education, ethnicity, BMI, cholesterol levels, or fasting glucose. At week six, there were no significant changes in weight, BMI, cholesterol levels, or fasting glucose. There was no significant difference in psychotic, negative or depressive symptoms. QTc significantly increased at week 2 but not at week 6.

Conclusions—The addition of 160 mg/day of ziprasidone was well tolerate but did not produce significant improvement in fasting glucose, insulin resistance, hyperlipidemia or lead to weight loss in olanzapine- or clozapine-treated subjects with schizophrenia or schizoaffective disorder.

Keywords

clozapine; olanzapine; ziprasidone; weight; lipids; insulin resistance; schizophrenia

Introduction

Clozapine, an atypical antipsychotic agent, remains the most effective psychopharmaceutical agent for treatment-resistant schizophrenia. Unfortunately, the side effects of clozapine are often difficult for patients to tolerate, particularly sedation and weight gain. Although some clozapine patients are able to remain out of the hospital, they continue to have significant

psychiatric symptoms despite adequate doses of clozapine. In other symptomatic patients, the clozapine dose is limited by significant side effects. Treatment options for patients who do not fully respond to clozapine have not been fully elucidated. Several years ago, Henderson et al. (Henderson and Goff, 1996) reported that chronic schizophrenia patients demonstrated an improved response to clozapine when risperidone was utilized as an adjunctive agent. In this open trial, significant improvements in ratings of positive, negative, and depressive symptoms were found. Furthermore, a more recent study found significant improvement in the disorganized thought subscale of the PANNS (Freudenreich et al., 2007). However, the metabolic sequelae of combination therapy have not been fully investigated, though elevations of the combination lead to significant elevations in prolactin compared to treatment with clozapine alone (Henderson et al., 2001).

Numerous reports of clozapine- and olanzapine-associated insulin resistance, hyperglycemia diabetic ketoacidosis associated with clozapine and olanzapine have emerged (Henderson et al., 2006a, Ananth et al., 2002, Baptista et al., 2002, Caro et al., 2002, Colli et al., 1999, Gianfrancesco et al., 2002, Hagg et al., 1998, Henderson et al., 2000b, Kato and Goodnick, 2001, Newcomer et al., 2002, Koller et al., 2001, Koller and Doraiswamy, 2002, Henderson et al., 2005). By increasing a patient's risk of obesity, antipsychotic agents may be placing patients at risk for associated morbidity and mortality (Pi-Sunyer, 1993). Patients who gain greater than 10% of their total body weight are at risk for developing weight associated conditions such as hypertension and type 2 diabetes mellitus. Henderson et al. found a high rate of diabetes in this cohort as 30 of 82 patients (36.6%) treated with clozapine developed diabetes mellitus over a 60 month period (Henderson et al., 2000a). Olanzapine has also been associated with significant weight gain (>7% total body weight), insulin resistance, hyperlipidemia, and new onset DM (Wilson et al., 2003, Opp and Hildebrandt, 2002, Seaburg et al., 2001, Melkersson et al., 2000, Cohen et al., 2003). While switching to an agent associated with less weight gain (such as ziprasidone or aripiprazole) offers the greatest opportunity for resolution of clozapine- or olanzapine-associated diabetes mellitus, many clinicians and patients are reluctant to do so. Because clozapine is reserved for treatment resistant schizophrenia patients, switching to another antipsychotic agent may not be feasible. The best intervention for clozapine patients may be to add an agent associated with less weight gain and eventually lower the dose of clozapine. In this setting, weight loss and improvements in glucose metabolism may occur.

Additionally, when switching patients treated with antipsychotic agents, many times clinicians make the switch rapidly and provide little time for overlap. Patients treated in this manner may have a higher risk of relapse which is often considered a drug trial failure, but in fact may be that the second agent did not have time to take hold. Determining whether combination antipsychotic agents therapy results in an improvement in metabolic parameters and psychopathology would be helpful to clinicians in their decision-making regarding pharmacotherapy. The results of this study may provide clinicians and patients with a potentially effective intervention to counteract the weight and metabolic effects of clozapine and olanzapine, while also showing the benefits of continued or improved efficacy. It may also allow clinicians and patients to be more comfortable with a much slower and safer switch to ziprasidone as the medical morbidity benefits may begin as soon as the drug is

started. If a patient experiences early medical benefits from ziprasidone, the speed of switching can be much slower while also significantly reducing the risk of relapse.

Ziprasidone is an atypical antipsychotic agent with high affinity for dopamine D₂ and 5-HT_{2A} receptors where it acts as an antagonist. Clinical trials indicate that ziprasidone is effective against positive, negative and affective symptoms in schizophrenia and schizoaffective disorder with minimal motor, cognitive, weight, prolactin related, or anticholinergic side effects (Daniel and Copeland, 2000). Ziprasidone also acts as an agonist of the 5-HT_{1A} receptor and moderately inhibits the re-uptake of serotonin and norepinephrine. The low liability of ziprasidone with respect to weight gain may have significance for patients even beyond the cardiovascular and other health effects (Cohen et al., 2003, Spivak et al., 2002, Alao et al., 2002, Wetterling, 2001, Allison et al., 1999). The distressing side effect of weight gain frequently leads to patient-driven decisions to switch or discontinue medications.

Ziprasidone was chosen as its side effect profile greatly differs from both clozapine and olanzapine and there was already pilot data examining aripiprazole (Henderson et al., 2006b). Different receptor affinities may play a role in weight gain and the development of diabetes mellitus in patients on antipsychotic medication. Aripiprazole may have partial agonist properties at 5-HT_{1A}, 5-HT_{2A}, and 5-HT_{2C} serotonin receptors (Shapiro et al., 2003), whereas ziprasidone is a D₂ and 5-HT_{2A} antagonist, and an agonist at 5-HT_{1A} receptors. Clozapine and olanzapine, which offer the greatest risk of weight gain, are structurally similar and both have binding capacities for serotonin 5-HT_{2C}, histamine H₁, and muscarinic M₁ receptors (Bymaster et al., 1996, Millan et al., 1998). A study by Kroeze et al. (Kroeze et al., 2003) found that affinities for histamine H₁, alpha(1A) adrenergic, 5-HT_{2C}, and 5-HT₆ receptors were most strongly correlated with weight gain when screening 17 typical antipsychotics. While neither aripiprazole nor ziprasidone have high affinities for the histamine H₁ receptors, the affinity of ziprasidone is lower than that of aripiprazole (Kroeze et al., 2003). This discrepancy may be a factor in each drug's ability to counteract clozapine and olanzapine associated weight gain.

Additionally, 5-HT_{2C} receptors have been implicated in the control of appetite (Vickers et al., 1999) and a variant of the 5-HT_{2C} receptor gene (-759C/T) was associated with less weight gain in a study of first episode schizophrenia patients (Reynolds et al., 2003). Therefore, it is conceivable that the minimal weight gain associated with aripiprazole may be due to its moderate binding affinity at 5-HT_{2C} receptors. Ziprasidone does not share this capacity for 5-HT_{2C} receptors. Accordingly, effects on clozapine and olanzapine associated weight gain were not demonstrated with the addition of ziprasidone comparable to aripiprazole.

In the present pilot study, we investigated the benefits for metabolic response, weight loss, and efficacy for positive and negative symptoms of ziprasidone 160 mg/day added to a stable dose of clozapine- or olanzapine-treated subjects with schizophrenia or schizoaffective disorder over a six week period.

Aims of the study

We investigated the efficacy of ziprasidone for weight loss in clozapine or olanzapine treated schizophrenia subjects during a 6-week open label trial. Based on our previous study with aripiprazole six weeks was determined to be adequate to see significant weight loss and reduction in lipids (Henderson et al., 2006b). The hypotheses were that the addition of ziprasidone 160 mg/day to stable clozapine- and olanzapine-treated schizophrenia or schizoaffective disorder subjects with diabetes mellitus (DM), impaired fasting glucose or insulin resistance would result in significant weight loss, and improvements in glucose and lipid metabolism over a 6 week period.

Methods

This six-week open label trial was conducted in the adult outpatient clinic of an urban community mental health center. The study was approved by the Institutional Review Board of the Massachusetts Department of Mental Health. After providing written informed consent all participants underwent a diagnostic evaluation by a research psychiatrist using the Structured Clinical Interview for DSM-IV (SCID) (Spitzer RL, 1992).

Subjects on clozapine or olanzapine were recruited for the study if they met the following criteria: diagnosis of schizophrenia or schizoaffective disorder, age 18–65 years, capable of providing informed consent, treatment with clozapine or olanzapine for at least one year and with a stable dose being administered for at least one month. Subjects were required to have a history of diabetes mellitus (DM), impaired fasting glucose or insulin resistance to participate in the study. Impaired fasting glucose was defined as a fasting glucose of greater than or equal to 100 mg/dL and less than 126 mg/dL. Insulin resistance was defined as fasting insulin greater or equal to 15 mU/L or a Homeostasis Model Assessment-Insulin Resistance (HOMA-IR) greater or equal to 2. Subjects were excluded from the study if they were unable to provide informed consent, had a significant unstable medical illness such as unstable cardiac disease, a current substance abuse problem, treatment with medications that significantly prolonged QTc or history of prolongation of QTc interval (>450msec) on EKG, clinically significant EKG abnormalities, hepatic or renal impairment, cancer, poorly-controlled seizure disorder, previous treatment with ziprasidone or treatment with more than one antipsychotic agent. Thirty patients were screened and 24 consented for the study. Of the 6 subjects that did not consent, three did not meet the BMI criteria and three decided not to participate. Two subjects consented but were lost to follow-up and one subject withdrew consent after receiving one dose of ziprasidone. The target for this pilot study was to have 20 subjects complete 6 weeks of the study (10 clozapine and 10 olanzapine).

Subjects were treated with open label ziprasidone 40 mg twice daily for the first two weeks. After two weeks, ziprasidone was increased to 80 mg twice daily as tolerated. Clozapine or olanzapine doses remained unchanged throughout the study. Patients that chose to remain on ziprasidone after the completion of the six-week trial were assessed at week ten.

Assays

Fasting blood samples were assayed for a complete blood count and concentrations of plasma glucose, cholesterol (total, HDL and LDL), and triglycerides at baseline, week 4 and week 6 using standard laboratory procedures. Insulin immunometric assays were performed using an Immulite Analyzer (Diagnostic Product Corporation, Los Angeles, CA) with an intraassay coefficient of variation of 4.2–7.6%. HOMA-IR was calculated from fasting glucose and insulin values at baseline and week six (Kissebah et al., 1982). The same assays were completed at week 10 for all subjects who chose to remain on Ziprasidone.

Subjects were assessed with a battery of symptom rating scales at baseline, weeks 2, 4 and 6. The assessment battery included The Positive and Negative Syndrome Scale (PANSS) (Kay et al., 1987), Scale for Assessment of Negative Symptoms (SANS), Hamilton Depression Rating Scale (HAM-D), Fatigue Scale Inventory (FSI) (Hann et al., 2000), the Quality of Life Scale (QOL), the Simpson-Angus Scale, Barnes Akathisia Scale, and the Abnormal Involuntary Movement Scale (AIMS). A single rater performed all assessments throughout the trial. A physical examination and medical history was performed at baseline and measurement of vital signs, weight, and waist/hip circumference at each visit. Diet and exercise interventions were not performed during the trial.

Statistical Analysis

Two-tailed paired samples t-tests were conducted to compare baseline and week 6 values for body weight, BMI, fasting lipids, fasting glucose, fasting insulin, HOMA-IR, fasting lipids, and clinical symptoms. Scores from week 4 (last observation carried forward (LOCF) method) were used for two subjects with missing end point measurements. Separate analyses were performed for the clozapine group and the olanzapine group. Additionally, analysis of subjects with DM and subjects without DM was performed. For all analysis, P-values of less than 0.05 were considered significant.

Results

Twenty-four subjects consented to participate in the study. Two subjects were lost to follow-up prior to starting the study medication; and one subject discontinued the medication after one dose. The remaining twenty-one subjects completed the six-week trial and are included in all analyses. Eleven patients (52%) were receiving clozapine and ten patients (48%) were receiving olanzapine.

The demographic data are summarized in Table 1. The mean age of the subjects was 49 ± 8 years and 17 (81%) were male. Fifteen subjects (71%) were Caucasian, 4 (19%) were Black, and 1 (5%) was Hispanic. There were no significant differences among the clozapine and olanzapine treated groups for race, marital status, employment status, or family history of hypertension or diabetes. Weight, BMI, cholesterol, and waist and hip measurements did not differ significantly between the clozapine treated group and the olanzapine treated group at baseline. Within the clozapine-treated group, 9 subjects (82%) were smokers, compared to 4 subjects (40%) in the olanzapine treated group ($p=0.049$). Eight subjects (38%) were treated for type 2 diabetes mellitus at the time of consent for the study.

Table 2 shows anthropometric changes from baseline to week 6 for each group and the entire sample. Comparing baseline to week 6, there was no significant difference in weight, BMI, waist circumference, or waist/hip ratio. The mean weight for the entire sample was 230 lbs $\pm$ 35 at baseline, and 229 lbs $\pm$ 35 at week 6 ($P=0.73$). There were no significant changes in total cholesterol, triglycerides, HDL-cholesterol or LDL-cholesterol. For the entire sample, the mean total cholesterol was 186 mg/dL $\pm$ 40 at baseline, and 185mg/dL $\pm$ 34 at week 6 ($P = 0.929$). Triglyceride levels decreased from 256 mg/dL $\pm$ 159 and 231 mg/dL $\pm$ 34 but was not statistically significant ($P =0.24$). There was no significant difference in any of the above outcome measures comparing baseline to week 6 when analyzing subgroups for clozapine or olanzapine alone. In the clozapine treated group, fasting plasma glucose and fasting serum insulin did not change significantly (Table 2). The mean fasting plasma glucose changed from 125 mg/dL $\pm$ 43 at baseline to 130 mg/dL $\pm$ 46 at week 6 ($P=0.719$) in the clozapine treated group. For non-diabetic subjects receiving either olanzapine or clozapine, there were similarly no significant changes in fasting plasma glucose from 104 mg/dL $\pm$ 13 at baseline to 102 mg/dL $\pm$ 18 at week 6 ($P=0.577$), or fasting insulin 16.3 $\pm$ 8.4 vs. 15.2 $\pm$ 11.06 ($p=.654$). Furthermore, in non-diabetic subjects, HOMA-IR decreased from 4.2 $\pm$ 2.2 at baseline to 3.9 $\pm$ 3.4 at week 6 but was not significant ($P =0.695$).

In addition, ziprasidone produced no significant differences between baseline and week 6 on the PANSS total scores and subscale scores, the SANS, or the Ham-D ($P_s > 0.129$) (Table 3). There were no serious adverse events as a result of treatment with ziprasidone. Four subjects (18%) experienced constipation, 2 subjects (9%) experienced diarrhea and 2 subjects (9%) experienced tremors while on ziprasidone. From baseline to week 2, the mean QTc increased from 417 $\pm$ 15 msec to 430 $\pm$ 16 ($P= 0.002$); however, the change in mean QTc from 417 $\pm$ 15 msec at baseline to 420 $\pm$ 21 at week 6 was not statistically significant ($P=0.507$).

Discussion

In this open label trial, the addition of ziprasidone to a steady dose of olanzapine or clozapine resulted in no significant benefit in weight loss or other metabolic parameters in patients with schizophrenia or schizoaffective disorder. Previous research has demonstrated the efficacy of ziprasidone to significantly improve weight and plasma lipids when switching treatment from another atypical antipsychotic medication. A study by Weiden et al. estimated weight loss of 10.3% of mean initial weight and total cholesterol decrease of 9.2% over 58 weeks for patients when switched from olanzapine to ziprasidone (Weiden et al., 2007). However, the lack of similar effects when used in combination with clozapine and olanzapine suggest that ziprasidone is ineffectual as an adjuvant therapy for weight and metabolism.

These results differ from a previous study, where the addition of aripiprazole to clozapine treatment resulted in significant decreases in weight ($P=0.003$), BMI ($P=0.004$), total cholesterol ($P=0.002$), total triglycerides ($P=0.040$), and HDL cholesterol ($P = 0.020$) (Henderson et al., 2006b).

The sample size in this study may not have been adequate to demonstrate the effectiveness of ziprasidone as an adjuvant to clozapine or olanzapine, and resulted in a type II error. It is also plausible that the receptor binding profile of ziprasidone does not significantly counteract the mechanisms of weight gain and other metabolic disturbances associated with clozapine and olanzapine.

In the present study, ziprasidone did not worsen extrapyramidal side effects or psychotic symptoms. It also did not result in any significant adverse events or increases in resting heart rate or blood pressure. There was a non-clinically relevant increase in QTc from baseline to week 2 which was not evident at week 6.

Conclusions

The addition of 160 mg/day of ziprasidone was well tolerate but did not produce significant improvement in fasting glucose, insulin resistance, hyperlipidemia or lead to weight loss in olanzapine- or clozapine-treated subjects with schizophrenia or schizoaffective disorder. This combination may not be of benefit to combat the medical morbidity associated with antipsychotic drugs. Investigation into new intervention strategies to control antipsychotic associated weight gain and metabolic disturbances is needed.

Acknowledgments

Funding Source: An investigator initiated grant from Pfizer Pharmaceuticals.

This study was funded by an investigator initiated grant from Pfizer Pharmaceuticals. Dr David C. Henderson is supported by a research grant from Solvay, Takeda and is a recipient of honorariums from Bristol-Myers Squibb, Janssen L.P., Pfizer, Inc., and Solvay Pharmaceuticals, and COVANCE, Primedia, Reed Medical Education. The research of Dr Donald C. Goff is supported by Pfizer, Cephalon, Janssen and he is a recipient of honorariums from Xenoport, Dainippon Sumitomo, Solvay/Wyeth, Bristol Meyer Squibb, Organon, Proteus, Genactics, Forest Laboratories, Xytis, MedReviews, LLC, and Vanda Pharmaceuticals, Primedia, Reed Medical Education. Dr Paul M. Copeland is a recipient of honorariums from Eli Lilly, Merck, Takeda, Sanofi-Aventis. The research of Dr Xiadou Fan is supported by Eli Lilly and he is a recipient of honorarium from Eli Lilly. The research of Dr Oliver Freudenreich is supported by Cephalon and he is a recipient of honorariums from Primedia, Reed Medical Education. The research of Dr A. Eden Evins is supported by Pfizer, GSK, NIDA CDDC. Drs Ryan Boxill, Corey Cather, and Bikash Sharma and Ms Christina P. Borba, Kate Miley, and Sharon I. Forstbauer report no disclosures. A copy of the research protocol may be requested from David C. Henderson, M.D. at dchenderson@partners.org. The statistical analysis was performed by Dr Xiadou Fan of the Massachusetts General Hospital Schizophrenia Research Program.

References

- ALAO AO, MALHOTRA K, DEWAN MJ. Comparing the side effect profile of the atypical antipsychotics. *West Afr J Med*. 2002; 21:313–5. [PubMed: 12665274]
- ALLISON DB, MENTORE JL, HEO M, CHANDLER LP, CAPPELLERI JC, INFANTE MC, WEIDEN PJ. Antipsychotic-induced weight gain: a comprehensive research synthesis. *Am J Psychiatry*. 1999; 156:1686–96. [PubMed: 10553730]
- ANANTH J, VENKATESH R, BURGOYNE K, GUNATILAKE S. Atypical antipsychotic drug use and diabetes. *Psychother Psychosom*. 2002; 71:244–54. [PubMed: 12207104]
- BAPTISTA T, KIN NM, BEAULIEU S, DE BAPTISTA EA. Obesity and Related Metabolic Abnormalities during Antipsychotic Drug Administration: Mechanisms, Management and Research Perspectives. *Pharmacopsychiatry*. 2002; 35:205–19. [PubMed: 12518268]
- BYMASTER FP, CALLIGARO DO, FALCONE JF, MARSH RD, MOORE NA, TYE NC, SEEMAN P, WONG DT. Radioreceptor binding profile of the atypical antipsychotic olanzapine. *Neuropsychopharmacology*. 1996; 14:87–96. [PubMed: 8822531]

- CARO JJ, WARD A, LEVINTON C, ROBINSON K. The risk of diabetes during olanzapine use compared with risperidone use: a retrospective database analysis. *J Clin Psychiatry*. 2002; 63:1135–9. [PubMed: 12523873]
- COHEN S, FITZGERALD B, OKOS A, KHAN S, KHAN A. Weight, lipids, glucose, and behavioral measures with ziprasidone treatment in a population with mental retardation. *J Clin Psychiatry*. 2003; 64:60–2. [PubMed: 12590625]
- COLLI A, COCCIOLO M, FRANCOBANDIERA F, ROGANTIN F, CATTALINI N. Diabetic ketoacidosis associated with clozapine treatment. *Diabetes Care*. 1999; 22:176–7. [PubMed: 10333925]
- DANIEL DG, COPELAND LF. Ziprasidone: comprehensive overview and clinical use of a novel antipsychotic. *Expert Opin Investig Drugs*. 2000; 9:819–28.
- FREUDENREICH O, HENDERSON DC, WALSH JP, CULHANE MA, GOFF DC. Risperidone augmentation for schizophrenia partially responsive to clozapine: a double-blind, placebo-controlled trial. *Schizophr Res*. 2007; 92:90–4. [PubMed: 17321111]
- GIANFRANCESCO FD, GROGG AL, MAHMOUD RA, WANG RH, NASRALLAH HA. Differential effects of risperidone, olanzapine, clozapine, and conventional antipsychotics on type 2 diabetes: findings from a large health plan database. *J Clin Psychiatry*. 2002; 63:920–30. [PubMed: 12416602]
- HAGG S, JOELSSON L, MJORNDAL T, SPIGSET O, OJA G, DAHLQVIST R. Prevalence of diabetes and impaired glucose tolerance in patients treated with clozapine compared with patients treated with conventional depot neuroleptic medications. *J Clin Psychiatry*. 1998; 59:294–9. [PubMed: 9671341]
- HANN DM, DENNISTON MM, BAKER F. Measurement of fatigue in cancer patients: further validation of the Fatigue Symptom Inventory. *Qual Life Res*. 2000; 9:847–54. [PubMed: 11297027]
- HENDERSON D, GOFF D. Risperidone as an adjunct to clozapine therapy in chronic schizophrenics. *Journal of Clinical Psychiatry*. 1996; 57:395–397. [PubMed: 9746446]
- HENDERSON DC, CAGLIERO E, COPELAND PM, BORBA CP, EVINS E, HAYDEN D, WEBER MT, ANDERSON EJ, ALLISON DB, DALEY TB, SCHOENFELD D, GOFF DC. Glucose Metabolism in Patients With Schizophrenia Treated With Atypical Antipsychotic Agents: A Frequently Sampled Intravenous Glucose Tolerance Test and Minimal Model Analysis. *Arch Gen Psychiatry*. 2005; 62:19–28. [PubMed: 15630069]
- HENDERSON DC, CAGLIERO E, GRAY C, NASRALLAH RA, HAYDEN DL, SCHOENFELD DA, GOFF DC. Clozapine, diabetes mellitus, weight gain, and lipid abnormalities: A five-year naturalistic study. *American Journal of Psychiatry*. 2000a; 157:975–981. [PubMed: 10831479]
- HENDERSON DC, CAGLIERO E, GRAY C, NASRALLAH RA, HAYDEN DL, SCHOENFELD DA, GOFF DC. Clozapine, diabetes mellitus, weight gain, and lipid abnormalities: A five-year naturalistic study. *Am J Psychiatry*. 2000b; 157:975–81. [PubMed: 10831479]
- HENDERSON DC, COPELAND PM, BORBA CP, DALEY TB, NGUYEN DD, CAGLIERO E, EVINS AE, ZHANG H, HAYDEN DL, FREUDENREICH O, CATHER C, SCHOENFELD DA, GOFF DC. Glucose metabolism in patients with schizophrenia treated with olanzapine or quetiapine: a frequently sampled intravenous glucose tolerance test and minimal model analysis. *J Clin Psychiatry*. 2006a; 67:789–97. [PubMed: 16841629]
- HENDERSON DC, GOFF DC, CONNOLLY CE, BORBA CP, HAYDEN D. Risperidone added to clozapine: impact on serum prolactin levels. *J Clin Psychiatry*. 2001; 62:605–8. [PubMed: 11561931]
- HENDERSON DC, KUNKEL L, NGUYEN DD, BORBA CP, DALEY TB, LOUIE PM, FREUDENREICH O, CATHER C, EVINS AE, GOFF DC. An exploratory open-label trial of aripiprazole as an adjuvant to clozapine therapy in chronic schizophrenia. *Acta Psychiatr Scand*. 2006b; 113:142–7. [PubMed: 16423166]
- KATO MM, GOODNICK PJ. Antipsychotic medication: effects on regulation of glucose and lipids. *Expert Opin Pharmacother*. 2001; 2:1571–82. [PubMed: 11825300]
- KAY SR, OPLER LA, LINDENMAYER JP. Reliability and validity of the positive and negative syndrome scale for schizophrenics. *Psychiatry Research*. 1987; 23:99–110. [PubMed: 3363019]

- KISSEBAH A, VYDELINGUM N, MURRY R, EVENS D, HARTZ A, KALKHOFF R, ADAMS P. Relationship of body fat distribution to metabolic complications of obesity. *J Clin Endocrinol Metab.* 1982; 54:254–260. [PubMed: 7033275]
- KOLLER E, SCHNEIDER B, BENNETT K, DUBITSKY G. Clozapine-associated diabetes. *Am J Med.* 2001; 111:716–23. [PubMed: 11747852]
- KOLLER EA, DORAISWAMY PM. Olanzapine-associated diabetes mellitus. *Pharmacotherapy.* 2002; 22:841–52. [PubMed: 12126218]
- KROEZE WK, HUFSEISEN SJ, POPADAK BA, RENOCK SM, STEINBERG S, ERNSBERGER P, JAYATHILAKE K, MELTZER HY, ROTH BL. H1-histamine receptor affinity predicts short-term weight gain for typical and atypical antipsychotic drugs. *Neuropsychopharmacology.* 2003; 28:519–26. [PubMed: 12629531]
- MELKERSSON KI, HULTING AL, BRISMAR KE. Elevated levels of insulin, leptin, and blood lipids in olanzapine-treated patients with schizophrenia or related psychoses. *J Clin Psychiatry.* 2000; 61:742–9. [PubMed: 11078035]
- MILLAN MJ, SCHREIBER R, DEKEYNE A, RIVET JM, BEROETS K, MAVRIDIS M, SEBBAN C, MAUREL-REMY S, NEWMAN-TANCREDI A, SPEDDING M, MULLER O, LAVIELLE G, BROCCO M. S 16924 ((R)-2-[1-[2-(2,3-dihydro-benzo[1,4] dioxin-5-yloxy)-ethyl]-pyrrolidin-3yl]-1-(4-fluoro-phenyl)-ethanone), a novel, potential antipsychotic with marked serotonin (5-HT)1A agonist properties: II. Functional profile in comparison to clozapine and haloperidol. *J Pharmacol Exp Ther.* 1998; 286:1356–73. [PubMed: 9732399]
- NEWCOMER JW, HAUPT DW, FUCETOLA R, MELSON AK, SCHWEIGER JA, COOPER BP, SELKE G. Abnormalities in glucose regulation during antipsychotic treatment of schizophrenia. *Arch Gen Psychiatry.* 2002; 59:337–45. [PubMed: 11926934]
- OPP D, HILDEBRANDT C. Olanzapine-associated type 2 diabetes mellitus. *Schizophr Res.* 2002; 56:195–6. [PubMed: 12084434]
- PI-SUNYER FX. Medical hazards of obesity. *Ann Intern Med.* 1993; 119:655–60. [PubMed: 8363192]
- REYNOLDS GP, ZHANG Z, ZHANG X. Polymorphism of the promoter region of the serotonin 5-HT(2C) receptor gene and clozapine-induced weight gain. *Am J Psychiatry.* 2003; 160:677–9. [PubMed: 12668355]
- SEABURG HL, MCLENDON BM, DORAISWAMY PM. Olanzapine-associated severe hyperglycemia, ketonuria, and acidosis: case report and review of literature. *Pharmacotherapy.* 2001; 21:1448–54. [PubMed: 11714220]
- SHAPIRO DA, RENOCK S, ARRINGTON E, CHIODO LA, LIU LX, SIBLEY DR, ROTH BL, MAILMAN R. Aripiprazole, a novel atypical antipsychotic drug with a unique and robust pharmacology. *Neuropsychopharmacology.* 2003; 28:1400–11. [PubMed: 12784105]
- SPITZER RLWJ, GIBBON M, FIRST MB. The Structured Clinical Interview for DSM-III-R (SCID). I: History, rationale, and description. *Arch Gen Psychiatry.* 1992; 49:624–9. [PubMed: 1637252]
- SPIVAK B, ALAMY SS, JARSKOG LF, SHEITMAN BB, LIEBERMAN JA. Ziprasidone alternative for olanzapine-induced hyperglycemia. *Am J Psychiatry.* 2002; 159:1606. [PubMed: 12202289]
- VICKERS SP, CLIFTON PG, DOURISH CT, TECOTT LH. Reduced satiating effect of d-fenfluramine in serotonin 5-HT(2C) receptor mutant mice. *Psychopharmacology (Berl).* 1999; 143:309–14. [PubMed: 10353435]
- WEIDEN PJ, NEWCOMER JW, LOEBEL AD, YANG R, LEOVITZ HE. Long-Term Changes in Weight and Plasma Lipids during Maintenance Treatment with Ziprasidone. *Neuropsychopharmacology.* 2007
- WETTERLING T. Bodyweight gain with atypical antipsychotics. A comparative review. *Drug Saf.* 2001; 24:59–73. [PubMed: 11219487]
- WILSON DR, D'SOUZA L, SARKAR N, NEWTON M, HAMMOND C. New-onset diabetes and ketoacidosis with atypical antipsychotics. *Schizophr Res.* 2003; 59:1–6. [PubMed: 12413635]

Table 1

Demographic and clinical characteristics of 21 schizophrenia patients treated with clozapine or olanzapine.

Characteristic	Entire sample (N=21)	Clozapine (N=11)	Olanzapine (N=10)	Group Comparison	p value
Age	49 ± 8	48 ± 7	52 ± 8	t(19)= 1.15	0.263
Education	12 ± 2	12 ± 2	11 ± 2	t(19)= -0.53	0.603
Gender, N (%)				X ² (1)= 1.01	0.314
Male	17 (81)	8 (73)	9 (90)		
Female	4 (19)	3 (27)	1 (10)		
Race, N (%)				X ² (3)= 2.02	0.568
Caucasian	15 (71)	7 (64)	8 (80)		
Black	4 (19)	2 (18)	2 (20)		
Hispanic	1 (5)	1 (9)	0 (0)		
Other	1 (5)	1 (9)	0 (0)		
Marital Status, N (%)				X ² (4)= 2.29	0.682
Single	14 (67)	7 (64)	7 (70)		
Married	2 (10)	1 (9)	1 (10)		
Separated	1 (5)	0 (0)	1 (10)		
Divorced	3 (14)	2 (18)	1 (10)		
Widowed	1 (5)	1 (9)	0 (0)		
Employment status, N (%)				X ² (8)= 8.05	0.429
Employed	8 (38)	5 (46)	3 (30)		
Unemployed	13 (62)	6 (55)	7 (70)		
History of hypertension, N (%)				X ² (1)= 0.29	0.593
Yes	3 (14)	2 (18)	1 (10)		
No	18 (86)	9 (82)	9 (90)		
Smoking status, N (%)				X ² (1)= 3.88	0.049
Non-smoker	8 (38)	2 (18)	6 (60)		
Smoker	13 (62)	9 (82)	4 (40)		
Family history of diabetes				X ² (1)= 0.40	0.528
Yes	12 (57)	7 (64)	5 (50)		
No	9 (43)	4 (36)	5 (50)		

<i>Characteristic</i>	Entire sample (N=21)	Clozapine (N=11)	Olanzapine (N=10)	Group Comparison	<i>p</i> value
Presence of diabetes					
Yes	8 (38)	6 (55)	2 (20)		
No	13 (62)	5 (46)	8 (80)	X ² (1)= 2.65	0.104

Notes: Values are expressed as mean ± SD unless otherwise indicated.

Table 2

Anthropometric and metabolic measures over 6-week treatment of ziprasidone as an adjuvant for clozapine- or olanzapine-treated patients with schizophrenia (N=21).

<i>Measurement</i>	Baseline	Week 6	Within Group comparison	p value
Entire sample (N=21)				
Fasting plasma glucose (mg/dL)	123 ± 51	123 ± 48	t(20)= -0.06	0.956
Fasting serum insulin (μIU/L)	15.0 ± 7.7	15.0 ± 9.5	t(20)= -0.02	0.988
HOMA-IR (log transformed)	4.5 ± 3.0	4.7 ± 4.0	t(20)= -0.22	0.826
HBA1C	6.5 ± 1.2	6.6 ± 1.1	t(19)= -0.26	0.795
Leptin (ng/mL)	70.0 ± 64.6	66.5 ± 57.0	t(19)= 0.51	0.615
Total cholesterol (mg/dL)	186 ± 40	185 ± 34	t(19)= 0.09	0.929
Triglyceride (mg/dL)	256 ± 159	231 ± 124	t(18)= 1.21	0.241
HDL (mg/dL)	43 ± 11	44 ± 9	t(18)= -0.99	0.334
LDL (mg/dL)	104 ± 27	102 ± 27	t(15)= 0.59	0.561
Weight (lb)	229.7 ± 34.9	229.3 ± 35.1	t(20)= 0.35	0.730
BMI (kg/m ²)	34.8 ± 4.0	34.8 ± 4.1	t(20)= 0.32	0.754
Waist circumference (cm)	108.4 ± 17.2	112.1 ± 7.2	t(20)= -1.09	0.289
Widest hip circumference (cm)	114.7 ± 19.3	118.6 ± 9.8	t(20)= -1.14	0.267
Clozapine group (N=11)				
Fasting plasma glucose (mg/dL)	125 ± 43	130 ± 46	t(10)= -0.37	0.719
Fasting serum insulin (μIU/L)	14.8 ± 6.9	15.8 ± 11.4	t(10)= -0.36	0.728
HOMA-IR (log transformed)	4.9 ± 3.6	4.9 ± 3.6	t(10)= -0.03	0.980
HBA1C	6.9 ± 1.2	7.0 ± 1.1	t(9)= -0.50	0.629
Leptin (ng/mL)	73.6 ± 63.8	72.4 ± 59.1	t(10)= 0.31	0.766
Total cholesterol (mg/dL)	183 ± 29	178 ± 33	t(10)= 0.58	0.574
Triglyceride (mg/dL)	293 ± 183	273 ± 129	t(10)= 0.57	0.580
HDL (mg/dL)	42 ± 13	42 ± 9	t(10)= -0.47	0.647
LDL (mg/dL)	95 ± 23	89 ± 23	t(7)= 0.88	0.406
Weight (lb)	230.1 ± 40.8	231.4 ± 40.8	t(10)= -1.15	0.278
BMI (kg/m ²)	36.1 ± 3.7	36.3 ± 3.7	t(10)= -1.22	0.251

<i>Measurement</i>	Baseline	Week 6	Within Group comparison	<i>p</i> value
Waist circumference (cm)	112.6 ± 9.5	113.1 ± 6.3	t(10)= - 0.29	0.779
Widest hip circumference (cm)	119.8 ± 11.8	119.6 ± 9.4	t(10)= 0.07	0.942
Olanzapine group (N=10)				
Fasting plasma glucose (mg/dL)	121 ± 61	116.5 ± 51.9	t(9)= 0.68	0.517
Fasting serum insulin (μIU/L)	15.1 ± 8.9	14.2 ± 7.5	t(9)= 0.33	0.747
HOMA-IR (log transformed)	4.1 ± 2.1	4.5 ± 4.6	t(9)= - 0.27	0.792
HBA1C	6.2 ± 1.2	6.2 ± 0.9	t(9)= 0.15	0.885
Leptin (ng/mL)	64.6 ± 69.2	59.2 ± 56.9	t(8)= 0.42	0.686
Total cholesterol (mg/dL)	189 ± 51	193 ± 36	t(8)= - 0.32	0.754
Triglyceride (mg/dL)	207 ± 110	173 ± 96	t(7)= 1.69	0.135
HDL (mg/dL)	45 ± 8	47 ± 9	t(7)= - 1.02	0.341
LDL (mg/dL)	114 ± 28	114 ± 26	t(7)= - 0.04	0.973
Weight (lb)	229.2 ± 29.3	227.0 ± 29.6	t(9)= 1.16	0.276
BMI (kg/m ²)	33.4 ± 4.0	33.1 ± 3.9	t(9)= 1.18	0.267
Waist circumference (cm)	103.9 ± 22.6	111.0 ± 8.2	t(9)= - 1.05	0.322
Widest hip circumference (cm)	109.2 ± 24.6	117.5 ± 10.7	t(9)= - 1.23	0.250

Notes: 1) Values are expressed as mean ± SD unless otherwise indicated; 2) BMI: body mass index; 3) Waist circumference is taken from iliac waist measures; 4) BMI: body mass index; 5) HOMA-IR: homeostasis model of assessment of insulin resistance.

Table 3

Psychopathology measures over a 6-week treatment of ziprasidone supplemented to clozapine- or olanzapine-treated subjects with schizophrenia (N=21).

Measurement	Baseline	Week 6	Within Group comparison	p value
Entire sample (N=21)				
HAM-D	11.9 ± 8.0	10.5 ± 8.2	t(20)= 1.40	0.177
PANSS – total	73.8 ± 15.3	72.1 ± 15.6	t(20)= 1.35	0.191
PANSS – positive	17.6 ± 6.6	16.7 ± 5.9	t(20)= 1.36	0.188
PANSS – negative	20.9 ± 4.9	21.4 ± 6.0	t(20)= – 0.64	0.532
PANSS – general	35.2 ± 8.9	34.0 ± 7.8	t(20)= 1.28	0.214
SANS – total	59.0 ± 17.6	59.0 ± 16.4	t(20)= – 0.03	0.981
Clozapine (N=11)				
HAM-D	8.6 ± 6.5	7.4 ± 5.2	t(10)= 0.96	0.358
PANSS – total	68.0 ± 16.2	64.9 ± 14.3	t(10)= 1.66	0.129
PANSS – positive	17.2 ± 6.7	15.5 ± 6.0	t(10)= 1.91	0.086
PANSS – negative	18.8 ± 4.1	18.6 ± 5.9	t(10)= 0.15	0.888
PANSS – general	32.0 ± 8.0	30.8 ± 5.7	t(10)= 1.04	0.322
SANS – total	53.1 ± 17.3	51.0 ± 14.0	t(10)= 0.72	0.488
Olanzapine (N=10)				
HAM-D	15.5 ± 8.1	13.9 ± 10.0	t(9)= 0.98	0.355
PANSS – total	80.1 ± 12.1	79.9 ± 13.4	t(9)= 0.12	0.907
PANSS – positive	18.1 ± 7.0	18.0 ± 5.9	t(9)= 0.09	0.927
PANSS – negative	23.2 ± 4.9	24.5 ± 5.0	t(9)= – 1.23	0.249
PANSS – general	38.8 ± 8.7	37.4 ± 8.6	t(9)= 0.79	0.448
SANS – total	65.4 ± 16.3	67.8 ± 14.7	t(9)= – 0.99	0.348

Notes : 1) Values are expressed as mean ± SD; 2) PANSS: Positive and Negative Syndrome Scale, including positive, negative, and general subscales; 3) HAM-D: Hamilton Depression Rating Scale; 4) SANS: Scale for the Assessment of Negative Symptoms