Published in final edited form as:

Front Biol (Beijing). 2012 June ; 7(3): 212–220. doi:10.1007/s11515-012-1222-2.

The role of GSK3 β in the development of the central nervous system

Jia Luo

Department of Internal Medicine, University of Kentucky College of Medicine, Lexington, KY 40536, USA

Abstract

Glycogen synthase kinase 3 β (GSK3 β) is a multifunctional serine/threonine kinase. It is particularly abundant in the developing central nervous system (CNS). Since GSK3 β has diverse substrates ranging from metabolic/signaling proteins and structural proteins to transcription factors, it is involved in many developmental events in the immature brain, such as neurogenesis, neuronal migration, differentiation and survival. The activity of GSK3 β is developmentally regulated and is affected by various environmental/cellular insults, such as deprivation of nutrients/trophic factors, oxidative stress and endoplasmic reticulum stress. Abnormalities in GSK3 β activity may disrupt CNS development. Therefore, GSK3 β is a critical signaling protein that regulates brain development. It may also determine neuronal susceptibility to damages caused by various environmental insults.

Keywords

development; differentiation; neurogenesis; proliferation

Introduction

Glycogen synthase kinase 3 (GSK3) is a multifunctional serine/threonine kinase. Originally found in mammals, and homologs have been found in all eukaryotes (Grimes and Jope, 2001; Doble and Woodgett, 2003). GSK3 was named for its ability to phosphorylate, and thereby inactivate glycogen synthase, a key regulatory molecule in the synthesis of glycogen. There are two highly homologous forms of GSK3 in mammals encoded by distinct genes, GSK3 α (51 kDa) and GSK3 β (47 kDa). GSK3 is an important component of diverse signaling pathways involved in the regulation of cell fate, protein synthesis, glycogen metabolism, cell mobility, transformation, proliferation and survival (Grimes and Jope, 2001; Doble and Woodgett, 2003; Luo, 2009a). Despite a high degree of similarity and functional overlap, these isoforms are not functionally identical and redundant. Between these isoforms, GSK3 β is better studied and highly expressed in the CNS; evidence indicates that GSK3 β plays a prominent role in the CNS. It is now known that GSK3 β regulates diverse early events of neuronal development, such as neurogenesis, neuronal migration,

differentiation and survival in the immature brain. This review discusses studies on the role of GSK3 β in the developing CNS.

GSK3 β signaling

Regulation of GSK3 β activity

GSK3 β is constitutively active in resting cells and undergoes a rapid and transient inhibition in response to a number of external signals (Grimes and Jope, 2001; Doble and Woodgett, 2003). GSK3 β activity is regulated by site-specific phosphorylation. Full activity of GSK3 β generally requires phosphorylation at tyrosine 216 (Tyr216), and conversely, phosphorylation at serine 9 (Ser9) inhibits GSK3 β activity. Phosphorylation of Ser9 is the most common and important regulatory mechanism. Many kinases are capable of phosphorylating Ser9, including p70 S6 kinase, extracellular signal-regulated kinases (ERKs), p90Rsk (also called MAP-KAP kinase-1), protein kinase B (also called Akt), certain isoforms of protein kinase C (PKC) and cyclic AMP-dependent protein kinase (protein kinase A, PKA) (Grimes and Jope, 2001; Kaytor and Orr, 2002). In opposition to the inhibitory modulation of GSK3 β that occurs by serine phosphorylation, tyrosine phosphorylation of GSK3 β increases the enzyme's activity. Studies of tyrosine phosphorylation of GSK3 β are relatively sparse. Stimulation of pGSK3 β (Tyr216) is reported to be mediated by alterations in intracellular calcium levels and a calcium-dependent tyrosine kinase, proline-rich tyrosine kinase 2 (PYK2) or by Fyn, a member of the Src tyrosine family (Hartigan and Johnson, 1999; Lesort et al., 1999; Hartigan et al., 2001; Sayas et al., 2006). pGSK3 β (Tyr216) is also subject to the regulation of mitogen-activated protein kinase kinase (MEK1/2) (Takahashi-Yanaga et al., 2004). It has been suggested that GSK3 β tyrosine phosphorylation may be regulated by autophosphorylation (Cole et al., 2004).

GSK3 β can be activated without apparent changes in phosphorylation of Tyr216 and Ser9 (Baltzis et al., 2007). Regulation of GSK3 β may also be mediated by its subcellular localization. Some substrates of GSK3 β , such as Tau, are cytosolic, whereas others, notably several transcription factors, are nuclear. Thus, GSK3 β must be located in both compartments to regulate these proteins. Diehl et al. (1998) report an increase in nuclear GSK3 β during the S phase of the cell cycle. Proapoptotic stimuli cause the translocation of Tyr216-phosphorylated GSK3 β to the nucleus (Bhat et al., 2000). Identification of a nuclear localization sequence (NLS) in GSK3 β shed light on the mechanisms underlying its shuttling between cytosolic and nuclear compartments (Meares and Jope, 2007).

Substrates of GSK3 β

More than 40 proteins are directly subjected to GSK3 β regulation. These proteins have diverse cellular functions, including glycogen metabolism, transcription, translation, cytoskeletal regulation, cell differentiation, proliferation, transformation and apoptosis (Grimes and Jope, 2001; Manoukian and Woodgett, 2002; Luo, 2009a). Substrates of GSK3 β can be divided into three major groups, namely, metabolic/signaling proteins, structural proteins and transcription factors. The proteins belonging to the first group include acetylCoA carboxylase, amyloid precursor protein, APC tumor suppressor protein, ATP-

citrate lyase, Axin, cyclic AMP-dependent protein kinase, cyclin D1, eIF2B, glycogen synthase, insulin receptor substrate-1 (IRS-1), myelin basic protein, NGF receptor, protein phosphatase 1, protein phosphatase inhibitor-2 and pyruvate dehydrogenase. The structural proteins include dynamin-like protein, microtubule-associated protein 1B (MAP1B), microtubule-associated protein 2 (MAP2), neural cell-adhesion protein (NCAM), neurofilaments, spindle-associated protein Astrin, ninein and Tau. GSK3 β -targeted transcription factors are AP-1 (Jun family), β -catenin, C/EBP α , CREB, glucocorticoid receptor, HSF-1, Myc, NFAT and NF- κ B.

β -catenin is probably the most well-known substrate of GSK3 β . Since β -catenin is an essential component of Wnt signaling, GSK3 β is critically involved in the Wnt system. The Wnts are a family of secreted, cysteine-rich and glycosylated protein ligands that signal by activating the Frizzled family of membrane-bound receptors. Wnt signal transduction causes nuclear translocation of β -catenin and ultimately results in the activation of genes regulated by the T cell factor (TCF)/lymphoid enhancer factor (LEF) family of transcription factors. In the absence of Wnt signals, free cytoplasmic β -catenin is incorporated into a cytoplasmic complex that includes Axin, GSK3 β and adenomatous polyposis coli (APC). This enables GSK3 β to phosphorylate β -catenin and results in ubiquitin-mediated degradation of β -catenin (Doble and Woodgett, 2003; Ille and Sommer, 2005). Wnt signaling inactivates GSK3 β and prevents it from phosphorylating β -catenin, thus stabilizing β -catenin in the cytoplasm. As β -catenin accumulates, it translocates into the nucleus where it binds to TCF/LEF and increases their transcriptional activity. Thus, GSK3 β must be inactivated for Wnt signaling to proceed, and the presence of active GSK3 β inhibits Wnt signaling. The Wnt signaling pathway is an integral component of embryonic development and is involved in morphogenesis and patterning, differentiation processes and lineage decision events during both central and peripheral nervous system development (Patapoutian and Reichardt, 2000; Ille and Sommer, 2005; Malaterre et al., 2007). Therefore, the effect of GSK3 β on development may be associated with Wnt signaling. Additionally recent studies indicate that GSK3 β is a crucial component of cell signaling pathways controlled by disrupted in schizophrenia 1 (DISC1), partitioning defective homolog 3 (PAR3) and PAR6. The role of these pathways in neural development and the involvement of GSK3 β have been reviewed elsewhere (Hur and Zhou, 2010).

GSK3 β in neuronal development

The expression of GSK3 β in the developing brain

Although GSK3 β is widely expressed in all tissues, it is particularly abundant in the CNS (Woodgett, 1990). In the developing brain, GSK3 β is predominantly expressed in neurons and barely detectable in astrocytes (Leroy and Brion, 1999; Takahashi et al., 2000). Leroy and Brion (1999) show that the expression of GSK3 β in the developing rat brain is highest from 18 days of embryonic life up to 10 days of postnatal life (PD10). The expression decreases thereafter and is lowest in the adult; strong GSK3 β immunoreactivity is localized in developing neurons, but is only weakly detected in layers containing neuroblasts. During development and in the adult, GSK3 β is detected in the perikarya and proximal part of dendrites. In the embryo, an intense GSK3 β immunoreactivity is also observed in axonal

tracts. This axonal immunoreactivity has markedly decreased by PD10 and is absent at PD20 as well as in the adult (Leroy and Brion, 1999). Similarly, Takahashi et al. (2000) show a high expression of GSK3 β in the rat brain during the first two weeks of postnatal life; GSK3 β levels peak at PD8–11, and its expression decreases significantly at PD20. Coyle-Rink et al. (2002) examine the expression of GSK3 β in the developing mouse brain. As in the rat brain, a strong GSK3 β expression is observed in the mouse brain during the first two postnatal weeks and then the level of GSK3 β drastically decreases after PD18.

GSK3 β and cell proliferation

GSK3 β is expressed more abundantly in postmitotic neurons than cycling neuroblasts in the developing brain (Leroy and Brion, 1999). Neuronal overexpression of a constitutively active GSK3 β induces microcephaly (Spittaels et al., 2000; 2002). This suggests that GSK3 β activity may negatively regulate the proliferation of neural precursors and neurogenesis. Cui et al. (1998) show the inhibition of GSK3 β can promote the proliferation of cerebellar granule neuron progenitors *in vitro*, suggesting the activation of GSK3 β may suppress the division of neuronal precursors. Knoepfler and Kenney (2006) also used an *in vitro* model of cerebellar granule neuron progenitors to show GSK3 β activation inhibits cell cycle progression, and the inhibitory effect of GSK3 β is mediated by phosphorylating and destabilizing N-Myc. Inhibition of GSK3 β enhances the proliferation of neural progenitor cells (Nedachi et al., 2011). GSK3 β is implicated in adult hippocampal neurogenesis (Boku et al., 2009). Lithium, an inhibitor of GSK3 enhances the proliferation of adult dentate gyrus-derived neural precursor cells in culture (Boku et al., 2009). GSK3 β inhibitor, SB216763 enhances hippocampal neurogenesis (Guo et al., 2011). Under most circumstances, the proliferation and differentiation of neural precursor cells are mutually exclusive during brain development. Fibroblast growth factor 2 (FGF2) promotes neural precursor cell proliferation and concurrently inhibits differentiation. FGF2 inactivates GSK3 β and induces β -catenin nucleus accumulation which enhances the proliferation of neural stem/precursor cells while concurrently inhibiting their differentiation in culture (Shimizu et al., 2008). Similarly, Jin et al. (2005) show that FGF2-stimulated proliferation of cortical neural progenitor cells is mediated by GSK3 β inactivation. According to some studies, however, GSK3 β activity appears necessary for cell cycle progression. Maurer et al. (2007) show that inhibition of GSK3 β by SB216763, a specific GSK3 β inhibitor, suppresses proliferation while promoting neuronal differentiation in neural stem cells isolated from the adult rat subventricular zone. GSK3 β activation by serum and potassium withdrawal induces the expression of several cell cycle regulating proteins, such as cyclin D1, cyclin E, transcription factor E2F-1 and the phosphorylation of retinoblastoma protein (Rb) in cultured cerebellar granule cells (Yeste-Velasco et al. 2007). In some tumor cells, inhibition of GSK3 β results in cell cycle arrest and apoptosis (Luo, 2009a).

Multiple mechanisms are involved in GSK3 β regulation of cell proliferation. First, GSK3 β may regulate mitogenic transcription factors. The activity of some mitogenic transcription factors, such as AP-1 and NF- κ B, is inhibited by GSK3 β activation (Ma et al., 2007; Luo, 2009a; Grimes and Jope, 2001). Secondly, GSK3 β may inhibit cell cycle progression by interacting with cell cycle regulatory proteins, such as cyclin D1 and D2 (Diehl et al., 1998; Huang et al., 2007). Thirdly, GSK3 β may directly regulate spindle microtubule assembly

and accurate chromosome segregation by interacting with the spindle-associated protein Astrin, a substrate for GSK3 β (Cheng et al., 2007).

GSK3 β and neural migration

During development of the CNS, neuronal cells undergo directional movements to specific sites in response to extracellular signals and establish proper connections to other cells. Cell migration in the direction of extracellular cues is mediated by actin and microtubule cytoskeletons. GSK3 β regulates the dynamics of microtubule cytoskeletons in response to extracellular cues (Ciani and Salinas, 2007; Barth et al., 2008), and therefore is an important signaling component that links extracellular signals to cytoskeletal components and governs neuronal migration. Evidence supports that GSK3 β plays an important role in neuronal migration. For example, Chen et al. (2009a) demonstrate that GSK3 β activity is essential for laminin-mediated neurite growth and neuronal migration in the developing cerebral cortex. Asada and Sanada (2010) show that GSK3 β is critical for centrosomal forward movement and neuronal migration in the developing neocortex.

GSK3 β regulates neuronal migration through its interaction with structural or signaling proteins. For instance, MAP1B, a neuron-specific microtubule-associated protein, is a substrate of GSK3 β ; it is implicated in the control of the dynamic stability of microtubules and in the cross-talk between microtubules and actin filaments in neurons (González-Billault et al., 2004). GSK3 β -dependent MAP1B phosphorylation is required for Reelin-regulated neuronal migration *in vivo* and *in vitro* (González-Billault et al., 2004). Platelet activating factor (PAF) induces apoptosis and inhibits the migration of cerebellar granule neurons in a GSK3 β -dependent manner; GSK3 β inhibitors block these effects of PAF (Tong et al., 2001). Focal adhesion kinase (FAK) is a non-receptor tyrosine kinase found in focal adhesion events and plays important roles in cell adhesion, spreading and migration (Hanks et al., 2003). It is shown that GSK3 β regulates cell spreading and migration through the phosphorylation of FAK at selected serine sites (Bianchi et al., 2005). Aspartyl (asparaginy)l-beta-hydroxylase (AAH) regulates cell motility by catalyzing post-translational hydroxylation of proteins involved in cell migration, such as Notch and Jagged (Carter et al., 2008). AAH is a substrate of GSK3 β and GSK3 β -mediated phosphorylation induces AAH degradation, resulting in retarded cell motility (Carter et al., 2008).

Microglia plays a prominent role in the brain's inflammatory response to injury or infection by migrating to affected locations and secreting inflammatory molecules. GSK3 inhibitors reduce the migration of microglia in cultured mouse hippocampal slices (Yuskaitis and Jope, 2009). The regulation of GSK3 β on microglia migration may underlie its role in inflammatory responses to injury in the CNS.

GSK3 β and neuronal differentiation

Neurons are highly polarized cells that contain a single long axon and multiple dendrites. Initial differentiation of neurons is to establish neuronal polarity. Polarization occurs when one of the multiple neurites emerging from the cell body initiates a phase of rapid elongation, becoming the axon; the remaining neurites will develop as dendrites. Rearrangement of the neuronal cytoskeleton provides support for the dramatic

morphological changes that occur during neuronal polarization and neurite outgrowth. GSK3 β plays a critical role in regulating neuronal polarity (Etienne-Manneville and Hall, 2003; Arévalo and Chao, 2005; Jiang et al., 2005; Yoshimura et al., 2005; Gärtner et al., 2006; Vohra et al., 2007). Several microtubule-associated proteins (MAPs) important for neuronal polarity and axonal outgrowth, such as Tau and MAP1B, are substrates of GSK3 β (Barth et al., 2008). The potential mechanisms for GSK3 β regulation of neuronal polarity have been discussed in a review by Hur and Zhou (2010).

GSK3 β also regulates the maturation and neurite outgrowth in neurons. Neuronal overexpression of a constitutively active GSK3 β causes a delayed postnatal maturation and differentiation of neurons in the mouse brain (Spittaels et al., 2000, 2002). In *in vitro* models of neuronal development, inhibition of GSK3 β by selective inhibitors or molecular approaches is shown to promote neurite outgrowth; in contrast, activation of GSK3 β causes neurite retraction (Muñoz-Montaña et al., 1999; Orme et al., 2003; Castelo-Branco et al., 2004; Sayas et al., 2006; Dill et al., 2008; Chen et al., 2009b). Lysophosphatidic acid induces neurite retraction in differentiated neuroblastoma cells via GSK3 β activation (Sun et al., 2011). WW domain-containing oxidoreductase (WWOX), a putative tumor suppressor induces neurite outgrowth and differentiation of SH-SY5Y neuroblastoma cells by repressing GSK3 β activity (Wang et al., 2011). In addition, inactivation of GSK3 β mediates a leptin-induced increase in the size of axonal growth cone in developing mouse cortical neurons (Valerio et al., 2006). GSK3 β also negatively regulates the differentiation of neural stem cells. Inhibition of GSK3 β by SB216763 inhibits proliferation while promoting neuronal differentiation in neural stem cells isolated from the adult rat subventricular zone (Maurer et al., 2007). Inactivation of GSK3 β is also beneficial for axonal regeneration after a spinal cord lesion (Dill et al., 2008).

On the other hand, some reports demonstrate that GSK3 β activity is necessary for neurite outgrowth and axonal extension. For example, lithium or other GSK3 β inhibitors are shown to reduce neurite outgrowth and axon elongation rates in primary cultures of rat hippocampal neurons and sensory neurons (Takahashi et al., 1999; Owen and Gordon-Weeks, 2003). The complex effects of GSK3 β on neuronal differentiation are demonstrated by studies using PC12 cells. PC12 cells are rat pheochromocytoma cells that are extensively used to study the cellular/molecular mechanisms of neuronal differentiation. PC12 cells differentiate into neuronal-like cells by extending neurites and expressing neurotransmitters in response to neurotrophic factors such as NGF and bFGF. There are conflicting reports on the effect of GSK3 β in PC12 cell differentiation; GSK3 β activity can either negatively or positively regulate neurite outgrowth of PC12 cells (Goold and Gordon-Weeks, 2001, 2005; Trivedi et al., 2005; Seng et al., 2006; Zhou et al., 2008; Zhang et al., 2009).

The effect of GSK3 β on neuronal differentiation appears to be mediated by its interaction with microtubule-associated proteins, such as MAP1B which is essential for neuronal polarization, migration neurite outgrowth and axon elongation (Gonzalez-Billault et al., 2001; Owen and Gordon-Weeks, 2003). GSK3 β phosphorylation of MAP1B acts as a molecular switch regulating the control that MAP1B exerts on microtubule dynamics in growing axons and growth cones (Trivedi et al., 2005). Besides MAP1B, GSK3 β may interact with other proteins that participate in regulating neuronal differentiation. For

example, E2F1 transcription factor is a key downstream target of the retinoblastoma tumor suppressor protein (pRB) and is involved in neuronal differentiation. NGF-induced GSK3 β /E2F1 interaction facilitates E2F1 degradation, increasing neurite outgrowth in PC12 cells (Zhou et al., 2008). NGF promotes the interaction between GSK3 β and MRP2, an actin binding protein in PC12 cells; the interaction enhances neurite outgrowth (Seng et al., 2006).

GSK3 β and neuronal survival

Apoptosis has been recognized to be an essential process during neural development and naturally occurring neuronal death plays a substantial developmental role in the building of neural circuitries (Madalosso et al., 2005). It is generally assumed that about half of the neurons produced during neurogenesis die before completion of CNS maturation; this process affects nearly all classes of neurons (Lossi and Merighi, 2003). GSK3 β is an important modulator of apoptosis (Grimes and Jope, 2001; Kaytor and Orr, 2002; Beurel and Jope, 2006). Lucas et al. (2001) use transgenic mice that conditionally overexpress GSK3 β to demonstrate the link between GSK3 β activity and neurodegeneration. They show that overexpression of GSK3 β to specific regions of the brain in mice results in neuronal death which is characteristic of apoptosis in these regions (Lucas et al., 2001). This is supported by studies using various *in vitro* models which show overexpression of GSK3 β in neurons is sufficient to trigger neuronal cell death; in contrast, expression of an inhibitory GSK3 β binding protein or a dominant interfering form of GSK3 β reduces neuronal apoptosis (Pap and Cooper, 1998; Crowder and Freeman, 2000; Hetman et al., 2000; Song et al., 2002). Although there is a lack of evidence regarding direct regulation of GSK3 β on naturally occurring neuronal death during brain development, *in vitro* studies suggest that GSK3 β plays a role in the death/survival of neural precursors or immature neurons. For example, the activation of GSK3 β promotes apoptotic signaling in cultured neural precursor cells (NPCs) derived from embryonic mouse brains subjected to two common apoptotic conditions, trophic factor withdrawal and genotoxic stress (Eom et al., 2007). Lithium, an inhibitor of GSK3 β , is shown to have an opposing effect on neuronal survival depending on the developmental status of the neuron; it induces apoptosis in immature cerebellar granule cells, but promotes survival of mature neurons (D'Mello et al., 1994).

GSK3 β may be activated and induces neuronal death in response to various environmental/cellular stresses, such as deprivation of trophic factors, oxidative stress and endoplasmic reticulum (ER) stress (Bhat et al., 2000; Hetman et al., 2000; Song et al., 2002; Chen et al., 2004; Koh et al., 2005; Brewster et al., 2006; Eom et al., 2007; Lee et al., 2007; Takadera et al., 2007). Relatively low levels of GSK3 β overexpression, which alone do not induce apoptosis, greatly facilitate pro-apoptotic signaling and promote apoptosis (Bijur et al., 2000). The expression of GSK3 β is developmentally regulated; therefore, the status of GSK3 β may play an important role in the susceptibility of the developing brain to environmental insults. For example, we have demonstrated as well as other studies that ethanol-induced death of neurons is mediated at least partially by the activation of GSK3 β in the developing brain, and lithium can protect neurons from ethanol-induced neurodegeneration (Liu et al., 2009; Luo, 2009b; Saito et al., 2010; Ke et al., 2011). We further show that manipulation of GSK3 β activity in neurons can alter neuronal susceptibility to ethanol toxicity (Liu et al., 2009).

GSK3 β activation during neuronal death is commonly mediated by deregulation of phosphorylation at Ser9. Bhat et al. (2000) report that pro-apoptotic stimuli (nerve growth factor withdrawal, ischemia or staurosporine treatment) increase Tyr216 phosphorylation of GSK3 β and induce its nuclear translocation. Multi-mechanisms underlie GSK3 β regulation of neuronal survival/death. It has been suggested that GSK3 β promotes cell death caused by the intrinsic apoptotic pathway, but inhibits the death receptor-mediated extrinsic apoptotic signaling pathway (Beurel and Jope, 2006). The intrinsic apoptotic signaling cascade can be induced by numerous stimuli that cause cell damage, such as DNA damage, oxidative stress and endoplasmic reticulum (ER) stress. The intrinsic apoptotic signaling pathway causes the disruption of mitochondria, leading to cell destruction. Other potential contributory mechanisms are also evident. It is suggested that GSK3 β -induced phosphorylation of Tau may destabilize microtubules which contributes to cytoskeletal collapse associated with apoptosis (Tsukane et al., 2007); GSK3 β -mediated phosphorylation of pyruvate dehydrogenase may impair Krebs cycle activity, reducing cell viability (Hoshi et al., 1996). Effects of GSK3 β on neuronal survival may be mediated by modulation of transcription factors. It is shown that GSK3 β activity is required for AP1-dependent expression of pro-apoptotic Bim, and inhibitors of GSK3 β block AP-1 activation and protect neurons from apoptosis (Hongisto et al., 2003). Activation of GSK3 β is shown to antagonize NF- κ B-mediated neuronal survival, resulting in decreased cell viability (Sui et al., 2006). GSK3 β binds to p53 and promotes p53-induced apoptosis (Watcharasit et al., 2003). Mixed lineage kinase 3 (MLK3) is a mitogen-activated protein kinase kinase kinase member that activates the c-Jun N-terminal kinase (JNK) pathway. GSK3 β -dependent MLK3 phosphorylation mediates neuronal death caused by NGF deprivation (Mishra et al., 2007). Additionally, the effect of GSK3 β on neuronal survival may be regulated by translational control. Translation initiation factor 2B (eIF2B) is a substrate of GSK3 β . GSK3 β phosphorylates and inhibits eIF2B, resulting in translational suppression and programmed cell death (Pap and Cooper, 2002). Finally, GSK3 β may directly modify the activity of pro-apoptotic proteins. For example, it has been demonstrated that GSK3 β directly phosphorylates pro-apoptotic Bax and promotes its mitochondrial localization during neuronal apoptosis (Linseman et al., 2004). In addition to the direct effect on neurons, GSK3 may contribute to neuronal injury by modulating brain inflammation. Excessive neuroinflammation contributes to neurodegeneration. GSK3 β activation promotes the production of inflammatory molecules and microglia migration, which together make GSK3 a powerful regulator of inflammation (Jope et al., 2007).

Conclusions

GSK3 β is primarily localized in neurons of the developing CNS and its expression is developmentally regulated. GSK3 β regulates diverse developmental events in the immature brain, including neurogenesis, neuronal migration, differentiation and survival. GSK3 β is sensitive to external signals and its activity is readily altered by various environmental/cellular insults, such as deprivation of nutrients/trophic factors, oxidative stress and endoplasmic reticulum stress. Therefore, GSK3 β not only plays an essential role in CNS development, it may also determine neuronal susceptibility to damages caused by various environmental insults.

Acknowledgements

I would like to thank Kimberly Bower for reading this manuscript. This research was supported by grants from the National Institutes of Health (AA015407, AA019693 and AA017226).

References

- Arévalo JC, Chao MV. Axonal growth: where neurotrophins meet Wnts. *Curr Opin Cell Biol.* 2005; 17(2):112–115. [PubMed: 15780585]
- Asada N, Sanada K. LKB1-mediated spatial control of GSK3 β and adenomatous polyposis coli contributes to centrosomal forward movement and neuronal migration in the developing neocortex. *J Neurosci.* 2010; 30(26):8852–8865. [PubMed: 20592207]
- Baltzis D, Pluquet O, Papadakis AI, Kazemi S, Qu LK, Koromilas AE. The eIF2 α kinases PERK and PKR activate glycogen synthase kinase 3 to promote the proteasomal degradation of p53. *J Biol Chem.* 2007; 282(43):31675–31687. [PubMed: 17785458]
- Barth AI, Caro-Gonzalez HY, Nelson WJ. Role of adenomatous polyposis coli (APC) and microtubules in directional cell migration and neuronal polarization. *Semin Cell Dev Biol.* 2008; 19(3):245–251. [PubMed: 18387324]
- Beurel E, Jope RS. The paradoxical pro- and anti-apoptotic actions of GSK3 in the intrinsic and extrinsic apoptosis signaling pathways. *Prog Neurobiol.* 2006; 79(4):173–189. [PubMed: 16935409]
- Bhat RV, Shanley J, Correll MP, Fieles WE, Keith RA, Scott CW, Lee CM. Regulation and localization of tyrosine216 phosphorylation of glycogen synthase kinase-3 β in cellular and animal models of neuronal degeneration. *Proc Natl Acad Sci USA.* 2000; 97(20):11074–11079. [PubMed: 10995469]
- Bianchi M, De Lucchini S, Marin O, Turner DL, Hanks SK, Villa-Moruzzi E. Regulation of FAK Ser-722 phosphorylation and kinase activity by GSK3 and PP1 during cell spreading and migration. *Biochem J.* 2005; 391(2):359–370. [PubMed: 15975092]
- Bijur GN, De Sarno P, Jope RS. Glycogen synthase kinase-3 β facilitates staurosporine- and heat shock-induced apoptosis. Protection by lithium. *J Biol Chem.* 2000; 275(11):7583–7590. [PubMed: 10713065]
- Boku S, Nakagawa S, Masuda T, Nishikawa H, Kato A, Kitaichi Y, Inoue T, Koyama T. Glucocorticoids and lithium reciprocally regulate the proliferation of adult dentate gyrus-derived neural precursor cells through GSK-3 β and beta-catenin/TCF pathway. *Neuropsychopharmacology.* 2009; 34(3):805–815. [PubMed: 19005466]
- Brewster JL, Linseman DA, Bouchard RJ, Loucks FA, Precht TA, Esch EA, Heidenreich KA. Endoplasmic reticulum stress and trophic factor withdrawal activate distinct signaling cascades that induce glycogen synthase kinase-3 β and a caspase-9-dependent apoptosis in cerebellar granule neurons. *Mol Cell Neurosci.* 2006; 32(3):242–253. [PubMed: 16765055]
- Carter JJ, Tong M, Silbermann E, Lahousse SA, Ding FF, Longato L, Roper N, Wands JR, de la Monte SM. Ethanol impaired neuronal migration is associated with reduced aspartyl-asparaginyl-beta-hydroxylase expression. *Acta Neuropathol.* 2008; 116(3):303–315. [PubMed: 18478238]
- Castelo-Branco G, Rawal N, Arenas E. GSK-3 β inhibition/beta-catenin stabilization in ventral midbrain precursors increases differentiation into dopamine neurons. *J Cell Sci.* 2004; 117(24):5731–5737. [PubMed: 15522889]
- Chen G, Bower KA, Ma C, Fang S, Thiele CJ, Luo J. Glycogen synthase kinase 3 β (GSK3 β) mediates 6-hydroxydopamine-induced neuronal death. *FASEB J.* 2004; 18(10):1162–1164. [PubMed: 15132987]
- Chen G, Bower KA, Xu M, Ding M, Shi X, Ke ZJ, Luo J. Cyanidin-3-glucoside reverses ethanol-induced inhibition of neurite outgrowth: role of glycogen synthase kinase 3 β . *Neurotox Res.* 2009b; 15(4):321–331. [PubMed: 19384566]
- Chen ZL, Haegeli V, Yu H, Strickland S. Cortical deficiency of laminin gamma1 impairs the AKT/GSK-3 β signaling pathway and leads to defects in neurite outgrowth and neuronal migration. *Dev Biol.* 2009a; 327(1):158–168. [PubMed: 19118544]

- Cheng TS, Hsiao YL, Lin CC, Yu CT, Hsu CM, Chang MS, Lee CI, Huang CY, Howng SL, Hong YR. Glycogen synthase kinase 3 β interacts with and phosphorylates the spindle-associated protein astrin. *J Biol Chem.* 2007; 283(4):2454–2464. [PubMed: 18055457]
- Ciani L, Salinas PC. c-Jun N-terminal kinase (JNK) cooperates with Gsk3 β to regulate Dishevelled-mediated microtubule stability. *BMC Cell Biol.* 2007; 8(1):27. [PubMed: 17608927]
- Cole A, Frame S, Cohen P. Further evidence that the tyrosine phosphorylation of glycogen synthase kinase-3 (GSK3) in mammalian cells is an autophosphorylation event. *Biochem J.* 2004; 377(1): 249–255. [PubMed: 14570592]
- Coyle-Rink J, Del Valle L, Sweet T, Khalili K, Amini S. Developmental expression of Wnt signaling factors in mouse brain. *Cancer Biol Ther.* 2002; 1(6):640–645. [PubMed: 12642687]
- Crowder RJ, Freeman RS. Glycogen synthase kinase-3 β activity is critical for neuronal death caused by inhibiting phosphatidylinositol 3-kinase or Akt but not for death caused by nerve growth factor withdrawal. *J Biol Chem.* 2000; 275(44):34266–34271. [PubMed: 10954722]
- Cui H, Meng Y, Bulleit RF. Inhibition of glycogen synthase kinase 3 β activity regulates proliferation of cultured cerebellar granule cells. *Brain Res Dev Brain Res.* 1998; 111(2):177–188.
- D'Mello SR, Anelli R, Calissano P. Lithium induces apoptosis in immature cerebellar granule cells but promotes survival of mature neurons. *Exp Cell Res.* 1994; 211(2):332–338. [PubMed: 8143780]
- Diehl JA, Cheng M, Roussel MF, Sherr CJ. Glycogen synthase kinase-3 β regulates cyclin D1 proteolysis and subcellular localization. *Genes Dev.* 1998; 12(22):3499–3511. [PubMed: 9832503]
- Dill J, Wang H, Zhou F, Li S. Inactivation of glycogen synthase kinase 3 promotes axonal growth and recovery in the CNS. *J Neurosci.* 2008; 28(36):8914–8928. [PubMed: 18768685]
- Doble BW, Woodgett JR. GSK-3: tricks of the trade for a multitasking kinase. *J Cell Sci.* 2003; 116(7):1175–1186. [PubMed: 12615961]
- Eom TY, Roth KA, Jope RS. Neural precursor cells are protected from apoptosis induced by trophic factor withdrawal or genotoxic stress by inhibitors of glycogen synthase kinase 3. *J Biol Chem.* 2007; 282(31):22856–22864. [PubMed: 17548347]
- Etienne-Manneville S, Hall A. Cdc42 regulates GSK-3 β and adenomatous polyposis coli to control cell polarity. *Nature.* 2003; 421(6924):753–756. [PubMed: 12610628]
- Gärtner A, Huang X, Hall A. Neuronal polarity is regulated by glycogen synthase kinase-3 (GSK-3 β) independently of Akt/PKB serine phosphorylation. *J Cell Sci.* 2006; 119(19):3927–3934. [PubMed: 16954147]
- Gonzalez-Billault C, Avila J, Cáceres A. Evidence for the role of MAP1B in axon formation. *Mol Biol Cell.* 2001; 12(7):2087–2098. [PubMed: 11452005]
- González-Billault C, Del Río JA, Ureña JM, Jiménez-Mateos EM, Barallobre MJ, Pascual M, Pujadas L, Simó S, Torre AL, Gavin R, Wandosell F, Soriano E, Avila J. A role of MAP1B in Reelin-dependent neuronal migration. *Cereb Cortex.* 2004; 15(8):1134–1145. [PubMed: 15590913]
- Goold RG, Gordon-Weeks PR. Microtubule-associated protein 1B phosphorylation by glycogen synthase kinase 3 β is induced during PC12 cell differentiation. *J Cell Sci.* 2001; 114(Pt 23): 4273–4284. [PubMed: 11739659]
- Goold RG, Gordon-Weeks PR. The MAP kinase pathway is upstream of the activation of GSK3 β that enables it to phosphorylate MAP1B and contributes to the stimulation of axon growth. *Mol Cell Neurosci.* 2005; 28(3):524–534. [PubMed: 15737742]
- Grimes CA, Jope RS. The multifaceted roles of glycogen synthase kinase 3 β in cellular signaling. *Prog Neurobiol.* 2001; 65(4):391–426. [PubMed: 11527574]
- Guo W, Murthy AC, Zhang L, Johnson EB, Schaller EG, Allan AM, Zhao X. Inhibition of GSK3 Improves Hippocampus-dependent Learning and Rescues Neurogenesis in a Mouse Model of Fragile X Syndrome. *Hum Mol Genet.* 2011
- Hanks SK, Ryzhova L, Shin NY, Brábek J. Focal adhesion kinase signaling activities and their implications in the control of cell survival and motility. *Front Biosci.* 2003; 8(1–3):d982–d996. [PubMed: 12700132]
- Hartigan JA, Johnson GV. Transient increases in intracellular calcium result in prolonged site-selective increases in Tau phosphorylation through a glycogen synthase kinase 3 β -dependent pathway. *J Biol Chem.* 1999; 274(30):21395–21401. [PubMed: 10409701]

- Hartigan JA, Xiong WC, Johnson GV. Glycogen synthase kinase 3beta is tyrosine phosphorylated by PYK2. *Biochem Biophys Res Commun*. 2001; 284(2):485–489. [PubMed: 11394906]
- Hetman M, Cavanaugh JE, Kimelman D, Xia Z. Role of glycogen synthase kinase-3beta in neuronal apoptosis induced by trophic withdrawal. *J Neurosci*. 2000; 20(7):2567–2574. [PubMed: 10729337]
- Hongisto V, Smeds N, Brecht S, Herdegen T, Courtney MJ, Coffey ET. Lithium blocks the c-Jun stress response and protects neurons via its action on glycogen synthase kinase 3. *Mol Cell Biol*. 2003; 23(17):6027–6036. [PubMed: 12917327]
- Hoshi M, Takashima A, Noguchi K, Murayama M, Sato M, Kondo S, Saitoh Y, Ishiguro K, Hoshino T, Imahori K. Regulation of mitochondrial pyruvate dehydrogenase activity by tau protein kinase I/glycogen synthase kinase 3beta in brain. *Proc Natl Acad Sci USA*. 1996; 93(7):2719–2723. [PubMed: 8610107]
- Huang W, Chang HY, Fei T, Wu H, Chen YG. GSK3 beta mediates suppression of cyclin D2 expression by tumor suppressor PTEN. *Oncogene*. 2007; 26(17):2471–2482. [PubMed: 17043650]
- Hur EM, Zhou FQ. GSK3 signalling in neural development. *Nat Rev Neurosci*. 2010; 11(8):539–551. [PubMed: 20648061]
- Ille F, Sommer L. Wnt signaling: multiple functions in neural development. *Cell Mol Life Sci*. 2005; 62(10):1100–1108. [PubMed: 15928805]
- Jiang H, Guo W, Liang X, Rao Y. Both the establishment and the maintenance of neuronal polarity require active mechanisms: critical roles of GSK-3beta and its upstream regulators. *Cell*. 2005; 120(1):123–135. [PubMed: 15652487]
- Jin L, Hu X, Feng L. NT3 inhibits FGF2-induced neural progenitor cell proliferation via the PI3K/GSK3 pathway. *J Neurochem*. 2005; 93(5):1251–1261. [PubMed: 15934945]
- Jope RS, Yuskaitis CJ, Beurel E. Glycogen synthase kinase-3 (GSK3): inflammation, diseases, and therapeutics. *Neurochem Res*. 2007; 32(4–5):577–595. [PubMed: 16944320]
- Kaytor MD, Orr HT. The GSK3 beta signaling cascade and neurodegenerative disease. *Curr Opin Neurobiol*. 2002; 12(3):275–278. [PubMed: 12049933]
- Ke Z, Liu Y, Wang X, Fan Z, Chen G, Xu M, Bower KA, Frank JA, Ou X, Shi X, Luo J. Cyanidin-3-glucoside ameliorates ethanol neurotoxicity in the developing brain. *J Neurosci Res*. 2011; 89(10):1676–1684. [PubMed: 21671257]
- Knoepfler PS, Kenney AM. Neural precursor cycling at sonic speed: N-Myc pedals, GSK-3 brakes. *Cell Cycle*. 2006; 5(1):47–52. [PubMed: 16322694]
- Koh SH, Lee YB, Kim KS, Kim HJ, Kim M, Lee YJ, Kim J, Lee KW, Kim SH. Role of GSK-3beta activity in motor neuronal cell death induced by G93A or A4V mutant hSOD1 gene. *Eur J Neurosci*. 2005; 22(2):301–309. [PubMed: 16045483]
- Lee KY, Koh SH, Noh MY, Park KW, Lee YJ, Kim SH. Glycogen synthase kinase-3beta activity plays very important roles in determining the fate of oxidative stress-inflicted neuronal cells. *Brain Res*. 2007; 1129(1):89–99. [PubMed: 17157278]
- Leroy K, Brion JP. Developmental expression and localization of glycogen synthase kinase-3beta in rat brain. *J Chem Neuroanat*. 1999; 16(4):279–293. [PubMed: 10450875]
- Lesort M, Jope RS, Johnson GV. Insulin transiently increases tau phosphorylation: involvement of glycogen synthase kinase-3beta and Fyn tyrosine kinase. *J Neurochem*. 1999; 72(2):576–584. [PubMed: 9930729]
- Linseman DA, Butts BD, Precht TA, Phelps RA, Le SS, Laessig TA, Bouchard RJ, Florez-McClure ML, Heidenreich KA. Glycogen synthase kinase-3beta phosphorylates Bax and promotes its mitochondrial localization during neuronal apoptosis. *J Neurosci*. 2004; 24(44):9993–10002. [PubMed: 15525785]
- Liu Y, Chen G, Ma C, Bower KA, Xu M, Fan Z, Shi X, Ke ZJ, Luo J. Over-expression of GSK3 beta sensitizes neuronal cells to ethanol toxicity. *J Neurosci Res*. 2009; 87(12):2793–2802. [PubMed: 19382207]
- Lossi L, Merighi A. *In vivo* cellular and molecular mechanisms of neuronal apoptosis in the mammalian CNS. *Prog Neurobiol*. 2003; 69(5):287–312. [PubMed: 12787572]

- Lucas JJ, Hernández F, Gómez-Ramos P, Morán MA, Hen R, Avila J. Decreased nuclear beta-catenin, tau hyperphosphorylation and neurodegeneration in GSK-3beta conditional transgenic mice. *EMBO J*. 2001; 20(1):27–39. [PubMed: 11226152]
- Luo J. Glycogen synthase kinase 3beta (GSK3beta) in tumorigenesis and cancer chemotherapy. *Cancer Lett*. 2009a; 273(2):194–200. [PubMed: 18606491]
- Luo J. GSK3beta in ethanol neurotoxicity. *Mol Neurobiol*. 2009b; 40(2):108–121. [PubMed: 19507062]
- Ma C, Wang J, Gao Y, Gao TW, Chen G, Bower KA, Odetallah M, Ding M, Ke Z, Luo J. The role of glycogen synthase kinase 3beta in the transformation of epidermal cells. *Cancer Res*. 2007; 67(16):7756–7764. [PubMed: 17699780]
- Madalosso SH, Pérez-Villegas EM, Armengol JA. Naturally occurring neuronal death during the postnatal development of Purkinje cells and their precerebellar afferent projections. *Brain Res Brain Res Rev*. 2005; 49(2):267–279. [PubMed: 16111555]
- Malaterre J, Ramsay RG, Mantamadiotis T. Wnt-Frizzled signalling and the many paths to neural development and adult brain homeostasis. *Front Biosci*. 2007; 12(1):492–506. [PubMed: 17127312]
- Manoukian AS, Woodgett JR. Role of glycogen synthase kinase-3 in cancer: regulation by Wnts and other signaling pathways. *Adv Cancer Res*. 2002; 84:203–229. [PubMed: 11883528]
- Maurer MH, Brömme JO, Feldmann RE Jr, Järve A, Sabouri F, Bürgers HF, Schelshorn DW, Krüger C, Schneider A, Kuschinsky W. Glycogen synthase kinase 3beta (GSK3beta) regulates differentiation and proliferation in neural stem cells from the rat subventricular zone. *J Proteome Res*. 2007; 6(3):1198–1208. [PubMed: 17330951]
- Meares GP, Jope RS. Resolution of the nuclear localization mechanism of glycogen synthase kinase-3: functional effects in apoptosis. *J Biol Chem*. 2007; 282(23):16989–17001. [PubMed: 17438332]
- Mishra R, Barthwal MK, Sondarva G, Rana B, Wong L, Chatterjee M, Woodgett JR, Rana A. Glycogen synthase kinase-3beta induces neuronal cell death via direct phosphorylation of mixed lineage kinase 3. *J Biol Chem*. 2007; 282(42):30393–30405. [PubMed: 17711861]
- Muñoz-Montaño JR, Lim F, Moreno FJ, Avila J, Díaz-Nido J. Glycogen synthase kinase-3 modulates neurite outgrowth in cultured neurons: possible implications for neurite pathology in Alzheimer's disease. *J Alzheimers Dis*. 1999; 1(6):361–378. [PubMed: 12214113]
- Nedachi T, Kawai T, Matsuaki T, Yamanouchi K, Nishihara M. Progranulin enhances neural progenitor cell proliferation through glycogen synthase kinase 3 β phosphorylation. *Neuroscience*. 2011; 185:106–115. [PubMed: 21540081]
- Orme MH, Giannini AL, Vivanco MD, Kypta RM. Glycogen synthase kinase-3 and Axin function in a beta-catenin-independent pathway that regulates neurite outgrowth in neuroblastoma cells. *Mol Cell Neurosci*. 2003; 24(3):673–686. [PubMed: 14664817]
- Owen R, Gordon-Weeks PR. Inhibition of glycogen synthase kinase 3beta in sensory neurons in culture alters filopodia dynamics and microtubule distribution in growth cones. *Mol Cell Neurosci*. 2003; 23(4):626–637. [PubMed: 12932442]
- Pap M, Cooper GM. Role of glycogen synthase kinase-3 in the phosphatidylinositol 3-Kinase/Akt cell survival pathway. *J Biol Chem*. 1998; 273(32):19929–19932. [PubMed: 9685326]
- Pap M, Cooper GM. Role of translation initiation factor 2B in control of cell survival by the phosphatidylinositol 3-kinase/Akt/glycogen synthase kinase 3beta signaling pathway. *Mol Cell Biol*. 2002; 22(2):578–586. [PubMed: 11756553]
- Patapoutian A, Reichardt LF. Roles of Wnt proteins in neural development and maintenance. *Curr Opin Neurobiol*. 2000; 10(3):392–399. [PubMed: 10851180]
- Saito M, Chakraborty G, Mao RF, Paik SM, Vadasz C, Saito M. Tau phosphorylation and cleavage in ethanol-induced neurodegeneration in the developing mouse brain. *Neurochem Res*. 2010; 35(4):651–659. [PubMed: 20049527]
- Sayas CL, Ariaens A, Ponsioen B, Moolenaar WH. GSK-3 is activated by the tyrosine kinase Pyk2 during LPA1-mediated neurite retraction. *Mol Biol Cell*. 2006; 17(4):1834–1844. [PubMed: 16452634]

- Seng S, Avraham HK, Jiang S, Venkatesh S, Avraham S. KLHL1/MRP2 mediates neurite outgrowth in a glycogen synthase kinase 3 β -dependent manner. *Mol Cell Biol.* 2006; 26(22):8371–8384. [PubMed: 16982692]
- Shimizu T, Kagawa T, Inoue T, Nonaka A, Takada S, Aburatani H, Taga T. Stabilized beta-catenin functions through TCF/LEF proteins and the Notch/RBP-Jkappa complex to promote proliferation and suppress differentiation of neural precursor cells. *Mol Cell Biol.* 2008; 28(24):7427–7441. [PubMed: 18852283]
- Song L, De Sarno P, Jope RS. Central role of glycogen synthase kinase-3 β in endoplasmic reticulum stress-induced caspase-3 activation. *J Biol Chem.* 2002; 277(47):44701–44708. [PubMed: 12228224]
- Spittaels K, Van den Haute C, Van Dorpe J, Geerts H, Mercken M, Bruynseels K, Lasrado R, Vandezande K, Laenen I, Boon T, Van Lint J, Vandenheede J, Moechars D, Loos R, Van Leuven F. Glycogen synthase kinase-3 β phosphorylates protein tau and rescues the axonopathy in the central nervous system of human four-repeat tau transgenic mice. *J Biol Chem.* 2000; 275(52):41340–41349. [PubMed: 11007782]
- Spittaels K, Van den Haute C, Van Dorpe J, Terwel D, Vandezande K, Lasrado R, Bruynseels K, Irizarry M, Verhoye M, Van Lint J, Vandenheede JR, Ashton D, Mercken M, Loos R, Hyman B, Van der Linden A, Geerts H, Van Leuven F. Neonatal neuronal overexpression of glycogen synthase kinase-3 β reduces brain size in transgenic mice. *Neuroscience.* 2002; 113(4):797–808. [PubMed: 12182887]
- Sui Z, Sniderhan LF, Fan S, Kazmierczak K, Reisinger E, Kovács AD, Potash MJ, Dewhurst S, Gelbard HA, Maggirwar SB. Human immunodeficiency virus-encoded Tat activates glycogen synthase kinase-3 β to antagonize nuclear factor- κ B survival pathway in neurons. *Eur J Neurosci.* 2006; 23(10):2623–2634. [PubMed: 16817865]
- Sun Y, Kim NH, Yang H, Kim SH, Huh SO. Lysophosphatidic acid induces neurite retraction in differentiated neuroblastoma cells via GSK-3 β activation. *Mol Cells.* 2011; 31(5):483–489. [PubMed: 21499833]
- Takadera T, Fujibayashi M, Kaniyu H, Sakota N, Ohyashiki T. Caspase-dependent apoptosis induced by thapsigargin was prevented by glycogen synthase kinase-3 inhibitors in cultured rat cortical neurons. *Neurochem Res.* 2007; 32(8):1336–1342. [PubMed: 17401651]
- Takahashi M, Tomizawa K, Ishiguro K. Distribution of tau protein kinase I/glycogen synthase kinase-3 β , phosphatases 2A and 2B, and phosphorylated tau in the developing rat brain. *Brain Res.* 2000; 857(1–2):193–206. [PubMed: 10700568]
- Takahashi M, Yasutake K, Tomizawa K. Lithium inhibits neurite growth and tau protein kinase I/glycogen synthase kinase-3 β -dependent phosphorylation of juvenile tau in cultured hippocampal neurons. *J Neurochem.* 1999; 73(5):2073–2083. [PubMed: 10537067]
- Takahashi-Yanaga F, Shiraishi F, Hirata M, Miwa Y, Morimoto S, Sasaguri T. Glycogen synthase kinase-3 β is tyrosine-phosphorylated by MEK1 in human skin fibroblasts. *Biochem Biophys Res Commun.* 2004; 316(2):411–415. [PubMed: 15020233]
- Tong N, Sanchez JF, Maggirwar SB, Ramirez SH, Guo H, Dewhurst S, Gelbard HA. Activation of glycogen synthase kinase 3 β (GSK-3 β) by platelet activating factor mediates migration and cell death in cerebellar granule neurons. *Eur J Neurosci.* 2001; 13(10):1913–1922. [PubMed: 11403684]
- Trivedi N, Marsh P, Goold RG, Wood-Kaczmar A, Gordon-Weeks PR. Glycogen synthase kinase-3 β phosphorylation of MAP1B at Ser1260 and Thr1265 is spatially restricted to growing axons. *J Cell Sci.* 2005; 118(5):993–1005. [PubMed: 15731007]
- Tsukane M, Yoshizaki C, Yamauchi T. Development and specific induction of apoptosis of cultured cell models overexpressing human tau during neural differentiation: Implication in Alzheimer's disease. *Anal Biochem.* 2007; 360(1):114–122. [PubMed: 17113559]
- Valerio A, Ghisi V, Dossena M, Tonello C, Giordano A, Frontini A, Ferrario M, Pizzi M, Spano P, Carruba MO, Nisoli E. Leptin increases axonal growth cone size in developing mouse cortical neurons by convergent signals inactivating glycogen synthase kinase-3 β . *J Biol Chem.* 2006; 281(18):12950–12958. [PubMed: 16522636]
- Vohra BP, Fu M, Heuckeroth RO. Protein kinase C ζ and glycogen synthase kinase-3 β control neuronal polarity in developing rodent enteric neurons, whereas SMAD specific E3 ubiquitin

- protein ligase 1 promotes neurite growth but does not influence polarity. *J Neurosci.* 2007; 27(35): 9458–9468. [PubMed: 17728459]
- Wang HY, Juo LI, Lin YT, Hsiao M, Lin JT, Tsai CH, Tzeng YH, Chuang YC, Chang NS, Yang CN, Lu PJ. WW domain-containing oxidoreductase promotes neuronal differentiation via negative regulation of glycogen synthase kinase 3 β . *Cell Death Differ.* 2011 Online Available December 23, 2011.
- Watcharasit P, Bijur GN, Song L, Zhu J, Chen X, Jope RS. Glycogen synthase kinase-3 β (GSK3 β) binds to and promotes the actions of p53. *J Biol Chem.* 2003; 278(49):48872–48879. [PubMed: 14523002]
- Woodgett JR. Molecular cloning and expression of glycogen synthase kinase-3/factor A. *EMBO J.* 1990; 9(8):2431–2438. [PubMed: 2164470]
- Yeste-Velasco M, Folch J, Trullàs R, Abad MA, Enguita M, Pallàs M, Camins A. Glycogen synthase kinase-3 is involved in the regulation of the cell cycle in cerebellar granule cells. *Neuropharmacology.* 2007; 53(2):295–307. [PubMed: 17612578]
- Yoshimura T, Kawano Y, Arimura N, Kawabata S, Kikuchi A, Kaibuchi K. GSK-3 β regulates phosphorylation of CRMP-2 and neuronal polarity. *Cell.* 2005; 120(1):137–149. [PubMed: 15652488]
- Yuskaitis CJ, Jope RS. Glycogen synthase kinase-3 regulates microglial migration, inflammation, and inflammation-induced neurotoxicity. *Cell Signal.* 2009; 21(2):264–273. [PubMed: 19007880]
- Zhang W, Smith A, Liu JP, Cheung NS, Zhou S, Liu K, Li QT, Duan W. GSK3 β modulates PACAP-induced neuritogenesis in PC12 cells by acting downstream of Rap1 in a caveolae-dependent manner. *Cell Signal.* 2009; 21(2):237–245. [PubMed: 18983912]
- Zhou F, Zhang L, Wang A, Song B, Gong K, Zhang L, Hu M, Zhang X, Zhao N, Gong Y. The association of GSK3 β with E2F1 facilitates nerve growth factor-induced neural cell differentiation. *J Biol Chem.* 2008; 283(21):14506–14515. [PubMed: 18367454]