Published in final edited form as:

Curr Opin Gastroenterol. 2015 January ; 31(1): 14–23. doi:10.1097/MOG.0000000000000134.

Autophagy and checkpoints for intracellular pathogen defense

Geraldine L.C. Paulus¹ and Ramnik J. Xavier^{1,2}

¹Broad Institute of Harvard University and Massachusetts Institute of Technology, Cambridge, MA, USA

²Gastrointestinal Unit, Center for the Study of Inflammatory Bowel Disease, and Center for Computational and Integrative Biology, Massachusetts General Hospital, Boston, MA, USA

Abstract

Purpose of review—Autophagy plays a crucial role in intracellular defense against various pathogens. Xenophagy is a form of selective autophagy that targets intracellular pathogens for degradation. In addition, several related yet distinct intracellular defense responses depend on autophagy-related (ATG) genes. This review gives an overview of these processes, pathogen strategies to subvert them, and their crosstalk with various cell death programs.

Recent findings—The recruitment of ATG proteins plays a key role in multiple intracellular defense programs, specifically xenophagy, LC3-associated phagocytosis (LAP), and the IFN γ -mediated elimination of pathogens such as *Toxoplasma gondii* and murine norovirus. Recent progress has revealed methods employed by pathogens to resist these intracellular defense mechanisms and/or persist in spite of them. The intracellular pathogen load can tip the balance between cell survival and cell death. Further, it was recently observed that LAP is indispensable for the efficient clearance of dying cells.

Summary—Autophagy-dependent and ATG gene-dependent pathways are essential in intracellular defense against a broad range of pathogens.

Keywords

autophagy; xenophagy; LC3-associated phagocytosis; IFN γ -mediated pathogen elimination; pathogen survival strategies

Introduction

Classical autophagy refers to an evolutionarily conserved catabolic process in eukaryotic cells in which cytoplasmic components are engulfed by a double-membraned autophagosome and degraded after fusion with a lysosome [1, 2]. The canonical autophagy machinery consists of more than 30 autophagy-related (ATG) genes, originally identified in yeast [2]. An important marker of autophagy is the recruitment of microtubule-associated

Corresponding author: Ramnik J. Xavier, Gastrointestinal Unit and Center for the Study of Inflammatory Bowel Disease, Massachusetts General Hospital, Richard B. Simches Research Center, 185 Cambridge Street, Boston, MA 02114, Phone: 617-643-3331, Fax: 617-643-3328, xavier@molbio.mgh.harvard.edu.

protein light-chain 3 (LC3), the mammalian orthologue of yeast Atg8, to the autophagosome [2].

Selective autophagy of pathogens, also referred to as xenophagy, is a key defense mechanism against a broad range of infections [3]. Additionally, genome-wide association studies have revealed associations between inflammatory bowel diseases (IBD) and single-nucleotide polymorphisms (SNPs) in genes involved in autophagy. Most notably, SNPs in *ATG16L1* and *IRGM* specifically increase the risk for Crohn's disease and genetic variants in *ATG4* and *FBNPIL* are associated with the presence of granulomas in Crohn's disease patients, a marker of a more aggressive disease course [1, 4, 5]. Other genes, like *SMURF1*, are found in risk loci for ulcerative colitis [5].

In addition, noncanonical forms of autophagy, which rely on select components of the autophagy machinery, are implicated in intracellular pathogen defense. For instance, in a process termed LC3-associated phagocytosis (LAP), a subset of ATG proteins assists in the digestion of phagocytosed pathogens. Central to this process is the conjugation of LC3 to the single-membraned phagosomes [6, 7]. Further, certain ATG proteins take part in the interferon gamma (IFN γ)-mediated defense against pathogens such as *Mycobacterium tuberculosis*, *Toxoplasma gondii*, and murine norovirus, whose human equivalent is the most common cause of viral gastroenteritis [8,9,10].

Autophagy in pathogen defense

The canonical macroautophagy pathway, hereafter referred to as autophagy, is the most studied and well-understood form of autophagy.

Different ATG protein conjugation complexes act at the different stages of the autophagy pathway [2,11]. First, an isolation membrane (phagophore) is generated *de novo* from the endoplasmic reticulum, Golgi apparatus, mitochondria, or other plasma membrane-derived endocytic organelles [11]. The initiation of the phagophore is mediated by the ULK1/FIP200/ATG101/ATG13 protein kinase complex and Vps34/Beclin-1/Vps15/ATG14 lipid kinase complex. These complexes induce the formation of the ATG5/ATG12/ATG16L1 complex (formed with the help of ATG7 and ATG10), which promotes the elongation of the phagophore that eventually closes around cytoplasmic cargo, forming a double-membraned vacuole called the autophagosome. Aided by ATG3 and ATG7, these complexes facilitate the addition of a phosphatidylethanolamine group to the mammalian LC3 homologues (LC3A, LC3B, LC3C, GABARAP, GABARAP-L1, GABARAP-L2) [12]), which are then recruited to the autophagosome. This conversion from the cytosolic to the lipid-bound form of LC3 is commonly referred to as LC3-I to LC3-II conversion. Finally, the autophagosome matures and fuses with the lysosome, forming a single-membraned vesicle termed the autolysosome, wherein the contents are degraded. This step is effectuated by several lysosomal membrane proteins, including the GTPase Rab7 and LAMP2, as well as by SNARE proteins such as syntaxin 17 and SNAP29. Another Beclin1/VPS34 complex assists in this final step, only here instead of ATG14 the UV radiation resistance-associated gene (UVRAG) is required.

Apart from their role in autophagosome formation, the LC3/GABARAP proteins provide docking sites for receptors that deliver cargo to phagophores during *selective* autophagy. To date, eight mammalian cargo receptors have been identified: p62, its paralogue NBR1, NDP52, its paralogue T6BP, optineurin [13, 14], Nix [15], NCOA4 [16, 17] and recently TRIM5 α [18]. They all share an LC3-interacting region (LIR), typically a WxxL sequence N-terminally preceded by negatively charged residues. Variations in their domain structure allow them to interact with specific members of the LC3/GABARAP family and specific pathogen-associated substrates [14, 19, 20]. Recent studies in yeast identified a LIR-independent interaction between ATG19 (a relative of mammalian p62 and NBR1) and ATG8 and ATG11 [21, 22]. ATG19 contains multiple binding sites for ATG8 and its interaction with ATG11 is achieved through kinase Hrr25 phosphorylation [22].

Xenophagy is a specific form of selective autophagy where the cargo captured by the autophagosome is an intracellular pathogen [23-26] (Fig. 1a). Xenophagy is an important immune response, through the direct and selective elimination of intracellular pathogens. Studies involving *M. tuberculosis* and *Salmonella* Typhimurium infection indicate that approximately 30% of intracellular bacteria are associated with autophagy markers [27, 28].

When bacteria enter the cell, they are engulfed by the host plasma membrane. The resulting single-membrane pathogen-containing vacuole can mature and fuse with the lysosome, forming a phagolysosome in a process termed phagocytosis. In order to evade this process and facilitate their survival and replication, most bacteria (e.g., *Coxiella*, *Chlamydia*, *Salmonella*, *Legionella*, *Mycobacterium*) adapt to or modulate the conditions of the phagosome, thereby establishing a vacuolar replication niche, and often concurrently block the fusion with the lysosome [29]. A percentage of these bacteria eventually become cytosolic because of failure to maintain vacuolar integrity, to obtain additional nutrients, or to spread to adjacent cells [27]. Other bacteria, including *Shigella*, *Listeria*, and *Francisella* quickly escape the vacuole and attempt to resist cytosolic host immune defense strategies.

When intracellular pathogens become cytosolic, their vacuole membrane ruptures, thereby exposing the previously inward-facing glycans to the host cytosol. The exposed glycans are damage-associated molecular patterns (DAMPs) that can be recognized by cargo receptors [3, 30]. For example, upon infection of epithelial cells with *S. Typhimurium*, galectin-8 interacts with exposed glycans on the ruptured membranes, accumulates, and recruits the cargo receptor NDP52 [31]. Additionally, xenophagy typically involves the poly-ubiquitination of either exposed phagosomemeglycans [32] or of cytosolic bacteria themselves [33, 34]. This process is governed by E3 ligases such as LRSAM1 [35, 36] and PARKIN [35]. Similarly, viruses are commonly targeted for ubiquitin-mediated xenophagy. The transmembrane protein STING (stimulator of interferon genes) can detect double-stranded DNA viruses such as herpes simplex virus 1 (HSV1) and human cytomegalovirus and target them for autophagy [37]. The E3 ligase SMURF1 causes the clearance of Sindbis virus and HSV1 [38].

Pathogen secretion products can also induce xenophagy. For instance, *M. tuberculosis* can permeabilize its vacuole and release extracellular DNA (eDNA) to assist bacterial adhesion to epithelial cells. STING can detect this eDNA and cause ubiquitination of the *M.*

tuberculosis-containing vacuole [9, 37, 39]. The resulting ubiquitin coat is subsequently recognized by cargo receptors NDP52, p62, NBR1, and optineurin [20].

In comparison to the insight in the recognition of pathogens as autophagic cargo, much less is understood about the signals inducing phagophore formation upon infection. One study reported that bacteria with access to the cytosol activate xenophagy by nutrient competition with the host [40]. Specifically, amino acid starvation in the host cell inhibits the mammalian target of rapamycin complex 1 (mTORC1), which negatively regulates the ULK1 complex. Additionally, nutrient starvation of the host cell can induce nuclear translocation of the transcription factor EB (TFEB), a master regulator of lysosome biogenesis, driving mTORC1-independent autophagy [41, 42].

Non-canonical autophagy programs in pathogen defense

Several intracellular defense mechanisms have been identified that depend on a subset of ATG genes.

LC3-associated phagocytosis

As discussed above, many microorganisms inside the phagosome have developed mechanisms to disrupt vesicular trafficking and avoid phagosome-lysosome fusion [29]. A complementary pathogen defense mechanism to xenophagy, coined LAP, involves the recruitment of LC3 to the still-intact phagosome (Fig. 1b) [6, 7, 43]. This process is independent of the ULK1 initiation complex, consistent with the absence of phagophore formation in LAP. LC3 is mobilized by TLRs or Fc γ receptors (for bacterial infections) [44] or CLEC7A (for fungal infections) [45]. The main function of LAP appears to be promoting rapid fusion of phagosomes with lysosomes, thereby accelerating conventional phagocytosis [46].

The discovery of LAP illustrates the importance of not solely relying on LC3 as a marker for autophagy. The mere presence of LC3 on pathogen-containing vacuoles cannot distinguish between canonical xenophagy and LAP. At present, the two mechanisms can be discerned using electron microscopy, which allows verification of whether LC3 is associated with a single or double membrane. Further, since LAP targets pathogens already contained in a phagosome, it does not depend on selective cargo receptors and is ULK1-independent. Moreover, in xenophagy the beclin-1 binding protein rubicon suppresses autophagosome maturation, whereas in LAP it positively regulates the generation of an oxidative burst, a key antimicrobial component of phagocytosis [47]. Regardless of the present discrimination methods, it would be beneficial to identify LAP-specific genetic markers.

IFN γ -mediated pathogen elimination

A related intracellular innate immune response involves the IFN γ -mediated control of pathogens, for instance the parasite *T. gondii* in macrophages [8]. In the acute stage of the infection, tachyzoites (fast-replicating forms of the parasite) actively penetrate the host cell at a moving junction and create a non-fusogenic vacuole (Fig. 1c). Inside this membrane-bound parasitophorous vacuole, the parasites replicate by endodyogeny. Meanwhile, extracellular profilin, a *T. gondii* protein crucial for its motility and host cell invasion is

recognized by TLR11 and TLR12 in murine dendritic cells. This causes the dendritic cells to secrete IL-12 in a MyD88-dependent manner, which promotes IFN γ production from natural killer cells. Once the latter is detected by the infected cell, it can halt *T.gondii* replication through multiple mechanisms. The induction of indoleamine 2,3 dioxygenase (IDO) results in reduced tryptophan levels, an essential amino acid without which *T.gondii* replication is decelerated. Second, IFN γ can produce inducible nitric oxide synthase (iNOS), resulting in decreased arginine levels and consequently increased nitric oxide (NO) levels. The former once more induces starvation of the parasite, whereas NO has direct microbicidal effects. Most importantly, IFN γ upregulates p47 GTPases, also known as IRGs, the murine equivalent of human IRGM. During the infection, the IRGs target the parasitophorous vacuole, leading to its disruption as well as the destruction of the parasite's plasma membrane. Subsequently, the stripped parasites are eliminated via lysosome-mediated degradation. Understanding the mechanism for the human immune response to *T. gondii* requires more work. First, humans lack functional TLR11. Supportive roles for TLR7 and TLR9 have been identified in mice, which could play a larger role in humans. Additionally, the human immune response can be mediated more strongly by neutrophils, not dendritic cells. Finally, human IRGM is not regulated by IFN γ , indicating a potentially larger role for NO/IDO-mediated pathogen destruction.

It was recently shown that the canonical autophagy initiation genes ATG14, ULK1, and ULK2 are not required for the IFN γ -mediated immune response to *T. gondii* [25]. However, it does require the downstream ATG5/ATG12/ATG16L1 complex and the presence of ATG3 and ATG7, responsible for LC3-I to LC3-II conversion. No autophagosomes are formed; however, LC3 associates directly with the parasitophorous vacuole membrane. This colocalization is required for the recruitment of the IFN γ -effectors to the parasitophorous vacuole during *T.gondii* infection, suggesting that the destruction of *T.gondii* requires same machinery as the aforementioned LAP. In fact, IFN γ -mediated LAP is observed as an alternative defense mechanism against *M. tuberculosis* [9].

The ATG5/ATG12/ATG16L1 complex also assists the IFN γ -mediated control of murine norovirus by inhibiting the formation of its viral replication complex. In contrast, LC3 is dispensable for this activity [10].

Non-canonical autophagosome formation

Certain studies report the manifestation of autophagosomes resulting in xenophagy in the absence of key ATG genes involved in phagophore formation. Notably, α -hemolysin is a toxin secreted by *Staphylococcus aureus* that is reported to induce a beclin 1-independent autophagy response [48]. Similarly, it was deduced that infection with the foot-and-mouth disease virus can activate autophagy in the absence of VPS34 [49].

The bigger picture: interaction of autophagy and ATG genes with other branches of intracellular immunity

There is accumulating evidence for complex cell type-specific interplay between ATG proteins and other cellular immune responses, yet mechanistic insight is often incomplete

[23, 50]. Pattern recognition receptors (PRRs) such as NOD-like receptors (NLRs), Toll-like receptors (TLRs), and RIG-I-like receptors (RLRs) are central players of innate immunity, capable of activating phagocytosis and various inflammatory processes, as well as chemokines and antimicrobial molecules. ATG16L1 is recruited by NOD1 and NOD2 to the bacterial entry site at the plasma membrane in epithelial cells, and silencing of NOD2 in monocyte-derived dendritic cells decreases LC3-II levels after treatment with the NOD2 ligand muramyl dipeptide [51]. Recent data show that infected bone marrow-derived macrophages lacking caspase-1 show increased autophagy, attributed to caspase-1's ability to cleave the TLR adaptor TRIF [52]. The study of *M. tuberculosis*-infected macrophages reveals how xenophagy can intricately modulate inflammatory responses and inflammasome signaling. The infection induces interleukin-1 β (IL-1 β), IFN γ , and tumor necrosis factor (TNF) levels, all T helper 1 (T_H1) cell-derived cytokines, which evoke xenophagy. On the other hand, T_H2 cytokines such as IL-4 and IL-13 have the opposite effect and inhibit starvation- and IFN γ -induced autophagosome formation [3, 53]. A recent study demonstrates the activation of the DNA damage-regulated autophagy modulator DRAM1 by the TLR/MYD88/NF- κ B signaling cascade, which subsequently activates autophagy [54]. Viri activate RLRs, which negatively regulate xenophagy, likely to prevent excessive IFN levels and control inflammation [55].

Autophagy assists in the secretion of antimicrobial peptides and proteins (AMPs) and intracellular viral restriction factors [23, 50]. For instance, an intestinal epithelial-specific deletion of vitamin D3 receptor (VDR), an IBD risk gene that can activate the AMP LL-37, induces a significant reduction of both ATG16L1 and the antimicrobial enzyme lysozyme as well as an accumulation of p62 *in vivo* [56]. Moreover, recent data showed that the viral restriction factor TRIM5 α doubles as a viral autophagy cargo receptor [18]. Additionally, it was recently observed that TFEB induces the expression of multiple antimicrobial and autophagy proteins in *Caenorhabditis elegans* and activates several proinflammatory cytokines and chemokines in mice after infection with *S. aureus* [57].

Finally, xenophagy assists the adaptive immune response via the fusion of autophagosomes with antigen-loading compartments for major histocompatibility complex class II (MHC-II) molecules. Similarly, a study in macrophages and dendritic cells showed LC3-positive phagosomes enable prolonged antigen presentation on MHC-II molecules [58]. Here, the delay of the fusion of the LC3-decorated phagosome with the lysosome allows the mounting of an adaptive immune response against the sequestered pathogen. Lastly, deletions of key ATG proteins are associated with the reduced development, activity, and survival of lymphocytes [28, 59, 60].

Mouse models provide an opportunity to study the cell type-specific effects of xenophagy, either in isolation from one another (e.g., using a conditional knockout mouse) or in connection with one another (e.g., using a conventional knockout or knock-in mouse). For instance, consider the coding SNP Thr300Ala (T300A) in ATG16L1, which is associated with increased risk for Crohn's disease. Consistent with the human phenotype, Paneth cells of T300A knock-in mice show abnormal lysozyme distribution, accompanied by reduced secretion of their antimicrobial compounds. Additionally, the knock-in mice showed altered goblet cell morphology. These phenotypes contribute to reduced pathogen clearance.

Establishing a vicious cycle, nearby dendritic cells and macrophages display more paracrine signaling of pro-inflammatory molecules [28, 61, 62]. This is a plausible mechanism for the observed chronic infection and altered microbial ecology in patients with Crohn's disease.

Pathogen resistance and persistence to LC3-associated intracellular immunity

Disrupting a crucial step in the host defense pathway is a common pathogen strategy to resist autophagy programs [23, 29, 63-65]. For example, cytosolic *Shigella* produces IcsB that binds to ATG5, thereby averting phagophore elongation [23], and the viral phosphoprotein P of parainfluenza virus type 3 (HPIV3) binds to SNAP29, preventing the SNARE complex from ensuring autophagosome-lysosome fusion [64].

Alternatively, certain pathogens use key elements of xenophagy or LAP to their advantage. *Coxiella burnetii* for instance does not merely resist the lysosomal degradative environment, but exploits the acidic pH for its own metabolic activation [66]. In influenza A, a LIR motif in a viral protein can redirect LC3 to the plasma membrane, where the budding of stable viruses is then promoted [67]. Similarly, vesicular stomatitis virus customizes UVRAG to hijack and redirect virus-bearing endosomes to specific cellular locations where it can deliver its genetic material [68, 69]. Hepatitis C virus engages autophagy to degrade mitochondria (mitophagy). It is hypothesized that the resulting increase in cellular ATP levels causes the observed reduced HCV-induced apoptosis [70].

Some pathogens exploit other intracellular defense mechanisms to reduce xenophagy. Specifically, adherent Invasive *Escherichia coli* (AIEC), associated with Crohn's disease, use the infection-induced NF- κ B response to upregulate microRNAs in the host cell. This reduces ATG5 and ATG16L1 levels, leading to an increased bacterial load and inflammation level [71].

A persistent infection is characterized by a subpopulation of intracellular pathogens hiding from the immune system by slowing or halting their replication altogether, thus assuming a dormant or latent state. For instance, *Listeria monocytogenes* depend on a pore-forming toxin, listeriolysin O (LLO), to escape the phagosome and replicate in the cytosol. However, a fraction of the bacteria that infects murine macrophages is characterized by reduced LLO activity, trapping them inside the phagosome. The lower LLO levels are nevertheless sufficient to block acidification of the phagosome and its fusion with a lysosome, thereby establishing the formation of a spacious *Listeria*-containing phagosome (SLAP) [43]. Inside, the bacteria reduce their replication rate, presumably due to a lack of nutrients. SLAPs have also been observed in a mouse model of *L. monocytogenes* persistent infection.

Similarly, mice infected with *S. Typhimurium* displayed a significant subpopulation of non-replicating bacteria in several organs, even 24 hours after infection [72]. Since most antibiotics target the enzymes of metabolically active and dividing bacteria, persisters can survive antibiotic treatment [73]. Indeed, after 5 days of antibiotic treatment that started 24 hours after inoculation, a small subset of non-replicating persisters was retrieved from the mesenteric lymph nodes, a known site of persistence [72].

Studies in *E. coli* have attributed the ability to develop persistence to the toxin gene of the toxin/antitoxins (TA) operons such as hipA/B, RelE/B, MazF/E, YafQ/DinJ, vapB/C, phd/doc, MqsR/A, HigB/A, and CcdA/B [73, 74]. Either stochastically or due to an external circumstance, such as the presence of antibiotics, oxidative stress, or starvation, the bacterial stress alarmone ppGpp accumulates and mediates the degradation of the operon's antitoxin compound, thereby creating persistence. Analysis of the *Salmonella* genome revealed 14 TA operons, the majority of which contributed to the persister formation [72], and proteomic profiling of *M. tuberculosis*, also capable of establishing latent infection, has revealed multiple nutrient starvation responsive TA systems [75]. Since culturing under nutrient starvation can mediate antitoxin degradation, it is plausible that the nutrient-poor conditions inside bacteria-containing vacuoles induce persister formation. Furthermore, in light of this discussion, it would be interesting to verify whether LLO is part of a TA operon (or comparable system) in the *Listeria* genome.

It should be noted that analogous phenomena have been observed as well for fungi, parasites, and viruses [76]. After the trigger for persistence is removed (e.g., aggressive immune response or antibiotic treatment), the persisters start replicating again and form the basis of a new population, thereby strongly contributing to the occurrence of chronic and recurrent infections [73].

A life-or-death situation on the cellular level

It is insightful to understand how the level of cellular stress mediates the LC3-mediated immune response (Fig. 2). Below, only concepts observed directly in the context of intracellular defense or potentially applicable thereto are described.

Initially, xenophagy and its variants perform a cytoprotective function; however, prolonged and irreversible stress will lead to cell death [77-80]. There is complex crosstalk between autophagy systems and cell death programs. For example, in the absence of cellular stress, beclin-1 is bound to anti-apoptotic proteins of the Bcl-2 family and both autophagy and apoptosis are inactive. At low to moderate stress levels, activated autophagy genes typically present a barrier for programmed cell death (apoptosis). For instance, high-mobility group box 1 (HMGB1), a DAMP, disrupts the beclin-1/Bcl-2 complex. While this does not alter the anti-apoptotic potential of Bcl-2, the now free beclin-1 both induces autophagy and inhibits apoptosis [79, 81, 82]. Past a critical stress level, complex regulatory mechanisms induce programmed cell death, where the cell breaks into multiple apoptotic bodies. Once this happens, autophagy is inactivated in part due to the caspase- or calpain-mediated cleavage of essential autophagy proteins such as ATG3 and beclin-1 [79]. Further, truncated ATG proteins can exhibit pro-apoptotic properties. For instance, antimicrobial peptides can cause an influx of calcium, which allows calpain-mediated cleavage of ATG5, actively contributing to apoptosis [83]. Meanwhile, studies have shown the potential of key autophagy proteins involved in autophagosome formation induce apoptosis via caspase activation [79].

Autophagosome accumulation can result in programmed cell necrosis, characterized by a loss of cell membrane integrity and release of cellular contents in the cytoplasm [79].

Finally, recent work showed that autophagy triggers, such as starvation and autophagy-inducing peptides, can induce a form of cell death with distinct morphological features, including the increased presence of autophagosome/autolysosomes. This autophagic cell death is termed “autosis” and is regulated by the sodium-potassium pump Na^+/K^+ -ATPase [80].

LAP has been observed to efficiently clear apoptotic corpses and necrotic cells [10, 84-86]. Here, LAP is triggered by the engagement of TIM4, a receptor for phosphatidylserine, displayed by dying cells. LAP-defective macrophages produce more pro- and fewer anti-inflammatory cytokines [86], which could contribute to disease.

Conclusion

Multiple ATG gene-dependent intracellular defense mechanisms have been identified, although to date, studies have largely focused on xenophagy, the autophagic clearance of pathogens. Additionally, a clearer picture of the complex interplay between cytoprotective autophagy-dependent systems and cell death cassettes is emerging.

Pathogens continually evolve to develop strategies to resist and persist ATG-dependent destruction. This fact, combined with their roles in numerous human infectious diseases and gastrointestinal disorders, underscores the importance of mechanistic studies of the pathways and regulatory networks involved in both the immune response and pathogen's survival strategies. The recent successful feasibility studies of conducting genome-wide CRISPR-Cas9 knockout screens in human cells pave the way for their application in this area [87, 88]. Relevant mouse models can help piece together a holistic picture of the consequences of impaired immunity *in vivo*.

Acknowledgments

This work was supported by funding from the Crohn's and Colitis Foundation of America, the Leona M. and Harry B. Helmsley Charitable Trust, and NIH grants DK097485 and DK043351 (to R.J.X.).

We thank Natalia Nedelsky for assistance with figures.

References

Papers of particular interest, published within the annual period of review, have been highlighted as:

*of special interest

**of outstanding interest

1. Choi AM, Ryter SW, Levine B. Autophagy in human health and disease. *N Engl J Med*. 2013; 368:651–662. [PubMed: 23406030]
2. Mizushima N, Yoshimori T, Ohsumi Y. The role of Atg proteins in autophagosome formation. *Annu Rev Cell Dev Biol*. 2011; 27:107–132. [PubMed: 21801009]
3. Deretic V, Saitoh T, Akira S. Autophagy in infection, inflammation and immunity. *Nat Rev Immunol*. 2013; 13:722–737. [PubMed: 24064518]
- 4*. Brinar M, Vermeire S, Cleynen I, et al. Genetic variants in autophagy-related genes and granuloma formation in a cohort of surgically treated Crohn's disease patients. *J Crohns Colitis*. 2012; 6:43–50. This article describes associations found between autophagy-associated proteins ATG4 and

- FNBP1L and the formation of granulomas in Crohn's disease, indicative of a more aggressive disease course. [PubMed: 22261526]
- 5*. Van Limbergen J, Radford-Smith G, Satsangi J. Advances in IBD genetics. *Nat Rev Gastroenterol Hepatol*. 2014; 11:372–385. This work reviews associations between SNPs in autophagy genes and IBD. [PubMed: 24614343]
 6. Romao S, Munz C. LC3-associated phagocytosis. *Autophagy*. 2014; 10:526–528. [PubMed: 24413059]
 7. Lai SC, Devenish RJ. LC3-Associated Phagocytosis (LAP): Connections with Host Autophagy. *Cells*. 2012; 1:396–408. [PubMed: 24710482]
 8. Sturge CR, Yarovsky F. Complex immune cell interplay in the gamma interferon response during *Toxoplasma gondii* infection. *Infect Immun*. 2014; 82:3090–3097. [PubMed: 24866795]
 9. Bradfute SB, Castillo EF, Arko-Mensah J, et al. Autophagy as an immune effector against tuberculosis. *Curr Opin Microbiol*. 2013; 16:355–365. [PubMed: 23790398]
 10. Bestebroer J, V'Kovski P, Mauthe M, et al. Hidden behind autophagy: the unconventional roles of ATG proteins. *Traffic*. 2013; 14:1029–1041. [PubMed: 23837619]
 - 11*. Lamb CA, Yoshimori T, Tooze SA. The autophagosome: origins unknown, biogenesis complex. *Nat Rev Mol Cell Biol*. 2013; 14:759–774. This work summarizes the important protein complexes in autophagosome formation as well as the potential sources of the phagophore membrane. [PubMed: 24201109]
 12. Shpilka T, Weidberg H, Pietrokovski S, et al. Atg8: an autophagy-related ubiquitin-like protein family. *Genome Biol*. 2011; 12:226. [PubMed: 21867568]
 13. Lamark T, Kirkin V, Dikic I, et al. NBR1 and p62 as cargo receptors for selective autophagy of ubiquitinated targets. *Cell Cycle*. 2009; 8:1986–1990. [PubMed: 19502794]
 - 14*. Boyle KB, Randow F. The role of 'eat-me' signals and autophagy cargo receptors in innate immunity. *Curr Opin Microbiol*. 2013; 16:339–348. This work describes the role of cargo receptors, their interaction with galectin-8 and poly-ubiquitin, and how their LIR domains determine specificity. [PubMed: 23623150]
 15. Novak I, Kirkin V, McEwan DG, et al. Nix is a selective autophagy receptor for mitochondrial clearance. *EMBO Rep*. 2010; 11:45–51. [PubMed: 20010802]
 - 16*. Mancias JD, Wang X, Gygi SP, et al. Quantitative proteomics identifies NCOA4 as the cargo receptor mediating ferritinophagy. *Nature*. 2014; 509:105–109. This article describes how quantitative proteomics can detect proteins associated with autophagosomes. Specifically, a novel cargo receptor, NCOA4, is identified that specifically targets ferritin for autophagy-mediated degradation. [PubMed: 24695223]
 17. Goodall M, Thorburn A. Identifying specific receptors for cargo-mediated autophagy. *Cell Res*. 2014; 24:783–784. [PubMed: 24797431]
 - 18*. Mandell MA, Jain A, Arko-Mensah J, et al. Proteins Regulate Autophagy and Can Target Autophagic Substrates by Direct Recognition. *Dev Cell*. 2014; 30:394–409. An siRNA screen of various TRIM proteins reveals their role in autophagy. Notably, TRIM5α, a known retroviral restriction factor, is identified as a cargo receptor for viral capsid proteins. [PubMed: 25127057]
 19. Wild P, McEwan DG, Dikic I. The LC3 interactome at a glance. *J Cell Sci*. 2014; 127:3–9. [PubMed: 24345374]
 20. Randow F, Youle RJ. Self and nonself: how autophagy targets mitochondria and bacteria. *Cell Host Microbe*. 2014; 15:403–411. [PubMed: 24721569]
 - 21*. Sawa-Makarska J, Abert C, Romanov J, et al. Cargo binding to Atg19 unmasks additional Atg8 binding sites to mediate membrane-cargo apposition during selective autophagy. *Nat Cell Biol*. 2014; 16:425–433. This study in yeast identifies a LIR-independent interaction of cargo receptor ATG19 (the yeast orthologue of p62) with ATG8 (the yeast orthologue of LC3). This interaction is assumed to tightly bend the phagophores around the cargo, such that other cytoplasmic cargo is not accidentally included in the autophagosome. [PubMed: 24705553]
 - 22*. Pfaffenwimmer T, Reiter W, Brach T, et al. Hrr25 kinase promotes selective autophagy by phosphorylating the cargo receptor Atg19. *EMBO Rep*. 2014; 15:862–870. This work attributes the interaction between ATG8 and ATG19 (described in reference 21), to the action of the kinase Hrr25. [PubMed: 24968893]

23. Gomes LC, Dikic I. Autophagy in antimicrobial immunity. *Mol Cell*. 2014; 54:224–233. [PubMed: 24766886]
24. Duan Z, Chen Y, Huang W, et al. Linkage of autophagy to fungal development, lipid storage and virulence in *Metarhizium robertsii*. *Autophagy*. 2013; 9:538–549. [PubMed: 23380892]
- 25**. Choi J, Park S, Biering SB, et al. The parasitophorous vacuole membrane of *Toxoplasma gondii* is targeted for disruption by ubiquitin-like conjugation systems of autophagy. *Immunity*. 2014; 40:924–935. This article shows that ATG5/ATG12/ATG16L1 as well as ATG7 and ATG3 assist in the conjugation of LC3 to the parasitophorous vacuole membrane of *T. gondii*, suggesting these proteins are required for the efficient IFN γ -mediated destruction of the pathogen. [PubMed: 24931121]
- 26*. Richetta C, Faure M. Autophagy in antiviral innate immunity. *Cell Microbiol*. 2013; 15:368–376. This review gives a comprehensive overview of the role autophagy plays in antiviral immunity. [PubMed: 23051682]
27. Fredlund J, Enninga J. Cytoplasmic access by intracellular bacterial pathogens. *Trends Microbiol*. 2014; 22:128–137. [PubMed: 24530174]
- 28*. Conway KL, Kuballa P, Song JH, et al. Atg16l1 is required for autophagy in intestinal epithelial cells and protection of mice from *Salmonella* infection. *Gastroenterology*. 2013; 145:1347–1357. The authors of this work use mice with a conditional knockout of the IBD-associated gene *ATG16L1*, specifically in epithelial cells, to demonstrate its role in the regulation of autophagy in intestinal epithelial cells. *ATG16L1* is required for bacterial clearance and to prevent systemic infection of mice with enteric bacteria. [PubMed: 23973919]
29. Baxt LA, Garza-Mayers AC, Goldberg MB. Bacterial subversion of host innate immune pathways. *Science*. 2013; 340:697–701. [PubMed: 23661751]
30. Mostowy S, Sancho-Shimizu V, Hamon MA, et al. p62 and NDP52 proteins target intracytosolic *Shigella* and *Listeria* to different autophagy pathways. *J Biol Chem*. 2011; 286:26987–26995. [PubMed: 21646350]
- 31*. Li S, Wandel MP, Li F, et al. Sterical hindrance promotes selectivity of the autophagy cargo receptor NDP52 for the danger receptor galectin-8 in antibacterial autophagy. *Sci Signal*. 2013; 6:ra9. This article reports the crystal structure of the NDP52-galectin-8 complex, demonstrating the molecular basis for the specificity of NDP52 for galectin-8, to the exclusion of other galectins. [PubMed: 23386746]
- 32*. Fujita N, Morita E, Itoh T, et al. Recruitment of the autophagic machinery to endosomes during infection is mediated by ubiquitin. *J Cell Biol*. 2013; 203:115–128. This article examines the precise role for ubiquitin in xenophagy, demonstrating that ubiquitin directly binds ATG16L1 and connects bacteria-containing endosomes with autophagic machinery. [PubMed: 24100292]
- 33*. Rogov V, Dotsch V, Johansen T, et al. Interactions between autophagy receptors and ubiquitin-like proteins form the molecular basis for selective autophagy. *Mol Cell*. 2014; 53:167–178. This article discusses the recent findings that ubiquitin-like proteins such as Atg8/LC3/GABARAP and Atg5 directly engage regulatory and core autophagy proteins, and discusses the implications of these findings in our understanding of selective autophagosome formation. [PubMed: 24462201]
34. Hubber A, Kubori T, Nagai H. Modulation of the ubiquitination machinery by *Legionella*. *Curr Top Microbiol Immunol*. 2013; 376:227–247. [PubMed: 23918174]
- 35**. Manzanillo PS, Ayres JS, Watson RO, et al. The ubiquitin ligase parkin mediates resistance to intracellular pathogens. *Nature*. 2013; 501:512–516. This article demonstrates a role for parkin in ubiquitin-mediated autophagy of *M. tuberculosis*, suggesting a compelling link between mitophagy and infectious disease. [PubMed: 24005326]
36. Huett A, Heath RJ, Begun J, et al. The LRR and RING domain protein LRSAM1 is an E3 ligase crucial for ubiquitin-dependent autophagy of intracellular *Salmonella* Typhimurium. *Cell Host Microbe*. 2012; 12:778–790. [PubMed: 23245322]
- 37*. Barber GN. STING-dependent cytosolic DNA sensing pathways. *Trends Immunol*. 2014; 35:88–93. This article reviews the current understanding of STING function, including its interaction with autophagy and inflammation. [PubMed: 24309426]
38. Cao Y, Zhang L. A Smurf1 tale: function and regulation of an ubiquitin ligase in multiple cellular networks. *Cell Mol Life Sci*. 2013; 70:2305–2317. [PubMed: 23007848]

39. Watson RO, Manzanillo PS, Cox JS. Extracellular *M. tuberculosis* DNA targets bacteria for autophagy by activating the host DNA-sensing pathway. *Cell*. 2012; 150:803–815. [PubMed: 22901810]
40. Tattoli I, Sorbara MT, Vuckovic D, et al. Amino acid starvation induced by invasive bacterial pathogens triggers an innate host defense program. *Cell Host Microbe*. 2012; 11:563–575. [PubMed: 22704617]
41. Settembre C, Di Malta C, Polito VA, et al. TFEB links autophagy to lysosomal biogenesis. *Science*. 2011; 332:1429–1433. [PubMed: 21617040]
42. Roczniaik-Ferguson A, Petit CS, Froehlich F, et al. The transcription factor TFEB links mTORC1 signaling to transcriptional control of lysosome homeostasis. *Sci Signal*. 2012; 5:ra42. [PubMed: 22692423]
- 43*. Lam GY, Cemma M, Muise AM, et al. Host and bacterial factors that regulate LC3 recruitment to *Listeria monocytogenes* during the early stages of macrophage infection. *Autophagy*. 2013; 9:985–995. This article describes how a subpopulation of *Listeria* succeeds in evading the fusion of lysosomes with the LC3-associated phagosomes that contain them, by slowing their growth rate and thereby becoming persistent. [PubMed: 23584039]
44. Huang J, Canadien V, Lam GY, et al. Activation of antibacterial autophagy by NADPH oxidases. *Proc Natl Acad Sci U S A*. 2009; 106:6226–6231. [PubMed: 19339495]
45. Ma J, Becker C, Lowell CA, et al. Dectin-1-triggered recruitment of light chain 3 protein to phagosomes facilitates major histocompatibility complex class II presentation of fungal-derived antigens. *J Biol Chem*. 2012; 287:34149–34156. [PubMed: 22902620]
46. Sanjuan MA, Dillon CP, Tait SW, et al. Toll-like receptor signalling in macrophages links the autophagy pathway to phagocytosis. *Nature*. 2007; 450:1253–1257. [PubMed: 18097414]
47. Mehta P, Henault J, Kolbeck R, et al. Noncanonical autophagy: one small step for LC3, one giant leap for immunity. *Curr Opin Immunol*. 2014; 26:69–75. [PubMed: 24556403]
48. Mestre MB, Fader CM, Sola C, et al. Alpha-hemolysin is required for the activation of the autophagic pathway in *Staphylococcus aureus*-infected cells. *Autophagy*. 2010; 6:110–125. [PubMed: 20110774]
49. Berryman S, Brooks E, Burman A, et al. Foot-and-mouth disease virus induces autophagosomes during cell entry via a class III phosphatidylinositol 3-kinase-independent pathway. *J Virol*. 2012; 86:12940–12953. [PubMed: 22993157]
50. Ma Y, Galluzzi L, Zitvogel L, et al. Autophagy and cellular immune responses. *Immunity*. 2013; 39:211–227. [PubMed: 23973220]
- 51*. Salem M, Seidelin JB, Rogler G, et al. Muramyl dipeptide responsive pathways in Crohn's disease: from NOD2 and beyond. *Cell Mol Life Sci*. 2013; 70:3391–3404. This review highlights how muramyl dipeptide is involved in the activation of inflammatory pathways that have been implicated in Crohn's disease. [PubMed: 23275943]
- 52*. Jabir MS, Ritchie ND, Li D, et al. Caspase-1 cleavage of the TLR adaptor TRIF inhibits autophagy and beta-interferon production during *Pseudomonas aeruginosa* infection. *Cell Host Microbe*. 2014; 15:214–227. This article demonstrates how cleavage of TRIF by caspase 1 downregulates autophagy, reduces production of type I interferons, and activates the inflammasome. [PubMed: 24528867]
53. Goletti D, Petruccioli E, Romagnoli A, et al. Autophagy in *Mycobacterium tuberculosis* infection: a passepartout to flush the intruder out? *Cytokine Growth Factor Rev*. 2013; 24:335–343. [PubMed: 23395260]
- 54*. van der Vaart M, Korbee CJ, Lamers GE, et al. The DNA damage-regulated autophagy modulator DRAM1 links mycobacterial recognition via TLR-MYD88 to autophagic defense. *Cell Host Microbe*. 2014; 15:753–767. This paper uses both zebrafish and human macrophage models of mycobacterial infection to demonstrate that mycobacterial infection induces DRAM1 in a MyD88- and NF- κ B-dependent manner, resulting in DRAM1 localization to mycobacteria within macrophages. Furthermore, DRAM1 mediates autophagic flux and lysosomal maturation. [PubMed: 24922577]
55. Chiramel AI, Brady NR, Bartenschlager R. Divergent roles of autophagy in virus infection. *Cells*. 2013; 2:83–104. [PubMed: 24709646]

- 56**. Wu S, Zhang YG, Lu R, et al. Intestinal epithelial vitamin D receptor deletion leads to defective autophagy in colitis. *Gut*. 2014 This article reports how the vitamin D receptor interacts with autophagy and the microbiome in a manner that promotes intestinal homeostasis, with potential implications for IBD. This work highlights a link between autophagy proteins and antimicrobial peptides.
- 57**. Visvikis O, Ihuegbu N, Labeled SA, et al. Innate host defense requires TFEB-mediated transcription of cytoprotective and antimicrobial genes. *Immunity*. 2014; 40:896–909. This work reports that TFEB is a key transcription factor in the host response to infection, demonstrating that TFEB is induced by *S. aureus* infection and drives antimicrobial and autophagy genes. [PubMed: 24882217]
- 58**. Romao S, Gasser N, Becker AC, et al. Autophagy proteins stabilize pathogen-containing phagosomes for prolonged MHC II antigen processing. *J Cell Biol*. 2013; 203:757–766. This article reports on a link between LAP and adaptive immunity. The data suggest that human macrophages and dendritic cells infected with *Candida albicans* pulsed with antigen, allow for prolonged antigen presentation to T cells as a result of LAP. [PubMed: 24322427]
- 59*. Conway KL, Kuballa P, Khor B, et al. ATG5 regulates plasma cell differentiation. *Autophagy*. 2013; 9:528–537. This work identifies an important role of the autophagy gene *ATG5* in adaptive immunity. Notably, deletion of *ATG5* in B cells leads to decreased antibody secretion. [PubMed: 23327930]
60. Liu G, Bi Y, Wang R, et al. Self-eating and self-defense: autophagy controls innate immunity and adaptive immunity. *J Leukoc Biol*. 2013; 93:511–519. [PubMed: 23271703]
- 61*. Lassen KG, Kuballa P, Conway KL, et al. Atg16L1 T300A variant decreases selective autophagy resulting in altered cytokine signaling and decreased antibacterial defense. *Proc Natl Acad Sci U S A*. 2014; 111:7741–7746. This article highlights the cell-dependent effects of a T300A SNP in *ATG16L1*, associated with Crohn's disease, by generating a knock-in mouse model. Moreover, this work shows how the phenotypes caused by the SNP in various cell types are interconnected. [PubMed: 24821797]
62. Lassen KG, Xavier RJ. An alteration in *ATG16L1* stability in Crohn disease. *Autophagy*. 2014; 10
63. Huang J, Brumell JH. Bacteria-autophagy interplay: a battle for survival. *Nat Rev Microbiol*. 2014; 12:101–114. [PubMed: 24384599]
- 64*. Ding B, Zhang G, Yang X, et al. Phosphoprotein of human parainfluenza virus type 3 blocks autophagosome-lysosome fusion to increase virus production. *Cell Host Microbe*. 2014; 15:564–577. This work identifies the mechanism by which parainfluenza virus type 3 avoids autophagy: its phosphoprotein binds to a SNARE protein, thereby inhibiting the latter from mediating autophagosome-lysosome fusion. [PubMed: 24832451]
65. Rajsbaum R, Garcia-Sastre A. Viral evasion mechanisms of early antiviral responses involving regulation of ubiquitin pathways. *Trends Microbiol*. 2013; 21:421–429. [PubMed: 23850008]
66. van Schaik EJ, Chen C, Mertens K, et al. Molecular pathogenesis of the obligate intracellular bacterium *Coxiella burnetii*. *Nat Rev Microbiol*. 2013; 11:561–573. [PubMed: 23797173]
- 67**. Beale R, Wise H, Stuart A, et al. A LC3-interacting motif in the influenza A virus M2 protein is required to subvert autophagy and maintain virion stability. *Cell Host Microbe*. 2014; 15:239–247. This study describes how the Matrix 2 (M2) ion-channel protein of influenza A virus contains a LIR that interacts with LC3 and redirects it to the plasma membrane, which benefits virus budding and virion stability. [PubMed: 24528869]
68. Pirooz S, He S, O'Connell D, et al. Viruses customize autophagy protein for efficient viral entry. *Autophagy*. 2014; 10:1355–1356. [PubMed: 24905575]
- 69*. Pirooz SD, He S, Zhang T, et al. UVRAG is required for virus entry through combinatorial interaction with the class C-Vps complex and SNAREs. *Proc Natl Acad Sci U S A*. 2014; 111:2716–2721. This work demonstrates how the autophagy-related gene UVRAG regulates viral entry of negative-strand RNA viri by assembling a fusogenic SNARE complex. [PubMed: 24550300]
- 70**. Kim SJ, Syed GH, Khan M, et al. Hepatitis C virus triggers mitochondrial fission and attenuates apoptosis to promote viral persistence. *Proc Natl Acad Sci U S A*. 2014; 111:6413–6418. This study shows how hepatitis C virus can instruct mitophagy through phosphorylation of Drp1. The

resulting increase in intracellular ATP is hypothesized to prevent apoptosis, ensuring viral survival. [PubMed: 24733894]

- 71*. Nguyen HT, Dalmasso G, Muller S, et al. Crohn's disease-associated adherent invasive *Escherichia coli* modulate levels of microRNAs in intestinal epithelial cells to reduce autophagy. *Gastroenterology*. 2014; 146:508–519. AIEC bacteria induce NF- κ B signaling, which upregulates microRNAs that inhibit the expression of key ATG genes, thereby disrupting the autophagy pathway. This work is especially relevant since the same microRNAs and ATG genes are upregulated and reduced, respectively, in patients with Crohn's disease. [PubMed: 24148619]
- 72*. Helaine S, Cheverton AM, Watson KG, et al. Internalization of *Salmonella* by macrophages induces formation of nonreplicating persisters. *Science*. 2014; 343:204–208. This work illustrates that a subpopulation of *Salmonella* becomes persistent inside a vacuolar niche by halting their replication. [PubMed: 24408438]
73. Maisonneuve E, Gerdes K. Molecular mechanisms underlying bacterial persisters. *Cell*. 2014; 157:539–548. [PubMed: 24766804]
74. Cohen NR, Lobritz MA, Collins JJ. Microbial persistence and the road to drug resistance. *Cell Host Microbe*. 2013; 13:632–642. [PubMed: 23768488]
- 75*. Albrethsen J, Agner J, Piersma SR, et al. Proteomic profiling of *Mycobacterium tuberculosis* identifies nutrient-starvation-responsive toxin-antitoxin systems. *Mol Cell Proteomics*. 2013; 12:1180–1191. A proteomic study reveals multiple toxin/antitoxin operons under conditions of nutrient starvation, forming the basis of bacterial persistence. [PubMed: 23345537]
76. Zhang Y. Persisters, persistent infections and the Yin-Yang model. *Emerg Microbes Infect*. 2014; 3:e3.
77. Deegan S, Saveljeva S, Gorman AM, et al. Stress-induced self-cannibalism: on the regulation of autophagy by endoplasmic reticulum stress. *Cell Mol Life Sci*. 2013; 70:2425–2441. [PubMed: 23052213]
78. Mukhopadhyay S, Panda PK, Sinha N, et al. Autophagy and apoptosis: where do they meet? *Apoptosis*. 2014; 19:555–566. [PubMed: 24415198]
79. Marino G, Niso-Santano M, Baehrecke EH, et al. Self-consumption: the interplay of autophagy and apoptosis. *Nat Rev Mol Cell Biol*. 2014; 15:81–94. [PubMed: 24401948]
- 80*. Liu Y, Shoji-Kawata S, Sumpter RM Jr, et al. Autosis is a Na⁺,K⁺-ATPase-regulated form of cell death triggered by autophagy-inducing peptides, starvation, and hypoxia-ischemia. *Proc Natl Acad Sci U S A*. 2013; 110:20364–20371. This study demonstrates how autophagy triggers such as starvation and the autophagy-inducing peptide Tat-beclin 1 induce a form of cell death that is morphologically distinct from necrosis and apoptosis. Initially the dying cells are characterized by an increased amount of autophagosomes and autolysosomes. At later stages the cell wall appears to lyse, similar to what occurs in necrosis. However, necrotic and apoptotic epithelial cells float, whereas autotic cells better retain adhesion. [PubMed: 24277826]
81. Lee JS, Ha TK, Park JH, et al. Anti-cell death engineering of CHO cells: co-overexpression of Bcl-2 for apoptosis inhibition, Beclin-1 for autophagy induction. *Biotechnol Bioeng*. 2013; 110:2195–2207. [PubMed: 23436561]
82. Takacs-Vellai K, Vellai T, Puoti A, et al. Inactivation of the autophagy gene bec-1 triggers apoptotic cell death in *C. elegans*. *Curr Biol*. 2005; 15:1513–1517. [PubMed: 16111945]
- 83*. Shi M, Zhang T, Sun L, et al. Calpain, Atg5 and Bak play important roles in the crosstalk between apoptosis and autophagy induced by influx of extracellular calcium. *Apoptosis*. 2013; 18:435–451. Calpains are calcium-dependent proteases involved in apoptosis. This study shows that the influx of calcium (e.g., as a result of antimicrobial peptides) results in the calpain-mediated cleavage of ATG5. The truncated form of ATG5 both inhibits autophagy and simultaneously actively induces apoptosis. This work provides insight into the complex relationship between autophagy and apoptosis. [PubMed: 23242420]
- 84*. Green DR, Levine B. To be or not to be? How selective autophagy and cell death govern cell fate. *Cell*. 2014; 157:65–75. This review highlights the complex crosstalk between autophagy programs and cell death. Notably, LC3-associated phagocytosis is able to efficiently clear apoptotic and necrotic cells induced by TIM4. [PubMed: 24679527]
85. Huang S, Jia K, Wang Y, et al. Autophagy genes function in apoptotic cell corpse clearance during *C. elegans* embryonic development. *Autophagy*. 2013; 9:138–149. [PubMed: 23108454]

86. Martinez J, Almendinger J, Oberst A, et al. Microtubule-associated protein 1 light chain 3 alpha (LC3)-associated phagocytosis is required for the efficient clearance of dead cells. *Proc Natl Acad Sci U S A*. 2011; 108:17396–17401. [PubMed: 21969579]
- 87*. Wang T, Wei JJ, Sabatini DM, et al. Genetic screens in human cells using the CRISPR-Cas9 system. *Science*. 2014; 343:80–84. This study demonstrates the feasibility of performing a CRISPR-based genome-wide knockout screen. Completely knocked out genes likely produce a stronger and more reliable phenotype than silenced genes. Thus the application of CRISPR-based genome-wide studies may reveal novel autophagy-related genes and pathways. [PubMed: 24336569]
88. Shalem O, Sanjana NE, Hartenian E, et al. Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science*. 2014; 343:84–87. [PubMed: 24336571]

Key points

- Autophagy and ATG gene-dependent processes such as IFN γ -induced pathogen elimination and LC3-associated phagocytosis (LAP) are vital host immune responses against a broad range of intracellular pathogens.
- Pathogens can resist these processes by a variety of mechanisms, including disrupting key steps in the degradation pathway, adapting to the associated degradative conditions, and exploiting host immune responses to promote their replication and/or survival.
- A persistent infection is characterized by a subpopulation of intracellular pathogens slowing or halting their replication inside a vacuole, thereby evading both immune responses and antibiotic drug action.
- Autophagy and various cell death programs show mutual regulation; in addition, LAP is essential for the efficient clearance of apoptotic and necrotic cells.

Figure 1. Pathways for LC3-associated pathogen defense

(a) Xenophagy, (b) LC3-associated phagocytosis, (c) IFN γ -induced pathogen degradation illustrated for *T. gondii* infection.

Figure 2. Complex stress level-dependent interplay between xenophagy, LAP, apoptosis, autosis, and necrosis

At low stress levels, autophagy proteins such as beclin-1 inhibit apoptosis. At higher stress intensity, apoptosis negatively regulates autophagy by cleaving ATG proteins. The truncated ATG proteins either are non-functional or obtain pro-apoptotic properties. Genes involved in autophagosome formation can induce apoptosis by activating caspases. Autophagy triggers can activate a morphologically distinct form of cell death, termed autosis. Finally, an accumulation of autophagosomes can induce necrosis. LC3-associated phagocytosis (LAP) efficiently removes apoptotic and necrotic cells.