Published in final edited form as:

J Child Psychol Psychiatry. 2015 April ; 56(4): 432–443. doi:10.1111/jcpp.12307.

Homotopic connectivity in drug-naïve, first-episode, early-onset schizophrenia

Hui-Jie Li^{1,2}, Yong Xu³, Ke-Rang Zhang³, Matthew J. Hoptman^{4,5}, and Xi-Nian Zuo^{1,2}

¹Key Laboratory of Behavioral Science and Magnetic Resonance Imaging Research Center, Chinese Academy of Sciences, Beijing, China

²Laboratory for Functional Connectome and Development, Institute of Psychology, Chinese Academy of Sciences, Beijing, China

³Department of Psychiatry, First Hospital of Shanxi Medical University, Taiyuan, China

⁴Department of Psychiatry, New York University School of Medicine, New York, USA

⁵Schizophrenia Research Division, Nathan S. Kline Institute for Psychiatric Research, Orangeburg, USA

Abstract

Background—The disconnection hypothesis of schizophrenia has been extensively tested in adults. Recent studies have reported the presence of brain disconnection in younger patients, adding evidence to support the neurodevelopmental hypothesis of schizophrenia. Because of drug confounds in chronic and medicated patients, it has been extremely challenging for researchers to directly investigate abnormalities in the development of connectivity and their role in the pathophysiology of schizophrenia. The present study aimed to examine functional homotopy – a measure of interhemispheric connection – and its relevance to clinical symptoms in first-episode drug-naïve early-onset schizophrenia (EOS) patients.

Methods—Resting-state functional magnetic resonance imaging was performed in 26 first-episode drug-naïve EOS patients (age: 14.5 ± 1.94 , 13 males) and 25 matched typically developing controls (TDCs) (age: 14.4 ± 2.97 , 13 males). We were mainly concerned with the functional connectivity between any pair of symmetric inter-hemispheric voxels (i.e., functional homotopy) measured by voxel-mirrored homotopic connectivity (VMHC).

Results—EOS patients exhibited both global and regional VMHC reductions in comparison with TDCs. Reduced VMHC values were observed within the superior temporal cortex and postcentral gyrus. These interhemispheric synchronization deficits were negatively correlated with negative symptom of the Positive and Negative Syndrome Scale. Moreover, regions of interest analyses

Correspondence: Xi-Nian Zuo, Key Laboratory of Behavioral Science, Institute of Psychology, Chinese Academy of Sciences; No 16 Lincui Road, Chaoyang District, Beijing, 100101, China; zuoxn@psych.ac.cn or zuoxinian@gmail.com. X-N. Z and Y. X. contributed equally as corresponding authors.

Conflict of interest statement: No conflicts declared.

The authors have declared that they have no potential or competing conflicts of interests.

Supporting Information

Additional supporting information is provided along with the online version of this article.

Appendix S1. Additional materials and methods

based on left and right clusters of temporal cortex and postcentral gyrus revealed abnormal heterotopic connectivity in EOS patients.

Conclusions—Our findings provide novel neurodevelopmental evidence for the disconnection hypothesis of schizophrenia and suggest that these alterations occur early in the course of the disease and are independent of medication status.

Keywords

resting-state fMRI; early-onset schizophrenia; functional connectivity; interhemispheric connectivity

Introduction

Schizophrenia is a severe psychotic disorder that affects approximately 1% of the population (Lopez & Murray, 1998) and is increasingly considered a disconnection syndrome (Friston & Frith, 1995; Frith et al., 1995; Pettersson-Yeo, Allen, Benetti, McGuire, & Mechelli, 2011). Using a recently developed brain imaging technique, resting state functional magnetic resonance imaging (RFMRI), a large number of studies have shown that patients with schizophrenia exhibit deficits in functional connectivity (Fornito, Zalesky, Pantelis, & Bullmore, 2012; Schmitt, Hasan, Gruber, & Falkai, 2011; Stephan, Baldeweg, & Friston, 2006). Pettersson-Yeo et al. (2011) systematically reviewed the RFMRI literature on schizophrenia and found that most studies consistently report decreased functional connectivity in different stages of schizophrenia (high risk, early-onset, first-episode, chronic). This decreased connectivity appeared mainly between the frontal lobe, the fronto-temporal lobe, the corpus callosum, the anterior cingulate gyrus and other cortical and subcortical brain regions (Pettersson-Yeo et al., 2011). Of note, most of the studies in the review focused on adulthood-onset schizophrenia, with only a few studies investigating functional connectivity in early-onset schizophrenia (EOS) patients.

“Early onset” refers to an onset of psychosis before the 18th birthday (Rhinewine et al., 2005; Vourdas, Pipe, Corrigall, & Frangou, 2003). Early onset patients have been found to be more impaired and have poorer outcomes than adult schizophrenia patients (Rhinewine et al., 2005; White, Schmidt, Kim & Calhoun, 2011). The neurodevelopmental hypothesis of schizophrenia proposes that abnormal neurodevelopment causes the illness (Owen, O’Donovan, Thapar, & Craddock, 2011; Rapoport, Giedd, & Gogtay, 2012; Weinberger, 1987). This highlights the importance of investigating the neurodevelopment of EOS, especially first-episode drug-naïve patients. Until now, few studies have explored functional connectivity using RFMRI in EOS patients. Ke et al.(2009) first observed reduced functional connectivity between right frontal regions and other brain regions in EOS patients. EOS patients also exhibited decreased functional connectivity between the cingulate gyrus and a variety of areas including frontal, temporal, occipital, and subcortical regions (Zhou, Tan, Tang, Chen, 2010). Alexander-Bloch and colleagues reported that childhood-onset schizophrenia (COS, defined as onset before age 13 years) patients showed deficits in whole-brain network local organization and brain modular community structure (Alexander-Bloch et al., 2010, 2012, 2013). Of note, in these studies, most of the EOS and COS patients were medicated. These studies therefore involve a potential confounding effect

of medication on the differences in functional connectivity. Antipsychotic medication has been shown to influence functional brain connectivity in schizophrenia (Honey et al., 1999).

Increasing evidence suggests that the disconnection syndrome in schizophrenia might include disruption of interhemispheric connectivity, as found in previous behavioral (Barnett, Kirk, & Corballis, 2007; Mohr, Pulvermuller, Cohen, & Rockstroh, 2000) and psychophysiological studies (Mohr, Pulvermuller, Rockstroh, & Endrass, 2008; Morrison-Stewart, Velikonja, Corning, & Williamson, 1996). Much of this interaction is mediated by the corpus callosum, which primarily connects homotopic regions of the two hemispheres (Hoptman & Davidson, 1994). As a fundamental characteristic of intrinsic brain architecture (Stark et al., 2008), functional homotopy – voxel-mirrored homotopic connectivity (VMHC) undergoes remarkable dynamic changes during the lifespan (Zuo et al., 2010). Using the VMHC approach, two studies reported reductions in functional homotopy in adult schizophrenia (Hoptman et al., 2012; Guo et al., 2014c).

Up to now, few studies investigated the functional connectivity in EOS, and no studies explored the functional connectivity in unmedicated EOS patients. This highlights the need for investigations focused on brain connectivity in drug-naïve EOS. In the present study, we examined neuroimaging data from 26 drug-naïve first-episode EOS patients and 25 demographically matched typically developing controls (TDCs). We used the VMHC method to examine potentially abnormal functional homotopy in this early stage of schizophrenia.

Materials and methods

Participants

Thirty-two EOS patients were recruited from the first hospital of Shanxi Medical University. The clinical diagnosis was made by two consultant psychiatrists according to a Chinese version of the Modified Structured Clinical Interview for DSM-IV for patient version (SCID-I/P). The patients who have other Axis-I or Axis-II comorbidity disorders were excluded. The age of the EOS patients ranged from 9.0 to 17.9 years. All patients were in their first episode and were drug-naïve when they received the neuroimaging scanning. Clinical symptoms were rated using the Positive and Negative Syndrome Scale (PANSS) (Kay, Fiszbein, & Opler, 1987). These patients had no history of other major psychiatric illness, neurological illness, head injury, alcohol or drug abuse.

Thirty TDCs were recruited from advertisements posted in the community and by word of mouth. They were interviewed with the Structured Clinical Interview for DSM-IV for nonpatient version (SCID-I/NP) and were selected to match members of the EOS group for gender and age. The age of the healthy controls ranged from 7.5 to 17.5 years. None of TDCs had any significant prior medical diagnosis, substance abuse, neurological disorders, or major psychiatric disorders. Those who had a first-degree relative with a history of severe mental disorder or suicidal behavior were also excluded.

Written informed consent was obtained from each participant and consent from each participant's guardian was also obtained prior to data acquisition. This study was approved

by the Ethical Committee for Medicine of First Hospital of Shanxi Medical University, China.

MRI Acquisition

MRI scanning was performed on the 3.0T Siemens Trio™ Scanner (Erlangen, Germany) at the First Hospital of Shanxi Medical University. For each participant, both an eight minute and 52 second RFMRI scan with echo planar imaging sequence (EPI: TR = 2,500 ms, TE = 30 ms, flip angle = 90°, matrix = 64 × 64, FOV = 240 mm, 4 mm slice thickness, 32 axial slices, 212 acquisitions) and a T1-weighted structural MRI scan with magnetization prepared rapidly acquired gradient echo sequence (MPRAGE: TR = 2300 ms, TE = 2.95 ms, TI = 900 ms, flip angle = 9°, matrix = 240 × 256, FOV = 225 × 240 mm, slice thickness = 1.2 mm, 160 slices) were acquired. For the resting state scan, participants were instructed to close their eyes and remain awake while lying quietly.

Data Analysis

Image preprocessing was conducted with the Connectome Computation System (CCS: <http://lfc.d.psych.ac.cn/ccs.html>), including both structural and functional image preprocessing (Zuo et al., 2013). Briefly, the functional scans were slice-timing corrected, motion corrected, normalized with a 4D mean of 10,000, filtered with a temporal band-pass filter (0.01–0.1Hz) and finally registered with individual anatomical images and further with the standard symmetric template. Data quality control procedure (QCP) was conducted to ensure usable data for subsequent analyses. For each participant, VMHC was computed as the resting-state functional connectivity (RSFC) between every pair of symmetric inter-hemispheric voxels (Zuo et al., 2010). Online supplementary Appendix S1 provides further details about the data analysis, such as the VMHC computation and generation of the standard symmetric brain template.

Group-level analyses were conducted using FSL's ordinary least squares (OLS) model implemented in FLAME. Voxel-wise one-way ANCOVA tests (covariates: age, sex, the error of nonlinear registration, minimal cost of co-registration, root mean square of frame-wise displacement, global VMHC (gVMHC), voxel-mirrored homotopic morphometry (VMHM) of the gray matter density) were performed to examine the differences in regional VMHC between the two groups. This statistical procedure produced thresholded Z-statistic maps of clusters defined by a threshold of $Z = 2.3$ and a cluster-level threshold based on Gaussian Random Field theory with a cluster-level family-wise error (FWE) correction for multiple comparisons (corrected $p < 0.05$). Of note, the gVMHC was also evaluated with a similar ANCOVA model (but without the covariates of gVMHC and VMHM) to test group differences.

Results

Demographic and Clinical Characteristics

Demographic and clinical information is listed in Table 1. Two EOS patients did not complete the scanning, and 4 other patients' data did not pass the imaging data QCP. Data

from 5 TDCs was excluded for failure to pass the QCP. The final sample included 26 EOS patients and 25 TDCs.

Functional Homotopy in TDCs and EOS

The spatial patterns of VMHC across the whole brain are shown in Figure 1 for TDCs (Figure 1A) and EOS (Figure 1B). Both groups exhibited the greatest VMHC in the precuneus and posterior cingulate cortex (PCU/PCC), followed by the insula, precentral gyrus (PRG), postcentral gyrus (PCG), superior temporal cortex (STC), and middle occipital gyrus, as well as subcortical areas. The lateral prefrontal cortex showed relatively weaker VMHC in both groups. On visual inspection, although the two groups showed similar patterns in VMHC, TDCs seemed to have stronger VMHC in PCU, PCG, STC, and occipital cortex.

Group Differences in Functional Homotopy

In order to assess structural confounders to the VMHC, we investigated the grey matter density and cortical thickness differences between EOS patients and TDCs. No significant differences were detectable in these two metrics between the two groups. Moreover, the data preprocessing was performed with two approaches, with scrubbing and without scrubbing. The results were reproducible across these two approaches, and here we only present the results without scrubbing in following sections.

EOS patients showed significantly reduced gVMHC in comparison with TDCs, indicating a deficit of whole-brain interhemispheric synchronization in EOS patients (Table 1). As seen in Figure 2A, EOS patients exhibited significantly lower VMHC in a cluster overlapping the STC/PCG. We also compared VMHC in the 24 EOS patients who had PANSS scores and the TDCs, and observed a similar decrease of VMHC in EOS patients (Figure 2B).

To evaluate the effects of disrupted interhemispheric connectivity on the relevant resting state networks, we used clusters of left and right STC/PCG as two regions of interest (including all voxels in the clusters of the left or right STC/PCG) to investigate the patterns of intrinsic functional connectivity (iFC) with the whole brain in both EOS and TDC groups, as well as group differences in iFC. The clusters of the left and right STC/PCG were defined as the cluster showing the VMHC group differences between EOS patients and TDCs and its left-right flipped version. We found that in TDC participants, the left STC/PCG showed positive iFC with the precentral and postcentral gyri, supplemental motor area, superior and middle temporal gyri, middle frontal gyrus, anterior cingulate cortex and anterior cerebellum. The left STC/PCG was negatively connected with the frontal pole, PCC, PCU, lateral occipital cortex, superior frontal gyrus, and medial frontal gyrus (Figure 3A: the left column). The right STC/PCG exhibited similar iFC patterns (Figure 3A: the right column). The iFC patterns seeded in the bilateral STC/PCG in the EOS group (Figure 3B) were highly similar but weaker in strength than those for TDC group. The group comparisons of left STC/PCG iFC analysis revealed that the EOS patients showed iFC reductions in the bilateral PCG, PRG, supplemental motor area, STC and insula, as well as increased iFC in the bilateral caudate, medial prefrontal gyrus, thalamus, PCC and PCU when compared with

TDCs (Figure 4A). The right STC/PCG iFC analysis showed highly similar findings with the left STC/PCG iFC results (Figure 4B).

Correlations with Clinical Scores in early onset schizophrenia patients

Mean VMHC of the STC/PCG was negatively correlated with negative symptoms ($r = -.43$, $p = 0.037$) (figure 2C) and marginally correlated with total PANSS score ($r = -.36$, $p = 0.085$). No significant correlations were found between the gVMHC and PANSS total and subscale scores or the mean VMHC within the STC/PCG and other subscales scores. To gain insights into the item-specific contribution to these correlations, we further examined the correlation between the VMHC and each item score in PANSS and detailed the findings in Appendix S1. Correlational analyses of iFC findings and the PANSS items can also be found in Appendix S1.

Reproducibility

To examine the reproducibility of our findings reported, we performed both the split-half and leave-one-out validations. First, 13 EOS patients and 13 TDCs were randomly selected from the sample and entered into the same statistical comparisons. The findings were depicted as in Figure 5A and fundamentally replicated the findings from the full sample. Notably, the sample size was relatively small and the split-half analysis may significantly reduce the statistical power of the group comparisons. This limitation is reflected by the following seed-based iFC analysis in the other half of the sample. The overall previous findings of iFC (Figure 4) were largely limited to the spatial extent of the half sample (Figure 6).

To validate the reproducibility and robustness of these findings on VMHC without losing statistical power, we performed a leave-one-out validation. Specifically, we left one EOS patient out of the sample and performed the same group comparisons based upon the permuted sample (i.e., 25 EOS versus 25 TDC). This analysis led to total 26 two-sample t-test images, based on which, for each voxel, we calculated number of tests where this voxel exhibited significant group differences across the total 26 tests as the reproducibility of the VMHC differences between EOS and TDC. Figure 5B indicated highly reproducible patterns of VMHC reductions across these tests as well as the original test reported.

Discussion

The current observation that drug-naïve EOS patients show substantial deficits in interhemispheric coordination provides neurodevelopmental evidence for the disconnection hypothesis of schizophrenia. Thus, alterations in functional homotopy in EOS might occur early on in the course of the illness and be independent of medication status.

EOS patients exhibited reduction of global whole-brain interhemispheric low frequency synchronization in comparison with TDCs. The present findings are consistent with recent studies reporting that COS patients exhibited a disrupted functional organization in their brain connectomes. Using a graph theory approach, Alexander-Bloch and colleagues observed reduced modularity and local organization in brain functional networks in chronic and medicated COS patients (Alexander-Bloch et al., 2010, 2012). Guo et al. (2013)

investigated the brain-wide functional inter-hemispheric connections in adult schizophrenia, and found reduced brain-wide inter-hemispheric functional networks. Our work employed drug-naïve first-onset EOS patients and provides strong evidence that these disruptions in the global metrics of whole brain functional connectomes might be an indication of the reduction of global hemispheric communication occurring in patients during development.

The regionally specific reduction of VMHC in the EOS group was mainly detected in the STC and extended to the PCG. With the same approach, Guo et al. (2014c) found first-episode, drug-naïve paranoid schizophrenia patients showed decreased VMHC in STC. The STC is considered to be responsible for auditory and language processing (Bigler et al., 2007; Menon et al., 1995). These results are consistent with well-established deficits shown by patients with schizophrenia in auditory processing (Javitt, 2009). Previous studies have reported that this region has abnormal features of both gray matter and white matter in COS and EOS patients (Tang et al., 2012; Taylor et al., 2005). Meanwhile, task-based fMRI studies have reported that adult schizophrenia patients show decreases in STC activation during language processing (Kircher, Leube, Erb, Grodd, & Rapp, 2007; Simons et al., 2010), the auditory oddball task (Kiehl & Liddle, 2001) and verbal working memory (Pae et al., 2008) compared to healthy controls. Additionally, EOS patients showed decreased activation in the STC during encoding stage of working memory (Haenschel et al., 2007). Another study found that the encoding-related bilateral anterior cingulate and temporal lobe network were disrupted in EOS patients during a working memory task (White et al., 2011). Moreover, adult schizophrenia patients also showed reduced inter-hemispheric EEG coherence in the temporal lobe (Winterer et al., 2001). The PCG is considered to be a key structure for receiving and processing somatosensory information (Nelson & Chen, 2008; Taylor & Jones, 1997). The decreased VMHC in this region suggests that EOS patients may have deficits in primary somatosensory processing.

The seed-based analysis revealed abnormal non-homotopic connectivity in EOS patients. The strength of functional connectivity of the STC/PCG in the EOS group was weaker than in the TDC group for both negative and positive iFC. More interestingly, EOS patients showed similar abnormal iFC patterns for left and right seeds of the STC/PCG, suggesting that the lower VMHC in EOS patients contributes to the dysfunctions in both hemispheres, as has been found in adult patients (Hoptman et al., 2012). In comparison to TDCs, EOS patients showed decreased iFC in bilateral PCG and STC, these findings were consistent with the VMHC group comparisons. EOS patients also showed decreased iFC in supplemental motor area and insula in comparison with TDCs. Insula is considered to play an important role in various aspects of emotional and cognitive processing (Augustine, 1996), while supplemental motor area is involved in internal movement generation and movement sequence construction (Goldberg, 1995; Tanji, 1994). These results suggested that EOS patients might also show abnormalities in emotion, cognition, and movement processing. Increased iFC was found in medial prefrontal gyrus, PCC, and PCU in EOS patients, these regions located in the default mode network. The present results were consistent with previous findings in adult schizophrenia (Garrity et al., 2007), possibly representing the boundary between imagination and reality (Buckner, Andrews-Hanna, & Schacter, 2008) is disrupted in early stage of the schizophrenia. The thalamus plays an important role in relaying and receiving the sensory and motor signals (Crick, 1984), while

caudate nucleus is involved in the learning and memory (Grahn, Parkinson, Owen, 2008). Increased iFC between the STC/PCG and regions responsible for sensory and goal-directed processing might represent decreased functional segregation in EOS patients (Fair et al., 2009). Moreover, non-homotopic functional connectivity disruptions have also been found in adult schizophrenia (Fornito, Zalesky, Pantelis, & Bullmore, 2012; Schmitt, Hasan, Gruber, & Falkai, 2011; Stephan, Baldeweg, & Friston, 2006) and COS patients (Alexander-Bloch et al., 2010, 2012, 2013). Disrupted homotopic as well as non-homotopic connectivity in EOS patients suggested that the disconnectivity might be the more general symptom of schizophrenia.

To the best of our knowledge, the present work represents the first study to investigate functional connectivity with unmedicated EOS patients. Consistent with our previous work on adult chronic schizophrenia patients (Hoptman et al., 2012), only decreased functional homotopy was detected in EOS patients, although the regions showing decreased connectivity were different between the two studies, suggesting medication and/or age effects. Guo et al. (2014a) revealed that healthy controls had more long-range interhemispheric connections than patients with schizophrenia (62% vs. 52%). However, they found patients with schizophrenia showed increased mean connectivity strength for both short and long-range interhemispheric links. This study used the automated anatomical labeling (AAL) atlas to parcellate the brain into 90 regions of interest (45 for each hemisphere), whereas VMHC approach was adopted in our study. Moreover, in the study of Guo et al. (2014a), adult patients with schizophrenia were recruited, and three quarters of the patients were receiving antipsychotic medications. The potential effects of antipsychotic drugs on brain structure in psychosis are receiving increasing attention. Navari and Dazzan (2009) systematically reviewed 33 studies investigating the association of antipsychotic medication and brain structure changes and concluded that the influences of antipsychotic treatment on the brain changes in psychosis are mostly regional and not global. In particular, antipsychotic treatment has been associated with reduced grey matter volume in the frontal and temporal lobes (Smieskova et al., 2009). Even short-term (mean 8.5 weeks) antipsychotic treatment might cause grey matter reduction in frontal and temporal-insular areas and the precuneus (Dazzan et al., 2005), and long-term antipsychotic treatment with different doses or types of antipsychotics presents further challenges to the interpretation of brain changes. Importantly, antipsychotic medications have also been found to affect brain activity or functional connectivity in schizophrenia. Researchers explicitly tested the effect of antipsychotics on brain function in drug-naïve adult schizophrenia before and after 6 weeks of treatment and observed increased regional amplitudes of intrinsic brain activity of the bilateral prefrontal, parietal, left superior temporal cortex and right caudate nucleus, as well as widespread decreased functional connectivity in schizophrenia patients (Lui et al., 2010). A recent study employing healthy volunteers also provided evidence that antipsychotic medications might attenuate the functional connectivity patterns (Cole et al., 2013). The results we report here in drug-naïve EOS patients exclude the effects of antipsychotics and suggest abnormal interhemispheric functional coordination in schizophrenia, even in unmedicated patients.

Several possible mechanisms may underlie the reductions of VMHC in EOS patients. Altered brain structures might disrupt the functional synchrony between homotopic regions.

The splenium of the corpus callosum connects left and right parietal and temporal areas (Hofer & Frahm, 2006; Knyazeva, 2013). A recent study reported that first-episode drug-naïve adult schizophrenia patients showed reduced FA in the splenium, but not the genu, of the corpus callosum (Gasparotti et al., 2009). Because the fiber tracts in the splenium of the corpus callosum interconnect temporal cortical areas, an altered corpus callosum structure may disrupt the neural signal transmitted from one hemisphere to another, leading to failure of functional integration or disrupted VMHC in EOS patients. A recent study found that long-range connections increased, whereas local connections decreased in adolescence (Dosenbach et al., 2010). We found decreased VMHC in EOS patients in a specific type of long-range connection, interhemispheric connectivity, implying that these decreases might reflect disruptions in functional integration during development. The disruptions of functional integration are associated with myelination, axon terminal arborization, and synapse formation (Fair et al., 2007, 2009). Another possible factor underlying abnormal functional homotopy is asymmetry of hemispheric growth in EOS patients. The two hemispheres may show subtle differences or delays in the trajectory of growth in schizophrenia (Bleich-Cohen, Hendler, Kotler, & Strous, 2009; Crow, 1998), as was found in the present seed-based analyses.

Specifically, previous studies have demonstrated disrupted functional connectivity in siblings of schizophrenia. Patients with schizophrenia and their unaffected siblings were found to show impaired brain connectivity between brain networks as well as within networks (Repovs et al., 2011). Chang et al. (2014) reported that patients with schizophrenia and their siblings shared disconnection within the anterior DMN, as well as in the left and right fronto-parietal network subsystems. Increased functional connectivity between the bilateral inferior temporal gyri was also found in patients with schizophrenia and their unaffected siblings (Liu et al., 2012). With the same VMHC approach, Guo et al. (2014b) found siblings of schizophrenia patients (mean age=22.96 years old) showed lower VMHC in angular gyrus, lingual gyrus/cerebellum lobule VI than healthy controls. These results found in unaffected siblings suggest that the disconnection may precede the psychosis and represent a vulnerability towards the illness. Most previous studies focused on the adult schizophrenia and their adult siblings, and few studies investigated the functional connectivity in EOS patients and their adolescent siblings. Future studies about EOS patients and their adolescent siblings would provide further neurodevelopmental evidence for the disconnection theory of schizophrenia.

The findings further extend the association between clinical symptoms and functional connectivity in EOS patients. The mean VMHC of the STC/PCG was correlated negatively with negative symptoms in EOS patients, indicating a relationship between severity of the illness and reduced VMHC. Previous studies have consistently found that reduced volume in the STC is significantly correlated with clinical symptoms (Lui et al., 2009; Menon et al., 1995; Tang et al., 2012). Functional neuroimaging studies have also found significant correlations between STC activation and symptomatology in schizophrenia patients during language processing (Kircher et al., 2001; Plaze et al., 2006). Guo et al. (2014c) found similar results in adult schizophrenia; they showed significant negative correlations between VMHC in STG and the PANSS negative and total scores. Interestingly, our recent study also observed a negative correlation between PANSS total score and VMHC in the PCG in adult

chronic schizophrenia (Hoptman et al., 2012). The patients in the present study were younger and drug-naïve and thus extend our previous findings to include medication-free and young patients. This has important clinical significance and may suggest a role for VMHC in evaluating and predicting the severity of psychopathology.

There are several limitations to the present study. First, the sample size was relatively small, and a larger sample would increase the power of the study and the generalizability of the findings. Second, the present study employed a cross-sectional design and is thus not able to separate intra- and inter-subject variability. A follow-up design could provide longitudinal data to clarify the abnormal developmental changes of functional homotopy in EOS. On the other hand, the EOS patients had to be treated for ethical reasons, and thus, we cannot exclude the effects of medication in a follow-up study. Third, although we tried to match the age and sex between EOS and TDCs, the age range was still a little different. Two of the TDCs were under 10 years old (7.5 and 8 years old, respectively), while only one EOS was under 10 years old (9 years old). During group analysis, we controlled the age to reduce the influence of the age range differences. Moreover, because the EOS patients were first-episode and drug-naïve, they might have difficulties performing neuropsychological tasks. We did not collect neuropsychological data, especially tasks that required interhemispheric transfer of information. Further exploration of these tasks could clarify whether decreased functional homotopy is reflected in abnormal behaviors. Socioeconomic status and IQ were also not collected, and thus we could not explore the associations between these factors and dysfunctional homotopy in EOS. Finally, we employed a voxel-wise method rather than a region-wise approach. Region-based approaches show advantages for exploring brain structure or anatomy in both the clinical and normal populations. These advantages partially originate from the fact that well-established knowledge of brain structure and anatomy has been used to create brain parcellation templates used in the ROI approach. However, it is not an obvious advantage for functional analyses because a well-established functional parcellation of the human brain is currently lacking. Rich differences in local functional homogeneity exist (Zang et al., 2004; Zuo et al., 2013). This fact makes it problematic to extract the regional mean time series by applying these structural parcellation templates (Jiang et al., 2014). Notably, due to these variabilities, large-scale regional approaches for functional connectomes only exhibited moderate test-retest reliability (Wang et al., 2011). To address above challenges, the voxel-wise method proposed in Zuo et al. (2010), takes the advantage of the original high-resolution of the fMRI data and defines the functional connectivity between the ‘homotopic voxels’, which are one voxel and its left-right geometrically symmetric voxel. VMHC shows high test-retest reliability (Zuo et al., 2010). The region-wise assessment on the homotopic changes will be an interesting topic once a functional parcellation of the human brain is well established in future.

In summary, the present study found that EOS patients showed substantial deficits of interhemispheric coordination, providing neurodevelopmental evidence for the disconnection theory of schizophrenia. Moreover, decreased VMHC in the STC/PCG correlated with clinical symptoms. These findings suggest that functional homotopy in neurodevelopmental processes may play an important role in the functional changes underlying schizophrenia.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the Key Technologies R&D Program (2012BAI36B01, XNZ), Natural Science Foundation of China (81278412, 81000583, 801411204369, 81171409), the Special Initiating Funds for Winners of Excellent Doctorate Degree Paper and Presidential Prizes of Institute of Psychology, Chinese Academy of Sciences (Y0CX262B01, HJL) and the Hundred Talents Program and the Key Research Program (KSZD-EW-TZ-002, XNZ) of Chinese Academy of Sciences, Program for New Century Excellent Talents in University (YX). Author M.J.H acknowledges support from the NIMH (R01 MH064783, R21 MH084031). The authors acknowledge Xiao-Dong Hu, Jun-Yan Shi, and Lei Cao for data collection.

References

- Alexander-Bloch AF, Gogtay N, Meunier D, Birn R, Clasen L, Lalonde F, Lenroot R, Giedd J, Bullmore ET. Disrupted modularity and local connectivity of brain functional networks in childhood-onset schizophrenia. *Frontiers in Systems Neuroscience*. 2010; 4:147. [PubMed: 21031030]
- Alexander-Bloch AF, Lambiotte R, Roberts B, Giedd J, Gogtay N, Bullmore E. The discovery of population differences in network community structure: new methods and applications to brain functional networks in schizophrenia. *NeuroImage*. 2012; 59:3889–3900. [PubMed: 22119652]
- Alexander-Bloch AF, Vertes PE, Stidd R, Lalonde F, Clasen L, Rapoport J, Giedd J, Bullmore ET, Gogtay N. The anatomical distance of functional connections predicts brain network topology in health and schizophrenia. *Cerebral Cortex*. 2013; 23:127–138. [PubMed: 22275481]
- Augustine JR. Circuitry and functional aspects of the insularlobe in primates including humans. *Brain Research Reviews*. 1996; 22:229–244. [PubMed: 8957561]
- Barnett KJ, Kirk IJ, Corballis MC. Bilateral disadvantage: lack of interhemispheric cooperation in schizophrenia. *Consciousness and Cognition*. 2007; 16:436–444. [PubMed: 16877008]
- Bigler ED, Mortensen S, Neeley ES, Ozonoff S, Krasny L, Johnson M, Lu J, Provencal SL, McMahon W, Lainhart JE. Superior temporal gyrus, language function, and autism. *Developmental Neuropsychology*. 2007; 31:217–238. [PubMed: 17488217]
- Bleich-Cohen M, Hendler T, Kotler M, Strous RD. Reduced language lateralization in first-episode schizophrenia: an fMRI index of functional asymmetry. *Psychiatry Research*. 2009; 171:82–93. [PubMed: 19185468]
- Chang X, Shen H, Wang L, Liu Z, Xin W, Hu D, Miao D. Altered default mode and fronto-parietal network subsystems in patients with schizophrenia and their unaffected siblings. *Brain Research*. 2014; 1562:87–99. [PubMed: 24675026]
- Cole DM, Oei NY, Soeter RP, Both S, van Gerven JM, Rombouts SA, Beckmann CF. Dopamine-dependent architecture of cortico-subcortical network connectivity. *Cerebral Cortex*. 2013; 23:1509–1516. [PubMed: 22645252]
- Crick F. Function of the thalamic reticular complex: the searchlight hypothesis. *Proceedings of the National Academy of Sciences of the United States of America*. 1984; 81:4586–4590. [PubMed: 6589612]
- Crow TJ. Schizophrenia as a transcallosal misconnection syndrome. *Schizophrenia Research*. 1998; 30:111–114. [PubMed: 9549773]
- Dazzan P, Morgan KD, Orr K, Hutchinson G, Chitnis X, Suckling J, Fearon P, McGuire PK, Mallett RM, Jones PB. Different effects of typical and atypical antipsychotics on grey matter in first episode psychosis: the AESOP study. *Neuropsychopharmacology*. 2005; 30:765–774. [PubMed: 15702141]
- Dosenbach NUF, Nardos B, Cohen AL, Fair DA, Power JD, Church JA, Nelson SM, Wig GS, Vogel AC, Lessov-Schlaggar CN, Barnes KA, Dubis JW, Feczko E, Coalson RS, Pruett JR, Barch DM, Petersen SE, Schlaggar BL. Prediction of Individual Brain Maturity Using fMRI. *Science*. 2010; 329:1358–1361. [PubMed: 20829489]

- Fair DA, Cohen AL, Power JD, Dosenbach NU, Church JA, Miezin FM, Schlaggar BL, Petersen SE. Functional brain networks develop from a “local to distributed” organization. *PLOS Computational Biology*. 2009; 5:e1000381. [PubMed: 19412534]
- Fair DA, Dosenbach NU, Church JA, Cohen AL, Brahmbhatt S, Miezin FM, Barch DM, Raichle ME, Petersen SE, Schlaggar BL. Development of distinct control networks through segregation and integration. *Proceedings of the National Academy of Sciences of the United States of America*. 2007; 104:13507–13512. [PubMed: 17679691]
- Fornito A, Zalesky A, Pantelis C, Bullmore ET. Schizophrenia, neuroimaging and connectomics. *NeuroImage*. 2012; 62:2296–314. [PubMed: 22387165]
- Friston KJ, Frith CD. Schizophrenia: a disconnection syndrome? *Journal of Clinical Neuroscience*. 1995; 3:89–97.
- Frith CD, Friston KJ, Herold S, Silbersweig D, Fletcher P, Cahill C, Dolan RJ, Frackowiak RS, Liddle PF. Regional brain activity in chronic schizophrenic patients during the performance of a verbal fluency task. *The British Journal of Psychiatry*. 1995; 167:343–349. [PubMed: 7496643]
- Garrity AG, Pearlson GD, McKiernan K, Lloyd D, Kiehl KA, Calhoun VD. Aberrant “default mode” functional connectivity in schizophrenia. *The American Journal of Psychiatry*. 2007; 164:450–457. [PubMed: 17329470]
- Gasparotti R, Valsecchi P, Carletti F, Galluzzo A, Liserre R, Cesana B, Sacchetti E. Reduced fractional anisotropy of corpus callosum in first-contact, antipsychotic drug-naïve patients with schizophrenia. *Schizophrenia Research*. 2009; 108:41–48. [PubMed: 19103476]
- Goldberg G. Supplementary motor area structure and function: Rev and hypotheses. *Behavioral and Brain Sciences*. 1985; 8:567–616.
- Grahn JA, Parkinson JA, Owen AM. The cognitive functions of the caudate nucleus. *Progress in Neurobiology*. 2008; 86:141–155. [PubMed: 18824075]
- Guo S, Kendrick KM, Zhang J, Broome M, Yu R, Liu Z, Feng J. Brain-wide functional inter-hemispheric disconnection is a potential biomarker for schizophrenia and distinguishes it from depression. *Neuroimage Clinical*. 2013; 23:818–26. [PubMed: 24179833]
- Guo S, Palaniyappan L, Yang B, Liu Z, Xue Z, Feng J. Anatomical Distance Affects Functional Connectivity in Patients With Schizophrenia and Their Siblings. *Schizophrenia Bulletin*. 2014a; 40:449–459. [PubMed: 24282323]
- Guo W, Jiang J, Xiao C, Zhang Z, Zhang J, Yu L, Liu J, Liu G. Decreased resting-state interhemispheric functional connectivity in unaffected siblings of schizophrenia patients. *Schizophrenia Research*. 2014b; 152:170–175. [PubMed: 24325975]
- Guo W, Xiao C, Liu G, Wooderson SC, Zhang Z, Zhang J, Yu L, Liu J. Decreased resting-state interhemispheric coordination in first-episode, drug-naïve paranoid schizophrenia. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*. 2014c; 48:14–19. [PubMed: 24075897]
- Haenschel C, Bittner RA, Haertling F, Rotarska-Jagiela A, Maurer K, Singer W, Linden DE. Contribution of impaired early-stage visual processing to working memory dysfunction in adolescents with schizophrenia: a study with event-related potentials and functional magnetic resonance imaging. *Archives of General Psychiatry*. 2007; 64:1229–1240. [PubMed: 17984392]
- Hofer S, Frahm J. Topography of the human corpus callosum revisited? Comprehensive fiber tractography using diffusion tensor magnetic resonance imaging. *NeuroImage*. 2006; 32:989–994. [PubMed: 16854598]
- Honey GD, Bullmore ET, Soni W, Varatheesan M, Williams SC, Sharma T. Differences in frontal cortical activation by a working memory task after substitution of risperidone for typical antipsychotic drugs in patients with schizophrenia. *Proceedings of the National Academy of Sciences of the United States of America*. 1999; 96:13432–13437. [PubMed: 10557338]
- Hoptman MJ, Davidson RJ. How and why do the two cerebral hemispheres interact? *Psychological Bulletin*. 1994; 116:195–219. [PubMed: 7972590]
- Hoptman MJ, Zuo XN, D’Angelo D, Mauro CJ, Butler PD, Milham MP, Javitt DC. Decreased interhemispheric coordination in schizophrenia: a resting state fMRI study. *Schizophrenia Research*. 2012; 141:1–7. [PubMed: 22910401]
- Javitt DC. Sensory processing in schizophrenia: neither simple nor intact. *Schizophrenia Bulletin*. 2009; 35:1059–1064. [PubMed: 19833806]

- Jiang L, Xu T, He Y, Hou XH, Wang J, Cao XY, Wei GX, Yang Z, He Y, Zuo XN. Toward neurobiological characterization of functional homogeneity in the human cortex: Regional variation, morphological association and functional covariance network organization. *Brain Structure and Function*. 2014 in press. 10.1007/s00429-014-0795-8
- Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia Bulletin*. 1987; 13:261–276. [PubMed: 3616518]
- Ke M, Shen H, Fan J, Huang X, Zhou Z, Chen X, Hu D. Combining resting-state fMRI and DTI analysis for early-onset schizophrenia. *International Journal of Intelligent Systems*. 2009; 2:375–385.
- Kiehl KA, Liddle PF. An event-related functional magnetic resonance imaging study of an auditory oddball task in schizophrenia. *Schizophrenia Research*. 2001; 48:159–171. [PubMed: 11295369]
- Kircher TT, Leube DT, Erb M, Grodd W, Rapp AM. Neural correlates of metaphor processing in schizophrenia. *NeuroImage*. 2007; 34:281–289. [PubMed: 17081771]
- Kircher TT, Liddle PF, Brammer MJ, Williams SC, Murray RM, McGuire PK. Neural correlates of formal thought disorder in schizophrenia: preliminary findings from a functional magnetic resonance imaging study. *Archives of General Psychiatry*. 2001; 58:769–774. [PubMed: 11483143]
- Knyazeva MG. Splenium of corpus callosum: patterns of interhemispheric interaction in children and adults. *Neural Plasticity*. 2013; 2013:639430. [PubMed: 23577273]
- Liu H, Kaneko Y, Ouyang X, Li L, Hao Y, Chen EYH, Jiang T, Zhou Y, Liu Z. Schizophrenic Patients and Their Unaffected Siblings Share Increased Resting-State Connectivity in the Task-Negative Network but Not Its Anticorrelated Task-Positive Network. *Schizophrenia Bulletin*. 2012; 38:285–294. [PubMed: 20595202]
- Lopez AD, Murray CC. The global burden of disease, 1990–2020. *Nature Medicine*. 1998; 4:1241–1243.
- Lui S, Deng W, Huang X, Jiang L, Ma X, Chen H, Zhang T, Li X, Li D, Zou L, Tang H, Zhou XJ, Mechelli A, Collier DA, Sweeney JA, Li T, Gong Q. Association of cerebral deficits with clinical symptoms in antipsychotic-naïve first-episode schizophrenia: an optimized voxel-based morphometry and resting state functional connectivity study. *The American journal of psychiatry*. 2009; 166:196–205. [PubMed: 18981063]
- Lui S, Li T, Deng W, Jiang LJ, Wu QZ, Tang HH, Yue QA, Huang XQ, Chan RC, Collier DA, Meda SA, Pearlson G, Mechelli A, Sweeney JA, Gong QY. Short-term Effects of Antipsychotic Treatment on Cerebral Function in Drug-Naïve First-Episode Schizophrenia Revealed by “Resting State” Functional Magnetic Resonance Imaging. *Archives of General Psychiatry*. 2010; 67:783–792. [PubMed: 20679586]
- Menon RR, Barta PE, Aylward EH, Richards SS, Vaughn DD, Tien AY, Harris GJ, Pearlson GD. Posterior superior temporal gyrus in schizophrenia: grey matter changes and clinical correlates. *Schizophrenia Research*. 1995; 16:127–135. [PubMed: 7577766]
- Mohr B, Pulvermuller F, Cohen R, Rockstroh B. Interhemispheric cooperation during word processing: evidence for callosal transfer dysfunction in schizophrenic patients. *Schizophrenia Research*. 2000; 46:231–239. [PubMed: 11120435]
- Mohr B, Pulvermuller F, Rockstroh B, Endrass T. Hemispheric cooperation--a crucial factor in schizophrenia? Neurophysiological evidence. *NeuroImage*. 2008; 41:1102–1110. [PubMed: 18456515]
- Morrison-Stewart SL, Velikonja D, Corning WC, Williamson P. Aberrant interhemispheric alpha coherence on electroencephalography in schizophrenic patients during activation tasks. *Psychological Medicine*. 1996; 26:605–612. [PubMed: 8733218]
- Navari S, Dazzan P. Do antipsychotic drugs affect brain structure? A systematic and critical review of MRI findings. *Psychological Medicine*. 2009; 39:1763–1777. [PubMed: 19338710]
- Nelson AJ, Chen R. Digit somatotopy within cortical areas of the postcentral gyrus in humans. *Cerebral Cortex*. 2008; 18:2341–2351. [PubMed: 18245039]
- Owen MJ, O'Donovan MC, Thapar A, Craddock N. Neurodevelopmental hypothesis of schizophrenia. *The British Journal of Psychiatry*. 2011; 198:173–175. [PubMed: 21357874]

- Pae CU, Juh R, Yoo SS, Choi BG, Lim HK, Lee C, Paik IH, Jeun SS, Lee CU. Verbal working memory dysfunction in schizophrenia: an fMRI investigation. *International Journal of Neuroscience*. 2008; 118:1467–87. [PubMed: 18788030]
- Pettersson-Yeo W, Allen P, Benetti S, McGuire P, Mechelli A. Dysconnectivity in schizophrenia: Where are we now? *Neuroscience & Biobehavioral Reviews*. 2011; 35:1110–1124. [PubMed: 21115039]
- Plaze M, Bartres-Faz D, Martinot JL, Januel D, Bellivier F, De Beaurepaire R, Chanraud S, Andoh J, Lefaucheur JP, Artiges E, Pallier C, Paillere-Martinot ML. Left superior temporal gyrus activation during sentence perception negatively correlates with auditory hallucination severity in schizophrenia patients. *Schizophrenia Research*. 2006; 87:109–115. [PubMed: 16828542]
- Rapoport JL, Giedd JN, Gogtay N. Neurodevelopmental model of schizophrenia: update 2012. *Molecular Psychiatry*. 2012; 17:1228–38. [PubMed: 22488257]
- Repovs G, Csernansky JG, Barch DM. Brain Network Connectivity in Individuals with Schizophrenia and Their Siblings. *Biological Psychiatry*. 2011; 69:967–973. [PubMed: 21193174]
- Rhinewine JP, Lencz T, Thaden EP, Cervellione KL, Burdick KE, Henderson I, Bhaskar S, Keehlisen L, Kane J, Kohn N, Fisch GS, Bilder RM, Kumra S. Neurocognitive profile in adolescents with early-onset schizophrenia: clinical correlates. *Biological Psychiatry*. 2005; 58:705–712. [PubMed: 16023084]
- Schmitt A, Hasan A, Gruber O, Falkai P. Schizophrenia as a disorder of disconnectivity. *European Archives of Psychiatry and Clinical Neurosciences*. 2011; 261:150–154.
- Simons CJ, Tracy DK, Sanghera KK, O'Daly O, Gillean J, Dominguez MD, Krabbendam L, Shergill SS. Functional magnetic resonance imaging of inner speech in schizophrenia. *Biological Psychiatry*. 2010; 67:232–237. [PubMed: 19846064]
- Smieskova R, Fusar-Poli P, Allen P, Bendfeldt K, Stieglitz R, Drewe J, Radue E, McGuire P, Riecher-Rossler A, Borgwardt S. The effects of antipsychotics on the brain: What have we learnt from structural imaging of schizophrenia? A systematic review. *Current Pharmaceutical Design*. 2009; 15:2535–2549. [PubMed: 19689326]
- Stark DE, Margulies DS, Shehzad ZE, Reiss P, Kelly AM, Uddin LQ, Gee DG, Roy AK, Banich MT, Castellanos FX, Milham MP. Regional variation in interhemispheric coordination of intrinsic hemodynamic fluctuations. *The Journal of Neuroscience*. 2008; 28:13754–13764. [PubMed: 19091966]
- Stephan KE, Baldeweg T, Friston KJ. Synaptic plasticity and dysconnection in schizophrenia. *Biological Psychiatry*. 2006; 59:929–939. [PubMed: 16427028]
- Tang JS, Liao YH, Zhou B, Tan CL, Liu WQ, Wang D, Liu TQ, Hao W, Tan LW, Chen XG. Decrease in Temporal Gyrus Gray Matter Volume in First-Episode, Early Onset Schizophrenia: An MRI Study. *PloS one*. 2012; 7:e40247. [PubMed: 22802957]
- Tanji J. The supplementary motor area in the cerebral cortex. *Neuroscience Research*. 1994; 19:251–268. [PubMed: 8058203]
- Taylor JL, Blanton RE, Levitt JG, Caplan R, Nobel D, Toga AW. Superior temporal gyrus differences in childhood-onset schizophrenia. *Schizophrenia Research*. 2005; 73:235–241. [PubMed: 15653266]
- Taylor L, Jones L. Effects of lesions invading the postcentral gyrus on somatosensory thresholds on the face. *Neuropsychologia*. 1997; 35:953–961. [PubMed: 9226657]
- Vourdas A, Pipe R, Corrigall R, Frangou S. Increased developmental deviance and premorbid dysfunction in early onset schizophrenia. *Schizophrenia Research*. 2003; 62:13–22. [PubMed: 12765738]
- Wang J, Zuo XN, Gohel S, Milham MP, Biswal BB, He Y. Graph Theoretical Analysis of Functional Brain Networks: Test-Retest Evaluation on Short- and Long-Term Resting-State Functional MRI Data. *PLoS ONE*. 2011; 6:e21976. [PubMed: 21818285]
- Weinberger DR. Implications of normal brain development for the pathogenesis of schizophrenia. *Archives of General Psychiatry*. 1987; 44:660. [PubMed: 3606332]
- White T, Schmidt M, Kim DI, Calhoun VD. Disrupted functional brain connectivity during verbal working memory in children and adolescents with schizophrenia. *Cerebral Cortex*. 2011; 21:510–518. [PubMed: 20670970]

- Winterer G, Egan MF, Radler T, Hyde T, Coppola R, Weinberger DR. An association between reduced interhemispheric EEG coherence in the temporal lobe and genetic risk for schizophrenia. *Schizophrenia Research*. 2001; 49:129–143. [PubMed: 11343872]
- Zang Y, Jiang T, Lu Y, He Y, Tian L. Regional homogeneity approach to fMRI data analysis. *Neuroimage*. 2004; 22:394–400. [PubMed: 15110032]
- Zhou B, Tan C, Tang J, Chen X. Brain functional connectivity of functional magnetic resonance imaging of patients with early-onset schizophrenia. *Zhong Nan Da Xue Xue Bao Yi Xue Ban*. 2010; 35:17–24. [PubMed: 20130360]
- Zuo XN, Kelly C, Di Martino A, Mennes M, Margulies DS, Bangaru S, Grzadzinski R, Evans AC, Zang YF, Castellanos FX, Milham MP. Growing together and growing apart: regional and sex differences in the lifespan developmental trajectories of functional homotopy. *The Journal of Neuroscience*. 2010; 30:15034–15043. [PubMed: 21068309]
- Zuo XN, Xu T, Jiang L, Yang Z, Cao XY, He Y, Zang YF, Castellanos FX, Milham MP. Toward reliable characterization of functional homogeneity in the human brain: preprocessing, scan duration, imaging resolution and computational space. *NeuroImage*. 2013; 65:374–386. [PubMed: 23085497]

Key points

- Few studies investigated the disconnection hypothesis in early-onset schizophrenia.
- This study explored the homotopic connectivity in first-episode drug-naïve early-onset schizophrenia.
- Early-onset schizophrenia patients exhibited global and regional homotopic connectivity reductions. Lower homotopic connectivity values were observed with in superior temporal cortex and postcentral gyrus.
- The lower interhemispheric synchronization was correlated with higher levels of negative symptoms.
- The findings provide novel neurodevelopmental evidence for the disconnection hypothesis of schizophrenia.

Figure 1. Whole brain voxel-wise homotopic functional connectivity patterns in early-onset schizophrenia (EOS) patients (A) and typically developing controls (TDCs) (B).

Figure 2.

(A) Comparisons of whole brain VMHC between early-onset schizophrenia (EOS) and typically developing controls (TDCs). The cool colors indicate regions showing higher VMHC in TDCs than EOS patients. (B) Comparisons of whole brain VMHC in the 24 EOS patients who had PANSS scores and the TDCs. Each VMHC value here came from the cluster in A. (C) Correlation between the mean VMHC of the STC/PCG and negative symptoms in EOS patients.

Figure 3.

Intrinsic functional connectivity patterns for left and right STC/PCG seeds: (A) early-onset schizophrenia (EOS) patients; (B) typically developing controls (TDCs). The figure is presented in neurological convention (image left is the left side of the brain). The left and right panels corresponded to the left and right STC/PCG seeds, respectively.

Figure 4. Group differences in intrinsic functional connectivity between early-onset schizophrenia (EOS) and typically developing controls (TDCs): (A) left STC/PCG seed; (B) right STC/PCG seed.

Figure 5.

(A) Split-half sample validation. Comparisons of whole brain VMHC were performed between 13 early-onset schizophrenia (EOS) patients and 13 typically developing controls (TDCs). The cool colors indicate regions showing higher VMHC in TDCs than EOS patients. (B) Leave-one-out sample validation. The group comparisons based upon the permuted samples (i.e., 25 EOS versus 25 TDC) for total 26 times. For each voxel, the color indicates number of tests where this voxel exhibited significant group differences across the total 26 tests (i.e., the reproducibility).

Figure 6. Split-half validation derived group differences in intrinsic functional connectivity between 13 early-onset schizophrenia (EOS) patients and 12 typically developing controls (TDCs): (A) left STC/PCG seed; (B) right STC/PCG seed.

Table 1**Participant Information**

	EOS (N=26)	TDC (N=25)
Age	14.5 (1.94)	14.4 (2.97)
Sex (male/female)	13/13	13/12
gVMHC*	0.2165 (0.0447)	0.2418 (0.0401)
errFNIRT ¹	0.0873 (0.0164)	0.0824 (0.0109)
mcBBR ^{2**}	0.5111 (0.0436)	0.5443 (0.0356)
rmsFD ³ (mm)	0.1147 (0.0562)	0.1121 (0.0477)
Total PANSS score	57.75 (13.74)	-
Positive symptoms	12.04 (3.94)	-
Negative symptoms	13.17 (4.93)	-
General psychopathology	28.58 (6.89)	-
Supplementary items	3.96 (1.57)	-

The values in brackets are standard deviations. EOS: Early-Onset Schizophrenia; TDC: Typical Development Control. PANSS scores were available for 24 EOS patients;

*
p < 0.05 and

**
p < 0.01.

¹ errFNIRT is the registration error of the individual spatial normalization to the MNI152 standard space (2mm isotropic voxels) measured by 1 – the spatial correlation between individual normalized T1 images and the standard template;

² mcBBR is the minimal cost of the intra-subject co-registration with the boundary-based registration;

³ rmsFD is the root mean square of the frame-wise displacement for in-scanner head motion.