EDITORIAL

Who Needs Sleep Apnea Treatment for Safety Critical Tasks—Are We There Yet?

Commentary on Karimi et al. Sleep apnea related risk of motor vehicle accidents is reduced by continuous positive airway pressure: Swedish traffic accident registry data. *SLEEP* 2015;38:341–349.

Mark E. Howard, PhD^{1,2,3}; Melinda L. Jackson, PhD^{1,4}; Mark Stevenson, PhD^{3,5}

¹*Institute for Breathing and Sleep, Austin Health, Heidelberg, Australia;* ²*Department of Medicine, University of Melbourne, Parkville, Australia;*

³*Co-operative Research Centre for Alertness, Safety and Productivity, Clayton, Australia;* ⁴*School of Health Sciences, RMIT University, Melbourne, Australia;* ⁵*Monash University Accident Research Centre, Monash Injury Research Institute, Clayton, Australia*

Normalizing cognitive function is key to optimizing road and workplace safety and is one of the main reasons why clinicians identify and treat obstructive sleep apnea (OSA). Whilst numerous studies have found that OSA increases the risk of motor vehicle crashes, few have applied ideal research designs to conclusively validate the relationship,¹ and importantly, to demonstrate that treatment of OSA reduces the risk of motor vehicle crashes.² In this issue of *SLEEP*, Karimi and colleagues³ attempt to redress the limitations of past research with respect to OSA and motor vehicle crash risk, for the first time assessing the impact of CPAP treatment adherence. Karimi et al. highlight an elevated risk of crash associated with OSA, reflecting previous findings.¹ The authors extend the current literature by using an objective measure of crash (police reports) and assess crash risk using two perspectives, both retrospective and prospective, among a cohort of patients referred for suspected OSA compared to an external cohort.

There are several challenges in optimally designing crash risk studies for OSA. While objective crash statistics are preferable, the low rate of police reported crashes necessitates a large sample. Clinical equipoise to leave a range of OSA patients untreated to assess crash risk is challenging because of safety concerns for severely sleepy patients.⁴ Bias may occur when a clinical population is used in that some patients may present because they have had a crash. Using a prospective community cohort, the Wisconsin cohort study found an increased road crash risk in men with OSA, but not women.⁵ Case (road crash drivers) control (drivers who have not had a crash) studies are an alternative, with assessment of OSA after the crash, and support increased crash risk in those with OSA.⁶ The findings of Karimi et al.³ are consistent with these previous studies. Some limitations must be acknowledged, though, including the potential for presentation bias that might increase the risk estimate in a clinical sample. OSA, sleepiness, and other confounders were not measured in the control cohort; hence it could have included some drivers with OSA and excessive sleepiness.³ Nevertheless, the study by Karimi is one of only a few controlled investigations to assess and identify a reduction in crash risk with CPAP treatment.^{2,7,8,3}

and it appears to be the first to demonstrate the importance of CPAP adherence (use ≥ 4 h/night). As for OSA crash risk, the designs of these studies have not been ideal (observational cohorts rather than randomized trials) due to the same challenges. The adherent CPAP users could have behaved differently to the non-adherent group (e.g., less speeding), which could be at least partly responsible for their reduced crash risk.

Disease severity was not related to crash risk within the OSA group of Karimi et al.,³ suggesting that respiratory indices (e.g., apnea hypopnea index [AHI]) may not be the best way of determining OSA-related cognitive impairment.^{9,10} Previous work has been divided on whether AHI predicts crash risk.^{1,5} Patient sleepiness may also be highly variable despite comparable OSA severity. Moreover, individuals vary in their susceptibility to sleep loss.¹¹ Some consistently and rapidly develop sleepiness and cognitive impairment during restricted sleep; others are more resistant. Recently, this concept has been explored in OSA patients; some maintain normal driving performance following restricted sleep, while others develop substantial impairment.¹² Supporting this concept both subjective and objective measures of sleepiness hold only modest relationships with respiratory indices of OSA severity.¹³ This is not surprising given that sleepiness can be influenced by many factors, including age, medications, and genetic factors.¹⁴ Given the poor relationship between OSA severity and crash risk, and the apparent variable impact of OSA on cognitive function and sleepiness, there is a need for alternative methods to assess crash risk in OSA patients. Severe subjective sleepiness is related to crash risk^{15,16}; however, this can be unreliable when licensing is at risk,¹⁷ particularly in occupational settings. Sleep latency testing is recommended when driving safety is in question and relates to driving performance,¹⁸ although its relationship to crash risk appears weak.¹⁹ Robust studies of current and alternative objective markers of sleepiness or cognitive function are required to better determine those OSA patients who require treatment in order to undertake safety critical tasks and adequacy of treatment.

The concept that individuals may have different cognitive responses to OSA and sleep loss raises the possibility that those in high-risk occupations might self-select to be more resistant to conditions that cause sleepiness.²⁰ Commercial vehicle drivers have a high prevalence of OSA.^{15,21} Their response to sleep loss appears to be similar to the general population²²; however, some large studies have not identified a relationship between OSA and crash risk.²³ Many other influences are evident here. Shift work, circadian misalignment, and sleep

Submitted for publication January, 2015

Accepted for publication January, 2015

Address correspondence to: Mark E. Howard, MD, Austin Hospital, Respiratory Department, Studley Road, Heidelberg, VIC 3084, Australia; Tel: (39) 4963688; Email: mark.howard@austin.org.au

restriction may be more important in this population as found by Karimi et al.³ and others.²⁴ Hence, it is important clinically to assess a range of individual factors associated with sleepiness and motor vehicle crash risk.²⁵ It is also important to consider that individuals operate in complex transport and workplace systems with many opportunities to intervene on key risk factors that impact on sleepiness at the system level. For example, improving heavy vehicle driver schedules may also improve sleep and crash risk.^{23,26,27}

Despite the challenges in designing ideal studies, OSA appears to increase crash risk with some indication that treatment of OSA may reduce crash risk. For the first time Karimi³ provides evidence that good CPAP adherence in particular is associated with reduced risk. This is not the first time, however, that OSA severity appears to be a poor discriminator of crash risk. Hence, there is an opportunity to improve stratification of risk and assessment of response to treatment through improved objective assessment of the cognitive impact of OSA, including sleepiness, a field of research that has not been sufficiently advanced in sleep medicine. It is important for sleep clinicians to address other factors that impact crash risk in their patient, in particular sleep duration and circadian effects in shift workers, and identify those activities and conditions that could help to mitigate this risk.

CITATION

Howard ME, Jackson ML, Stevenson M. Who needs sleep apnea treatment for safety critical tasks—are we there yet? *SLEEP* 2015;38(3):331–332.

DISCLOSURE STATEMENT

Dr. Howard has received research support from Resmed Foundation, EdanSafe, and the CRC for Alertness, Safety and Productivity; he has consulted for the National Transport Commission and Prevention Express. The other authors have indicated no financial conflicts of interest.

REFERENCES

1. Tregear S, Reston J, Schoelles K, Phillips B. Obstructive sleep apnea and risk of motor vehicle crash: systematic review and meta-analysis. *J Clin Sleep Med* 2009;5:573–81.
2. Tregear S, Reston J, Schoelles K, Phillips B. Continuous positive airway pressure reduces risk of motor vehicle crash among drivers with obstructive sleep apnea: systematic review and meta-analysis. *Sleep* 2010;33:1373–80.
3. Karimi M, Hedner J, Häbel H, Nerman O, Grote L. Sleep apnea related risk of motor vehicle accidents is reduced by continuous positive airway pressure: Swedish traffic accident registry data. *Sleep* 2015;38:341–9.
4. McEvoy RD, Anderson CS, Antic NA, et al. The sleep apnea cardiovascular endpoints (SAVE) trial: rationale and start-up phase. *J Thorac Dis* 2010;2:138–43.
5. Young T, Blustein J, Finn L, Palta M. Sleep-disordered breathing and motor vehicle accidents in a population-based sample of employed adults. *Sleep* 1997;20:608–13.
6. Teran-Santos J, Jimenez-Gomez A, Cordero-Guevara J. The association between sleep apnea and the risk of traffic accidents. Cooperative Group Burgos-Santander [see comments]. *N Engl J Med* 1999;340:847–51.
7. Findley L, Smith C, Hooper J, Dineen M, Suratt PM. Treatment with nasal CPAP decreases automobile accidents in patients with sleep apnea. *Am J Respir Crit Care Med* 2000;161:857–9.
8. George CF. Reduction in motor vehicle collisions following treatment of sleep apnoea with nasal CPAP. *Thorax* 2001;56:508–12.
9. Quan SF, Wright R, Baldwin CM, et al. Obstructive sleep apnea-hypopnea and neurocognitive functioning in the Sleep Heart Health Study. *Sleep Med* 2006;7:498–507.
10. Engleman HM, Kingshott RN, Martin SE, Douglas NJ. Cognitive function in the sleep apnea/hypopnea syndrome (SAHS). *Sleep* 2000;23(Suppl 4):S102–8.
11. Van Dongen HP, Baynard MD, Maislin G, Dinges DF. Systematic interindividual differences in neurobehavioral impairment from sleep loss: evidence of trait-like differential vulnerability. *Sleep* 2004;27:423–33.
12. Vakulin A, Catcheside PG, Baulk SD, et al. Individual variability and predictors of driving simulator impairment in patients with obstructive sleep apnea. *J Clin Sleep Med* 2014;10:647–55.
13. Bennett LS, Langford BA, Stradling JR, Davies RJ. Sleep fragmentation indices as predictors of daytime sleepiness and nCPAP response in obstructive sleep apnea. *Am J Respir Crit Care Med* 1998;158:778–86.
14. Viola AU, Archer SN, James LM, et al. PER3 polymorphism predicts sleep structure and waking performance. *Curr Biol* 2007;17:613–8.
15. Howard ME, Desai AV, Grunstein RR, et al. Sleepiness, sleep-disordered breathing, and accident risk factors in commercial vehicle drivers. *Am J Respir Crit Care Med* 2004;170:1014–21.
16. Lloberes P, Levy G, Descals C, et al. Self-reported sleepiness while driving as a risk factor for traffic accidents in patients with obstructive sleep apnoea syndrome and in non-apnoeic snorers. *Respir Med* 2000;94:971–6.
17. Engleman HM, Hirst WS, Douglas NJ. Under reporting of sleepiness and driving impairment in patients with sleep apnoea/hypopnoea syndrome. *J Sleep Res* 1997;6:272–5.
18. Philip P, Sagaspe P, Taillard J, et al. Maintenance of Wakefulness Test, obstructive sleep apnea syndrome, and driving risk. *Ann Neurol* 2008;64:410–6.
19. Kingshott RN, Cowan JO, Jones DR, et al. The role of sleep-disordered breathing, daytime sleepiness, and impaired performance in motor vehicle crashes—a case control study. *Sleep Breath* 2004;8:61–72.
20. Saksvik IB, Bjorvatn B, Hetland H, Sandal GM, Pallesen S. Individual differences in tolerance to shift work—a systematic review. *Sleep Med Rev* 2011;15:221–35.
21. Stoohs RA, Bingham LA, Itoi A, Guilleminault C, Dement WC. Sleep and sleep-disordered breathing in commercial long-haul truck drivers. *Chest* 1995;107:1275–82.
22. Howard ME, Jackson ML, Swann P, Berlowitz DJ, Grunstein RR, Pierce RJ. Deterioration in driving performance during sleep deprivation is similar in professional and nonprofessional drivers. *Traffic Inj Prev* 2014;15:132–7.
23. Stevenson MR, Elkington J, Sharwood L, et al. The role of sleepiness, sleep disorders, and the work environment on heavy-vehicle crashes in 2 Australian states. *Am J Epidemiol* 2014;179:594–601.
24. Philip P, Akerstedt T. Transport and industrial safety, how are they affected by sleepiness and sleep restriction? *Sleep Med Rev* 2006;10:347–56.
25. Connor J, Norton R, Ameratunga S, et al. Driver sleepiness and risk of serious injury to car occupants: population based case control study. *BMJ* 2002;324:1125.
26. Hanowski RJ, Hickman J, Fumero MC, Olson RL, Dingus TA. The sleep of commercial vehicle drivers under the 2003 hours-of-service regulations. *Accid Anal Prev* 2007;39:1140–5.
27. Jackson ML, Banks S, Belenky G. Investigation of the effectiveness of a split sleep schedule in sustaining sleep and maintaining performance. *Chronobiol Int* 2014;31:1218–30.