Current FDA-Related Drug Information

Approvals, Submission, and Important Labeling Changes for US Marketed Pharmaceuticals

Danial E. Baker, PharmD, FASHP, FASCP*

This monthly feature will help readers keep current on new drugs, new indications, dosage forms, and safety-related changes in labeling or use. Efforts have been made to ensure the accuracy of this information; however, if there are any questions, please let me know at danial.baker@wsu.edu.

Table 1. New drugs approved by the US Food and Drug Administration (FDA): November 16, 2014 through December 12, 2014

Blinatumomab – Blincyto (Amgen)	
<i>Comparative agents:</i>	None
<i>Indication:</i>	Treatment of Philadelphia chromosome–negative relapsed or refractory B-cell precursor acute lymphoblastic leukemia
<i>Mechanism of action:</i>	Bispecific CD19-directed CD3 T-cell engager
<i>Common adverse effects:</i>	Pyrexia, headache, peripheral edema, febrile neutropenia, nausea, hypokalemia, tremor, rash, and constipation
<i>Dosage form & strength:</i>	Injection; 36 mcg single-use vial
<i>Product labeling:</i>	http://www.accessdata.fda.gov/drugsatfda_docs/label/2014/125557lbl.pdf
Clindamycin/benzoyl peroxide – Onexton (Valeant Pharmaceuticals)	
<i>Comparative agents:</i>	Topical clindamycin and benzoyl peroxide–containing products
<i>Indication:</i>	Treatment of comedonal (noninflammatory) and inflammatory acne vulgaris in patients 12 and older
<i>Mechanism of action:</i>	Antibacterial and anti-inflammatory properties
<i>Common adverse effects:</i>	Burning sensation, contact dermatitis, pruritis, and rash
<i>Dosage form & strength:</i>	Topical gel; 1.2%/3.75%
<i>Product labeling:</i>	http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm?fuseaction=Search.Label_ApprovalHistory#labelinfo
Human papillomavirus 9-valent vaccine, recombinant – Gardasil 9 (Merck Sharp & Dohme)	
<i>Comparative agents:</i>	Human papillomavirus quadrivalent (types 6, 11, 16, 18) vaccine, recombinant
<i>Indication:</i>	Prevention of certain diseases caused by 9 types of human papillomavirus; for use in females, ages 9 through 26 years, and males, ages 9 through 15 years
<i>Mechanism of action:</i>	Immunization
<i>Common adverse effects:</i>	Injection site pain, swelling, redness, and headaches
<i>Dosage form & strength:</i>	Injection
<i>Product labeling:</i>	http://www.fda.gov/downloads/BiologicsBloodVaccines/Vaccines/ApprovedProducts/UCM426457.pdf

(continued)

*Director, Drug Information Center, College of Pharmacy, Washington State University Spokane, PO Box 1495, Spokane, WA 99210-1495.

Table 1. New drugs approved by the US Food and Drug Administration (FDA): November 16, 2014 through December 12, 2014 (CONT.)

Hydrocodone bitartrate – Hysingla ER (Purdue Pharma)	
<i>Comparative agents:</i>	Hydrocodone bitartrate
<i>Indication:</i>	Management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate
<i>Mechanism of action:</i>	Opioid agonist
<i>Common adverse effects:</i>	Constipation, nausea, vomiting, fatigue, upper respiratory tract infection, dizziness, headache, and somnolence
<i>Dosage form & strength:</i>	Extended-release tablets: 20, 30, 40, 60, 80, 100, and 120 mg
<i>Product labeling:</i>	http://www.accessdata.fda.gov/drugsatfda_docs/label/2014/206627s000lbl.pdf

Table 2. New dosage forms and indications approved by the FDA: November 16, 2014 through December 12, 2014

Generic name	Brand name (Company)	Indication and comments
New indications		
Ramucirumab	<i>Cyramza</i> (Lilly)	Treatment of patients with metastatic non-small-cell lung cancer (NSCLC)
Rifapentine	<i>Priftin</i> (Sanofi-Aventis)	Use in combination with isoniazid for the treatment of latent tuberculosis infection.
Ruxolitinib	<i>Jakafi</i> (Incyte Corporation)	Treatment of patients with polycythemia vera

Table 3. Agents pending FDA approval: November 16, 2014 through December 12, 2014

Generic name	Brand name (Company)	Indication and comments
Complete response letter: more information required		
Daclatasvir	(Bristol-Myers Squibb)	The application that was submitted to the FDA used a combination of daclatasvir and asunaprevir for the treatment of hepatitis C virus (HCV); since that time asunaprevir has been withdrawn by BMS. The FDA has requested data on the effectiveness and safety of daclatasvir with other antiviral agents.
Recommended for approval by an FDA advisory panel		
Ceftazidime-avibactam	(Actavis)	Treatment of complicated intra-abdominal infections and complicated urinary tract infections caused by gram-negative pathogens when little or no treatment options are available. The Committee did not recommend its approval for the treatment of hospital-acquired bacterial pneumonia or bacteremia.
Agents scheduled for review by an FDA advisory panel		
Desmopressin	<i>Nocdurna</i> (Ferring Pharmaceuticals)	Orally disintegrating sublingual tablets for the treatment of nocturia due to nocturnal polyuria in adults who awaken 2 or more times each night to void (meeting date: January 12, 2015)

(continued)

Current FDA-Related Drug Information

Table 3. Agents pending FDA approval: November 16, 2014 through December 12, 2014 (CONT.)

Generic name	Brand name (Company)	Indication and comments
EP2006	Sandoz	Proposed biosimilar to Amgen Inc.'s <i>Neupogen</i> (filgrastim). The proposed indications for this product are: (1) to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with nonmyeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a significant incidence of severe neutropenia with fever; (2) for reducing the time to neutrophil recovery and the duration of fever, following induction or consolidation chemotherapy treatment of adults with acute myeloid leukemia; (3) to reduce the duration of neutropenia and neutropenia-related clinical sequelae, eg, febrile neutropenia in patients with nonmyeloid malignancies undergoing myeloablative chemotherapy followed by marrow transplantation; (4) for the mobilization of hematopoietic progenitor cells into the peripheral blood for collection by leukapheresis; and (5) for chronic administration to reduce the incidence and duration of sequelae of neutropenia (eg, fever, infections, oropharyngeal ulcers) in symptomatic patients with congenital neutropenia, cyclic neutropenia, or idiopathic neutropenia (meeting date: January 7, 2015)
Isavuconazonium sulfate	(Astellas Pharma)	Treatment of invasive aspergillosis and mucormycosis (meeting date: January 22, 2015)

Table 4. New drug/biologics license applications filed by manufacturer: November 16, 2014 through December 12, 2014

Generic name Brand name (Company)	Comparative agents	Indication	Mechanism of action	Common adverse effects	Dosage form or route	Comments
Paliperidone palmitate (Janssen)	Paliperidone palmitate	Treatment for schizophrenia in adults	Mechanism of action of paliperidone is unknown; might be a combination of central dopamine Type 2 (D2) and serotonin Type 2 (5HT2A) receptor antagonism	Injection site reactions, somnolence/ sedation, dizziness, akathisia, and extrapyramidal disorder	Injectable; intramuscular	Injectable that would be administered every 3 months instead of monthly
Trabectedin <i>Yondelis</i> (Janssen)	Doxorubicin, pegylated liposomal doxorubicin, epirubicin, ifosfamide, taxanes	Treatment of patients with advanced soft tissue sarcoma, including liposarcoma and leiomyosarcoma subtypes, who have received prior chemotherapy including an anthracycline	Alkylating agent	Nausea, vomiting, fatigue, weight loss, neutropenia, thrombocytopenia, and increased liver function tests	Intravenous	Original derived from the sea squirt

Table 5. Significant labeling changes or “Dear Health Professional” letters related to safety^a

Generic name Brand name (Company)	Warning
Ampicillin sodium/ sulbactam sodium <i>Unasyn</i> (Pfizer)	<p>CONTRAINDICATIONS</p> <ul style="list-style-type: none"> <i>Unasyn</i> is contraindicated in patients with a previous history of cholestatic jaundice/hepatic dysfunction associated with <i>Unasyn</i>. <p>WARNINGS</p> <p>Hepatotoxicity</p> <ul style="list-style-type: none"> Hepatic dysfunction, including hepatitis and cholestatic jaundice, has been associated with the use of <i>Unasyn</i>. Hepatic toxicity is usually reversible; however, deaths have been reported. Hepatic function should be monitored at regular intervals in patients with hepatic impairment. <p>ADVERSE REACTIONS</p> <ul style="list-style-type: none"> Cholestatic hepatitis and cholestasis and phlebitis <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm426034.htm</p>
Anagrelide hydrochloride <i>Agrylin</i> (Shire)	<p>WARNINGS AND PRECAUTIONS</p> <p>Cardiovascular Toxicity</p> <ul style="list-style-type: none"> [Added]...Hepatic impairment increases anagrelide exposure and could increase the risk of QTc prolongation. Monitor patients with hepatic impairment for QTc prolongation and other cardiovascular adverse reactions. The potential risks and benefits of anagrelide therapy in a patient with mild and moderate hepatic impairment should be assessed before treatment is commenced. Reduce <i>Agrylin</i> dose in patients with moderate hepatic impairment. Use of <i>Agrylin</i> in patients with severe hepatic impairment has not been studied. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm175918.htm</p>
Asenapine <i>Saphris</i> (Forest Laboratories)	<p>WARNINGS AND PRECAUTIONS</p> <p>Metabolic Changes</p> <ul style="list-style-type: none"> Atypical antipsychotic drugs have been associated with metabolic changes that may increase cardiovascular/cerebrovascular risk. These metabolic changes include hyperglycemia, dyslipidemia, and body weight gain. While all of the drugs in the class have been shown to produce some metabolic changes, each drug has its own specific risk profile. Dyslipidemia section added <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm271083.htm</p>
Bosutinib <i>Bosulif</i> (Wyeth)	<p>WARNINGS AND PRECAUTIONS</p> <p>Renal Toxicity</p> <ul style="list-style-type: none"> An on-treatment decline in estimated glomerular filtration rate (eGFR) has occurred in patients treated with <i>Bosulif</i>. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm426240.htm</p>
Brentuximab vedotin <i>Adcetris</i> (Seattle Genetics)	<p>WARNINGS AND PRECAUTIONS</p> <p>Increased Toxicity in the Presence of Severe Renal Impairment</p> <ul style="list-style-type: none"> The frequency of grade 3 adverse reactions and deaths was greater in patients with severe renal impairment compared to patients with normal renal function. Grade 3 adverse reactions may be more frequent in patients with severe renal impairment compared to patients with normal renal function. Use of <i>Adcetris</i> in patients with severe renal impairment [creatinine clearance (CLcr) less than 30 mL/min] should be avoided. <p>Increased Toxicity in the Presence of Moderate or Severe Hepatic Impairment</p> <ul style="list-style-type: none"> The frequency of grade 3 adverse reactions and deaths was greater in patients with moderate and severe hepatic impairment compared to patients with normal hepatic function. Use of <i>Adcetris</i> in patients with moderate (Child-Pugh B) or severe (Child-Pugh C) hepatic impairment should be avoided.

(continued)

Current FDA-Related Drug Information

Table 5. Significant labeling changes or “Dear Health Professional” letters related to safety^a (CONT.)

Generic name Brand name (Company)	Warning
	<p>Hepatotoxicity</p> <ul style="list-style-type: none"> Serious cases of hepatotoxicity, including fatal outcomes, have occurred in patients receiving <i>Adcetris</i>. Cases were consistent with hepatocellular injury, including elevations of transaminases and/or bilirubin. Cases have occurred after the first dose of <i>Adcetris</i> or after <i>Adcetris</i> rechallenge. Preexisting liver disease, elevated baseline liver enzymes, and concomitant medications may also increase the risk. Monitor liver enzymes and bilirubin. Patients experiencing new, worsening, or recurrent hepatotoxicity may require a delay, change in dose, or discontinuation of <i>Adcetris</i>. <p>Serious Dermatologic Reactions</p> <ul style="list-style-type: none"> Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), including fatal outcomes, have been reported with <i>Adcetris</i>. If SJS or TEN occurs, discontinue <i>Adcetris</i> and administer appropriate medical therapy. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm426241.htm</p>
Cinacalcet <i>Sensipar</i> (Amgen)	<p>WARNINGS AND PRECAUTIONS</p> <p>Hypocalcemia</p> <ul style="list-style-type: none"> <i>Sensipar</i> lowers serum calcium and, therefore, patients should be carefully monitored for the occurrence of hypocalcemia during treatment. Life-threatening events and fatal outcomes associated with hypocalcemia have been reported in patients treated with <i>Sensipar</i>, including pediatric patients. Potential manifestations of hypocalcemia include paresthesias, myalgias, muscle cramping, tetany, and convulsions. <i>Sensipar</i> is not indicated for patients with chronic kidney disease (CKD) not on dialysis. In patients with secondary hyperparathyroidism (HPT) and CKD not on dialysis, the long-term safety and efficacy of <i>Sensipar</i> have not been established. <p>ADVERSE REACTIONS</p> <p>Hypocalcemia</p> <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm268351.htm</p>
Collagenase clostridium histolyticum <i>Xiaflex</i> (Auxilium Pharmaceuticals)	<p>CONTRAINDICATIONS</p> <p><i>Xiaflex</i> is contraindicated in the treatment of Peyronie’s plaques that involve the penile urethra due to potential risk to this structure and in patients with a history of hypersensitivity to <i>Xiaflex</i> or to collagenase used in any other therapeutic application or application method.</p> <p>WARNINGS AND PRECAUTIONS</p> <p>Tendon Rupture or Other Serious Injury to the Injected Finger/Hand in the Treatment of Dupuytren’s Contracture</p> <ul style="list-style-type: none"> Information regarding the risk of skin laceration was added. The possible increased risk for 2 concurrent injections over a single injection is described, as well as postmarketing experience with serious skin lacerations requiring skin graft. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm422930.htm</p>
Daptomycin <i>Cubicin</i> (Cubist)	<p>WARNINGS AND PRECAUTIONS</p> <p>Potential Nervous System and/or Muscular System Effects in Pediatric Patients Younger than 12 Months</p> <ul style="list-style-type: none"> Avoid use of <i>Cubicin</i> in pediatric patients younger than 12 months due to the risk of potential effects on muscular, neuromuscular, and/or nervous systems (either peripheral and/or central) observed in neonatal dogs with intravenous daptomycin. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm225842.htm</p>
Dimethyl fumarate <i>Tecfidera</i> (Biogen Idec)	<p>WARNINGS</p> <p>Warning will be added after a patient with multiple sclerosis treated with <i>Tecfidera</i> (dimethyl fumarate) progressive multifocal leukoencephalopathy (PML) and died.</p> <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/SafetyAlertsforHumanMedicalProducts/ucm424752.htm?source=govdelivery&utm_medium=email&utm_source=govdelivery</p>

(continued)

Table 5. Significant labeling changes or “Dear Health Professional” letters related to safety^a (CONT.)

Generic name Brand name (Company)	Warning
Docetaxel <i>Docetaxel</i> (Caraco Pharmaceutical Laboratories)	<p>WARNINGS AND PRECAUTIONS</p> <p>Eye Disorders</p> <ul style="list-style-type: none"> Cystoid macular edema (CME) has been reported in patients treated with <i>Docetaxel</i>. Patients with impaired vision should undergo a prompt and comprehensive ophthalmologic examination. If CME is diagnosed, <i>Docetaxel</i> treatment should be discontinued and appropriate treatment initiated. Alternative non-taxane cancer treatment should be considered. <p>Alcohol Content</p> <ul style="list-style-type: none"> Cases of intoxication have been reported with some formulations of docetaxel due to the alcohol content. The alcohol content in a dose of <i>Docetaxel</i> injection may affect the central nervous system and should be taken into account for patients in whom alcohol intake should be avoided or minimized. Consideration should be given to the alcohol content in <i>Docetaxel</i> injection on the ability to drive or use machines immediately after the infusion. Each administration of <i>Docetaxel</i> injection at 100 mg/m² delivers 1.425 g/m² of ethanol. For a patient with a body surface area (BSA) of 2.0 m², this would deliver 2.85 g of ethanol. Other docetaxel products may have a different amount of alcohol. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm422805.htm</p>
Docetaxel <i>Taxotere</i> (Sanofi-Aventis)	<p>WARNINGS AND PRECAUTIONS</p> <p>Alcohol Content</p> <ul style="list-style-type: none"> Cases of intoxication have been reported with some formulations of docetaxel due to the alcohol content. The alcohol content in a dose of <i>Taxotere</i> injection may affect the central nervous system and should be taken into account for patients in whom alcohol intake should be avoided or minimized. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm212079.htm</p>
Duloxetine HCl <i>Cymbalta</i> (Lilly)	<p>WARNINGS AND PRECAUTIONS</p> <p>Orthostatic Hypotension, Falls, and Syncope</p> <ul style="list-style-type: none"> Orthostatic hypotension, falls, and syncope have been reported with therapeutic doses of <i>Cymbalta</i>. Syncope and orthostatic hypotension tend to occur within the first week of therapy but can occur at any time during <i>Cymbalta</i> treatment, particularly after dose increases. The risk of falling appears to be related to the degree of orthostatic decrease in blood pressure as well as other factors that may increase the underlying risk of falls. In an analysis of patients from all placebo-controlled trials, patients treated with <i>Cymbalta</i> reported a higher rate of falls compared to patients treated with placebo. Risk appears to be related to the presence of orthostatic decrease in blood pressure. The risk of blood pressure decreases may be greater in patients taking concomitant medications that induce orthostatic hypotension (such as antihypertensives) or are potent CYP1A2 inhibitors and in patients taking <i>Cymbalta</i> at doses above 60 mg daily. Consideration should be given to dose reduction or discontinuation of <i>Cymbalta</i> in patients who experience symptomatic orthostatic hypotension, falls, and/or syncope during <i>Cymbalta</i> therapy. Risk of falling also appeared to be proportional to a patient's underlying risk for falls and appeared to increase steadily with age. As elderly patients tend to have a higher underlying risk for falls due to a higher prevalence of risk factors such as use of multiple medications, medical comorbidities, and gait disturbances, the impact of increasing age by itself is unclear. Falls with serious consequences including bone fractures and hospitalizations have been reported. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm319241.htm</p>

(continued)

Table 5. Significant labeling changes or “Dear Health Professional” letters related to safety^a (CONT.)

Generic name Brand name (Company)	Warning
Duloxetine hydrochloride <i>Cymbalta</i> (Lilly)	<p>BOXED WARNING: Suicidal Thoughts and Behaviors</p> <ul style="list-style-type: none"> Antidepressants increased the risk of suicidal thoughts and behavior in children, adolescents, and young adults in short-term studies. These studies did not show an increase in the risk of suicidal thoughts and behavior with antidepressant use in patients over age 24; there was a reduction in risk with antidepressant use in patients aged 65 and older. In patients of all ages who are started on antidepressant therapy, monitor closely for worsening and for emergence of suicidal thoughts and behaviors. Advise families and caregivers of the need for close observation and communication with the prescriber. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm319241.htm</p>
Gemfibrozil <i>Lopid</i> (Pfizer)	<p>WARNINGS AND PRECAUTIONS</p> <p>Concomitant Anticoagulants</p> <ul style="list-style-type: none"> Caution should be exercised when warfarin is given in conjunction with <i>Lopid</i>. The dosage of warfarin should be reduced to maintain the prothrombin time at the desired level to prevent bleeding complications. Frequent prothrombin determinations are advisable until it has been definitely determined that the prothrombin level has stabilized. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm175927.htm</p>
Goserelin acetate <i>Zoladex</i> (AstraZeneca)	<p>WARNINGS AND PRECAUTIONS</p> <p>Effect on QT/QTc Interval</p> <ul style="list-style-type: none"> Androgen deprivation therapy may prolong the QT/QTc interval. Providers should consider whether the benefits of androgen deprivation therapy outweigh the potential risks in patients with congenital long QT syndrome, congestive heart failure, frequent electrolyte abnormalities, and in patients taking drugs known to prolong the QT interval. Electrolyte abnormalities should be corrected. Consider periodic monitoring of electrocardiograms and electrolytes. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm182245.htm</p>
Isosorbide dinitrate <i>Dilatrate SR</i> (Auxilium Pharmaceuticals)	<p>CONTRAINDICATIONS</p> <ul style="list-style-type: none"> Do not use <i>Dilatrate SR</i> in patients who are taking certain drugs for erectile dysfunction (phosphodiesterase inhibitors), such as avanafil, sildenafil, tadalafil, vardenafil. Concomitant use can cause severe hypotension, syncope, or myocardial ischemia. Do not use <i>Dilatrate SR</i> in patients who are taking the soluble guanylate cyclase stimulator riociguat. Concomitant use can cause hypotension. <p>PRECAUTIONS</p> <p>Drug Interactions</p> <ul style="list-style-type: none"> Concomitant use of <i>Dilatrate SR</i> with phosphodiesterase inhibitors in any form is contraindicated. Concomitant use of <i>Dilatrate SR</i> with riociguat, a soluble guanylate cyclase stimulator, is contraindicated. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm422810.htm</p>
Isosorbide mononitrate <i>Monoket</i> (Kremers Urban Pharmaceuticals)	<p>CONTRAINDICATIONS</p> <ul style="list-style-type: none"> [Added] Do not use <i>Monoket</i> in patients who are taking certain drugs for erectile dysfunction (phosphodiesterase inhibitors), such as sildenafil, tadalafil, or vardenafil. Concomitant use can cause severe hypotension, syncope, or myocardial ischemia. Do not use <i>Monoket</i> in patients who are taking the soluble guanylate cyclase stimulator riociguat. Concomitant use can cause hypotension. <p>PRECAUTIONS</p> <p>Drug Interactions</p> <ul style="list-style-type: none"> [Added] Concomitant use of <i>Monoket</i> with phosphodiesterase inhibitors in any form is contraindicated. Concomitant use of <i>Monoket</i> with riociguat, a soluble guanylate cyclase stimulator, is contraindicated. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm422928.htm</p>

(continued)

Table 5. Significant labeling changes or “Dear Health Professional” letters related to safety^a (CONT.)

Generic name Brand name (Company)	Warning
Leflunomide <i>Arava</i> (Sanofi)	<p>CONTRAINDICATIONS</p> <ul style="list-style-type: none"> <i>Arava</i> is contraindicated in patients with known hypersensitivity to leflunomide, teriflunomide, or any of the other components of <i>Arava</i>. <p>WARNINGS</p> <p>Skin Reactions</p> <ul style="list-style-type: none"> Drug reactions with eosinophilia and systemic symptoms (DRESS) have been reported in patients receiving <i>Arava</i>. If a patient taking <i>Arava</i> develops any of these conditions, <i>Arava</i> therapy should be stopped, and a drug elimination procedure is recommended. <p>PRECAUTIONS</p> <p>General</p> <ul style="list-style-type: none"> Co-administration of teriflunomide with leflunomide is not recommended, as leflunomide is the parent compound of teriflunomide. <p>ADVERSE REACTIONS</p> <p>Skin and Appendages</p> <ul style="list-style-type: none"> Drug reaction with DRESS <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm230245.htm</p>
Leuprolide acetate for injectable suspension <i>Eligard</i> (Tolmar Pharmaceuticals)	<p>WARNINGS AND PRECAUTIONS</p> <p>Effect on QT/QTc Interval</p> <ul style="list-style-type: none"> Androgen deprivation therapy may prolong the QT/QTc interval. Providers should consider whether the benefits of androgen deprivation therapy outweigh the potential risks in patients with congenital long QT syndrome, congestive heart failure, frequent electrolyte abnormalities, and in patients taking drugs known to prolong the QT interval. Electrolyte abnormalities should be corrected. Consider periodic monitoring of electrocardiograms and electrolytes. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm232194.htm</p>
Levetiracetam <i>Keppra</i> (UCB)	<p>WARNINGS AND PRECAUTIONS [following sections updated with additional data from studies using an oral formulation]</p> <p>Behavioral Abnormalities and Psychotic Symptoms; Somnolence and Fatigue; Hematologic Abnormalities</p> <p>Increase in Blood Pressure [section added]</p> <ul style="list-style-type: none"> In a randomized, placebo-controlled study in patients 1 month to less than 4 years of age using an oral formulation of <i>Keppra</i>, a significantly higher risk of increased diastolic blood pressure was observed in the <i>Keppra</i>-treated patients (17%) compared to placebo-treated patients (2%). There was no overall difference in mean diastolic blood pressure between the treatment groups. This disparity between the <i>Keppra</i> and placebo treatment groups was not observed in the studies of older children or in adults. Monitor patients 1 month to less than 4 years of age for increases in diastolic blood pressure. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm413465.htm</p>

(continued)

Table 5. Significant labeling changes or “Dear Health Professional” letters related to safety^a (CONT.)

Generic name Brand name (Company)	Warning
Methoxy polyethylene glycol-epoetin beta <i>Mircera</i> (Hoffmann-La Roche)	<p>BOXED WARNING [edited]</p> <p>WARNINGS: Erythropoiesis-stimulating agents (ESAs) increase the risk of death, myocardial infarction, stroke, venous thromboembolism, thrombosis of vascular access, and tumor progression or recurrence.</p> <p>Chronic Kidney Disease</p> <ul style="list-style-type: none"> • In controlled trials, patients experienced greater risks for death, serious adverse cardiovascular reactions, and stroke when administered ESAs to target a hemoglobin level of greater than 11 g/dL. • No trial has identified a hemoglobin target level, ESA dose, or dosing strategy that does not increase these risks. • Use the lowest <i>Mircera</i> dose sufficient to reduce the need for red blood cell (RBC) transfusions. • <i>Mircera</i> is not indicated and is not recommended for the treatment of anemia due to cancer chemotherapy. A dose-ranging study of <i>Mircera</i> was terminated early because of more deaths among patients receiving <i>Mircera</i> than another ESA. • ESAs have shown shortened overall survival and/or increased the risk of tumor progression or recurrence in clinical studies in patients with breast, non-small-cell lung, head and neck, lymphoid, and cervical cancers. <p>CONTRAINDICATIONS</p> <p><i>Mircera</i> is contraindicated in patients with:</p> <ul style="list-style-type: none"> • Uncontrolled hypertension • Pure red cell aplasia that begins after treatment with <i>Mircera</i> or other erythropoietin protein drugs • History of serious or severe allergic reactions to <i>Mircera</i> (eg, anaphylactic reactions, angioedema, bronchospasm, skin rash, and urticaria). <p>WARNINGS AND PRECAUTIONS [Significant edits to the following sections]</p> <p>Increased Mortality, Myocardial Infarction, Stroke, and Thromboembolism; Increased Mortality and/or Increased Risk of Tumor Progression or Recurrence; Hypertension; Seizures; Pure Red Cell Aplasia; Serious Allergic Reactions; Dialysis Management; and Laboratory Monitoring.</p> <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm423232.htm</p>
Metoprolol succinate extended release / hydrochlorothiazide <i>Dutoprol</i> (Covis Pharmaceuticals)	<p>WARNINGS AND PRECAUTIONS</p> <p>Heart Failure</p> <ul style="list-style-type: none"> • Worsening cardiac failure may occur during up-titration of beta-blockers. If such symptoms occur, increase diuretics and restore clinical stability (compensated heart failure) before advancing the dose of <i>Dutoprol</i>. It may be necessary to lower the dose of <i>Dutoprol</i> or temporarily discontinue it. Such episodes do not preclude subsequent successful titration of <i>Dutoprol</i>. <p>Impaired Hepatic Function</p> <ul style="list-style-type: none"> • Consider initiating <i>Dutoprol</i> therapy at doses lower than those recommended; gradually increase dosage to optimize therapy, while monitoring closely for adverse events. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm249753.htm</p>

(continued)

Table 5. Significant labeling changes or “Dear Health Professional” letters related to safety^a (CONT.)

Generic name Brand name (Company)	Warning
Minocycline for injection <i>Minocin</i> intravenous (Rempex Pharms)	<p>WARNINGS</p> <p>Intracranial Hypertension</p> <ul style="list-style-type: none"> Intracranial hypertension (IH, pseudotumor cerebri) has been associated with the use of tetracyclines including <i>Minocin</i>. Clinical manifestations of IH include headache, blurred vision, diplopia, and vision loss; papilledema can be found on funduscopy. Women of childbearing age who are overweight or have a history of IH are at greater risk for developing tetracycline-associated IH. Concomitant use of isotretinoin and <i>Minocin</i> should be avoided because isotretinoin is also known to cause pseudotumor cerebri. Although IH typically resolves after discontinuation of treatment, the possibility for permanent visual loss exists. If visual disturbance occurs during treatment, prompt ophthalmologic evaluation is warranted. Since intracranial pressure can remain elevated for weeks after drug cessation, patients should be monitored until they stabilize. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm422931.htm</p>
Moxifloxacin HCl <i>Avelox</i> (Bayer HealthCare Pharmaceuticals)	<p>WARNINGS AND PRECAUTIONS</p> <p>Blood Glucose Disturbances</p> <ul style="list-style-type: none"> As with all fluoroquinolones, disturbances in blood glucose, including both hypoglycemia and hyperglycemia, have been reported with <i>Avelox</i>. In <i>Avelox</i>-treated patients, dysglycemia occurred predominantly in elderly diabetic patients receiving concomitant treatment with an oral hypoglycemic agent (eg, sulfonylurea) or with insulin. In diabetic patients, careful monitoring of blood glucose is recommended [see Adverse Reactions (6.2)]. If a hypoglycemic reaction occurs, <i>Avelox</i> should be discontinued and appropriate therapy should be initiated immediately and the addition of cyclosporine to the list of drugs that do not affect moxifloxacin concentration. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm319318.htm</p>
Nitroglycerin <i>Nitro-Dur</i> transdermal infusion system (Merck & Co.)	<p>CONTRAINDICATIONS [added]</p> <ul style="list-style-type: none"> Do not use <i>Nitro-Dur</i> in patients who are taking phosphodiesterase inhibitors (such as sildenafil, tadalafil, or vardenafil) for erectile dysfunction or pulmonary arterial hypertension. Concomitant use can cause severe drops in blood pressure. Do not use <i>Nitro-Dur</i> in patients who are taking the soluble guanylate cyclase stimulator riociguat. Concomitant use can cause hypotension. <p>PRECAUTIONS [added]</p> <ul style="list-style-type: none"> Concomitant use of <i>Nitro-Dur</i> with phosphodiesterase inhibitors in any form is contraindicated. Concomitant use of <i>Nitro-Dur</i> with soluble guanylate cyclase stimulators is contraindicated. <p>Drug Interactions [added]</p> <ul style="list-style-type: none"> Concomitant use of <i>Nitro-Dur</i> with phosphodiesterase inhibitors in any form is contraindicated. <p>Concomitant use of <i>Nitro-Dur</i> with soluble guanylate cyclase stimulators is contraindicated.</p> <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm422803.htm</p>

(continued)

Current FDA-Related Drug Information

Table 5. Significant labeling changes or “Dear Health Professional” letters related to safety^a (CONT.)

Generic name Brand name (Company)	Warning
Nitroglycerin 5% in dextrose for injection (various)	<p>CONTRAINDICATIONS [added]</p> <ul style="list-style-type: none"> Do not use nitroglycerin in 5% dextrose injection in patients who are taking certain drugs for erectile dysfunction (phosphodiesterase inhibitors) such as sildenafil, tadalafil, or vardenafil. Concomitant use can cause severe hypotension, syncope, or myocardial ischemia. Do not use nitroglycerin in 5% dextrose in patients who are taking the soluble guanylate cyclase stimulator riociguat. Concomitant use can cause hypotension. <p>PRECAUTIONS [added]</p> <ul style="list-style-type: none"> Concomitant use of nitroglycerin in 5% dextrose injection with phosphodiesterase inhibitors in any form is contraindicated. Concomitant use of nitroglycerin in 5% dextrose with riociguat, a soluble guanylate cyclase stimulator, is contraindicated. <p>Drug Interactions [added]</p> <ul style="list-style-type: none"> Concomitant use of nitroglycerin in 5% dextrose injection with phosphodiesterase inhibitors in any form is contraindicated. Concomitant use of nitroglycerin in 5% dextrose with riociguat, a soluble guanylate cyclase stimulator, is contraindicated. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm422804.htm</p>
Pazopanib HCl <i>Votrient</i> (GlaxoSmithKline)	<p>WARNINGS AND PRECAUTIONS</p> <p>Cardiac Dysfunction</p> <ul style="list-style-type: none"> Cardiac dysfunction, such as decreased left ventricular ejection fraction (LVEF) and congestive heart failure, have been reported. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm303649.htm</p>
Romidepsin <i>Istodax</i> (Celgene Corporation)	<p>WARNINGS AND PRECAUTIONS</p> <p>Infections</p> <ul style="list-style-type: none"> Fatal and serious infections, including pneumonia, sepsis, and viral reactivation, including Epstein Barr and hepatitis B viruses, have been reported in clinical trials with <i>Istodax</i>. These can occur during treatment and within 30 days after treatment. The risk of life-threatening infections may be greater in patients with a history of prior treatment with monoclonal antibodies directed against lymphocyte antigens and in patients with disease involvement of the bone marrow. Reactivation of hepatitis B virus infection has occurred in 1% of peripheral T-cell lymphoma (PTCL) patients in clinical trials in Western populations. In patients with evidence of prior hepatitis B infection, consider monitoring for reactivation and consider antiviral prophylaxis. Reactivation of Epstein Barr viral infection leading to liver failure has occurred in a trial of patients with relapsed or refractory extranodal NK/T-cell lymphoma. In one case, ganciclovir prophylaxis failed to prevent Epstein Barr viral reactivation. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm360070.htm</p>
Sofosbuvir <i>Sovaldi</i> (Gilead)	<p>Pregnancy: Use with Ribavirin or Peginterferon Alfa/Ribavirin</p> <ul style="list-style-type: none"> When <i>Sovaldi</i> is used in combination with ribavirin or peginterferon alfa/ribavirin, women of childbearing potential and their male partners must use 2 forms of effective contraception during treatment and for at least 6 months after treatment has concluded. Routine monthly pregnancy tests must be performed during this time. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm426234.htm</p>

(continued)

Table 5. Significant labeling changes or “Dear Health Professional” letters related to safety^a (CONT.)

Generic name Brand name (Company)	Warning
Sweet vernal, orchard, perennial rye, timothy, and Kentucky blue grass mixed pollens allergen extract <i>Oralair</i> (Stallergenes)	<p>CONTRAINDICATIONS</p> <p><i>Oralair</i> is contraindicated in patients with:</p> <ul style="list-style-type: none"> • Severe, unstable or uncontrolled asthma • History of any severe systemic allergic reaction • History of any severe local reaction to sublingual allergen immunotherapy • A history of eosinophilic esophagitis • Hypersensitivity to any of the inactive ingredients (mannitol, microcrystalline cellulose, croscarmellose sodium, colloidal anhydrous silica, magnesium stearate and lactose monohydrate) contained in this product <p>WARNINGS AND PRECAUTIONS</p> <p>Eosinophilic Esophagitis</p> <ul style="list-style-type: none"> • Eosinophilic esophagitis has been reported in association with sublingual tablet immunotherapy. Discontinue <i>Oralair</i> and consider a diagnosis of eosinophilic esophagitis in patients who experience severe or persistent gastro-esophageal symptoms including dysphagia or chest pain. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm422927.htm</p>
Teriflunomide <i>Aubagio</i> (Genzyme Corporation)	<p>WARNINGS AND PRECAUTIONS</p> <p>Hepatotoxicity</p> <ul style="list-style-type: none"> • [Updated study data] In placebo-controlled trials, ALT greater than 3 times the upper limit of normal (ULN) occurred in 61/1,045 (5.8%) and 62/1,002 (6.2%) of patients receiving <i>Aubagio</i> 7 mg and 14 mg, respectively, and 38/997 (3.8%) of patients receiving placebo, during the treatment period. <p>Bone Marrow Effects/Immunosuppression Potential/Infections [updated study data in the following sections]</p> <ul style="list-style-type: none"> • White blood cell count decrease • Risk of infection/tuberculosis screening <p>Peripheral Neuropathy</p> <ul style="list-style-type: none"> • [Updated study data] The incidence of peripheral neuropathy confirmed by nerve conduction studies was 1.4% (13 patients) and 1.9% (17 patients) of patients receiving 7 mg and 14 mg of <i>Aubagio</i>, respectively, compared with 0.4% receiving placebo (4 patients). Treatment was discontinued in 0.7% (8 patients) with confirmed peripheral neuropathy (3 patients receiving <i>Aubagio</i> 7 mg and 5 patients receiving <i>Aubagio</i> 14 mg). Five of them recovered following treatment discontinuation. Not all cases of peripheral neuropathy resolved with continued treatment. Peripheral neuropathy also occurred in patients receiving leflunomide. <p>Increased Blood Pressure</p> <ul style="list-style-type: none"> • [Updated study data] In placebo-controlled studies, the mean change from baseline to the end of study in systolic blood pressure was +2.3 mm Hg and +2.7 mm Hg for <i>Aubagio</i> 7 mg and 14 mg, respectively, and -0.6 mm Hg for placebo. The change from baseline in diastolic blood pressure was +1.4 mm Hg and +1.9 mm Hg for <i>Aubagio</i> 7 mg and 14 mg, respectively, and -0.3 mm Hg for placebo. Hypertension was an adverse reaction in 3.1% and 4.3% of patients treated with 7 mg or 14 mg of <i>Aubagio</i> compared with 1.8% for placebo. <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm423101.htm</p>

(continued)

Current FDA-Related Drug Information

Table 5. Significant labeling changes or “Dear Health Professional” letters related to safety^a (CONT.)

Generic name <i>Brand name</i> (Company)	Warning
Vemurafenib <i>Zelboraf</i> (Genentech)	<p>WARNINGS AND PRECAUTIONS</p> <p>Hepatotoxicity</p> <ul style="list-style-type: none"> • Liver injury leading to functional hepatic impairment, including coagulopathy or other organ dysfunction, can occur with <i>Zelboraf</i>. <p>ADVERSE REACTIONS</p> <ul style="list-style-type: none"> • Panniculitis and neutropenia <p>http://www.fda.gov/Safety/MedWatch/SafetyInformation/ucm364374.htm</p>
Ziprasidone <i>Geodon</i> (Pfizer) & generics	<p>WARNING</p> <ul style="list-style-type: none"> • Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) has been reported with ziprasidone therapy. Patients who have a fever with a rash and/or swollen lymph glands should seek urgent medical care. Health care professionals should immediately stop treatment with ziprasidone if DRESS is suspected. <p>http://www.fda.gov/Drugs/DrugSafety/ucm426391.htm?source=govdelivery&utm_medium=email&utm_source=govdelivery</p>

^aPractitioners are encouraged to check the US Food and Drug Administration’s MEDWATCH Web site (<http://www.fda.gov/medwatch/safety.htm>) for updated information.