Published in final edited form as:

Neurorehabil Neural Repair. 2008 ; 22(3): 217–227. doi:10.1177/1545968308315600.

The Predictive Brain State: Timing Deficiency in Traumatic Brain Injury?

Jamshid Ghajar, MD, PhD, Richard B. Ivry, PhD, and the Cognitive and Neurobiological Research Consortium

Brain Trauma Foundation and the Department of Neurological Surgery, Weill Medical College of Cornell University, New York, New York (JG), and the Department of Psychology, University of California, Berkeley (RBI)

Abstract

Attention and memory deficits observed in traumatic brain injury (TBI) are postulated to result from the shearing of white matter connections between the prefrontal cortex, parietal lobe, and cerebellum that are critical in the generation, maintenance, and precise timing of anticipatory neural activity. These fiber tracts are part of a neural network that generates predictions of future states and events, processes that are required for optimal performance on attention and working memory tasks. The authors discuss the role of this anticipatory neural system for understanding the varied symptoms and potential rehabilitation interventions for TBI.

Preparatory neural activity normally allows the efficient integration of sensory information with goal-based representations. It is postulated that an impairment in the generation of this activity in traumatic brain injury (TBI) leads to performance variability as the brain shifts from a predictive to reactive mode. This dysfunction may constitute a fundamental defect in TBI as well as other attention disorders, causing working memory deficits, distractibility, a loss of goal-oriented behavior, and decreased awareness.

“The future is not what is coming to meet us, but what we are moving forward to meet.”

—Jean-Marie Guyau¹

Keywords

Attention; mild traumatic brain injury; head injury; concussion; post-concussive symptoms; diffuse axonal injury; neuropsychological tests

TBI is a global health problem in terms of incidence, cost, and impact on daily living.² Worldwide, an estimated 57 million individuals have been hospitalized with TBI.³ The Glasgow Coma Scale (GCS) has come to serve as a standardized assessment instrument to classify the severity of TBI, with scores ranging from a low of 3 (comatose) to 15 (awake and following commands). The GCS score is a reliable measure of the level of consciousness, but it is of questionable utility for assessing current cognitive status and

predicting longer-term cognitive sequelae. Additional assessments are especially critical given that more than 80% of individuals classified as mild TBI or concussion have GCS scores between 13 and 15. In the United States, medical center emergency rooms report approximately 1.74 million cases of mild TBI.⁴ These estimates fail to include unrecognized or unreported TBI cases that may number up to 3 million^{3,5} for sports-related injuries and up to 40% for personnel participating in current military engagements.^{6,7}

Mild TBI or concussion with a normal GCS score implies little or no brain injury; indeed, CT imaging studies reveal traumatic lesions in less than 15% of patients.⁸ Cognitive sequelae, which are difficult to measure, are often ascribed to the traumatic event or pre-morbid factors.⁹ However, we postulate that subtle brain white matter tract lesions that are not detected by standard forms of neuroimaging may lead to chronic difficulties in complex cognitive tasks. These white matter tract lesions are located predominantly in anterior cerebral areas, which are selectively vulnerable to rotational shearing injury. Furthermore, these fiber tracts are part of a neural network that generates predictions of future states and events, processes that are required for optimal performance on attention and working memory tasks. We hypothesize that impairments in this anticipatory process likely contribute to the fact that approximately 20% of individuals with TBI end up being classified as chronically disabled.¹⁰

Unless otherwise specified, we use the term “TBI” to denote mild TBI patients and survivors of more severe TBI who are awake and responsive. Although we expect the general ideas advanced here are relevant for understanding the cognitive deficits in all forms of TBI, much of the discussion in the literature has centered on accounting for the cognitive deficits that persist even in individuals with relatively mild overt symptoms.

THE PREDICTIVE BRAIN STATE

Currently, there is no causative explanation for the attention deficits and persistence of symptoms in TBI patients.^{11,12} An extensive review of TBI by Alexander¹³ finds no single psychological, physiologic, somatic, or demographic factor that can account for persistent postconcussive symptoms (PCS). Many PCS symptoms are equally compatible with diagnoses of depression, anxiety, and chronic pain as they are with disorders of cognition. The lack of a clear understanding of the mechanisms underlying concussive symptoms has important consequences for the treatment of these individuals. For example, clear assessment guidelines are lacking to help determine when it is appropriate for athletes with concussive symptoms to resume sports, in large part because there is no consensus on objective measures to assess the level of cognitive function at which there remains a high vulnerability for reinjury.¹⁴

A widely accepted view is that frontal lobe damage underlies the cognitive deficits associated with TBI. This view is consistent, to some degree, with the central role attributed to this brain region in functions related to attention and working memory. However, frontal lobe lesion volumes in TBI patients do not correlate with neuropsychological deficits.¹⁵ This observation, coupled with more sophisticated models of cognition, suggests that these complex processes involve distributed neural networks and that the effects of TBI reflect

disruption of these networks.¹⁶ While TBI certainly results in frontal-like symptoms, one goal in this review is to encourage a broader conceptualization of how TBI might disrupt control processes. In particular, we outline a network-based hypothesis that emphasizes an impairment in anticipatory control, a form of prediction. Rather than attribute this process to a single neural region such as the frontal lobe, we consider how this predictive aspect of brain function emerges through the interactions of frontal and parietal regions, as well as the cerebellum.

A core premise we develop in this article is that one aspect of attention emerges from the generation of moment-to-moment expectancies about the immediate future. To be clear, we argue that the causal link here is not that sensory expectancy requires attention; rather, we propose that attention is the consequence of, or rather, is manifest as expectancies of the future. These expectancies are a core feature of goal-oriented behavior, providing predictions to be compared with sensory feedback. These expectancies, based on prior experience, are essential for ensuring the fluid operation of cognitive functions, allowing them to be anticipatory rather than reactive. One consequence of successful anticipation is that performance becomes less variable. By accurately anticipating the future, the performer is less likely to miss task-relevant information or be distracted by irrelevant information, 2 prominent sources of errors. *By anticipation, the individual is “paying attention,” creating a future-oriented brain state.*

Anticipatory Neural Network

Prediction is a general feature of cognition, operating over multiple time scales. When driving, we may anticipate the actions of an oncoming car as well as think about the location of a lunch-stop, 100 miles down the highway. Within the framework of our model of TBI-related cognitive deficits, we limit our discussion to the saliency of short-range predictions that require the comparison of expectancies and feedback with real-time properties. We hypothesize that these temporal predictions emerge from the interactions of a network involving frontal, parietal, and cerebellar areas (Table 1). Could one prominent source of performance variability in TBI be related to an impaired ability to generate and/or maintain predictive states? Our premise is that disruption of the connectivity between these areas affects this predictive capability and, as such, results in deficits in goal-oriented behavior arising from problems with attention and working memory.

One way to examine the neural underpinnings of prediction is to study the activation and coherence of activation in and between cortical areas that are required for anticipatory activity in attention and working memory. Studies of spatial attention frequently use tasks in which a cue, indicating either the probable location of a stimulus or serving solely as an alerting signal, is followed after a short delay by an imperative stimulus. Neuroimaging studies consistently reveal that, following the cue, a network encompassing the dorsolateral prefrontal cortex (PFC) and parietal lobe is engaged, most prominently in the right hemisphere.^{17–19} In addition to this cerebral cortical network, anticipation also engages the cerebellum; functional magnetic resonance imaging (fMRI) studies consistently show activation of the cerebellum during the interval following attentional cues (Figure 1).^{20–22}

This activation is not dependent on the presence of movement, although it may require the preparation of a potential response.

Electrophysiological markers of anticipation associated with the PFC include the readiness potential (RP)²³ on tasks in which the movement is self-initiated and contingent negative variation (CNV)²⁴ when the response is initiated following an imperative stimulus. In TBI patients, the CNV amplitude is decreased. Reaction times become more variable, and errors in performance correlate with the amplitude of the CNV.²⁵ There are also reported reductions or absence of the RP in TBI patients in self-initiated movements (Figure 2),²⁶ similar to that observed in patients with lesions of the cerebellar dentate nucleus.²⁷ The cerebellum (and basal ganglia) are assumed to contribute to the RP and CNV through thalamic relays.²⁸ Similar findings of decreased frontal preparatory electroencephalography have been reported to predict deficits in attention in normal volunteers.²⁹

The relationship between parietal and PFC anticipatory activity has been examined in a variety of ways.³⁰ Within the parietal lobe, anticipatory markers are observed in inferior parietal lobule (IPL), intraparietal sulcus, and the posterior parietal area.^{19,31,32} A recent functional connectivity study identified 2 distinct networks. A fronto-parietal-cerebellar network was consistently engaged following cues indicating the onset of experimental tasks; in contrast, a cingulo-opercular network was selectively associated with the maintenance of internal goals.³³ The former was most active in predictable situations, when the conditions allow the participants to know when a stimulus will appear rather than when the stimuli occur at random points in time. This observation underscores the temporal property of this attentional network. Goal maintenance or intention is generally independent of this temporal constraint.³⁴

The frontal-parietal network is also activated in working memory tasks.³⁵ Variability in antecedent activity in IPL can predict on a trial-to-trial basis performance on a working memory task.³⁶ Interestingly, initial encoding in a working memory task does not correlate with performance unless there is a high level of sustained activity in PFC and IPL during the maintenance period before retrieval.³⁷

Polysynaptic tracers have been used to identify cerebral cortical targets of the deep cerebellar nuclei in the monkey. This work has shown that, in addition to projections to primary and secondary motor areas, there are significant cerebello-thalamo-cortical projections to the prefrontal and parietal cortex.^{38,39} These anatomical connections could be essential for refining anticipatory neural activity in the cortex. Within the domain of sensorimotor control, it has been hypothesized that the cerebellum generates forward models related to the consequences of planned actions and the resultant changes in sensory feedback.⁴⁰ Extending this idea to a more general role in anticipation that includes the prefrontal and parietal cortex, the specialized role of the cerebellum may be to provide the precise timing for these predictions. As such, increases in performance variability could be a consequence of poorly formed predictions.^{41,42} We recognize that a lack of preparation can arise from many causes. For example, performance will be variable if an alert person is unable to maintain the current goal. Less consideration has been given to preparatory problems that may result from failures of anticipation, or predictions that are mistimed.

These findings suggest that the PFC, IPL, and cerebellum interact to maintain information during delay periods associated with anticipatory activity, either related to the expectation of an imperative stimulus or the retrieval of information from working memory. Models of these interactions suggest a synchronization process that enables an expectancy to be maintained and compared with incoming sensory information (Figure 3).^{40,43}

Performance Variability, Predictive Timing, and Self-Agency

In most studies, performance variability is considered at the group level. Individual variability tends to be ignored, with the various observations in each condition collapsed into a single mean. However, in terms of the study of attention and anticipation, intra-individual variability (IIV) is likely to prove quite informative. Interestingly, variability has frequently been the primary dependent variable in studies of temporal processing. A recent review⁴⁴ finds that high IIV is a prominent feature of TBI.^{45,46} In addition to an increase in IIV and their resultant errors,⁴⁷ TBI patients have longer reaction times,^{25,48} are more distractible,⁴⁹ and experience difficulty in sustaining attention.⁵⁰ Similar findings have been reported in ADHD^{51,52} and in aging studies,^{53–56} with increasing IIV accounting for most of the group variance.⁵⁷ In fact, IIV is one of the better predictors of cognitive decline.⁵⁸

The capability to anticipate the future, as well as the variability in this process, may have implications for awareness. Awareness is, of course, a component of paying attention; moreover, the qualia of object awareness is likely a by-product of self-agency that results from the predictive state.⁵⁹ Disruption of the predictive state, either artificially induced by altering the timing of sensory feedback⁶⁰ or arising as part of clinical syndromes such as schizophrenia⁶¹ and autism,⁶² will result in decreased awareness of self-agency. The impact of variability on awareness and self-agency is also evident in right parietal lesion patients with left-sided neglect—less attention or awareness is directed toward the left side with high performance variability.⁶³ Similarly, the IIV that occurs in TBI⁶⁴ is associated with decreased awareness and, in high variability states, a lack of awareness and self-agency.

Accurate predictive timing, the ability to synchronize what is expected with what is observed, produces low variability and a heightened focus of attention and self-agency. This ability is a prerequisite for performance that will be less prone to interference or distraction. The better one attends, the more aware one can be of the object of attention.

ANTERIOR AXONAL SHEARING AND SYMPTOMS

The preceding discussion outlines a general model in which well-timed predictions can yield attentional control that results in decreased IIV and reduced distractibility, and consequently, improved performance and awareness. This predictive process may be selectively disrupted in TBI due to the rotational shearing vulnerability of white matter tracts that support this preparatory activity. In addition, compensation for variable predictive timing by increased prefrontal cortical recruitment may explain many of the “secondary” post-concussive symptoms of TBI.

Diffuse Axonal Injury

Diffuse axonal injury (DAI), also known as traumatic axonal injury when lesions are more focal, resulting from rotational shear forces^{65–67} is common in TBI⁶⁸ and may account for persistent cognitive deficits and symptoms.^{69,70} This pattern of injury, distinguished from focal contusion without shearing, is characterized by damage to anterior axons at the gray/white matter junction of the cerebral hemispheres, corpus callosum, and in severe cases, dorsolateral midbrain. These patterns of shearing are often accompanied by similar damage in the superior cerebellar peduncles.^{71,72} Deep white matter abnormalities on magnetic resonance imaging (MRI) are associated with poorer neuropsychological test performances and poor long-term outcome.⁷³ The severity of deep white matter tract injury may explain many TBI-induced cognitive deficits, ranging from mild concussion to the vegetative state.^{74,75}

The normal CT or MRI scans observed in the majority of cases of mild TBI may belie microscopic axonal pathology. The sensitivity of most MRI structural studies is insufficient to detect white matter tract disruption or small areas of edema. Postmortem analysis, when conducted within a few hours of severe TBI, shows extensive microscopic hemorrhages and severing of nerve fibers in pons, midbrain, and corpus callosum, consistent with shearing forces.⁷⁶ Even in TBI cases classified as mild (in which the patients died of other causes), white matter microhemorrhages in the brain stem, cerebral hemispheres, and corpus callosum are evident.⁷⁷

DAI and Deficits in Attention and Working Memory

MRI diffusion tensor imaging (DTI) uses the characteristics of water diffusion to assess the integrity of white matter pathways.⁷⁸ This method has been used to better quantify the extent of DAI damage^{79–82} and clinical outcome¹¹ following TBI. Deficits in processing speed, working memory, and attention have been attributed to DAI.^{83,84} In addition, DAI patients have broader cortical activation in working memory tasks.⁸⁵ More recent studies suggest that DAI can cause chronic post-concussive symptoms in executive function and memory dysfunction.^{69,81,86–88} Compared to conventional structural studies, DTI appears to be more sensitive and accurate⁸⁹ in correlating pathology with attention and memory deficits (Figure 4).^{90,111} These observations suggest that white matter shear injury in TBI plays a prominent causal role in the observed cognitive deficits.

As revealed by DTI, the predominant sites for mild TBI axonal shearing are located in anterior white matter tracts connecting PFC with posterior cortical regions and subcortical structures.^{91,111} Even in normal patients, DTI imaging shows a significant correlation between working memory function and maturation of white matter tract connectivity between the PFC and the parietal lobe.⁹² PFC and IPL have reciprocal connections with the cerebellum with input passing through the internal capsule and pons and output through the superior cerebellar peduncle and thalamus. Shearing anywhere along this pathway could disrupt preparatory neural processes, as well as comparison processes in which these predictive representations are compared to sensory input.

The diffuse character of TBI has traditionally discouraged rigorous studies of structure-function relationships. However, recent results from our research group report a significant dissociation between cognitive impairments in TBI individuals with predominant pathology in the anterior corona radiata or uncinate fasciculus. The former connects the lateral PFC with other cortical areas, thalamus, and cerebellum; the latter serves as a primary pathway between the lateral PFC and parahippocampal cortex. Patients with prominent shearing in the uncinate fasciculus exhibited deficits in working memory tasks. In contrast, patients with prominent shearing in the anterior corona radiata exhibited deficits in tasks of executive attention.¹¹¹ Others have reported significant DTI changes in tracts that support working memory in TBI patients.⁹⁰ The direction of rotational shear forces at the time of TBI may lead to shearing in certain pathways due to differences in tract orientation.

TBI Symptoms: Predictive Timing Deficits and Compensation?

The most common postconcussive symptoms encountered after TBI^{10,13,93} are traditionally grouped into 3 categories: cognitive, somatic complaints, and affective complaints (Table 2). Cognitive function is a major predictor of poor outcome in TBI patients.⁹⁴ In particular, working memory is vulnerable^{95,96} and likely underlies many of the prominent problems associated with daily life activities that require planning and problem solving.⁹⁷ Similar neuropsychological deficiencies are seen in children with TBI.⁹⁸ Some symptoms of TBI may result from damage to core cognitive processes, whereas others may be due to compensatory mechanisms recruited to minimize the ensuing performance variability.

The performance of individuals with TBI is especially vulnerable to tasks that tax attention, such as when task-relevant and task-irrelevant stimuli are intermixed,⁹⁶ or if there is an increase in task difficulty. Activation patterns in chronic patients with TBI tend to be abnormal on tasks of executive function. These abnormalities may be manifest as more diffuse patterns of cortical recruitment⁸⁵ or as greater increases in prefrontal activity despite performance levels that are below that observed in controls.⁹⁹ Even when performance is matched to controls, TBI patients with diffuse axonal injury exhibit elevated pre-frontal activation on working memory tasks (Figure 5).¹⁰⁰ These findings of increased cortical activation during cognitive tasks in TBI patients is similar to those reported for aging,¹⁰¹ stroke,¹⁰² and Alzheimer's disease.¹⁰³ These effects have been related to increased variability in reaction time⁴⁵ and inconsistency in performance.¹⁰⁴

Could the increase in cortical recruitment in TBI reflect a compensatory mechanism arising because anticipatory neural activity is disrupted? Consider an analogy to cerebellar dysmetria. This disorder results in increased cortical control to correct trajectory deviations, a compensatory process that can correct for impairments in the feed-forward processes dependent on the cerebellum.¹⁰⁵ An increase in prefrontal activity may be a response to the disruption of predictive mechanisms for cognition, perhaps triggered by an increase in performance variability and errors in performance. Evidence in support of this hypothesis comes from a developmental fMRI study of working memory. Compared to adults, children exhibited weaker PFC-IPL coherence and greater activation of PFC during the delay period. These neural changes were accompanied by higher performance variability and distractibility.¹⁰⁶ Similar studies need to be done in TBI patients to investigate if a similar

dissociation is observed between measures of PFC-IPL-cerebellar coherence and local activation levels.

Classifying symptoms based on the framework outlined in this article would posit that postconcussive symptoms encountered after a TBI could be grouped into primary and secondary causes (Table 3). Primary symptoms arise from TBI axonal shearing in white matter tracts connecting the PFC-IPL-cerebellar network leading to decreased PFC-IPL coherence (Figure 6). The resulting increase in variability would be manifest as increased distractibility, working memory deficits, and problems with balance and coordination. An inability to anticipate sensory events, resulting in a temporal mismatch of sensory expectation to actual sensory input, could lead to dizziness, tinnitus, and sensory hypersensitivity. Self-agency would be abnormal due to the desynchronization between actions and their resulting consequences.

Secondary symptoms might be a consequence of heightened activation of the PFC. This might arise from an increase in error signals and performance variability. PFC recruitment, serving as a compensatory mechanism, could help bridge the moment-to-moment temporal discrepancies. The expended effort might underlie fatigue, headache, irritability, anxiety, and when prolonged, depression.

Variability As a Metric of Prediction

If a consequence of accurate anticipation is a reduction in performance variability, then variability can be used as a metric to assess this aspect of attention. By definition, attention varies over time. Relatively continuous and extended measures of performance should be used to gauge moment-to-moment fluctuations in performance within individuals. Tasks used to assess IIV have typically involved relatively discrete measures such as reaction time,^{25,45} allowing measurements, at best, every few seconds. Such measures may be relatively insensitive to lapses in attention.

We have employed a task in which we measure circular smooth pursuit eye movements during a predictive tracking task, lasting 2.5 seconds using a high-resolution camera sampling performance at 500 Hz over blocks of 20 trials. The predictive nature of this task engages the PFC-parietal-cerebellar network, similar to that observed in studies of spatial attention. The variability in the difference between eye and target position, as well as velocity error (the difference between target velocity and eye velocity), both correlate with performance on a working memory task.^{107,108} In addition, TBI patients with chronic cognitive symptoms had higher variability and a persistent lag in eye position compared to control individuals.

Further studies have demonstrated that adding a concurrent secondary task (short-term verbal memory) during the smooth pursuit tracking task produces different effects on velocity error variability in control and TBI patients. In TBI patients, there is a significant increase in variability as the difficulty of the secondary task increases. In contrast, control individuals showed a decrease in velocity error variability when the length of the word list was increased, a counterintuitive finding, yet one that is also observed in other attention-loading studies with normal individuals (Ghajar J, Suh M, Kolster RA, et al. Smooth pursuit

eye movement variability under dual-task load predicts white matter shear injury in mild TBI patients. Society for Neuroscience 37th Annual Meeting. San Diego 2007) (Figure 7).¹⁰⁹ We suggest that the secondary task results in increased recruitment of the PFC, with the consequence of greater coherence across the prefrontal-parietal-cerebellar network required for tracking. In TBI, a similar increase in PFC activity^{85,100,110} can support secondary task performance but fails to result in benefits on the tracking task. Indeed, by diverting resources from the tracking task, performance becomes more variable.

SUMMARY

The inability to efficiently and consistently use predictive mechanisms, even with increased prefrontal cortical activation, is posited to be a major cause of disability in TBI, and possibly other disorders of attention. Damage to a prefrontal-parietal-cerebellum network, both from direct neural insult and shearing of the white matter tracts linking these areas, impinges on this predictive capability. The cerebellum is posited to be involved in producing the requisite timing required to match predictions and the consequences of actions. In the absence of this timing, the PFC must be reactive in integrating expectancies and input. A breakdown in anticipatory behavior will result in an increase in performance variability and a host of profound disturbances of cognition, some of which are the direct result of the pathology and some of which arise as part of compensatory processes.

Current accounts of TBI that emphasize a loss of top-down control of attention dependent on the prefrontal cortex are insufficient to explain the deficits in TBI. A perspective that focuses on how the brain generates predictive states through the interaction of the prefrontal and parietal cortex, with feed-forward timing from the cerebellum, may offer a more complete account. Presently, there are not strong data to support either model. However, the predictive brain state model offers a testable scientific construct, as well as a framework that can be useful in diagnostic and therapeutic interventions that emphasize individual variability in performance.

Acknowledgments

We thank Susan Fitzpatrick, PhD, for helpful comments on the manuscript and Akiyo Kadera for assistance in preparation of the manuscript. Supported by the Rachel L. Mellon TBI Research Fund and a collaborative grant from the James S. McDonnell Foundation to the Brain Trauma Foundation, with lead investigators from Weill Medical College of Cornell University, University of California, San Francisco, and University of California, Berkeley.

References

1. Guyau, J-M. La genèse de l'idée de temps. Paris: Félix Alcan; 1890.
2. Ghajar J. Traumatic brain injury. *Lancet*. 2000; 356:923–929. [PubMed: 11036909]
3. Langlois JA, Rutland-Brown W, Wald MM. The epidemiology and impact of traumatic brain injury: a brief overview. *J Head Trauma Rehabil*. 2006; 21:375–378. [PubMed: 16983222]
4. Fife D. Head injury with and without hospital admission: comparisons of incidence and short-term disability. *Am J Public Health*. 1987; 77:810–812. [PubMed: 2954475]
5. Goldberg LD, Dimeff RJ. Sideline management of sport-related concussions. *Sports Med Arthrosc*. 2006; 14:199–205. [PubMed: 17135969]
6. Taber KH, Warden DL, Hurley RA. Blast-related traumatic brain injury: what is known? *J Neuropsychiatry Clin Neurosci*. 2006; 18:141–145. [PubMed: 16720789]

7. Warden D. Military TBI during the Iraq and Afghanistan wars. *J Head Trauma Rehabil.* 2006; 21:398–402. [PubMed: 16983225]
8. Stein SC, Burnett MG, Glick HA. Indications for CT scanning in mild traumatic brain injury: a cost-effectiveness study. *J Trauma.* 2006; 61:558–566. [PubMed: 16966987]
9. Dikmen SS, Ross BL, Machamer JE, Temkin NR. One year psychosocial outcome in head injury. *J Int Neuropsychol Soc.* 1995; 1:67–77. [PubMed: 9375211]
10. Dikmen S, McLean A, Temkin N. Neuropsychological and psychosocial consequences of minor head injury. *J Neurol Neurosurg Psychiatry.* 1986; 49:1227–1232. [PubMed: 3794728]
11. Schaefer PW, Huisman TA, Sorensen AG, Gonzalez RG, Schwamm LH. Diffusion-weighted MR imaging in closed head injury: high correlation with initial Glasgow coma scale score and score on modified Rankin scale at discharge. *Radiology.* 2004; 233:58–66. [PubMed: 15304663]
12. Huisman TA, Schwamm LH, Schaefer PW, et al. Diffusion tensor imaging as potential biomarker of white matter injury in diffuse axonal injury. *AJNR Am J Neuroradiol.* 2004; 25:370–376. [PubMed: 15037457]
13. Alexander MP. Mild traumatic brain injury: pathophysiology, natural history, and clinical management. *Neurology.* 1995; 45:1253–1260. [PubMed: 7617178]
14. Concussion (mild traumatic brain injury) and the team physician: a consensus statement. *Med Sci Sports Exerc.* 2006; 38:395–399. [PubMed: 16531912]
15. Scheibel RS, Newsome M, Steinberg J, Pearson D, Rauch R, Mao H, Levin H. Altered brain activation during cognitive control in patients with moderate to severe traumatic brain injury. *Neurorehabil Neural Repair.* 2007; 21:36–45. [PubMed: 17172552]
16. Knight RT. Neuroscience. Neural networks debunk phrenology. *Science.* 2007; 316:1578–1579. [PubMed: 17569852]
17. Posner MI, Walker JA, Friedrich FJ, Rafal RD. Effects of parietal injury on covert orienting of attention. *J Neurosci.* 1984; 4:1863–1874. [PubMed: 6737043]
18. Thiel CM, Zilles K, Fink GR. Cerebral correlates of alerting, orienting and reorienting of visuospatial attention: an event-related fMRI study. *Neuroimage.* 2004; 21:318–328. [PubMed: 14741670]
19. Naghavi HR, Nyberg L. Common fronto-parietal activity in attention, memory, and consciousness: shared demands on integration? *Conscious Cogn.* 2005; 14:390–425. [PubMed: 15950889]
20. Kim SG, Ugurbil K, Strick PL. Activation of a cerebellar output nucleus during cognitive processing. *Science.* 1994; 265:949–951. [PubMed: 8052851]
21. Tomasi D, Ernst T, Caparelli EC, Chang L. Practice-induced changes of brain function during visual attention: a parametric fMRI study at 4 Tesla. *Neuroimage.* 2004; 23:1414–1421. [PubMed: 15589105]
22. Macar F, Anton JL, Bonnet M, Vidal F. Timing functions of the supplementary motor area: an event-related fMRI study. *Brain Res Cogn Brain Res.* 2004; 21:206–215. [PubMed: 15464352]
23. Cunnington R, Windischberger C, Deecke L, Moser E. The preparation and readiness for voluntary movement: a high-field event-related fMRI study of the Bereitschafts-BOLD response. *Neuroimage.* 2003; 20:404–412. [PubMed: 14527600]
24. Walter WG, Cooper R, Aldridge VJ, McCallum WC, Winter AL. Contingent negative variation: an electric sign of sensorimotor association and expectancy in the human brain. *Nature.* 1964; 203:380–384. [PubMed: 14197376]
25. Segalowitz SJ, Dywan J, Unsal A. Attentional factors in response time variability after traumatic brain injury: an ERP study. *J Int Neuropsychol Soc.* 1997; 3:95–107. [PubMed: 9126851]
26. Di Russo F, Incoccia C, Formisano R, Sabatini U, Zoccolotti P. Abnormal motor preparation in severe traumatic brain injury with good recovery. *J Neurotrauma.* 2005; 22:297–312. [PubMed: 15716635]
27. Kitamura J, Shabasaki H, Terashi A, Tashima K. Cortical potentials preceding voluntary finger movement in patients with focal cerebellar lesion. *Clin Neurophysiol.* 1999; 110:126–132. [PubMed: 10348331]
28. Paradiso G, Cunic D, Chen R. Involvement of subcortical structures in the preparation of self-paced movement. *J Psychophysiol.* 2004; 18:130–139.

29. Padilla ML, Wood RA, Hale LA, Knight RT. Lapses in a pre-frontal-extrastriate preparatory attention network predict mistakes. *J Cogn Neurosci*. 2006; 18:1477–1487. [PubMed: 16989549]
30. Nagai Y, Critchley HD, Featherstone E, Fenwick PB, Trimble MR, Dolan RJ. Brain activity relating to the contingent negative variation: an fMRI investigation. *Neuroimage*. 2004; 21:1232–1241. [PubMed: 15050551]
31. Knight RT. Distributed cortical network for visual attention. *J Cogn Neurosci*. 1997; 9:75–91. [PubMed: 23968181]
32. Wiese H, Stude P, Nebel K, Forsting M, de Greiff A. Prefrontal cortex activity in self-initiated movements is condition-specific, but not movement-related. *Neuroimage*. 2005; 28:691–697. [PubMed: 16054840]
33. Dosenbach NU, Fair DA, Miezin FM, et al. Distinct brain networks for adaptive and stable task control in humans. *Proc Natl Acad Sci U S A*. 2007; 104:11073–11078. [PubMed: 17576922]
34. Manly T, Owen AM, McAvinue L, et al. Enhancing the sensitivity of a sustained attention task to frontal damage: convergent clinical and functional imaging evidence. *Neurocase*. 2003; 9:340–349. [PubMed: 12925947]
35. LaBar KS, Gitelman DR, Parrish TB, Mesulam M. Neuroanatomic overlap of working memory and spatial attention networks: a functional MRI comparison within subjects. *Neuroimage*. 1999; 10:695–704. [PubMed: 10600415]
36. Sapir A, d'Avossa G, McAvoy M, Shulman GL, Corbetta M. Brain signals for spatial attention predict performance in a motion discrimination task. *Proc Natl Acad Sci U S A*. 2005; 102:17810–17815. [PubMed: 16306268]
37. Pessoa L, Gutierrez E, Bandettini P, Ungerleider L. Neural correlates of visual working memory: fMRI amplitude predicts task performance. *Neuron*. 2002; 35:975–987. [PubMed: 12372290]
38. Dum RP, Strick PL. An unfolded map of the cerebellar dentate nucleus and its projections to the cerebral cortex. *J Neurophysiol*. 2003; 89:634–639. [PubMed: 12522208]
39. Middleton FA, Strick PL. Cerebellar projections to the prefrontal cortex of the primate. *J Neurosci*. 2001; 21:700–712. [PubMed: 11160449]
40. Blakemore SJ, Sirigu A. Action prediction in the cerebellum and in the parietal lobe. *Exp Brain Res*. 2003; 153:239–245. [PubMed: 12955381]
41. Mauk MD, Medina JF, Nores WL, Ohyama T. Cerebellar function: coordination, learning or timing? *Curr Biol*. 2000; 10:R522–R525. [PubMed: 10898992]
42. Mehta B, Schaal S. Forward models in visuomotor control. *J Neurophysiol*. 2002; 88:942–953. [PubMed: 12163543]
43. Volker, S.; Gross, H-M. Neural anticipative architecture for expectation driven perception. *Proceedings of the 2001 IEEE Systems, Man, and Cybernetics Conference*; 2001;
44. MacDonald SW, Nyberg L, Backman L. Intra-individual variability in behavior: links to brain structure, neurotransmission and neuronal activity. *Trends Neurosci*. 2006; 29:474–480. [PubMed: 16820224]
45. Stuss DT, Pogue J, Buckle L, Bondar J. Characterization of stability of performance in patients with traumatic brain injury: variability and consistency on reaction time test. *Neuropsychologia*. 1994; 8:316–324.
46. Collins LF, Long CJ. Visual reaction time and its relationship to neuropsychological test performance. *Arch Clin Neuropsychol*. 1996; 11:613–623. [PubMed: 14588445]
47. Dockree PM, Bellgrove MA, O'Keeffe FM, et al. Sustained attention in traumatic brain injury and healthy controls: enhanced sensitivity with dual-task load. *Exp Brain Res*. 2006; 168:218–219. [PubMed: 16044297]
48. Hetherington CR, Stuss DT, Finlayson MA. Reaction time and variability 5 and 10 years after traumatic brain injury. *Brain Inj*. 1996; 10:473–486. [PubMed: 8806008]
49. Mangels JA, Craik FI, Levine B, Schwartz ML, Stuss DT. Effects of divided attention on episodic memory in chronic traumatic brain injury: a function of severity and strategy. *Neuropsychologia*. 2002; 40:2369–2385. [PubMed: 12417466]
50. Whyte J, Polansky M, Fleming M, Coslett HB, Cavallucci C. Sustained arousal and attention after traumatic brain injury. *Neuropsychologia*. 1995; 33:797–813. [PubMed: 7477808]

51. Castellanos FX, Sonuga-Barke EJ, Scheres A, Di Martino A, Hyde C, Walters JR. Varieties of attention-deficit/hyperactivity disorder-related intra-individual variability. *Biol Psychiatry*. 2005; 57:1416–1423. [PubMed: 15950016]
52. Klein C, Wendling K, Huettner P, Ruder H, Peper M. Intra-subject variability in attention-deficit hyperactivity disorder. *Biol Psychiatry*. 2006; 60:1088–1097. [PubMed: 16806097]
53. Hultsch DF, MacDonald SW, Dixon RA. Variability in reaction time performance of younger and older adults. *J Gerontol B Psychol Sci Soc Sci*. 2002; 57:P101–P115. [PubMed: 11867658]
54. Martin M, Hofer SM. Intraindividual variability, change, and aging: conceptual and analytical issues. *Gerontology*. 2004; 50:7–11. [PubMed: 14654720]
55. Nesselroade JR, Salthouse TA. Methodological and theoretical implications of intraindividual variability in perceptual-motor performance. *J Gerontol B Psychol Sci Soc Sci*. 2004; 59:P49–P55. [PubMed: 15014087]
56. Gorus E, De Raedt R, Mets T. Diversity, dispersion and inconsistency of reaction time measures: effects of age and task complexity. *Aging Clin Exp Res*. 2006; 18:407–417. [PubMed: 17167305]
57. Strauss E, MacDonald SW, Hunter M, Moll A, Hultsch DF. Intraindividual variability in cognitive performance in three groups of older adults: cross-domain links to physical status and self-perceived affect and beliefs. *J Int Neuropsychol Soc*. 2002; 8:893–906. [PubMed: 12405540]
58. MacDonald SW, Hultsch DF, Dixon RA. Performance variability is related to change in cognition: evidence from the Victoria Longitudinal Study. *Psychol Aging*. 2003; 18:510–523. [PubMed: 14518812]
59. Jeannerod M. The mechanism of self-recognition in humans. *Behav Brain Res*. 2003; 142:1–15. [PubMed: 12798261]
60. Blakemore SJ, Frith CD, Wolpert DM. Spatio-temporal prediction modulates the perception of self-produced stimuli. *J Cogn Neurosci*. 1999; 11:551–559. [PubMed: 10511643]
61. Posada A, Franck N, Georgieff N, Jeannerod M. Anticipating incoming events: an impaired cognitive process in schizophrenia. *Cognition*. 2001; 81:209–225. [PubMed: 11483170]
62. Schmitz C, Martineau J, Barthélemy C, Assaiante C. Motor control and children with autism: deficit of anticipatory function? *Neurosci Lett*. 2003; 348:17–20. [PubMed: 12893415]
63. Anderson B, Mennemeier M, Chatterjee A. Variability not ability: another basis for performance decrements in neglect. *Neuropsychologia*. 2000; 38:785–796. [PubMed: 10689054]
64. Prigatano GP. Disturbances of self-awareness and rehabilitation of patients with traumatic brain injury: a 20-year perspective. *J Head Trauma Rehabil*. 2005; 20:19–29. [PubMed: 15668568]
65. Gennarelli TA, Graham DI. Neuropathology of the head injuries. *Semin Clin Neuropsychiatry*. 1998; 3:160–175. [PubMed: 10085204]
66. Povlishock JT, Katz DI. Update of neuropathology and neurological recovery after traumatic brain injury. *J Head Trauma Rehabil*. 2005; 20:76–94. [PubMed: 15668572]
67. Smith DH, Meaney DF, Shull WH. Diffuse axonal injury in head trauma. *J Head Trauma Rehabil*. 2003; 18:307–316. [PubMed: 16222127]
68. Medana IM, Esiri MM. Axonal damage: a key predictor of outcome in human CNS diseases. *Brain*. 2003; 126:515–530. [PubMed: 12566274]
69. Salmond CH, Menon DK, Chatfield DA, et al. Diffusion tensor imaging in chronic head injury survivors: correlations with learning and memory indices. *Neuroimage*. 2006; 29:117–124. [PubMed: 16084738]
70. Schmithorst VJ, Wilke M, Dardzinski BJ, Holland SK. Cognitive functions correlate with white matter architecture in a normal pediatric population: a diffusion tensor MRI study. *Hum Brain Mapp*. 2005; 26:139–147. [PubMed: 15858815]
71. Gennarelli TA, Thibault LE, Adams JH, Graham DI, Thompson CJ, Marcincin RP. Diffuse axonal injury and traumatic coma in the primate. *Ann Neurol*. 1982; 12:564–574. [PubMed: 7159060]
72. Raghupathi R, Margulies SS. Traumatic axonal injury after closed head injury in the neonatal pig. *J Neurotrauma*. 2002; 19:843–853. [PubMed: 12184854]
73. Wilson JT, Pettigrew LE, Teasdale GM. Structured interviews for the Glasgow Outcome Scale and the extended Glasgow Outcome Scale: guidelines for their use. *J Neurotrauma*. 1998; 15:573–585. [PubMed: 9726257]

74. Ross BL, Temkin NR, Newell D, Dikmen SS. Neuropsychological outcome in relation to head injury severity. Contributions of coma length and focal abnormalities. *Am J Phys Med Rehabil.* 1994; 73:341–347. [PubMed: 7917164]
75. Kampfl A, Schmutzhard E, Franz G, et al. Prediction of recovery from post-traumatic vegetative state with cerebral magnetic-resonance imaging. *Lancet.* 1998; 351:1763–1767. [PubMed: 9635948]
76. Oppenheimer DR. Microscopic lesions in the brain following head injury. *J Neurol Neurosurg Psychiatry.* 1968; 31:299–306. [PubMed: 4176429]
77. Blumbergs PC, Scott G, Manavis J, Wainwright H, Simpson DA, McLean AJ. Staining of amyloid precursor protein to study axonal damage in mild head injury. *Lancet.* 1994; 344:1055–1056. [PubMed: 7523810]
78. Basser PJ, Pierpaoli C. Microstructural and physiological features of tissues elucidated by quantitative-diffusion-tensor MRI. *J Magn Reson B.* 1996; 111:209–219. [PubMed: 8661285]
79. Arfanakis K, Haughton VM, Carew JD, Rogers BP, Dempsey RJ, Meyerand ME. Diffusion tensor MR imaging in diffuse axonal injury. *AJNR Am J Neuroradiol.* 2002; 23:794–802. [PubMed: 12006280]
80. Huisman TA. Diffusion-weighted imaging: basic concepts and application in cerebral stroke and head trauma. *Eur Radiol.* 2003; 13:2283–2297. [PubMed: 14534804]
81. Huisman TA, Schwamm LH, Schaefer PW, et al. Diffusion tensor imaging as potential biomarker of white matter injury in diffuse axonal injury. *AJNR Am J Neuroradiol.* 2004; 25:370–376. [PubMed: 15037457]
82. Lee ZI, Byun WM, Jang SH, Ahn SH, Moon HK, Chang Y. Diffusion tensor magnetic resonance imaging of microstructural abnormalities in children with brain injury. *Am J Phys Med Rehabil.* 2003; 82:556–559. [PubMed: 12819543]
83. Wallesch CW, Curio N, Galazky I, Jost S, Synowitz H. The neuropsychology of blunt head injury in the early postacute stage: effects of focal lesions and diffuse axonal injury. *J Neurotrauma.* 2001; 18:11–20. [PubMed: 11200246]
84. Wallesch CW, Curio N, Kutz S, Jost S, Bartels C, Synowitz H. Outcome after mild-to-moderate blunt head injury: effects of focal lesions and diffuse axonal injury. *Brain Inj.* 2001; 15:401–412. [PubMed: 11350654]
85. Christodoulou C, DeLuca J, Ricker JH, et al. Functional magnetic resonance imaging of working memory impairment after traumatic brain injury. *J Neurol Neurosurg Psychiatry.* 2001; 71:161–168. [PubMed: 11459886]
86. Fork M, Bartels C, Ebert AD, Grubich C, Synowitz H, Wallesch CW. Neuropsychological sequelae of diffuse traumatic brain injury. *Brain Inj.* 2005; 19:101–108. [PubMed: 15841754]
87. Himanen L, Portin R, Isoniemi H, Helenius H, Kurki T, Tenovuo O. Cognitive functions in relation to MRI findings 30 years after traumatic brain injury. *Brain Inj.* 2005; 19:93–100. [PubMed: 15841753]
88. Himanen L, Portin R, Isoniemi H, Helenius H, Kurki T, Tenovuo O. Longitudinal cognitive changes in traumatic brain injury: a 30-year follow-up study. *Neurology.* 2006; 66:187–192. [PubMed: 16434651]
89. Mac Donald CL, Dikranian K, Song SK, Bayly PV, Holtzman DM, Brody DL. Detection of traumatic axonal injury with diffusion tensor imaging in a mouse model of traumatic brain injury. *Exp Neurol.* 2007; 205:116–131. [PubMed: 17368446]
90. Salmond CH, Menon DK, Chatfield DA, et al. Diffusion tensor imaging in chronic head injury survivors: correlations with learning and memory indices. *Neuroimage.* 2005; 29:117–124. [PubMed: 16084738]
91. Inglese M, Makani S, Johnson G, et al. Diffuse axonal injury in mild traumatic brain injury: a diffusion tensor imaging study. *J Neurosurg.* 2005; 103:298–303. [PubMed: 16175860]
92. Olesen PJ, Nagy Z, Westerberg H, Klingberg T. Combined analysis of DTI and fMRI data reveals a joint maturation of white and grey matter in a fronto-parietal network. *Brain Res Cogn Brain Res.* 2003; 18:48–57. [PubMed: 14659496]
93. Martinez P, Richters JE. The NIMH community violence project: II. Children's distress symptoms associated with violence exposure. *Psychiatry.* 1993; 56:22–35. [PubMed: 8488209]

94. Kaplan CP, Corrigan JD. The relationship between cognition and functional independence in adults with traumatic brain injury. *Arch Phys Med Rehabil.* 1994; 75:643–647. [PubMed: 8002762]
95. Levin HS, Gary HE Jr, Eisenberg HM, et al. Neurobehavioral outcome 1 year after severe head injury. Experience of the Traumatic Coma Data Bank. *J Neurosurg.* 1990; 73:699–709. [PubMed: 2213159]
96. Stuss DT, Ely P, Hugenholtz H, et al. Subtle neuropsychological deficits in patients with good recovery after closed head injury. *Neurosurgery.* 1985; 17:41–47. [PubMed: 4022286]
97. Drake AI, Gray N, Yoder S, Pramuka M, Llewellyn M. Factors predicting return to work following mild traumatic brain injury: a discriminant analysis. *J Head Trauma Rehabil.* 2000; 15:1103–1112. [PubMed: 10970931]
98. Ewing-Cobbs L, Fletcher JM, Levin HS, Francis DJ, Davidson K, Miner ME. Longitudinal neuropsychological outcome in infants and preschoolers with traumatic brain injury. *J Int Neuropsychol Soc.* 1997; 3:581–591. [PubMed: 9448371]
99. Levine B, Cabeza R, McIntosh AR, Black SE, Grady CL, Stuss DT. Functional reorganisation of memory after traumatic brain injury: a study with H(2)(15)O positron emission tomography. *J Neurol Neurosurg Psychiatry.* 2002; 73:173–181. [PubMed: 12122177]
100. Maruishi M, Miyatani M, Nakao T, Muranaka H. Compensatory cortical activation during performance of an attention task by patients with diffuse axonal injury: a functional magnetic resonance imaging study. *J Neurol Neurosurg Psychiatry.* 2007; 78:168–173. [PubMed: 16952916]
101. Cabeza R, Grady CL, Nyberg L, et al. Age-related differences in neural activity during memory encoding and retrieval: a positron emission tomography study. *J Neurosci.* 1997; 17:391–400. [PubMed: 8987764]
102. Lotze M, Grodd W, Rodden F, Schonle P, Katdatzke B, Cohen LG. Neuroimaging patterns associated with motor control in traumatic brain injury. *Neurorehabil Neural Repair.* 2006; 20:14–23. [PubMed: 16467275]
103. Woodard JL, Grafton ST, Votaw JR, Green RC, Dobraski ME, Hoffman JM. Compensatory recruitment of neural resources during overt rehearsal of word lists in Alzheimer's disease. *Neuropsychology.* 1998; 12:491–504. [PubMed: 9805319]
104. Bleiberg J, Garmoe WS, Halpern EL, Reeves DL, Nadler JD. Consistency of within-day and across-day performance after mild brain injury. *Neuropsychiatr Neuropsychol Behav Neurol.* 1997; 10:247–253.
105. Timmann D, Citron R, Watts S, Hore J. Increased variability in finger position occurs throughout overarm throws made by cerebellar and unskilled subjects. *J Neurophysiol.* 2001; 86:2690–2702. [PubMed: 11731529]
106. Olesen PJ, Macoveanu J, Tegner J, Klingberg T. Brain activity related to working memory and distraction in children and adults. *Cereb Cortex.* 2007; 17:1047–1054. [PubMed: 16801377]
107. Suh M, Kolster R, Sarkar R, McCandliss B, Ghajar J. Deficits in predictive smooth pursuit after mild traumatic brain injury. *Neurosci Lett.* 2006; 401:108–113. [PubMed: 16554121]
108. Suh M, Basu S, Kolster R, Sarkar R, McCandliss B, Ghajar J. Increased oculomotor deficits during target blanking as an indicator of mild traumatic brain injury. *Neurosci Lett.* 2006; 410:203–207. [PubMed: 17055156]
109. Stuss DT, Murphy KJ, Binns MA, Alexander MP. Staying on the job: the frontal lobes control individual performance variability. *Brain.* 2003; 126:2363–2380. [PubMed: 12876148]
110. McAllister TW, Saykin AJ, Flashman LA, et al. Brain activation during working memory 1 month after mild traumatic brain injury: a functional MRI study. *Neurology.* 1999; 53:1300–1308. [PubMed: 10522888]
111. Niogi SN, Mukherjee P, Ghajar J, Johnson C, Kolster RA, Sarkar R, Lee H, Meeker M, Zimmerman RD, Manley GT, McCandliss BD. Extent of Microstructural White Matter Injury in Postconcussive Syndrome Correlates with Impaired Cognitive Reaction Time: A 3T Diffusion Tensor Imaging Study of Mild Traumatic Brain Injury. *AJNR Am J Neuroradiol.* 2008 Feb 26. Epub ahead of print.

Figure 1.

Schematic of predictive brain state network. Predictive neural network that is activated in cued, learned attention tasks in the time period before task presentation. The sensory cue (CUE), with a visual cue as an example, activates cerebellum (Cblm) to delay motor or cognitive action to coincide with onset of the cued task. This learned delay, mediated by the cerebellum, synchronizes expectancy with actual sensory input. The timed expectancy is manifested by a mainly right-sided coherence (large gray arrowheads) between the dorsolateral prefrontal cortex (PFC) and inferior parietal lobe (IPL).

Figure 2.

Movement-related cortical potentials in control group and severe traumatic brain injury (TBI) group with good recovery before voluntary movement of left finger. BP is the readiness potential that appears 2.5 seconds before movement in normal individuals but is absent in TBI patients (NS is the rapid negative slope of the BP). MP = Movement potential; RAP = Reafferent positivity. Reproduced from Di Russo F, Incoccia C, Formisano R, Sabatini U, Zoccolotti P. Abnormal motor preparation in severe traumatic brain injury with good recovery. *J Neurotrauma*. 2005;22:297–312. The publisher for this copyrighted material is Mary Ann Liebert, Inc, publishers.

Figure 3.

Diagram showing how anticipatory timing can facilitate performance in a simple cueing task when the time of a stimulus occurs after a known interval. Predictive timing allows the participant to anticipate the stimulus and generate an expectancy of the resulting feedback. Accurate predictive timing ensures synchronization of the predictions, in this example, generated in anticipation of the “Go” stimulus and feedback from the response.

Figure 4.

Magnetic resonance imaging–diffusion tensor imaging images in a normal control individual and a mild traumatic brain injury (TBI) patient. The patient shows reduced white matter integrity in the right anterior corona radiata (ACR) shown by the circled region, compared to the normal individual's circled left and right ACR. Courtesy of Niogi S, Weill Medical College of Cornell University.

Figure 5.

Compensatory right prefrontal cortex (PFC) activation in traumatic brain injury (TBI) patients in an attention-demanding task. A) Target-related activation in the frontal and parietal areas in normal individuals. B) Target-related activation in TBI patients with pure diffuse axonal injury (DAI) injury. The TBI patient shows target-related activation in the right PFC as well as increased activation in the left PFC. Target-related parietal activity is not evident in the TBI group. Reproduced from Maruishi M, Miyatani M, Nakao T, Muranaka H. Compensatory cortical activation during performance of an attention task by patients with diffuse axonal injury: a functional magnetic resonance imaging study. *J Neurol Neurosurg Psychiatry*. 2007;78:168–173. The publisher for this copyrighted material is BMJ Publishing Group Ltd.

Figure 6.
Figure showing predominant focal areas of diffuse axonal injury (DAI).

Figure 7.

Variability of tracking velocity error (VE) during circular smooth pursuit task. The task was performed under 4 levels of attentional load in a task requiring rehearsal of a list of 0, 1, 5, or 9 words during the 30-second tracking trial. Traumatic brain injury (TBI) patients show an increase in variability with increases in demands on working memory, whereas controls become less variable over the lower levels of increased load. Reproduced from Ghajar J, Suh M, Kolster RA, et al. Smooth pursuit eye movement variability under dual-task load predicts white matter shear injury in mild TBI patients. Paper presented at: Society for Neuroscience 37th Annual Meeting.

Table 1**Attention Network for Expectancy**

-
- Anticipatory Neural Activity: A feed-forward process that reduces performance variability by generating real-time predictions of anticipated sensory signals.
 - Predominant right hemispheric dorsolateral prefrontal cortex (PFC) and inferior parietal lobe (IPL) attention network with timing mediated by the cerebellum.
 - Neural correlates of this activity involve:
 1. Coherence of PFC-IPL to synchronize expectancy with sensory input.
 2. Feed-forward timing of PFC-IPL coherence is mediated by cerebellum.
 3. Associative capabilities of cerebellum learn appropriate time delays for prediction.
-

Table 2

Post-Concussive Symptoms Scale

Cognitive	Somatic	Affective
<ul style="list-style-type: none"> • Memory difficulty[*] • Attention deficit • Decreased concentration[*] 	<ul style="list-style-type: none"> • Headache[*] • Fatigue[*] • Insomnia • Balance and coordination problem • Dizziness[*] • Tinnitus • Sleep disturbances[*] • Sensitivity to noise or light[*] 	<ul style="list-style-type: none"> • Depression • Irritability[*] • Anxiety

^{*} Three of any of these symptoms persisting for more than 3 months defines persistent post-concussive syndrome. Alexander MP. Mild traumatic brain injury: pathophysiology, natural history, and clinical management. *Neurology*. 1995;45:1253–1260.

Table 3

Post-Concussive Symptoms

Primary Symptoms – Related to Predictive Timing Deficit	Secondary Symptoms – Related To PFC Compensation and Error Signaling
<ul style="list-style-type: none">• Decreased attention• Decreased memory• Decreased concentration• Balance and coordination problem• Decreased awareness• Dizziness• Tinnitus• Sensory sensitivity	<ul style="list-style-type: none">• Fatigue• Headache• Insomnia• Irritability• Anxiety• Depression