Published in final edited form as:

Expert Rev Mol Med. ; 12: e18. doi:10.1017/S1462399410001493.

Gene therapy for bone healing

Christopher H. Evans

Center for Advanced Orthopaedic Studies, Beth-Israel Deaconess Medical Center, Harvard Medical School, 330, Brookline Avenue, RN-115, Boston, MA 02215, USA and Collaborative Research Center, AO Foundation. cevans@bidmc.harvard.edu

Abstract

Clinical problems in bone healing include large segmental defects, nonunion and delayed union of fractures, and spinal fusions. Gene-transfer technologies have the potential to aid healing by permitting the local delivery and sustained expression of osteogenic gene products within osseous lesions. Key questions for such an approach include the choice of transgene, vector and gene-transfer strategy. Most experimental data have been obtained using cDNAs encoding osteogenic growth factors such as bone morphogenetic protein-2 (BMP-2), BMP-4 and BMP-7, in conjunction with both nonviral and viral vectors using in vivo and ex vivo delivery strategies. Proof of principle has been convincingly demonstrated in small-animal models. Relatively few studies have used large animals, but the results so far are encouraging. Once a reliable method has been developed, it will be necessary to perform detailed pharmacological and toxicological studies, as well as satisfy other demands of the regulatory bodies, before human clinical trials can be initiated. Such studies are very expensive and often protracted. Thus, progress in developing a clinically useful gene therapy for bone healing is determined not only by scientific considerations, but also by financial constraints and the ambient regulatory environment.

Despite the remarkable, intrinsic ability of bone to regenerate without scarring, there are a number of clinical conditions where healing is impaired. In approximately 10% of fractures, for example, bone healing is either very slow (delayed union) or the bone does not heal at all (nonunion). The reasons for this are unclear, although it is more common in smokers (Ref. 1), alcohol abusers (Ref. 2) and diabetics (Ref. 3). Bone healing is also an issue for the elderly (Ref. 4) and patients with osteoporosis (Ref. 5). Healing is also difficult under conditions where a large segment of bone is lost because of injury, infection or tumour resection; indeed, segmental defects beyond a certain 'critical size' never heal spontaneously, even in young, healthy individuals (Ref. 6). Problems in bone healing also arise for iatrogenic reasons; for instance, when the surgeon needs to fuse two vertebrae together as a way of treating back pain (Ref. 7). An improved ability to form bone would also be useful for the treatment of osteonecrosis and for increasing bone stock around prosthetic joints, with the possibility of preventing or treating aseptic loosening. Moreover, there are numerous potential applications in the craniofacial, orthodontic and dental areas.

The most reliable way to form bone under these conditions is by the surgical implantation of some of the patient's own living bone (autograft). This is usually harvested surgically from the iliac crest of the pelvis. Although this method has a high success rate, its usefulness is restricted by the limited amounts of bone available for autografting, and side effects,

especially pain, at the harvest site (Ref. 8). As an alternative or supplement to autograft bone, it is possible to use bone recovered from cadavers (allograft) (Ref. 9). Although allograft bone is available in almost unlimited quantities, there are concerns about disease transmission, and because processing kills endogenous cells, it is essentially dead bone. Therefore, unlike autograft, it is unable to participate actively in the formation of new bone and thus serves as an inert filler. If the allograft is implanted into areas where it needs to bear load, it frequently fails, because unlike living bone, it cannot remodel.

In recent decades, the search for new ways to promote bone healing has increasingly turned to biology (Ref. 10). In general, there are two routes through which the body forms bone: endochondral and intramembranous (Ref. 11). The former process involves the local differentiation of progenitor cells into chondrocytes that lay down a cartilaginous matrix. The cartilage is then replaced by bone through a process known as endochondral ossification, during which the cartilaginous matrix is degraded, blood vessels invade the cartilage and chondrocytes are replaced by osteoblasts. The intramembranous formation of bone involves the differentiation of progenitor cells directly into osteoblasts without a cartilaginous intermediate. Both the endochondral and intramembranous routes lead to the formation of immature, woven bone that undergoes remodeling into mature, lamellar bone (Ref. 11).

Study of osteogenesis has identified a number of growth factors with the ability to stimulate one or more of the steps involved in endochondral and intramembranous bone formation (Refs 12, 13). The best characterised of these factors are the bone morphogenetic proteins (BMPs). Several members of this large family, including BMP-2, BMP-4, BMP-6, BMP-7 and BMP-9, are able to induce the formation of bone when implanted experimentally into the muscles of laboratory animals (Ref. 14). Recombinant, human BMP-2 and BMP-7 are available clinically as the active components of the products Infuse® and OP-1® (osteogenic protein-1), respectively. Despite the powerful osteogenic properties of these proteins in laboratory animals, their clinical performance is mixed (Ref. 15). For instance, they appear to be much more effective for spinal fusion than for long-bone healing, and need to be administered at extremely high doses of several milligrammes – many orders of magnitude greater than the levels at which they occur naturally in bone. The use of such high doses not only raises safety concerns, but also enormously increases the cost. In the USA, for example, a single dose of Infuse® can cost around \$5000. There is considerable optimism that gene-therapy approaches can be harnessed to improve the effectiveness of osteogenic factors, such as BMP-2, while lowering both costs and the potential for side effects.

As its name suggests, gene therapy involves the transfer of genes, or more usually cDNA (complementary DNA), to patients for therapeutic purposes. Gene therapy was originally developed as a means of curing genetic diseases, but in recent years its potential use in treating nongenetic disorders has become increasingly appreciated and a range of nonmendelian diseases, including various cancers (Ref. 16), arthritis (Ref. 17) and Parkinson disease (Ref. 18), have been the subject of clinical trials. In the context of bone healing, the aim is to deliver to the fracture site cDNAs encoding osteogenic proteins, such as BMPs. Successful gene delivery and expression leads to the continuous, local, focal synthesis of the

osteogenic protein, which is likely to have undergone authentic post-translational modification and, unlike the recombinant product, to be uncontaminated by inactive, misfolded products that can trigger immune responses. Safety is further enhanced by gene transfer because the local production of osteogenic proteins at approximately physiological concentrations is less likely than the bolus application of a large amount of recombinant protein to lead to the systemic spread of these proteins to ectopic sites.

As with most other applications of gene therapy, the key questions for bone healing are: which gene(s) to transfer? Where to transfer them? How to get them there? Does it work? Is it safe? Much of the remainder of this article addresses these questions, bearing in mind that different clinical applications, such as fracture nonunions versus segmental defects, and different parts of the skeleton, such as maxillofacial bones versus long bones, might require different approaches. Gene therapy for bone healing has been recently reviewed by Carofino and Lieberman (Ref. 19)

Which genes to transfer?

There are a large number of growth factors known to stimulate one or more of the processes involved in osteogenesis and bone healing. Most of these either stimulate the differentiation of progenitor cells into chondrocytes or osteoblasts, or enhance the bone-forming activities of mature osteoblasts. Because bone is highly vascularised, vascular endothelial growth factor (VEGF) and other angiogenic factors are also important, especially for intramembranous bone formation; chondrogenesis, the first step in endochondral ossification, does not require a blood supply. The importance of angiogenesis in bone healing has been nicely demonstrated by Peng and colleagues (Ref. 20) who used gene transfer to show that VEGF enhanced, and a VEGF antagonist (sFlt1) inhibited, repair of cranial defects in mice. Because individual growth factors act at different stages of osteogenesis, combinations of different factors should promote bone healing more potently than either factor alone. This has been confirmed in animal models using gene delivery of BMP-2 and BMP-7 (Refs 21, 22); BMP-4 and VEGF (Ref. 20); BMP-4 and transforming growth factor- β (TGF- β) (Ref. 23).

An alternative approach to the delivery of cDNAs encoding growth factors involves the delivery of transcription factors associated with osteogenesis, such as RUNX2 (Ref. 24) and osterix (Transcription factor Sp7) (Ref. 25). Because these are intracellular proteins, they are not easily delivered by traditional methods of protein delivery, but are well suited to gene transfer. Their safety might be greater than that of growth factors, because they are not secreted and do not circulate. This is a disadvantage in terms of efficiency, however, because gene transfer would have to be very effective to modify most of the progenitor cells in the vicinity of the fracture. Timing of gene transfer would then be important, to synchronise with the entry of progenitor cells into the defect. In the case of a secreted gene product, by contrast, there is only the need to modify adequate numbers of cells to create a sufficiently high local concentration of the secreted factor for a sufficient period of time. LIM mineralisation protein-1 (LMP-1) is another intracellular, osteogenic molecule of interest. It was discovered during a screen of transcripts that are induced during osteogenesis, and its function is still unclear (Ref. 26). It was originally thought to be a

transcription factor, but that now seems not to be the case (REF?). However, remarkably potent and dramatic preclinical results in models of spinal fusion have been reported using plasmid DNA-liposome as the vector in rats (Ref. 27) and adenovirus vectors in rabbits (Ref. 28). Vectors expressing a related protein, LMP-3, are also osteogenic in animal models (Refs 29, 30, 31)

A slightly different approach is indicated by data which show that the knockdown of the BMP inhibitors noggin (Ref. 32) and chordin (Ref. 33) also stimulates osteogenesis. The delivery of siRNA molecules is possible by RNA transfection or the delivery of a vector that encodes the inhibitory RNA molecule. There do not appear to have been any experimental attempts to enhance fracture repair by inhibition of osteoclast function using gene transfer, although this might be a promising short-term approach to increase the net accumulation of bone.

Where to transfer genes?

Although it is possible to deliver osteogenic genes to sites where secreted gene products are able to circulate systemically, such strategies are considered only for conditions such as osteoporosis, where there is disseminated loss of bone (Refs 34, 35, 36). There is some evidence that mesenchymal stem cells (MSCs) can home to fracture sites and deliver a genetic payload (Ref. 37), but this has not been widely studied. Instead, because fractures occur locally, gene-therapy strategies are based upon the direct, local delivery of genes to cells within or around the fracture site. This has the advantage of achieving high local concentrations of the gene product, with minimal exposure of other organs. As noted earlier, this introduces an important safety component.

How to transfer genes to fractures?

Basic principles

A variety of different vectors have been developed for gene transfer. These can be divided into nonviral and viral gene delivery; nonviral gene transfer is known as transfection, and viral gene transfer is known as transduction. Each has advantages and disadvantages. Viral vectors are much more complicated to manufacture and they raise greater safety concerns, but continue to be used for the majority of human gene therapy trials because they are so much more efficient than nonviral vectors. For most gene-transfer applications, viruses are engineered to reduce any pathogenicity they might have had, while retaining their ability to infect target cells. For nearly all purposes, the viruses are attenuated, so that after they have delivered their genetic payload to the cells they infect, they cannot replicate. A variety of different viruses have been engineered in this way, and a number have found use in human clinical trials (www.wiley.co.uk/genetherapy/clinical/). There is not space here to describe them all in detail (for review see Ref. 38), but some general comments can be made about the vectors used in studies of bone healing.

Oncoretroviruses (usually referred to as retroviruses) and lentiviruses (also members of the retrovirus family) are RNA viruses that integrate their genetic material into the chromosomal DNA of the cells they infect, thus providing the potential for long-term

transgene expression. Although this is highly desirable when treating a genetic disease for the lifetime of the patient, it is probably not necessary for healing a fracture, which might be expected to take 6–8 weeks. The continued expression of a transgene beyond this time would probably be counterproductive. Another disadvantage of integrating retrovirus and lentivirus vectors is their potential for causing insertional mutagenesis. This has led to leukaemia in children being treated for severe combined immunodeficiency disease (SCID) (Ref. 39). Retrovirus, but not lentivirus, has the additional disadvantage of requiring host-cell division for successful transduction (Ref. 40). Lentivirus vectors are being engineered to prevent integration (Ref. 41), but it still remains doubtful that such viruses could be used in humans to treat a non-life-threatening condition, such as an osseous nonunion, especially as the most commonly used lentiviral vectors come with the ‘psychological baggage’ of having been derived from HIV.

Adenovirus and recombinant (unlike wild-type) adenoassociated virus (AAV) are DNA viruses whose genomes do not integrate with high frequency into host-cell DNA. Nevertheless, they are capable of maintaining transgene expression for lengths of time compatible with the needs of bone healing. Among the advantages of adenovirus is the relative ease with which recombinant vector can be produced at high titre, and its high infectivity towards many cell types, regardless of whether they are dividing or not (Ref. 42). As reflected in Tables 1 and 2, they have been the most widely used vectors for studies of gene therapy for bone healing. A perceived disadvantage of adenovirus is its high antigenicity (Ref. 42). Most of the population have circulating, neutralising antibodies to adenovirus serotype 5, the serotype most commonly used for gene transfer, which might reduce the effectiveness of adenovirus-based gene therapy. Moreover, the inflammatory response to adenovirus may be detrimental to healing.

AAV is perceived to be safe, and the death in 2007 of a subject receiving intra-articular injections of recombinant AAV in an arthritis gene-therapy trial was not attributed to the vector (Ref. 43). Although it has a small carrying capacity, most of the growth factors of interest have small cDNAs that fit comfortably into the recombinant AAV genome. The single-stranded DNA genome of AAV was an earlier limitation for many purposes, because transgene expression required the host cell to synthesise the second strand of DNA and this was sometimes very inefficient. This limitation was overcome by the development of recombinant, selfcomplementing AAV with double-stranded DNA genomes (Ref. 44) that give much higher levels of transgene expression much more quickly and reliably. Use of AAV remains constrained by the technological challenges of producing vector in large quantities and the need for relatively high multiplicities of infection.

Nonviral vectors (Ref. 45) can be as simple as naked plasmid DNA. The efficiency of gene transfer can be increased by associating the DNA with a carrier, such as a liposome, polymer or nanoparticle, or by using a physical stimulus such as electroporation or sonication.

Regardless of the vector, there are two basic strategies for gene transfer: ex vivo and in vivo (Fig. 1). For ex vivo delivery, cells are removed from the host, genetically modified outside the body and then reimplanted. This was very common in the early days of gene therapy, because the first useful vectors were retroviruses that require cell division for effective

transduction. Usually the cells are expanded in tissue culture, which, for human application, is very cumbersome and expensive. As described later, when host-cell division is not required, cell culture may not be necessary and tissue can be removed, genetically modified and reimplanted intraoperatively in an expedited *ex vivo* fashion (Ref. 46). For *in vivo* delivery, the vector is introduced directly into the host, often by injection or by implantation in association with a matrix (Fig. 1).

For certain applications, such as when treating a disease like rheumatoid arthritis that has flares and remissions, it might be desirable to regulate the level of transgene expression. There are a number of strategies for this, most of which involve the use of inducible promoters. This has not been widely investigated in the context of bone healing because, as far as we know, the levels of the most commonly used growth factors do not need to be closely regulated, and many of the commonly used vectors express for suitable lengths of time without the need for modification. Nevertheless, it has been confirmed that BMP-2 expression and consequent osteogenesis can be regulated *in vivo* using a TetON inducible expression system (Ref. 47).

Applications in bone healing

Four strategies have emerged for gene transfer to osseous lesions; two of them are *in vivo* and two are *ex vivo* (Fig. 1). All have shown promise, and it is not yet clear which of them will find eventual clinical use in human or veterinary medicine.

Ex vivo strategies

Traditional *ex vivo* gene transfer—As noted above, *ex vivo* gene transfer to culture-expanded cells was the first gene-therapy strategy, and the first to be used in human clinical trials (Ref. 48). It was also among the first experimental approaches to be used to enhance bone healing in experimental animal models. Lieberman and co-workers (Refs 49, 50) used a recombinant adenovirus vector carrying the cDNA for human *BMP2* (Ad.BMP-2) to transduce cultures of bone marrow stromal cells. The transduced cells were seeded onto collagen scaffolds and implanted into critical-sized defects in the femora of rats. All defects healed within 12 weeks of implanting the genetically modified cells, and showed better histological integrity than control defects treated with recombinant human BMP-2. Because rat marrow stromal cells differentiate into osteoblasts in response to BMP-2, the transduced cells have the potential to serve both as a provider of BMP-2 and also a source of osteoprogenitors, thereby enhancing the reparative response. Subsequent investigators have confirmed the utility of this strategy in experimental animals using osteoprogenitor cells derived from additional sources, such as periosteum (Ref. 51), fat (Refs 52, 53) and muscle (Ref. 54). Despite the success of these methods, it is still unclear whether the use of osteoprogenitor cells as vehicles for *ex vivo* gene delivery confers much of an advantage. It has been difficult to identify donor cells in the healed bone (Ref. 54), and some investigators report success with skin (Refs 29, 55) and gingival (Ref. 56) fibroblasts, which are not noted for their osteogenic potential. In agreement with this, the abilities of different cell types to promote the formation of heterotopic bone intramuscularly in mice was compared, but it was concluded that the cell type was not important (Ref. 57).

Although most investigators use adenovirus vectors for experiments of this kind, gene expression usually lasts for only 2–3 weeks in vivo. Virk and colleagues (Ref. 58) compared lentivirus, which expresses BMP-2 within osseous lesions for 8 weeks in vivo, with adenovirus. Although both vectors achieved bony union in the femoral critical-size-defect model, the repaired bone formed in response to the lentivirus vector had better mechanical and histological properties. In contrast to this, Gazit and colleagues (Ref. 59) have achieved remarkable success in murine models of bone healing using nonviral delivery of plasmids containing a human *BMP2* cDNA. Under these conditions, BMP-2 expression is low and transient. There has been very little research into species differences when using gene transfer approaches, and it is possible that the murine model is especially responsive to human BMP-2.

Expedited ex vivo gene transfer—Ex vivo strategies have the safety advantage that vectors are not introduced directly into the body. However, there are disadvantages to the traditional approaches described above: they require two invasive procedures, one to harvest cells and one to implant them, and they incur the high costs of expanding cells under GMP (good-manufacturing-practice) conditions. To obviate these problems, expedited ex vivo methods are being developed in which tissues are removed from the body, genetically modified and reimplanted within a single operative period (Ref. 60). The first use of this approach in bone healing was in a rabbit model of spinal fusion (Ref. 28). Marrow was withdrawn and the buffy coats isolated, transduced with a recombinant adenovirus vector encoding LMP-1 and implanted locally during a single operative session. Full spinal fusion occurred in each rabbit receiving the transgene in this fashion.

Skeletal muscle and fat have also been evaluated as vehicles for an expedited, ex vivo gene-based technology, known as ‘facilitated endogenous repair’, for the healing of lesions in bone, cartilage and other connective tissues (Refs 46, 60). Muscle (Ref. 61) and fat (Ref. 62) were selected because they contain osteoprogenitor cells, can serve as space-filling scaffolds, and they can be harvested, genetically modified and reimplanted in a single sitting. Using these methods in conjunction with Ad.BMP-2, it was possible to heal rat critical-size femoral defects more rapidly and reliably than with recombinant human BMP-2 (Ref. 46). Implantation of genetically modified muscle led to the rapid formation of cartilage within the lesion, presumably by the efficient differentiation of progenitor cells within the implant into chondrocytes, which underwent rapid endochondral ossification. Tracking of the donor cells confirmed that at least some of the newly formed bone was derived from the implanted muscle (Ref. 46). The expedited ex vivo approach also eliminates the humoral response to the adenovirus vector, which is an advantage for safety, efficacy and redosing.

In vivo strategies

In vivo strategies include direct injection of genes and gene-activated matrices and have been comprehensively reviewed recently (Ref. 63).

Direct injection of vector—Among the simplest of all gene-delivery strategies is to inject the vector directly into the osseous defect; this can often be performed percutaneously. Adenovirus vectors have been widely used for this purpose, and success has been reported in

rats (Refs 64, 65, 66), rabbits (Refs 67, 68) and horses (Ref. 69) using BMP-2 (Refs 64, 65, 66, 67, 69) and BMP-6 (Refs 68, 69) as transgenes. Bone healing in a rat femoral defect model was improved by delaying the administration of the virus for 5–10 days after creating the defect (Ref. 64). This could reflect the time it takes for osteoprogenitor cells to enter the defect, and also the possibility that the coxsackie virus adenovirus receptor (CAR), through which adenovirus enters cells, is maximally induced at the lesion site at this time (Ref. 70).

There also appear to be species differences in the osteogenic response to the direct injection of adenovirus. As noted above, very promising results have been reported in rabbits, rats and horses. In the sheep, the results have been quite disappointing. First experiments were conducted with a tibial osteotomy model (Ref. 71). This lesion spontaneously heals, and the administration of Ad.BMP-2 delayed, rather than accelerated, healing. Antibodies against both adenovirus and human BMP-2 were detected in the recipient sheep (Ref. 71), suggesting that immune reactions were responsible for the poor healing response. In agreement with this conclusion, the administration of Ad.BMP-2 to sheep that had been treated with steroids enhanced the healing of the tibial osteotomy (Ref. 72). It is not known whether the sheep immune system is especially sensitive to adenovirus, or whether the sheep, which spend much of their lives outdoors on a farm, had been previously exposed to adenovirus. Regardless of the explanation, it highlights again the importance of considering species differences and the immune system in studies of this sort.

Gene-activated matrices—Gene-activated matrices (GAMs) were developed to provide an off-the-shelf, nonviral, gene-based method for healing bone. In the original GAM formulation, plasmid DNA was incorporated into a collagen sponge and fitted into an osseous defect. The concept was to transfect in situ host reparative cells that infiltrated the GAM after implantation. The transfected cells would then secrete the transgene locally, triggering reparative responses. Using parathyroid hormone (PTH) 1–34 as the transgene, impressive results were reported in large segmental defects in rats (Ref. 73) and dogs (Ref. 74). However, with the exception of a GAM containing DNA encoding VEGF that showed activity in a rabbit segmental defect model (Ref. 75), no further progress has been reported using plasmid DNA and a collagen matrix.

Some investigators have attempted to improve the efficiency of GAMs by using different scaffolds (Refs 76, 77, 78, 79) or by using GAMs containing viral vectors. Once such GAM containing recombinant adenovirus carrying a cDNA encoding platelet-derived growth factor (PDGF) B has shown preliminary success in clinical trials for healing diabetic skin ulcers (www.t-r-co.com), and has shown promise in growing periodontal bone in animal models (Ref. 80). The fact that is already in clinical trials for wound healing will facilitate its adaptation for bone healing in humans. There is evidence that a collagen scaffold reduces the immune response to adenovirus (Ref. 81).

Schwarz and colleagues have developed a modified GAM process in which allograft bone serves as the scaffold and AAV as the vector (Ref. 82). Efficacy was first demonstrated in a murine segmental-defect model. Allograft bone which, as described earlier in this article, contains no living cells, was ‘revitalised’ by coating it with two different AAV vectors carrying NF- κ B ligand (RANKL), to promote osteoclastogenesis, and VEGF, to promoted

angiogenesis. When the construct was implanted into the mouse femoral defect, AAV transduced the host cells surrounding the implant, leading to the remodelling of the allograft and its replacement with host bone (Ref. 82). Similar responses were noted using a constitutively active *ALK2* cDNA (Ref. 83). This encodes a type I BMP receptor bearing the same mutation as found in the heterotopic bone-deposition disease fibrodysplasia ossificans progressiva (Ref. 84). Because the AAV-coated construct is relatively stable, it can form the basis of an off-the-shelf material for bone healing.

Does it work?

Table 1 lists the extensive literature confirming the ability of gene-transfer approaches to heal osseous defects in small laboratory animals, and Figure 2 shows one example of successful bone healing in a rat model. Far fewer studies have been performed in large-animal models (Table 2), but the results are encouraging, especially when using the traditional *ex vivo* approach in horses (Ref. 55), goats (Refs 103, 104, 105) and pigs (Ref. 106). Results using GAM containing *PTH* 1-34 DNA were more equivocal (Refs 74, 85). Data from the *in vivo* delivery of Ad.BMP-2 and Ad.BMP-6 to osseous defects in horses were encouraging (Ref. 69), whereas similar experiments sheep gave disappointing results (Refs 71, 72). Further research is clearly indicated.

Is it safe?

Over 1500 human gene-therapy clinical studies have taken place in the 20 years since the first properly authorised clinical gene-transfer study (Ref. 48) (www.wiley.co.uk/genetherapy/clinical/). Despite a common perception that gene therapy is unsafe, only two human deaths have been attributed to gene transfer (Ref. 86). Nevertheless, safety issues continue to burden the clinical development of gene therapy. The 1999 death of Jesse Gelsinger (Ref. 87) was widely reported and halted what had been a decade of spectacular growth in the number of clinical gene therapy trials. His death resulted from an uncontrollable activation of the innate immune system following the *in vivo* infusion of a very high dose of recombinant adenovirus vectors into the hepatic portal vein. The second death occurred in Paris, where haematopoietic stem cells of children with SCID were treated *ex vivo* with a retrovirus vector. Several subjects developed leukaemia because of insertional mutagenesis, the retroviral genome having been inserted adjacent to the LMO oncogene (Ref. 39). One of the affected individuals died, but the leukaemia was successfully treated in the other cases. The 2007 death of a subject who received an AAV vector by intraarticular injection in an arthritis gene-therapy trial was not attributed to gene transfer (Refs 43, 86).

Safety issues are of particular concern when gene therapy is used for a nonlethal condition, such as bone healing, especially if alternative, nongenetic treatments are available. For this reason, vectors based upon integrating viruses, such as retrovirus and lentivirus, are unlikely to see early clinical application. Because the efficacy of nonviral delivery systems remains in doubt, adenovirus and AAV emerge as the vectors most likely to be used in the first clinical trials. Although recombinant adenovirus vectors were responsible for the death of Jesse Gelsinger, any application in bone healing will use far less virus and the vector will be

delivered locally into an osseous lesion, rather than systemically into a major vessel. Moreover, it is likely that adenovirus will be used in an *ex vivo* fashion, which means that no virions enter the patient and the immune response to the virus is attenuated.

Progress towards clinical trials

Bringing gene-therapy protocols into the clinic is a lengthy and very expensive process. Most of the experimental data concerning bone healing, although very promising, have been generated with rodents and rabbits (Table 1). Efficient and reliable bone healing would need to be established in large-animal models (Table 2). Detailed pharmacological and toxicological studies are also mandated. These include, among other things, extensive biodistribution studies using very sensitive PCR-based methods, conducted under GMP conditions, to determine the possible spread of vector genomes throughout the body and their persistence. Demonstration of safety and efficacy in animal models is a necessary prelude to Phase I (safety, small number), II (efficacy, small number) and III (safety and efficacy, large number) studies in humans. Only if these are successful can the therapeutic enter clinical use. The total cost of such investigations is enormous. Because they lie beyond the typical means of an academic investigator, the clinical development of gene therapies for bone healing will almost certainly require the sustained interest of a commercial entity. Whether this occurs could determine the fate of the field. The other dominating factor is the ambient regulatory environment, which invariably tightens after a serious adverse event in a human gene-therapy trial. This makes a gene-therapy programme vulnerable to events occurring elsewhere, over which the investigator has absolutely no control. Nevertheless, GAMs containing adenovirus carrying *PDGF* cDNA are already undergoing clinical trials for wound healing and are thus favourably positioned to enter trials for bone healing.

In addition to human medicine, there are many veterinary uses for a gene-based approach to bone healing, and the route to veterinary application is less burdensome from the regulatory point of view. Demonstration of safety and efficacy in veterinary medicine should facilitate the application of a gene therapeutic for bone healing in humans.

Research in progress and outstanding research questions

There is a compelling logic to the strategy of healing osseous defects by delivering osteogenic cDNAs locally to sites of injury. However, further research is needed to determine which transgene and which gene delivery system are the most appropriate for taking forward into advanced preclinical studies. One advantage of using transgenes encoding BMP-2, BMP-7 and PDGF, which are already used clinically as recombinant proteins, is the large body of accumulated data concerning the efficacy, pharmacology and toxicology of these substances in humans. This considerably smoothes the regulatory pathway to clinical application. The use of cDNAs encoding novel or esoteric osteogenic products, by contrast, although scientifically more exciting, will have a much more tortuous path to the clinic. The same is true for the simultaneous use of multiple transgenes. Thus, much of the current research uses cDNAs encoding well-studied proteins.

Similar considerations influence the choice of gene-delivery strategy and vector. *Ex vivo* gene delivery has many attractions, especially related to safety, and considerable research

activity surrounds the evaluation of different types of cells for their suitability as vehicles of ex vivo gene delivery. However, the traditional route of using monolayers of cells that have been serially expanded is very expensive, especially if the cells need to be autologous. The use of allografted cells would reduce the cost and complexity of this type of ex vivo gene therapy, and there is much interest in the possibility that MSCs can be successfully allografted because they produce immunosuppressive factors (Ref. 88). However, femoral defects in rats could only be healed using allografted MSCs if the animals were immunosuppressed (Ref. 89). Nevertheless, the successful healing of fractures in immunocompetent, osteoporotic rats using xenografts of retrovirally transduced, irradiated, human chondrocytes expressing BMP-2 has been reported (Ref. 90). The ability of MSCs to differentiate into osteoblasts promises to be an advantage but, as noted earlier, it is still not clear to what degree transplanted MSCs become osteoblasts in animal models of bone healing. Moreover, other easily obtainable cells of mesenchymal origin, such as skin fibroblasts, seem able to differentiate into osteoblasts. MSCs are of additional interest because of their purported ability to home to sites of osseous damage, which, if true, would considerably simplify delivery.

Related to the choice of cell is the choice of scaffold upon which to seed the genetically modified cells. Discussion of scaffolds is beyond the scope of this review, but they are key adjuncts to traditional ex vivo methods of gene delivery to bone and the subject of much research. Expedited ex vivo strategies obviate the need to identify suitable cell types and scaffolds. Instead, the main research focus is to establish whether marrow, muscle, fat or some other tissue is the most appropriate one to use.

There is still no consensus concerning the most appropriate vectors to develop for clinical use. Although retrovirus and lentivirus have shown utility for demonstrating proof of principle, safety concerns will make them difficult to take forward into the clinic. Among the viral vectors, adenovirus and AAV seem to hold the most promise, because they are effective and the safety issues are manageable. They are thus the focus of much current investigation. The development of effective nonviral vectors would greatly accelerate the clinical development of the field, but these have not proved very effective. Nevertheless, there is much research in this area.

Finally, there is the need to move beyond the small-animal models that most of us use for our research. Because most of the skeleton bears load, the scale-up from a rodent to a large animal has important biomechanical consequences, superimposed upon which are species differences in responses to osteogenic stimuli. Although large-animal studies are expensive and slow, they need to be done if the field is to continue to move forward towards clinical application. Development of translational outcome measures of bone healing is central to the design of feasible gene-therapy studies in large animals and clinical trials. Novel, noninvasive techniques are being developed to determine mechanical strength and bone quality. Such techniques include cone-beam computed tomography, positron emission spectroscopy, dynamic-contrast-enhanced magnetic-resonance imaging, union ratio, connectivity and bone mineral density.

Acknowledgments

The author's work in this area has been supported by NIH grant R01 AR 050243 from NIAMS and the AO Foundation. I thank Elisabeth Ferreira, PhD, for reviewing an earlier draft of this manuscript. The suggestions of the four anonymous reviewers are also appreciated.

References

1. Kwiatkowski TC, Hanley EN Jr, Ramp WK. Cigarette smoking and its orthopedic consequences. *American Journal of Orthopedics* (Belle Mead NJ). 1996; 25:590–597.
2. Chakkalakal DA. Alcohol-induced bone loss and deficient bone repair. *Alcoholism Clinical and Experimental Research*. 2005; 29:2077–2090.
3. Gandhi A, et al. Diabetic fracture healing. *Foot and Ankle Clinics*. 2006; 11:805–824. [PubMed: 17097518]
4. Gruber R, et al. Fracture healing in the elderly patient. *Experimental Gerontology*. 2006; 41:1080–1093. [PubMed: 17092679]
5. Giannoudis P, et al. Fracture healing in osteoporotic fractures: is it really different? A basic science perspective. *Injury*. 2007; 38:S90–99. [PubMed: 17383490]
6. McKee MD. Management of segmental bony defects: the role of osteoconductive orthobiologics. *Journal of the American Academy of Orthopaedic Surgeons*. 2006; 14:S163–167. [PubMed: 17003191]
7. Daniels AH, et al. Adverse events associated with anterior cervical spine surgery. *Journal of the American Academy of Orthopaedic Surgeons*. 2008; 16:729–738. [PubMed: 19056921]
8. Kim DH, et al. Prospective study of iliac crest bone graft harvest site pain and morbidity. *Spine Journal*. 2009; 9:886–892. [PubMed: 19540168]
9. Delloye C, et al. Bone allografts: What they can offer and what they cannot. *Journal of Bone and Joint Surgery. British Volume*. 2007; 89:574–579.
10. Watson JT. Overview of biologics. *Journal of Orthopaedic Trauma*. 2005; 19:S14–S16. [PubMed: 16479217]
11. Shapiro F. Bone development and its relation to fracture repair. The role of mesenchymal osteoblasts and surface osteoblasts. *European Cells and Materials*. 2008; 15:53–76. [PubMed: 18382990]
12. Simpson AH, Mills L, Noble B. The role of growth factors and related agents in accelerating fracture healing. *Journal of Bone and Joint Surgery. British Volume*. 2006; 88:701–705.
13. Lieberman JR, Daluiski A, Einhorn TA. The role of growth factors in the repair of bone. *Biology and clinical applications. Journal of Bone and Joint Surgery. American Volume*. 2002; 84:1032–1044.
14. Tsumaki N, Yoshikawa H. The role of bone morphogenetic proteins in endochondral bone formation. *Cytokine and Growth Factor Reviews*. 2005; 16:279–285. [PubMed: 15869898]
15. Kwong FN, Harris MB. Recent developments in the biology of fracture repair. *Journal of the American Academy of Orthopaedic Surgeons*. 2008; 16:619–625. [PubMed: 18978283]
16. Gillet JP, et al. The development of gene therapy: from monogenic recessive disorders to complex diseases such as cancer. *Methods in Molecular Biology*. 2009; 542:5–54. [PubMed: 19565894]
17. Evans CH, et al. Gene transfer to human joints: progress toward a gene therapy of arthritis. *Proceedings of the National Academy of Sciences of the United States of America*. 2005; 102:8698–8703. [PubMed: 15939878]
18. Fiandaca M, Forsayeth J, Bankiewicz K. Current status of gene therapy trials for Parkinson's disease. *Experimental Neurology*. 2008; 209:51–57. [PubMed: 17920059]
19. Carofino BC, Lieberman JR. Gene therapy applications for fracture-healing. *Journal of Bone and Joint Surgery. American Volume*. 2008; 90:99–110.
20. Peng H, et al. Synergistic enhancement of bone formation and healing by stem cell-expressed VEGF and bone morphogenetic protein-4. *Journal of Clinical Investigation*. 2002; 110:751–759. [PubMed: 12235106]

21. Koh JT, et al. Combinatorial gene therapy with BMP2/ 7 enhances cranial bone regeneration. *Journal of Dental Research*. 2008; 87:845–849. [PubMed: 18719211]
22. Zhu W, et al. Combined bone morphogenetic protein-2 and -7 gene transfer enhances osteoblastic differentiation and spine fusion in a rodent model. *Journal of Bone and Mineral Research*. 2004; 19:2021–2032. [PubMed: 15537446]
23. Li BC, et al. Treatment of rabbit femoral defect by firearm with BMP-4 gene combined with TGF-beta1. *Journal of Trauma*. 2009; 66:450–456. [PubMed: 19204520]
24. Zhao Z, et al. Healing cranial defects with AdRunx2-transduced marrow stromal cells. *Journal of Dental Research*. 2007; 86:1207–1211. [PubMed: 18037657]
25. Tu Q, et al. Osterix overexpression in mesenchymal stem cells stimulates healing of critical-sized defects in murine calvarial bone. *Tissue Engineering*. 2007; 13:2431–2340. [PubMed: 17630878]
26. Boden SD, et al. LMP-1, a LIM-domain protein, mediates BMP-6 effects on bone formation. *Endocrinology*. 1998; 139:5125–5134. [PubMed: 9832452]
27. Boden SD, et al. Lumbar spine fusion by local gene therapy with a cDNA encoding a novel osteoinductive protein (LMP-1). *Spine (Phila. PA 1976)*. 1998; 23:2486–2492. [PubMed: 9854747]
28. Viggewarapu M, et al. Adenoviral delivery of LIM mineralization protein-1 induces new-bone formation in vitro and in vivo. *Journal of Bone and Joint Surgery. American Volume*. 2001; 83:364–376.
29. Lattanzi W, et al. Ex vivo-transduced autologous skin fibroblasts expressing human Lim mineralization protein-3 efficiently form new bone in animal models. *Gene Therapy*. 2008; 15:1330–1343. [PubMed: 18633445]
30. Parrilla C, et al. Ex vivo gene therapy using autologous dermal fibroblasts expressing hLMP3 for rat mandibular bone regeneration. *Head and Neck*. 2009; 32:310–318. [PubMed: 19626643]
31. Pola E, et al. Efficient bone formation by gene transfer of human LIM mineralization protein-3. *Gene Therapy*. 2004; 11:683–693. [PubMed: 14724674]
32. Wan DC, et al. Noggin suppression enhances in vitro osteogenesis and accelerates in vivo bone formation. *Journal of Biological Chemistry*. 2007; 282:26450–26459. [PubMed: 17609215]
33. Kwong FN, Richardson SM, Evans CH. Chordin knockdown enhances the osteogenic differentiation of human mesenchymal stem cells. *Arthritis Research and Therapy*. 2008; 10:R65. [PubMed: 18533030]
34. Baltzer AW, et al. Gene therapy for osteoporosis: evaluation in a murine ovariectomy model. *Gene Therapy*. 2001; 8:1770–1776. [PubMed: 11803396]
35. Bolon B, et al. Adenoviral delivery of osteoprotegerin ameliorates bone resorption in a mouse ovariectomy model of osteoporosis. *Molecular Therapy*. 2001; 3:197–205. [PubMed: 11237676]
36. Kostenuik PJ, et al. Gene therapy with human recombinant osteoprotegerin reverses established osteopenia in ovariectomized mice. *Bone*. 2004; 34:656–664. [PubMed: 15050896]
37. Shen FH, et al. Systemically administered mesenchymal stromal cells transduced with insulin-like growth factor-I localize to a fracture site and potentiate healing. *Journal of Orthopaedic Trauma*. 2002; 16:651–659. [PubMed: 12368646]
38. Young LS, et al. Viral gene therapy strategies: from basic science to clinical application. *Journal of Pathology*. 2006; 208:299–318. [PubMed: 16362990]
39. Kohn DB, Sadelain M, Glorioso JC. Occurrence of leukaemia following gene therapy of X-linked SCID. *Nature Reviews Cancer*. 2003; 3:477–488.
40. Kurian KM, Watson CJ, Wyllie AH. Retroviral vectors. *Molecular Pathology*. 2000; 53:173–176. [PubMed: 11040938]
41. Sarkis C, et al. Non-integrating lentiviral vectors. *Current Gene Therapy*. 2008; 8:430–437. [PubMed: 19075626]
42. Douglas JT. Adenoviral vectors for gene therapy. *Molecular Biotechnology*. 2007; 36:71–80. [PubMed: 17827541]
43. Frank KM, et al. Investigation of the cause of death in a gene-therapy trial. *New England Journal of Medicine*. 2009; 361:161–169. [PubMed: 19587341]

44. McCarty DM, Monahan PE, Samulski RJ. Self-complementary recombinant adeno-associated virus (scAAV) vectors promote efficient transduction independently of DNA synthesis. *Gene Therapy*. 2001; 8:1248–1254. [PubMed: 11509958]
45. Glover DJ, Lipps HJ, Jans DA. Towards safe, non-viral therapeutic gene expression in humans. *Nature Reviews Genetics*. 2005; 6:299–310.
46. Evans CH, et al. Use of genetically modified muscle and fat grafts to repair defects in bone and cartilage. *European Cells and Materials*. 2009; 18:96–111. [PubMed: 20073015]
47. Gafni Y, et al. Gene therapy platform for bone regeneration using an exogenously regulated, AAV-2-based gene expression system. *Molecular Therapy*. 2004; 9:587–595. [PubMed: 15093189]
48. Rosenberg SA, et al. Gene transfer into humans--immunotherapy of patients with advanced melanoma, using tumor-infiltrating lymphocytes modified by retroviral gene transduction. *New England Journal of Medicine*. 1990; 323:570–578. [PubMed: 2381442]
49. Lieberman JR, et al. The effect of regional gene therapy with bone morphogenetic protein-2-producing bone-marrow cells on the repair of segmental femoral defects in rats. *Journal of Bone and Joint Surgery. American Volume*. 1999; 81:905–917.
50. Lieberman JR, et al. Regional gene therapy with a BMP-2-producing murine stromal cell line induces heterotopic and orthotopic bone formation in rodents. *Journal of Orthopaedic Research*. 1998; 16:330–339. [PubMed: 9671928]
51. Breitbart AS, et al. Gene-enhanced tissue engineering: applications for bone healing using cultured periosteal cells transduced retrovirally with the BMP-7 gene. *Annals of Plastic Surgery*. 1999; 42:488–495. [PubMed: 10340856]
52. Peterson B, et al. Healing of critically sized femoral defects, using genetically modified mesenchymal stem cells from human adipose tissue. *Tissue Engineering*. 2005; 11:120–129. [PubMed: 15738667]
53. Hsu WK, et al. Stem cells from human fat as cellular delivery vehicles in an athymic rat posterolateral spine fusion model. *Journal of Bone and Joint Surgery. American Volume*. 2008; 90:1043–1052.
54. Lee JY, et al. Effect of bone morphogenetic protein-2-expressing muscle-derived cells on healing of critical-sized bone defects in mice. *Journal of Bone and Joint Surgery. American Volume*. 2001; 83:1032–1039.
55. Ishihara A, et al. Dermal fibroblast-mediated BMP2 therapy to accelerate bone healing in an equine osteotomy model. *Journal of Orthopaedic Research*. 2010; 28:403–411. [PubMed: 19777486]
56. Shin JH, et al. Ex vivo bone morphogenetic protein-2 gene delivery using gingival fibroblasts promotes bone regeneration in rats. *Journal of Clinical Periodontology*. 2010; 37:305–311. [PubMed: 20041973]
57. Gugala Z, et al. Osteoinduction by ex vivo adenovirus-mediated BMP2 delivery is independent of cell type. *Gene Therapy*. 2003; 10:1289–1296. [PubMed: 12883525]
58. Virk MS, et al. Influence of short-term adenoviral vector and prolonged lentiviral vector mediated bone morphogenetic protein-2 expression on the quality of bone repair in a rat femoral defect model. *Bone*. 2008; 42:921–931. [PubMed: 18295562]
59. Gazit D, et al. Engineered pluripotent mesenchymal cells integrate and differentiate in regenerating bone: a novel cell-mediated gene therapy. *Journal of Gene Medicine*. 1999; 1:121–133. [PubMed: 10738576]
60. Evans CH, et al. Facilitated endogenous repair: making tissue engineering simple, practical, and economical. *Tissue Engineering*. 2007; 13:1987–1993. [PubMed: 17518747]
61. Bosch P, et al. Osteoprogenitor cells within skeletal muscle. *Journal of Orthopaedic Research*. 2000; 18:933–944. [PubMed: 11192254]
62. Dragoo JL, et al. Tissue-engineered bone from BMP-2-transduced stem cells derived from human fat. *Plastic and Reconstructive Surgery*. 2005; 115:1665–1673. [PubMed: 15861072]
63. Pelled G, et al. Direct Gene Therapy for Bone Regeneration: Gene Delivery, Animal Models and Outcome Measures. *Tissue Engineering Part A*. 2009 [Epub ahead of print: 21 Jul 2009].

64. Betz OB, et al. Delayed administration of adenoviral BMP-2 vector improves the formation of bone in osseous defects. *Gene Therapy*. 2007; 14:1039–1044. [PubMed: 17460719]
65. Betz OB, et al. Direct percutaneous gene delivery to enhance healing of segmental bone defects. *Journal of Bone and Joint Surgery. American Volume*. 2006; 88:355–365.
66. Betz VM, et al. Healing of segmental bone defects by direct percutaneous gene delivery: effect of vector dose. *Human Gene Therapy*. 2007; 18:907–915. [PubMed: 17910523]
67. Baltzer AW, et al. Genetic enhancement of fracture repair: healing of an experimental segmental defect by adenoviral transfer of the BMP-2 gene. *Gene Therapy*. 2000; 7:734–739. [PubMed: 10822299]
68. Bertone AL, et al. Adenoviral-mediated transfer of human BMP-6 gene accelerates healing in a rabbit ulnar osteotomy model. *Journal of Orthopaedic Research*. 2004; 22:1261–1270. [PubMed: 15475207]
69. Ishihara A, et al. Osteogenic gene regulation and relative acceleration of healing by adenoviral-mediated transfer of human BMP-2 or -6 in equine osteotomy and ostectomy models. *Journal of Orthopaedic Research*. 2008; 26:764–771. [PubMed: 18241059]
70. Ito T, et al. Coxsackievirus and adenovirus receptor (CAR)-positive immature osteoblasts as targets of adenovirus-mediated gene transfer for fracture healing. *Gene Therapy*. 2003; 10:1623–1628. [PubMed: 12907955]
71. Egernann M, et al. Effect of BMP-2 gene transfer on bone healing in sheep. *Gene Ther*. 2006; 13:1290–1299. [PubMed: 16642029]
72. Egernann M, et al. Direct adenoviral transfer of bone morphogenetic protein-2 cDNA enhances fracture healing in osteoporotic sheep. *Human Gene Therapy*. 2006; 17:507–517. [PubMed: 16716108]
73. Fang J, et al. Stimulation of new bone formation by direct transfer of osteogenic plasmid genes. *Proceedings of the National Academy of Sciences of the United States of America*. 1996; 93:5753–5758. [PubMed: 8650165]
74. Bonadio J, et al. Localized, direct plasmid gene delivery in vivo: prolonged therapy results in reproducible tissue regeneration. *Nature Medicine*. 1999; 5:753–759.
75. Geiger F, et al. Vascular endothelial growth factor gene-activated matrix (VEGF165-GAM) enhances osteogenesis and angiogenesis in large segmental bone defects. *Journal of Bone and Mineral Research*. 2005; 20:2028–2035. [PubMed: 16234976]
76. Endo M, et al. Bone regeneration by modified gene-activated matrix: effectiveness in segmental tibial defects in rats. *Tissue Engineering*. 2006; 12:489–497. [PubMed: 16579682]
77. Huang YC, et al. Bone regeneration in a rat cranial defect with delivery of PEI-condensed plasmid DNA encoding for bone morphogenetic protein-4 (BMP-4). *Gene Therapy*. 2005; 12:418–426. [PubMed: 15647766]
78. Ono I, et al. Combination of porous hydroxyapatite and cationic liposomes as a vector for BMP-2 gene therapy. *Biomaterials*. 2004; 25:4709–4718. [PubMed: 15120517]
79. Valimaki VV, et al. Molecular biological evaluation of bioactive glass microspheres and adjunct bone morphogenetic protein 2 gene transfer in the enhancement of new bone formation. *Tissue Engineering*. 2005; 11:387–394. [PubMed: 15869418]
80. Chang PC, et al. Adenovirus encoding human platelet-derived growth factor-B delivered to alveolar bone defects exhibits safety and biodistribution profiles favorable for clinical use. *Human Gene Therapy*. 2009; 20:486–496. [PubMed: 19199824]
81. Sonobe J, et al. Osteoinduction by bone morphogenetic protein 2-expressing adenoviral vector: application of biomaterial to mask the host immune response. *Human Gene Therapy*. 2004; 15:659–668. [PubMed: 15242526]
82. Ito H, et al. Remodeling of cortical bone allografts mediated by adherent rAAV-RANKL and VEGF gene therapy. *Nature Medicine*. 2005; 11:291–297.
83. Koefoed M, et al. Biological effects of rAAV-caAlk2 coating on structural allograft healing. *Molecular Therapy*. 2005; 12:212–218. [PubMed: 16043092]
84. Shore EM, et al. A recurrent mutation in the BMP type I receptor ACVR1 causes inherited and sporadic fibrodysplasia ossificans progressiva. *Nature Genetics*. 2006; 38:525–527. [PubMed: 16642017]

85. Backstrom KC, et al. Response of induced bone defects in horses to collagen matrix containing the human parathyroid hormone gene. *American Journal of Veterinary Research*. 2004; 65:1223–1232. [PubMed: 15478769]
86. Evans CH, Ghivizzani SC, Robbins PD. Arthritis gene therapy's first death. *Arthritis Research and Therapy*. 2008; 10:110. [PubMed: 18510784]
87. Raper SE, et al. Fatal systemic inflammatory response syndrome in a ornithine transcarbamylase deficient patient following adenoviral gene transfer. *Molecular Genetics and Metabolism*. 2003; 80:148–158. [PubMed: 14567964]
88. Xu G, et al. Immunosuppressive properties of cloned bone marrow mesenchymal stem cells. *Cell Research*. 2007; 17:240–248. [PubMed: 17325691]
89. Tsuchida H, et al. Engineered allogeneic mesenchymal stem cells repair femoral segmental defect in rats. *Journal of Orthopaedic Research*. 2003; 21:44–53. [PubMed: 12507579]
90. Yi Y, et al. Irradiated human chondrocytes expressing bone morphogenetic protein 2 promote healing of osteoporotic bone fracture in rats. *Tissue Engineering Part A*. 2009; 15:2853–2863. [PubMed: 19254139]
91. Tarkka T, et al. Adenoviral VEGF-A gene transfer induces angiogenesis and promotes bone formation in healing osseous tissues. *Journal of Gene Medicine*. 2003; 5:560–566. [PubMed: 12825195]
92. Li R, et al. Effect of cell-based VEGF gene therapy on healing of a segmental bone defect. *Journal of Orthopaedic Research*. 2009; 27:8–14. [PubMed: 18634016]
93. Shen HC, et al. Structural and functional healing of critical-size segmental bone defects by transduced muscle-derived cells expressing BMP4. *Journal of Gene Medicine*. 2004; 6:984–991. [PubMed: 15352071]
94. Hsu WK, et al. Lentiviral-mediated BMP-2 gene transfer enhances healing of segmental femoral defects in rats. *Bone*. 2007; 40:931–938. [PubMed: 17236835]
95. Guo X, et al. Bone regeneration with active angiogenesis by basic fibroblast growth factor gene transfected mesenchymal stem cells seeded on porous beta-TCP ceramic scaffolds. *Biomedical Materials*. 2006; 1:93–99. [PubMed: 18458388]
96. Hidaka C, et al. Enhancement of spine fusion using combined gene therapy and tissue engineering BMP-7-expressing bone marrow cells and allograft bone. *Spine (Phila. PA 1976)*. 2003; 28:2049–2057. [PubMed: 14501913]
97. Hasharoni A, et al. Murine spinal fusion induced by engineered mesenchymal stem cells that conditionally express bone morphogenetic protein-2. *Journal of Neurosurgery Spine*. 2005; 3:47–52. [PubMed: 16122022]
98. Riew KD, et al. Induction of bone formation using a recombinant adenoviral vector carrying the human BMP-2 gene in a rabbit spinal fusion model. *Calcified Tissue International*. 1998; 63:357–360. [PubMed: 9744997]
99. Park J, et al. Bone regeneration in critical size defects by cell-mediated BMP-2 gene transfer: a comparison of adenoviral vectors and liposomes. *Gene Therapy*. 2003; 10:1089–1098. [PubMed: 12808439]
100. Southwood LL, et al. Evaluation of Ad-BMP-2 for enhancing fracture healing in an infected defect fracture rabbit model. *Journal of Orthopaedic Research*. 2004; 22:66–72. [PubMed: 14656661]
101. Ashinoff RL, et al. Bone morphogenetic protein-2 gene therapy for mandibular distraction osteogenesis. *Annals of Plastic Surgery*. 2004; 52:585–591. [PubMed: 15166991]
102. Alden TD, et al. The use of bone morphogenetic protein gene therapy in craniofacial bone repair. *Journal of Craniofacial Surgery*. 2000; 11:24–30. [PubMed: 11314095]
103. Rundle CH, et al. In vivo bone formation in fracture repair induced by direct retroviral-based gene therapy with bone morphogenetic protein-4. *Bone*. 2003; 32:591–601. [PubMed: 12810166]
104. Rundle CH, et al. Retroviral-based gene therapy with cyclooxygenase-2 promotes the union of bony callus tissues and accelerates fracture healing in the rat. *Journal of Gene Medicine*. 2008; 10:229–241. [PubMed: 18088065]

105. Strohbach CA, et al. LMP-1 retroviral gene therapy influences osteoblast differentiation and fracture repair: a preliminary study. *Calcified Tissue International*. 2008; 83:202–211. [PubMed: 18709396]
106. Dai KR, et al. Repairing of goat tibial bone defects with BMP-2 gene-modified tissue-engineered bone. *Calcified Tissue International*. 2005; 77:55–61. [PubMed: 16007479]
107. Xu XL, et al. Immune response and effect of adenovirus-mediated human BMP-2 gene transfer on the repair of segmental tibial bone defects in goats. *Acta Orthopaedica*. 2005; 76:637–646. [PubMed: 16263609]
108. Tang TT, et al. Treatment of osteonecrosis of the femoral head with hBMP-2-gene-modified tissue-engineered bone in goats. *Journal of Bone and Joint Surgery. British Volume*. 2007; 89:127–129.
109. Chang SC, et al. Large-scale bicortical skull bone regeneration using ex vivo replication-defective adenoviral-mediated bone morphogenetic protein-2 gene-transferred bone marrow stromal cells and composite biomaterials. *Neurosurgery*. 2009; 65:75–83. [PubMed: 19935005]

Further reading, resources and contacts

The following website compiles information on human gene therapy clinical trials. <http://www.wiley.co.uk/genetherapy/clinical/>

This website, within the Office of Biotechnology Activities of the US National Institutes of Health, provides data on human gene transfer trials in the USA. http://oba.od.nih.gov/rdna/oba_gemcris_public.html

Websites of relevant learned societies: American Society of Gene and Cell Therapy. <http://www.asgct.org/>

British Society of Gene Therapy. <http://www.bsgt.org/>

European Society of Gene and Cell Therapy. <http://www.esgct.eu/congress/>

Figure 1.

Strategies for gene transfer to defects in bone. There are two general strategies: in vivo and ex vivo. For in vivo gene delivery, the vector is introduced directly into the site of the osseous lesion, either as a free suspension (top right) or incorporated into a gene-activated matrix (GAM) (bottom right). For ex vivo delivery, vectors are not introduced directly into the defect. Instead they are used for the genetic modification of cells, which are subsequently implanted. Traditional ex vivo methods (top left) usually involve the establishment of cell cultures, which are genetically modified in vitro. The modified cells are then introduced into the lesion, often after seeding onto an appropriate scaffold. Expedited ex vivo methods (bottom left) avoid the need for cell culture by genetically modifying tissues such as marrow, muscle and fat, intraoperatively and inserting them into the defect during a single operative session.

Figure 2.

Healing of a rat femoral segmental defect following in vivo delivery of an adenovirus vector encoding *BMP2*. Representative radiographic images of segmental bone defects after direct injection of adenoviral vectors encoding human BMP-2 (Ad.BMP-2) or luciferase cDNA (Ad.luc). Most defects that had been treated with Ad.BMP-2 displayed bone formation within the defect by 4 weeks (a) and complete union by 8 weeks (c). Control defects treated with Ad.luc (and untreated defects) did not display appreciable signs of healing within this time (b and d). Images reproduced from Ref. 65.

Table 1

Use of gene therapy to treat osseous defects in small-animal models

Transgene product	Vector	Model	Ref.
Traditional ex vivo			
BMP-2	Adenovirus	Rodent long bone	49, 50, 52
IGF-1	Plasmid ^a		37
VEGF	Plasmid		92
BMP-4	Retrovirus		93
BMP-2	Lentivirus		94
FGF-2	Plasmid		95
BMP-2	Retrovirus ^b		90
BMP-2, BMP-7	Retrovirus, adenovirus	Rodent cranial defect	51, 54
BMP-4 and VEGF	Adenovirus		20
Cbfa1	Adenovirus		24
Osterix	Retrovirus		25
LMP-1	Plasmid, liposomes	Rat spine fusion	27
BMP-7	Adenovirus		96
BMP-2	Adenovirus		53
BMP-2	Adenovirus	Mouse spine fusion	97
BMP-2	Adenovirus	Rabbit spine fusion	98
BMP-2	Liposomes, adenovirus	Rat mandibular defect	99
LMP-3	Adenovirus		29
In vivo			
BMP-2	Adenovirus	Rabbit femur	67
BMP-2	Adenovirus	Infected rabbit femur	100
BMP-2	Adenovirus	Rat femur	64, 65, 66
BMP-2	Adenovirus	Distraction osteogenesis –rat mandible	101
BMP-6	Adenovirus	Rabbit ulna	68
BMP-2, BMP-9	Adenovirus	Rat mandible	102
BMP-4, COX-2	Retrovirus	Rat femoral fracture	103, 104
VEGF	Adenovirus	Rat femur drilling	91
BMP-2 and BMP-7	Adenovirus	Rat spine fusion	22
BMP-2	Adenovirus	Rabbit spine fusion	99
LMP-1	Retrovirus	Rat femoral fracture	105
Gene-activated matrix (GAM)			
PTH 1-34	Plasmid, collagen	Rat femur	73
VEGF	Plasmid, collagen	Rabbit radius	75
BMP-2	Plasmid, collagen-CaP	Rat tibia	76
BMP-2	Liposomes, hydroxyapatite	Rabbit cranium	78
BMP-2	Bioglass, adenovirus	Rat tibia	79
BMP-4	Plasmid, poly(ethylenimine)	Rat cranial defect	77

Transgene product	Vector	Model	Ref.
PDGF	Adenovirus	Rat peridontal	80
VEGF + RANKL	AAV	Mouse femur	82
caAlk-2	AAV	Mouse femur	83
Facilitated ex vivo delivery			
LMP-1	Adenovirus, marrow buffy coats	Rabbit spine fusion	28
BMP-2	Adenovirus, muscle, fat	Rat femur	46

^aStably transfected cell line.

^bTransduced, irradiated line of human chondrocytes; osteoporotic rats.

Abbreviations: AAV, attenuated adenovirus; BMP, bone morphogenic protein; COX-2, cyclooxygenase-2; CaP, calcium phosphate; caAlk-2, constitutively active Alk-2; FGF, fibroblast growth factor; IGF, insulin-like growth factor; LMP, LIM mineralisation protein; PDGF, platelet-derived growth factor; PTH, parathyroid hormone; RANKL, receptor activator of nuclear factor kappa-B ligand; VEGF, vascular endothelial growth factor.

Table 2

Use of gene therapy to treat osseous defects in large-animal models

Gene product	Vector	Strategy	Model	Ref.
BMP2	Adenovirus	Traditional ex vivo	Goat tibial defect	106, 107
BMP2	Adenovirus	Traditional ex vivo	Horse metacarpus/metatarsus	55
BMP2	Adenovirus	Traditional ex vivo	Goat, osteonecrosis femoral head	108
BMP2	Adenovirus	Traditional ex vivo	Pig cranial defect	109
PTH 1-34	Plasmid-collagen	GAM	Dog femur Horse carpus	74
BMP2, BMP6	Adenovirus	In vivo	Horse metatarsal	69
BMP2	Adenovirus	In vivo	Sheep tibia	71, 72

Abbreviations: BMP, bone morphogenic protein; GAM, gene-activated matrix; PTH, parathyroid hormone.