Published in final edited form as:

Ann Surg. 2015 July ; 262(1): 171–178. doi:10.1097/SLA.0000000000000883.

Surgeon and Facility Variation in the Use of Minimally Invasive Breast Biopsy in Texas

Nina P. Tamirisa, MD^{1,2}, Kristin M. Sheffield, PhD¹, Abhishek D. Parmar, MD^{1,2}, Christopher J. Zimmermann, MS¹, Deepak Adhikari, MS, Gabriela M. Vargas, MD¹, Yong-Fang Kuo, PhD³, James S. Goodwin, MD¹, and Taylor S. Riall, MD, PhD¹

¹Departments of Surgery, The University of Texas Medical Branch, Galveston, Texas

²The University of California, San Francisco-East Bay, Oakland, California

³Department of Internal Medicine, The University of Texas Medical Branch, Galveston, Texas

Abstract

OBJECTIVE AND SUMMARY BACKGROUND DATA—Minimally invasive breast biopsy (MIBB) rates remain well below guideline recommendations of >90% and varies across geographic areas. Our aim was to determine the variation in use attributable to the surgeon and facility and determine the patient, surgeon, and facility characteristics associated with use of MIBB.

METHODS—We used 100% Texas Medicare claims data (2000–2008) to identify women >66 years with a breast biopsy (open or minimally invasive) and subsequent breast cancer diagnosis/operation within 1 year. The percentage of patients undergoing MIBB as the first diagnostic modality was estimated for each surgeon and facility. Three-level hierarchical generalized linear models (patients clustered within surgeons within facilities) were used to evaluate variation in MIBB use.

RESULTS—22,711 patients underwent a breast cancer operation by 1,226 surgeons at 525 facilities. MIBB was the initial diagnostic modality in 62.4% of cases. Only 7.0% of facilities and 12.9% of surgeons used MIBB in >90% of patients. In 3-level models adjusted for patient characteristics, the percentage of patients who received MIBB ranged from 7.5%–96.0% across facilities (mean 50.1%, median 49.2%) and from 8.0% to 87.0% across surgeons (mean 50.3%, median 50.9%). 28.8% of the variance in MIBB was attributable to the facility and 15.4% to the surgeon. Lower surgeon and facility volume, longer surgeon years in practice and smaller facility bed size were associated with lower rates of MIBB use.

CONCLUSION—Identification of surgeon and facility characteristics associated with low use of MIBB provides potential targets for interventions to improve MIBB rates and decrease variation in use.

Keywords

Minimally invasive breast biopsy; surgeon characteristics; hierarchical models

INTRODUCTION

For patients presenting with palpable breast masses or mammographic abnormalities, minimally invasive breast biopsy (MIBB) offers several advantages over open surgical biopsy. Diagnostic accuracy is similar for both procedures, but women who undergo MIBB experience less peri-procedural pain and have lower rates of post-procedure complications.¹ In cases of malignancy, an MIBB approach is more cost-effective and reduces the overall number of surgical procedures.^{2,3} The minimally invasive approach also provides clinicians with the opportunity for multidisciplinary planning prior to surgical intervention. The 2009 National Comprehensive Cancer Network (NCCN) guidelines recommend MIBB as the gold standard and first line approach to the diagnosis of suspicious breast masses, citing a target of >90% MIBB rates for women presenting with palpable breast masses or mammographic abnormalities requiring biopsy.⁴

Despite these guidelines and the advantages associated with MIBB, population-based studies have demonstrated that MIBB rates are significantly below the NCCN target of >90%.^{5–9} While these studies demonstrate an improvement in MIBB rates since the initial 2001 NCCN consensus statement, the use of open biopsy remains unacceptably high, with open biopsy rates exceeding 20%–30% reported in observational studies through 2008.^{7–11} Furthermore, a recent population-based study from Texas found variation in the use of MIBB across geographic areas (hospital service areas) with similar demographic characteristics and access to MIBB. In addition, improvement in the use of MIBB across geographic regions was variable over time.⁶ These findings suggest that physician and facility practice patterns may explain some of the observed geographic variation in the use of MIBB.^{6,9}

The aim of this study was to evaluate the proportion of variance in MIBB use attributable to the surgeon and facility and to evaluate the surgeon and facility characteristics associated with low MIBB use. We hypothesize that a large portion of the variance in MIBB use across geographic regions can be explained by facility and physician practice patterns. Identifying characteristics of facilities and physicians associated with low MIBB use will highlight potential targets to improve the delivery of patient care and meet the NCCN target guidelines of >90% MIBB rates.

METHODS

The study was reviewed by the Institutional Review Board at the University of Texas Medical Branch, Galveston and granted exemption, as it was not considered human subjects research.

Data Source

This study used enrollment and claims data for 100% of Medicare beneficiaries in the state of Texas from 2000 to 2008. Demographic and enrollment information for each beneficiary was obtained from the Denominator File. Race/ethnicity was assigned based on the Medicare enrollment race variable for patients who did not have Part D data available. The

Outpatient Standard Analytic File (OUTSAF), which contains claims submitted by institutional outpatient providers, and the Carrier Standard Analytic File, which contains claims submitted by non-institutional providers, were used to identify outpatient facility services and physician services. The Medicare Provider Analysis and Review (MEDPAR) files were used to obtain inpatient hospital admissions and claims data.

U.S. Census data for the year 2000 provided ZIP code level education and population estimates. The ZIP code level income was obtained from the 2006 ZIP Code Tabulation Area population estimations produced by the Dartmouth Atlas of Health Care.

Cohort Selection (Figure 1)

The study cohort selection is summarized in Figure 1. We used Medicare claims data to identify all minimally invasive and open breast biopsies performed between 2001 and 2008 for any reason, including breast masses and mammographic abnormalities as previously described; and the CPT codes for minimally invasive and open biopsy are shown in Table 1.⁶ Current Procedural Terminology codes 10021 and 10022 identify FNA done for any reason. Therefore, to identify FNAs done specifically for breast lesions, we chose only those associated with a diagnosis of breast mass, benign or malignant (ICD-9 diagnosis codes 174.0 to 174.9, 217, 233.0, 238.3, 239.3, 610.0 to 610.9, 611.0 to 611.9).

In order to identify the treating surgeon and facility, we then limited our cohort to women with a diagnosis of breast cancer defined by identification of ICD-9-CM diagnosis codes for breast cancer with a subsequent breast cancer operation in the year after the initial biopsy, also shown Table 1. If a woman had more than one breast cancer over the time period, we included the first episode only. We also excluded patients in whom we could not identify the surgeon (N=1,031) or the facility (N=4,377) and those who had operations in facilities outside of Texas (N=426). The final study cohort included 22,711 women with surgically treated breast cancers.

Biopsy

For each patient, the initial breast biopsy in the episode of care was classified as either minimally invasive or open as previously described.⁶ MIBB included fine needle aspiration and core-needle biopsies. Open surgical procedures included incisional and excisional biopsies.

Facility Identification

The facility where the breast cancer operation was performed was identified by the facility ID on the OUTSAF or MEDPAR claim. Facility ID was linked to the Provider of Service file from CMS to obtain facility characteristics, including type of hospital (non-profit, profit, government), bed size (<200, 200–324, 325–500, >500), and medical school affiliation (major, limited, graduate, and none). For procedures done at ambulatory surgical centers (N=1,164) we were unable to determine facility characteristics.

Surgeon Identification

The surgeon who performed the breast cancer operation was identified by the Unique Provider Identification Number (UPIN, 2001–2007) or National Provider Identifier (NPI, 2008) number on the OUTSAF or Carrier claim for the operation. Surgeon UPIN was linked to the American Medical Association Physician Masterfile to obtain surgeon age, sex, year of medical school graduation, graduation from U.S. or foreign medical school, specialty in surgery (surgical oncology vs. general surgery vs. others), and board certification status. Years in practice were calculated from Medical School graduation date. As age and years in practice were collinear, only years in practice was used in the multivariable models. For procedures done at ambulatory surgical centers (ASCs) we could not identify individual surgeons for breast cancer operations because an ASC UPIN is used on these claims.

Patient Characteristics

Patient covariates included age, race/ethnicity (white, black, Hispanic, and other), year of biopsy, and size of the Metropolitan Statistical Area (<10,000, 10,000–250,000, 250,001 to 1 million, and ≥1 million population). Beneficiary race/ethnicity was obtained from the Part D Denominator File (2006–2008), which uses first and last name algorithms to designate race/ethnicity.¹² Race/ethnicity was assigned based on the Medicare enrollment race variable for patients who did not have Part D data available. Median income and education levels (percentage of residents with <12 years of education) in the ZIP code were stratified into quartiles.

Statistical Analysis

Unadjusted differences in the use of MIBB and open biopsy were compared by patient, surgeon, and facility characteristics using chi-square tests for categorical variables and t-tests for continuous variables. From the initial cohort of 22,711 patients, 525 facilities, and 1,226 surgeons, we only included facilities or surgeons that performed >5 cases to create stable estimates of variation in unadjusted and multilevel models. There were 295 facilities performing >5 cases (N=22,331 patients) and 793 surgeons performing >5 cases (N=21,902 patients). We calculated the percentage of the total number of patients treated by each facility or surgeon who underwent MIBB as the first diagnostic modality and evaluated the unadjusted range of MIBB across the 295 facilities and 793 surgeons.

We used multilevel hierarchical modeling to evaluate the variation in the use of MIBB in Texas Medicare beneficiaries treated for breast cancer (2000–2008). In multilevel analyses, we adjusted for patient characteristics including age, race/ethnicity, education, size of the Metropolitan Statistical Area (MSA), and year of biopsy. With various levels of data clustered within each other, hierarchical modeling allows for the estimation and partitioning of variance in MIBB use between the patient, surgeon, and facility levels. We used both 2-level [patients (level 1) clustered within facilities (level 2), N=22,331] and 3-level [patients (level 1) clustered within surgeons (level 2) and surgeons clustered within facilities (level 3), N=17,076] hierarchical models. To address issues with cross-classification in the 3-level model, surgeons who operated in more than one facility were assigned to the facility where they did the most number of cases. Cases performed outside the facility where they did the greatest number of cases (N=4,921) and cases done by surgeons or facilities with <5 cases

(N=714) were removed. The final cohort for the 3-level model included 17,076 patients undergoing breast cancer operations by 792 surgeons at 229 facilities. A 2-level model of patients clustered within facilities was performed with the smaller, cross-classified cohort as well. We estimated adjusted variation in surgeon and hospital MIBB use in both the 2- and 3-level models. To prevent overestimation of the true variation, we generated empirical Bayes “shrunk” estimates.¹⁴ To account for chance in these models, the unstructured covariance of random effects was used to allow for each variance and covariance to be distinct.¹⁵

We estimated the intraclass correlation coefficients (ICC) in each model using the threshold technique that is appropriate for dichotomous outcomes.¹³ The residual ICCs represent the percentage of the total variance in MIBB use attributable to each level of the model. The 2-level model was performed to determine the percent of the variance in MIBB use attributable to the facility alone. The 3-level model was used to determine the percentage of the variance attributable to the surgeon as well as the percentage of the facility variance that was explained by the surgeon. The change in the ICC for facility levels between the 2 and 3-level models can be interpreted as the amount of facility variation explained by surgeon characteristics. The 2 and 3-level models were used to calculate adjusted facility and surgeon rates of MIBB, represented graphically in caterpillar plots. In these plots, each surgeon- and facility-specific rate was adjusted toward the mean of the overall rate as a factor of panel size. For surgeons or facilities with low caseloads, the rates were less reliable and had more adjustment toward the mean and for those with high caseloads, less adjustment was required.

To determine surgeon and facility characteristics associated with the use of MIBB, additional 3-level hierarchical generalized models were estimated including patient, surgeon, and facility-level predictors of MIBB. Since we did not have facility characteristics for ASCs, cases performed at ASCs along with surgeons and facilities performing <5 cases were removed (N=1,260) for a total cohort of 16,530 cases within this model. Statistical significance was accepted at the $p < 0.05$ level. All analyses were performed with were performed with SAS version 9.3 (SAS, Inc.) and STATA 13. All multilevel models and empirical Bayes estimates were performed with STATA 13.

RESULTS

Patient, Surgeon, and Facility Characteristics

22,711 patients undergoing breast biopsy met our inclusion criteria. Table 2 demonstrates overall cohort characteristics and the results of the unadjusted analysis demonstrating MIBB rates by patient, surgeon, and facility characteristics. MIBB was the initial diagnostic modality for 14,171 patients (62.4%). MIBB was performed by a radiologist in 70.0% of cases, by surgeons in 24.9% of cases, and other physicians of other specialties in 5.1%. 85.9% of open biopsies were performed by surgeons. Of note, in 86.6% of the 8,254 who underwent open biopsy as the initial treatment modality, the surgeon who performed the definitive operation also performed the breast biopsy.

In an unadjusted analysis, patient characteristics associated with lower use of MIBB included Hispanic race, lower education, and residence in non-metropolitan or smaller areas. Surgeon characteristics associated with low use of MIBB included male gender, training outside of the United States, and lower case volume. Facility characteristics associated with low use of MIBB included non teaching hospitals, <200 beds, and low case volume (Table 2).

Unadjusted Variation in MIBB Use

There was significant variation in the use of MIBB as the initial diagnostic modality across facilities and surgeons. Only 7.0% of facilities and 12.9% of surgeons performed MIBB in >90% of patients. MIBB use across 295 facilities ranged from 0% to 100% with a mean of 50.9% (median=53.2%, IQR = 27.1%–74.2%). The unadjusted MIBB use across 793 surgeons also ranged from 0 to 100%. Of note, the unadjusted mean percentage of patients receiving MIBB across surgeons was 56.4% (median=60.0%, IQR 32.0%–80.9%).

Percentage of Variance in MIBB Use Attributable to Facility and Surgeon (Table 3)

In the 2-level hierarchical model (patients clustered within facilities) adjusted for patient characteristics, 33.6% of the variance in MIBB use was attributable to the facility (ICC=33.6%, Table 3, Model 1). Of note, when the 2-level model of patients clustered within facilities was performed with the cohort excluding cross-classified cases, the ICC was similar at 35.8% for facilities (ICC=35.8%, Table 3, Model 2).

We then identified the variance in MIBB use at the surgeon and facility levels using a 3-level model of patients clustered within surgeons clustered within facilities (Table 3, Model 3). In the 3-level model, the ICC for the surgeon contribution to variance in MIBB was 15.4% (Table 3, Model 3). The percentage of variance in MIBB use attributable to the facility decreased from 33.6% in the 2-level model (Table 3, Model 1) to 28.8% in the 3-level model (Table 3, Model 3) demonstrating that approximately 15% of the facility-level variance was explained by the surgeon. These models are depicted graphically.

From the 2-level model of patients clustered within facilities (Table 3, Model 1) the adjusted rates of MIBB across facilities were plotted by rank in MIBB use from lowest to highest, ranging from 5.0% to 96.3% (Figure 2A). Of these 295 facilities, 24.1% (N=71) had rates of MIBB use significantly below and 28.1% (N=83) had rates significantly above the adjusted mean facility rate of 51.7% (Figure 2a, $p<0.05$). Figures 2b and 3 reflect the 3-level model of patients clustered within surgeons clustered within facilities. Using the 3-level model to evaluate facility variance, 6.1% of facilities had rates significantly lower than the mean MIBB use of 50.1%; 13.5% of facilities had rates significantly higher than mean ($p<0.05$). Facility variation in the 3-level model is shown in Figure 2b. Variability in MIBB use across facilities decreased to 7.5% – 96.0% (median = 49.2, IQR = 33.4%–68.3%) after adjusting for surgeon factors. Comparing Figure 2a and Figure 2b, facility variation in MIBB use diminished after accounting for surgeons in the model. In other words, a portion of the variation among facilities was due to variation in practice patterns among surgeons.

We then used the 3-level model to evaluate adjusted surgeon variation. Adjusted MIBB use across 792 surgeons ranged from 8.0% to 87.0% (Figure 3). One hundred and nine surgeons

(13.8%) had rates of MIBB use significantly different than the mean; 6.9% were significantly lower and 6.8% significantly higher than the mean of 50.3% (median = 50.9, IQR =39.8%–61.2%).

Patient, Surgeon, and Facility Characteristics Associated with MIBB Use (Table 4)

Surgeon and facility characteristics were added to the 3-level hierarchical model above to determine the patient, surgeon, and facility factors associated with high MIBB use (N=16,530, ambulatory surgery centers excluded). Surgeons with more years in practice and surgeons with lower case volume were less likely to use MIBB. Facility factors associated with lower MIBB use included smaller facility bed size and low facility volume.

DISCUSSION

To our knowledge, ours is the first study to use multilevel hierarchical models to evaluate surgeon- and facility-level variation in the use of MIBB as the initial diagnostic modality in a woman presenting with a palpable breast mass or mammographic abnormality. Consistent with previous population-based studies,^{6,10,11} MIBB use in our cohort fell well below the NCCN target rate of 90%, with only 62.4% of women undergoing MIBB as the initial diagnostic modality. We identified only 7.0% of facilities and 13% of surgeons performing MIBB in >90% of their breast cancer cases. We observed wide variation in the use of MIBB across surgeons (8.0% to 87.0%) and facilities (7.5% to 96.0%). In our multilevel models, over 28% of the variance in MIBB use was attributable to the facility and 15% attributable to the surgeon. Among the 295 facilities that performed > 5 cases, 279 facilities had the capacity to perform MIBB, therefore availability of technology was not a driver of variation in our study. Finally, we also identified specific surgeon and facility characteristics associated with a low use of MIBB.

The role of physician characteristics in practice variation across a variety of disease processes has been previously evaluated.^{16–21} In a retrospective cohort study, Feinstein et al. used hierarchical linear regression models to demonstrate that surgeons accounted for as high as 15.7% of the variance in a patient's receipt of radiation therapy after breast conserving surgery.²¹ This is similar to the findings in our study where the percentage of variance in MIBB use attributable to the surgeon was 15.4%. When the amount of practice variation attributable to the physician is high (as in our study), efforts directed at changing physician behavior would potentially have a significant impact on outcomes. As a result, our findings suggest that interventions targeted at changing physician behaviors could potentially improve MIBB rates dramatically. This idea is supported by a single-institution quality improvement initiative by Lovrics et al., where surgeon directed strategies resulted in the increase of MIBB rates from 73% to 92%.²²

Previous studies have demonstrated similar findings of facility variation in practice patterns.^{23,24} In a study evaluating blood transfusion and patient outcomes in cardiac surgery, 30% of the variance in transfusion practices was attributable to the facility despite specific recommendations from evidence based guidelines.²⁴ In such cases, efforts directed at implementing a hospital wide quality improvement program improved guideline adherence rates.²⁵ In our study, a comparable percentage of the variation in MIBB use was

attributable to the facility at 28.8%. We identified lower volume facilities and smaller facilities as having lower rates of MIBB use. Quality improvement initiatives targeted at low-volume, smaller hospitals may improve MIBB rates at the facility level, leading to decreased variation.

Finally, we observed that U.S.-trained surgeons with high volumes were more likely to use MIBB. We also found an inverse correlation between surgeon years in practice and use of MIBB. This may represent a higher proportion of young surgeons with specialized training and practices devoted to breast disease or an increased willingness of young surgeons to adopt current guidelines. A previous study using Medicare claims data also identified an association between high surgeon volume and adherence to guidelines, specifically adequate lymphadenectomy in early stage breast cancer.²⁶ In a single-institution, retrospective study of 465 breast cancer patients, Clarke-Pearson et al. identified differences in MIBB rates between academic breast surgeons (91%), private practice breast surgeons on the clinical faculty (74%), and general surgeons performing breast cases (58%) as part of their practice.⁷ While we were not able to determine the proportion of a surgeon's practice dedicated to breast disease or surgeon's who were breast fellowship trained, we also observed a volume-outcomes relationship in the use of MIBB. Our findings support that interventions to improve physician use of MIBB may be directed towards underperforming surgeons with longer years in practice and lower volumes.

Our study assigns patients to the surgeon or hospital that did the definitive breast cancer operation. However, previous data demonstrate that over 70% of minimally invasive biopsies are performed by radiologists and not the surgeon doing the definitive breast cancer operation.⁶ Likewise, it is possible that surgeons who did the definitive operation but not the original open biopsy would be incorrectly classified as having made this decision, when instead the patient was referred after the open biopsy had occurred. However, 86.6% of the open biopsies performed were performed by the same surgeon that did the biopsy. 356 of the 792 surgeons in the cohort operated on one or more of the 1,146 patients in whom the open biopsy was performed by a different surgeon. As such, most surgeons only had few cases that were inappropriately classified. Conversely, many patients may have minimally invasive biopsy before being referred to a surgeon, who might have otherwise done the biopsy open. We chose the surgeon doing the breast cancer operation as the unit of analysis because this person is easy to identify in the claims data, but we feel this represents a larger systemic issue and reflects both referral and practice patterns and should be evaluated in this context.

In many settings, such as large academic, multidisciplinary breast centers, mammography and biopsy occur without surgeon involvement and surgeons only evaluate patients after a tissue diagnosis is established and multidisciplinary consultation is underway. It is likely that in this setting, patients referred without biopsy would be sent for minimally invasive biopsy as this is standard practice. This is evidenced by reports of MIBB exceeding 90% in academic centers.²⁷ In private settings, surgeons may be referred patients with breast masses prior to diagnosis and open breast biopsy may represent a significant proportion of a surgeon's practice. In the study by Clarke-Pearson and colleagues, diagnostic excisional biopsy made up 10% of academic breast surgeon practices, but 35% of private practice

breast surgeon practices, and 37% of general surgeon practices. Increasing referring physician awareness as well as surgeon awareness of the current recommendations may lead to improved MIBB rates.

Our study has several limitations. Our study cohort was limited to women who were Medicare beneficiaries >66 years in the state of Texas, and results may not be generalizable to other age groups or geographic regions. In addition, we included only patients with breast cancer. Rates of open biopsy have been shown to be higher in patients with benign disease,⁶ so we may have actually overestimated rates of MIBB use. In addition, we limited our cohort to patients who had a cancer operation after biopsy, in order to identify the surgeon and the facility. Patients with breast cancer may be more likely to be treated by a breast specialist or at a multidisciplinary center. Variability in the use of MIBB across surgeons and facilities may differ for a cohort of women that includes both benign and malignant disease. Finally, we were able to identify surgeon specialty as surgical oncology vs. general surgery using the Medicare claims. However, these data are self-reported and have not been validated. We were unable to identify surgeons that are breast fellowship trained or have practices devoted to breast disease.

In conclusion, we used multilevel hierarchical modeling techniques to demonstrate that a large amount of the observed variation in MIBB use was attributable to both surgeon and facility. In addition, we identified several surgeon and facility factors associated with low MIBB use. These data points can be used as specific targets for intervention to achieve MIBB rates of >90% for breast cancer patients in Texas.

ACKNOWLEDGEMENTS

This study was supported by grants from the National Institute of Health (T32 DK007639), the UTMB Clinical and Translational Science Award UL1TR000071, the Cancer Prevention Research Institute of Texas (RP101207-P03), and the Agency for Healthcare Research and Quality (1R24HS022134). The empirical bayes estimations were generated with the assistance of Dr. Nai Wei Chen from UTMB Galveston.

Funding: Cancer Prevention Research Institute of Texas Grant # RP101207-P03, UTMB Clinical and Translational Science Award #UL1TR000071, NIH T-32 Grant # T32DK007639, AHRQ Grant # 1R24HS022134.

REFERENCES

1. Verkooijen HM, Buskens E, Peeters PH, et al. Diagnosing non-palpable breast disease: short-term impact on quality of life of large-core needle biopsy versus open breast biopsy. *Surg Oncol.* 2002; 10:177–181. [PubMed: 12020672]
2. Hatmaker AR, Donahue RM, Tarpley JL, Pearson AS. Cost-effective use of breast biopsy techniques in a Veterans health care system. *Am J Surg.* 2006; 192:e37–e41. [PubMed: 17071179]
3. Bruening W, Fontanarosa J, Tipton K, Treadwell JR, Launder J, Schoelles K. Systematic review: comparative effectiveness of core-needle and open surgical biopsy to diagnose breast lesions. *Ann Intern Med.* 2010; 152:238–246. [PubMed: 20008742]
4. Bevers TB, Anderson BO, Bonaccio E, et al. NCCN clinical practice guidelines in oncology: breast cancer screening and diagnosis. *J Natl Compr Canc Netw.* 2009; 7:1060–1096. [PubMed: 19930975]
5. Williams RT, Yao K, Stewart AK, et al. Needle versus excisional biopsy for noninvasive and invasive breast cancer: report from the National Cancer Data Base, 2003–2008. *Ann Surg Oncol.* 2011; 18:3802–3810. [PubMed: 21630122]

6. Zimmermann CJ, Sheffield KM, Duncan CB, et al. Time trends and geographic variation in use of minimally invasive breast biopsy. *J Am Coll Surg.* 2013; 216:814–824. discussion 824-817. [PubMed: 23376029]
7. Clarke-Pearson EM, Jacobson AF, Boolbol SK, et al. Quality assurance initiative at one institution for minimally invasive breast biopsy as the initial diagnostic technique. *J Am Coll Surg.* 2009; 208:75–78. [PubMed: 19228506]
8. Breslin TM, Caughran J, Pettinga J, et al. Improving breast cancer care through a regional quality collaborative. *Surgery.* 2011; 150:635–642. [PubMed: 22000174]
9. Holloway CM, Saskin R, Brackstone M, Paszat L. Variation in the use of percutaneous biopsy for diagnosis of breast abnormalities in Ontario. *Ann Surg Oncol.* 2007; 14:2932–2939. [PubMed: 17619931]
10. Gutwein LG, Ang DN, Liu H, et al. Utilization of minimally invasive breast biopsy for the evaluation of suspicious breast lesions. *Am J Surg.* 2011; 202:127–132. [PubMed: 21295284]
11. Frieze CR, Neville BA, Edge SB, Hassett MJ, Earle CC. Breast biopsy patterns and outcomes in Surveillance, Epidemiology, and End Results-Medicare data. *Cancer.* 2009; 115:716–724. [PubMed: 19152430]
12. Agency for Healthcare Research and Quality. [Accessed July 20, 2013] Creation of new race ethnicity codes and socioeconomic status (SES) indicators for Medicare beneficiaries. Available at: <http://www.ahrq.gov/libux.utmb.edu/qual/medicareindicators/medicareindicators.pdf>.
13. Snijders, TA.; Bosker, RJ. Multilevel analysis: an introduction to basic and advanced multilevel modeling. Thousand Oaks: SAGE Publications; 1999.
14. Seymour CW, Iwashyna TJ, Ehlenbach WJ, Wunsch H, Cooke CR. Hospital-level variation in the use of intensive care. *Health Serv Res.* 2012; 47:2060–2080. [PubMed: 22985033]
15. [Accessed May 2014] <http://www.stata.com/manuals13/metoc.pdf>.
16. Beaulieu MD, Blais R, Jacques A, Battista RN, Lebeau R, Brophy J. Are patients suffering from stable angina receiving optimal medical treatment? *Qjm.* 2001; 94:301–308. [PubMed: 11391028]
17. Cowen ME, Strawderman RL. Quantifying the physician contribution to managed care pharmacy expenses: a random effects approach. *Med Care.* 2002; 40:650–661. [PubMed: 12187179]
18. Hofer TP, Hayward RA, Greenfield S, Wagner EH, Kaplan SH, Manning WG. The unreliability of individual physician "report cards" for assessing the costs and quality of care of a chronic disease. *Jama.* 1999; 281:2098–2105. [PubMed: 10367820]
19. Shahinian VB, Kuo YF, Freeman JL, Goodwin JS. Determinants of androgen deprivation therapy use for prostate cancer: role of the urologist. *J Natl Cancer Inst.* 2006; 98:839–845. [PubMed: 16788157]
20. Sixma HJ, Spreeuwenberg PM, van der Pasch MA. Patient satisfaction with the general practitioner: a two-level analysis. *Med Care.* 1998; 36:212–229. [PubMed: 9475475]
21. Feinstein AJ, Soulos PR, Long JB, et al. Variation in receipt of radiation therapy after breast-conserving surgery: assessing the impact of physicians and geographic regions. *Med Care.* 2013; 51:330–338. [PubMed: 23151590]
22. Lovrics P, Hodgson N, O'Brien MA, et al. The implementation of a surgeon-directed quality improvement strategy in breast cancer surgery. *Am J Surg.* 2013
23. Goodnough LT, Johnston MF, Toy PT. The variability of transfusion practice in coronary artery bypass surgery. Transfusion Medicine Academic Award Group. *Jama.* 1991; 265:86–90. [PubMed: 1984129]
24. Rogers MA, Blumberg N, Saint S, Langa KM, Nallamotheu BK. Hospital variation in transfusion and infection after cardiac surgery: a cohort study. *BMC Med.* 2009; 7:37. [PubMed: 19646221]
25. Brevig J, McDonald J, Zelinka ES, Gallagher T, Jin R, Grunkemeier GL. Blood transfusion reduction in cardiac surgery: multidisciplinary approach at a community hospital. *Ann Thorac Surg.* 2009; 87:532–539. [PubMed: 19161774]
26. Gilligan MA, Neuner J, Sparapani R, Laud PW, Nattinger AB. Surgeon characteristics and variations in treatment for early-stage breast cancer. *Arch Surg.* 2007; 142:17–22. [PubMed: 17224496]

27. Linebarger JH, Landercasper J, Ellis RL, et al. Core needle biopsy rate for new cancer diagnosis in an interdisciplinary breast center: evaluation of quality of care 2007–2008. *Ann Surg.* 2012; 255:38–43. [PubMed: 22167007]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.

Cohort Selection. We identified all breast biopsies for women age 66 and older in Texas, 2001–2008. Unique episodes of care and unique patients were included. Only women with a breast cancer diagnosis and primary operation for breast cancer were included. N=22,711.

Figure 2.

a. Empirical Bayes shrunk estimates of facility variation in the adjusted rates of MIBB based on the 2-level hierarchical model adjusted for patient characteristics (N=22,331). 295 facilities performed >5 surgeries and were ranked from lowest to highest MIBB use with a range of 5.0%–96.3%. The horizontal line represents the overall mean rate of MIBB use. Error bars represent 95% confidence intervals for the MIBB rates of individual facilities. Black error bars represent facilities that have rates statistically significantly ($p < 0.05$) above or below the mean rate of 51.7% (median = 51.0%; IQR = 31.4%–73.1%) and light gray bars represent rates that are not different from the mean rate.

b. Empirical Bayes shrunk estimates of facility variation in the adjusted rates based on 3-level hierarchical model (patients clustered with surgeons and surgeons clustered with facilities) adjusted for patient characteristics (N=17,076). 229 facilities performed >5 surgeries and were ranked in the 3-level model from lowest to highest MIBB use with a range of 7.5%–96.0%. The horizontal line represents the overall mean rate of MIBB use. Error bars represent 95% confidence intervals for the MIBB rates of individual facilities. Black error bars represent facilities that have rates statistically significantly ($p < 0.05$) above or below the mean rate of 50.1% (median = 49.2%; IQR = 33.4%–68.3%) and light gray bars represent rates that are not different from the mean rate.

Figure 3.

Empirical Bayes shrunk estimates of surgeon variation in the adjusted rates of MIBB use for 792 surgeons doing >5 cases were ranked from lowest to highest MIBB use with a range of 8.0%–87.0% (N=17,076). Rates were generated from the 3-level hierarchical model (adjusted for patient characteristics). The horizontal line represents the overall mean rate of MIBB use. Error bars represent 95% confidence intervals for the MIBB rates of individual surgeons. Black error bars represent facilities that have rates statistically significantly ($p < 0.05$) above or below the mean rate of 50.3% (median = 50.9%; IQR = 39.8%–61.2%) and light gray bars represent rates that are not different from the mean rate.

Table 1

Biopsy Modality and Cancer Operation (N=22,711).

TO IDENTIFY ALL BREAST BIOPSIES FOR ANY REASON	
CPT or ICD-9 Code	Procedure
Core Biopsy	
19100	Breast biopsy; percutaneous, needle core, not using imaging guidance
19102	Breast biopsy, percutaneous, needle core with imaging guidance
19103	Biopsy of breast; percutaneous, automated vacuum assisted or rotating biopsy device, using imaging guidance
Open	
19101	Breast biopsy; open, incisional
19120	Excision of cyst, fibroadenoma, or other benign or malignant tumor, aberrant breast tissue, duct lesion, nipple or areolar lesion, open, one or more lesions
19125	Excision of breast lesion identified by preoperative placement of radiological marker, open; single lesion
FNA	
10021 *	Fine-needle aspiration; not using imaging guidance
10022 *	Fine-Needle aspiration; using imaging guidance
TO IDENTIFY PATIENTS WITH BREAST CANCER AFTER BIOPSY	
Breast Cancer	
174.0 **	Malignant neoplasm of nipple and areola of female breast
233.0 **	Carcinoma in situ of breast
19110	Nipple exploration, with or without excision of a solitary lactiferous duct or a papilloma lactiferous duct
19120	Excision of cyst, fibroadenoma, or other benign or malignant tumor, aberrant breast tissue, duct lesion, nipple or areolar lesion, open, one or more lesions
19125	Excision of breast lesion identified by preoperative placement of radiological marker, open; single lesion
19126	Excision of breast lesion identified by preoperative placement of radiological marker, open; each additional lesion separately identified by a preoperative radiological marker
19160	Mastectomy, partial
19162	Mastectomy, partial, with axillary lymphadenectomy
19301	Mastectomy, partial (eg, lumpectomy, tylectomy, quadrantectomy, segmentectomy)
19302	Mastectomy, partial (eg, lumpectomy, tylectomy, quadrantectomy, segmentectomy); with axillary lymphadenectomy
19180	Mastectomy, simple, complete
19182	Mastectomy, subcutaneous
19200	Mastectomy, radical, including pectoral muscles, axillary lymph nodes
19220	Mastectomy, radical, including pectoral muscles, axillary, and internal mammary lymph node
19240	Mastectomy, modified radical, including axillary lymph nodes, with or without pectoralis minor muscle, but excluding pectoralis major muscle
19303	Mastectomy, simple, complete
19304	Mastectomy, subcutaneous
19305	Mastectomy, radical, including pectoral muscles, axillary lymph nodes
19306	Mastectomy, radical, including pectoral muscles, axillary and internal mammary lymph nodes
19307	Mastectomy, radical, including pectoral muscles, axillary and internal mammary lymph nodes

* CPT codes 10021 and 10022 identify fine-needle aspiration done for any reason. Specifically for breast lesions, we chose only those associated with a diagnosis of breast mass, benign or malignant (ICD-9 codes: 174.0–174.9, 217, 233.0, 238.3, 239.3, 610.0–610.9, 611.0–611.9).

** To identify facilities and surgeons we chose breast cancers associated with an operation in the year after the initial biopsy (ICD-9-CM codes: 85.22–85.26, 85.4–85.48; CPT codes: 19110, 19120, 19125–6, 19160, 19162, 19301–2, 19180, 19182, 19200, 19220, 19240, 19303–7).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Percent Minimally Invasive Breast Biopsy by Patient, Surgeon, and Facility Characteristics (Total N=22,711)

Patient Characteristics	Overall N	MIBB N (% MIBB)	P value
Overall Cohort	22,711	14,171 (62.4)	
Race			<0.0001
White	18,609	11,773 (63.3)	
Black	1,692	1,064 (62.9)	
Hispanic	2,200	1,189 (54.1)	
Other	210	145 (69.1)	
Age			0.4200
66–74	11,705	7,333 (62.7)	
75+	11,006	6,838 (62.1)	
Area of Residence			<0.0001
Metropolitan	17,696	11,484 (64.9)	
Non Metropolitan	4,579	2,433 (53.1)	
Rural	436	254 (58.3)	
Education Level* (Quartile)			<0.0001
Lowest	5,688	3,199 (56.2)	
2	5,675	3,393 (59.8)	
3	5,443	3,481 (64.0)	
4 -Highest	4,995	3,575 (71.6)	
Unknown	910	523 (57.5)	
MSA**			<0.0001
> 1 million	11,419	7,973 (69.8)	
250,000–1 million	2,406	1,136 (47.2)	
10,000 –250,000	3,871	2,375 (61.4)	
<10,000	5,015	2,687 (53.6)	
Surgeon Characteristics			
Age			<0.0001
Mean (SD)	54.45(9.5)	53.74 (9.6)	
Median	54.0	53.0	
Years of practice			<0.0001
Mean (SD)	27.81(10.0)	27.03 (10.1)	
Median	27.0	26.0	
Sex			<0.0001
Male	18,280	10,905 (59.7)	
Female	4,431	3,266 (73.7)	
US Trained			<0.0001
Yes	20,292	13,123 (64.7)	

Patient Characteristics	Overall N	MIBB N (% MIBB)	P value
No	2,419	1,048 (43.3)	
Specialty			<0.0001
General Surgery	20,170	12,472 (61.8)	
Surgical Oncology	1,315	1,031 (78.4)	
Others	1,226	668 (54.5)	
Volume of Surgeon (Quartile)			<0.0001
1-Lowest	5,604	2,994 (53.4)	
2	5,634	3,096 (55.0)	
3	5,891	3,649 (62.0)	
4-Highest	5,582	4,432 (79.4)	
Facility Characteristics ^ψ			
Type of facility			<0.0001
Non-profit	12,030	7,737 (64.3)	
Profit	6,525	3,957 (60.6)	
Government	3,165	1,902 (60.1)	
Teaching Hospital			<0.0001
Yes	8,227	5,652 (68.7)	
No	13,493	7,944 (58.9)	
Facility Bed Size			<0.0001
<200 Beds	5,769	2,749 (47.7)	
200–324 beds	3,714	2,982 (53.4)	
325–500 beds	5,051	3,567 (70.6)	
>500 beds	7,186	5,298 (73.7)	
Volume of Facility (Quartile)			<0.0001
1-Lowest	5,099	2,346 (46.0)	
2	5,652	3,394 (60.1)	
3	5,280	3,343 (63.3)	
4 –Highest	5,689	4,513 (79.3)	
ASC only	991	575 (58.0)	<0.0001

* = Lowest level of >12 years of schooling

** = Metropolitan Statistical Area

^ψ = Hospital patients only

Table 3

Hierarchical Multilevel Models: Percentage of Variance in MIBB use Attributable to Patient, Surgeon, and Facility

Characteristics	Percentage of Variance in MIBB use [*]
	Adjusted for patient characteristics
2-level model (Patient, Facility)	
MODEL 1	22,331
Number of Patients	295
Number of Facilities	33.6%
Residual ICC (% variance) - Facility ^{*†}	
2-level model (Patient, Facility)	
MODEL 2	17,716
Number of Patients	249
Number of Facilities	35.8%
Residual ICC (% variance) - Facility ^{*†}	
3-level model (Patient, Surgeon, Facility)	
MODEL 3	17,076
Number of Patients	792
Number of Surgeons	229
Number of Facilities	15.4%
Residual ICC (% variance) - Surgeon ^{*†}	28.7%
Residual ICC (% variance) - Facility ^{*†}	

Model 1. For the 2-level hierarchical models, only 295 facilities with ≥ 5 cases were included on the initial cohort of 525 facilities for a final N=22,331. **Model 2.** The 2-level hierarchical model was repeated with the cohort of cross-classified cases. Surgeons were assigned to facilities where they performed the greatest number cases, cases done at other facilities (N=4,921) and facilities with < 5 cases (N=714) were excluded resulting in 17,716 patients. **Model 3.** Surgeons were assigned to facilities where they performed the greatest number cases, cases done at other facilities (N=4,921) and facilities with < 5 cases (N=714) were excluded resulting in 17,076 patients. The percentage of variance in MIBB use attributable to the facility decreased from 33.6% in the 2-level model to 28.7% in the 3-level model.

^{*} Hierarchical generalized linear models: 2-level model “level 1” variables are patient characteristics and “level 2” variables are facility identifiers. 3-level model “level 1” variables are patient characteristics, “level 2” variables are surgeon identifiers, and “level 3” variables are facility identifiers. ICC = intraclass correlation coefficient.

[†] The percentage of variance attributable to the surgeon and facility are calculated with a threshold model, after simultaneous adjustment of all available patient characteristics. The denominator for the calculation of the percentage was composed of the variance attributable to the facility (2-level model) or surgeon and facility (3-level model), after adjustment for available patient characteristics, and the variance attributable to unexplained patient variables plus error.

Table 4

Three-Level Model (patients within surgeon within facilities) Adjusted for Patient, Surgeon and Facility Characteristics (N=16,530)

Variables	Odds Ratio	95% CI
Patient Characteristics		
Age (ref: <74) 75+	1.13	1.05–1.23
Biopsy Year (ref: 2001–2003) 2004–2006 2007–2008	1.90 3.44	1.74–2.08 3.05–3.87
Race (ref: White) Hispanic Black Others	0.91 0.97 0.85	0.77–1.07 0.82–1.15 0.55–1.31
Education (ref: Q1) Q2 Q3 Q4-Highest	1.06 0.99 0.98	0.93–1.20 0.87–1.13 0.86–1.12
MSA (ref: 1M) 0.25 – 1M 10,000 – 0.25M <10,000 vs. 1M	0.61 0.69 0.78	0.47–0.81 0.56–0.84 0.66–0.92
Surgeon Characteristics		
Year of Practice (ref: Q1) Q2 Q3 Q4-Highest	0.73 0.54 0.49	0.57–0.94 0.42–0.70 0.38–0.64
Surgeon Gender (ref: Male) Female	1.16	0.87–1.15
US Trained (ref: Yes) No	0.91	0.69–1.21
Specialty (ref: General Surgery) Others Surgical Oncology	0.99 1.25	0.68–1.44 0.64–2.44
Surgeon Volume (ref: Q1) Q2 Q3 Q4-Highest	1.17 1.54 2.14	0.94–1.45 1.19–1.99 1.48–3.11
Facility Characteristics		
Type of Hospital (ref: non-profit) Government Profit	1.27 1.10	0.83–1.94 0.70–1.75
Teaching Hospital (ref: Yes) No	0.99	0.61–1.60
Facility Bed Size (ref: <200 beds) 200–324 325–500 >500	1.54 3.40 3.15	0.92–2.56 1.89–6.13 1.54–5.95
Facility Volume (ref: Q1) Q2 Q3 Q4-Highest	1.48 1.24 2.60	0.92–2.36 0.67–2.29 1.09–5.98