Published in final edited form as:

J Asthma. 2013 April ; 50(3): 223–230. doi:10.3109/02770903.2012.761231.

Enhanced generation of suppressor T cells in patients with asthma taking oral contraceptives

A. Catalina Vélez-Ortega, M.S., Ph.D.c*,

Department of Microbiology, Immunology and Molecular Genetics. University of Kentucky

James Temprano, M.D.*,

Department of Medicine, Division of Allergy and Immunology. University of Kentucky

Mary Catherine Reneer, Ph.D.,

Department of Microbiology, Immunology and Molecular Genetics. University of Kentucky

Gavin I. Ellis, B.S., Ph.D.c,

Department of Microbiology, Immunology and Molecular Genetics. University of Kentucky

Andrea McCool, B.S.,

Department of Microbiology, Immunology and Molecular Genetics. University of Kentucky

Tonya Gardner, C.C.R.C.,

Department of Medicine, Division of Allergy and Immunology. University of Kentucky

Mehdi Khosravi, M.D., and

Department of Medicine, Division of Pulmonary and Critical Care Division. University of Kentucky

Francesc Marti, Ph.D.

Department of Microbiology, Immunology and Molecular Genetics. University of Kentucky

Lexington, KY 40536

Abstract

INTRODUCTION—A dysregulation of regulatory T cells (Tregs) could play a major role in the pathogenesis of bronchial asthma. Sex-dependent differences as well as the impact of hormonal changes in the incidence and severity of asthma are widely recognized. Emerging evidence suggests that asthma symptoms are alleviated in female patients taking hormone oral contraceptives (OCs). The impact of OCs on the generation of induced Tregs (iTregs) was assessed in a cohort of female patients with asthma.

METHODS—Thirteen patients were included in this pilot study. During three distinct phases of their menstrual cycles we measured exhaled nitric oxide (eNO) levels, forced expiratory volume at

Corresponding author: Francesc Marti, Ph.D., 800 Rose St – MN415 UK Medical Center, Phone: 323-7858; Fax: 323-8268, fmart3@uky.edu.

*Both authors contributed equally to this work

Present affiliation: James Temprano, M.D., Department of Internal Medicine, Saint Louis University School of Medicine

DECLARATION OF INTEREST STATEMENT

A.C.V., M.C.R., G.I.E., A.M., T.G., M.K. and F.M. declare no financial or commercial conflict of interest. J.T. receives research funding from Merck and Dyax.

1s (FEV1s), asthma control test (ACT) score, sex steroid hormone levels in serum, natural Tregs in peripheral blood and the ability of CD4⁺ T cells to generate iTregs *ex vivo*.

RESULTS—The luteal serum levels of estradiol and progesterone negatively correlated with the proportion of iTregs generated *ex vivo* in patients not taking OCs. In addition, physiological doses of estradiol and progesterone prevented the acquisition of a suppressor T cell phenotype *in vitro*. Interestingly, patients taking OCs had reduced serum sex hormone levels associated with higher iTreg induction, a better ACT score and a tendency toward lower eNO levels.

CONCLUSIONS—Our results identify an impact of sex hormones in the capacity of T cells to polarize towards a regulatory phenotype and suggest the regulation of peripheral T cell lineage plasticity as a potential mechanism underlying the beneficial effects of OCs in women with asthma.

Keywords

Induced regulatory T cells; menstrual cycle; estradiol; progesterone; ACT score; CD4 T cells

INTRODUCTION

The T helper type 1 (Th1) vs. Th2 dichotomy of T cell responses has been considered for many years a cornerstone of the adaptive immune system. An exaggerated Th2-driven response is believed to lead the inflammatory allergic reaction of asthma, a highly complex, multifactorial disease characterized by airway hyperresponsiveness, inflammation and infiltration of different immune cell types, including eosinophils, mast cells and T lymphocytes. Particularly, Th2 cytokines induce the production of immunoglobulin E and mast cells, activation and recruitment of eosinophils, and airway hyperreactivity and mucus secretion¹. Clinical observations and epidemiological studies suggest a role for sex hormones in the incidence and/or regulation of symptom severity in numerous Th2-related pathologies². Indeed, while the incidence of asthma is higher in boys before puberty, the manifestations of asthma are predominant in females after puberty^{3–7} and up to 30–40% of female patients experience perimenstrual worsening of symptoms^{8,9}. Moreover, the use of oral contraceptives (OCs) has been shown to improve symptoms in some women with asthma^{10–12}. The capacity of the sex steroid hormones estradiol and progesterone as contributing factors in asthma is being actively addressed,¹³ but the underlying immunoregulatory mechanisms have not been fully elucidated yet.

Additional T helper expression profiles that challenge the Th1/Th2 paradigm have been identified. The Th17 population has been implicated in neutrophilic asthma and resistance to steroids in mouse models of asthma^{14,15}. Of major importance are Tregs, which make up 1–5% of the T cells in the peripheral blood of adult mice and humans¹⁶. Tregs can inhibit or suppress the effector function of T helper cells and their tolerance to environmental allergens¹⁷ thus preventing the induction or reducing the severity of asthma. In fact, a perturbed Treg compartment has been associated with autoimmunity, inflammatory disorders and cancer development¹⁸. In addition to the “natural” Tregs (nTregs) that stem as a separate lineage in the thymus, induced Tregs (iTregs) can be generated in peripheral tissues from conventional CD4⁺ T cells. Interestingly, recent evidence supports the

prevailing relevance of iTregs in mucosal tissues and inflammatory processes^{19–21}. As such, any potential mechanism altering the generation, maintenance or function of iTregs might be relevant to the etiology, pathogenesis and/or evolution of asthma. In fact, the pathological Th2 response of asthma patients has been associated with the reduced number or suppressive activity of circulating CD4⁺ CD25⁺FoxP3⁺ Tregs^{22–24}. Consequently, there is a growing interest in the mechanisms that regulate iTreg function and differentiation to therapeutically manipulate their enormous potential in immune and inflammatory conditions, including asthma^{25,26}.

Here we investigate the impact of OC intake on the ability of T cells to generate suppressor iTregs in asthma patients.

METHODS

Patients with asthma and controls

Thirteen female patients with asthma were recruited at the University of Kentucky Clinics. Their asthma severity was mild to moderate persistent according to the National Asthma Education and Prevention Program (NAEPP) treatment guidelines²⁷. All patients were followed for two consecutive months during three specific phases of the menstrual cycle for a total of six study visits. The three time points studied were menses (days 1–5 of the menstrual cycle), mid-cycle (days 12–16) and luteal phase (days 23–27). One cycle (3 visits) from one of the patients was removed because pregnancy was suspected. In each visit we drew approximately 20 mL of a heparinized blood sample. In addition, prick skin testing was performed at the initial visit to assess for the presence of atopy. All study subjects had well-controlled asthma at the initial study visit according to NAEPP guidelines. Six patients were taking combination (estradiol and progesterone or equivalent) OCs and the remaining seven patients were not under any hormone- based contraceptive treatment (non-OC). Patients were non-smokers, did not exhibit other medical conditions and did not receive systemic steroid treatment 4 weeks prior to the study enrollment or during the study. The study protocol was approved by the Medical Institutional Review Board of the University of Kentucky (approval number 06-0578) and written informed consent was obtained from all study participants.

Fresh buffy coats from twenty-six healthy adult (mean age of 34, range 19 to 45) anonymous blood donors with no gender identification were obtained from the Kentucky Blood Center (Lexington, KY) and included for control purposes.

Asthma clinical parameters

All study subjects underwent spirometry in each visit according to American Thoracic Society criteria²⁸. Forced expiratory volume in 1 second (FEV1s) was measured with a MicroLab portable spirometer (Micro Direct, Inc.) at each visit. Asthma control test (ACT) questionnaires were filled out by every subject on each visit²⁹. Modified ACT questionnaires were constructed allowing reported symptoms to only reflect those experienced since the previous study visit; therefore, an assessment could be made on level of asthma symptoms during each phase of the menstrual cycle. Fractional exhaled nitric

oxide (eNO) levels were measured with the Aerocrine NIOX MINO system³⁰ in accordance with recommendations of the American Thoracic Society³¹ at a flow rate of 250 ml per second. Subjects were instructed to refrain from green leafy vegetables, processed meat products, and smoked meat 24 hours prior to visits to minimize interference with eNO measurements.

Serum hormone measurement

Serum levels of estradiol and progesterone were measured using the Immulite 1000 (Siemens Medical Solutions Diagnostics) following manufacturer's instructions.

Generation of iTregs ex vivo

The method for the polarization of CD4⁺ T cells towards a regulatory phenotype has been described in detail elsewhere^{32,33}. Briefly, negatively isolated (StemCell Technologies) CD4⁺ T cells were placed under polarizing conditions for 5 days: supplemented RPMI with 5ng/ml recombinant human IL-2, 2 ng/mL recombinant human TGF- β 1 (eBioscience) and plate-bound anti-CD3 antibodies (OKT3 clone, Bio X Cell). For some experiments, different concentrations of estradiol (17 β -Estradiol) or of the synthetic progesterone analog Norgestrel (Cayman Chemical) were added to the polarizing media at the beginning of the assay.

Flow cytometry

Cells were stained with antibodies against CD25, CD45RA (Miltenyi), CD4, FOXP3 and CTLA-4 (BD). All antibodies were monoclonal and directly conjugated with fluorescent markers. Data were acquired using a FACSCalibur flow cytometer (BD) and analyzed using FlowJo software (Tree Star) by an examiner blind to the group assignments of the patients and the menstrual time points. Live cells were gated based on their forward and side scatter profiles.

Statistics

Data were analyzed using a two-way ANOVA with repeated measurements, the Student's *t* test or the Mann-Whitney test using Prism 5 (GraphPad Software). *P* values for the correlation analyses were calculated from an *F* test. Values of *P* < 0.05 were considered to be statistically significant. Data are presented as mean $\pm$ SEM.

RESULTS

CD4 T cells from female patients with asthma taking OCs exhibit improved ability to generate iTregs ex vivo

Demographic data of the patients with asthma included in this study is shown in Table 1. Data were collected at three different points of the menstrual cycle: follicular, mid-cycle and luteal phases, for two consecutive cycles. Averaged clinical parameters obtained in their first visit of each menstrual cycle are also shown in Table 1. As expected in non-OC patients, serum levels of the steroid sex hormones estradiol and progesterone peaked at mid-cycle and luteal phases, respectively (Figure 1). Patients using OCs showed an overall decrease in the

serum levels of estradiol and progesterone and the absence of pronounced peaks at any of the menstrual cycle phases (Figure 1).

We first looked for alterations in the frequency of nTregs in the patients' peripheral blood by flow cytometry. We used a stringent CD4⁺CD25^{hi} gating strategy and found similar levels of Tregs between OC and non-OC patients (Figures 2A and 2B), which resembled the values observed in our cohort of normal donors (data not shown). Moreover, no significant differences were observed in the frequencies of nTregs at the different phases of the menstrual cycle (Figure 2B).

Next, we purified peripheral blood CD4⁺ T cells to assess their ability to adopt a regulatory phenotype *ex vivo* using a previously established strategy. Upon Treg polarization, we are able to gate on CD45RA⁻CD25^{hi} T cells to obtain a population enriched with FoxP3⁺/CTLA-4⁺/CD127⁻ cells that exhibit suppressor activity³²⁻³⁴ (Supp. Fig 1 and data not shown). After 5 days in culture under these Treg polarizing conditions, T cells from non-OC patients with asthma generated a lower percentage of CD45RA⁻CD25^{hi}FoxP3⁺ iTregs than T cells from patients taking OCs (Figures 2C and D). Of note, the majority of the cells using this gating strategy expressed CTLA-4 ($87.1 \pm 1.92\%$), further supporting the *bona fide* regulatory phenotype of these cells³⁵. The levels of iTregs generated *ex vivo* did not correlate with the frequency of CD4⁺CD25^{hi} nTregs present in the initial T cell pool (Supp. Fig 2), which argues against an expansion of nTregs *in vitro*. Interestingly, T cells from healthy donors cultured under the same conditions yielded a frequency of CD45RA⁻CD25^{hi}FoxP3⁺ iTregs ($11.2\% \pm 2.39$, n=22) that resembled that observed for women with asthma taking OCs (Figure 2D).

High estradiol and progesterone levels in serum during the luteal phase associate with reduced ability to generate iTregs

Since a higher proportion of iTregs was observed in the patients that exhibited lower levels of sex hormones in serum (*i.e.* patients taking OCs), we wondered whether estradiol or progesterone affected the ability of CD4⁺ T cells to polarize towards a regulatory phenotype. We compared serum hormone levels at each menstrual cycle phase with the percentages of CD45RA⁻CD25^{hi} iTregs generated *ex vivo* in both groups of patients. We found that, for non-OC patients, the percentage of iTregs generated in culture negatively correlated with the serum levels of estradiol ($r = -0.96$, $p < 0.001$) and progesterone ($r = -0.86$, $p < 0.05$) during the luteal phase (Figure 3). No correlations between hormone serum levels and the frequency of iTregs were observed in the follicular phase, in menses or for patients taking OCs (data not shown).

Estradiol and progesterone prevent the generation of iTregs *in vitro*

We further investigated the direct impact of estradiol and progesterone in the differentiation of iTregs by adding these hormones to our *in vitro* system in doses that resembled the levels observed in the patients' sera: 10, 50 and 250 pg/mL estradiol (Figure 4B) or 0.5 and 2 ng/mL progesterone (Figure 4C). Similar to our *in vivo* findings, when CD4⁺ T cells from healthy donors were cultured for 5 days under Treg polarizing conditions in the presence of

either hormone, the ability to generate CD45RA⁻CD25^{hi} iTregs was reduced in a dose-dependent manner (Figure 4D).

Oral contraceptives influence on asthma parameters

We next wondered whether the relationship between serum hormone levels and iTreg development correlated with changes in clinical parameters in the patients. The asthma severity was assessed by evaluating the levels of eNO, FEV1s and ACT score. When comparing the measurements obtained during the luteal phase, where a clear correlation between serum hormone levels and iTreg generation had been established, we found that female patients with asthma using OCs exhibited significantly higher ACT scores (Figure 5G) and significantly lower eNO levels (Figure 5H) when compared to non-OC patients. In contrast, we found similar FEV1s levels for both groups of patients (Figure 5I). Similar results were observed at the midcycle phase of the menstrual cycle (Figures 5D–F).

DISCUSSION

We evaluated the ability of peripheral blood CD4⁺ T cells from a small cohort of women with well-controlled mild to moderate persistent asthma to generate iTregs *ex vivo*. We found a negative correlation between luteal serum levels of estrogen and progesterone and the frequency of iTregs generated in culture. The ability of T cells to differentiate into iTregs also declined when we cultured CD4⁺ T cells isolated from healthy donors with physiological concentrations of estrogen or progesterone. These results suggest that the presence of female sex hormones may directly contribute to the perturbed development of iTregs and could perpetuate the asthmatic inflammation in an at-risk female population. Consistent with this possibility, the lower serum levels of circulating estrogen and progesterone in women with asthma using OCs were associated with higher percentages of *ex vivo*-generated iTregs when compared with non-OC patients. Interestingly, a reduction in the frequency of peripheral blood Tregs was reported during the second trimester of pregnant women which we replicated *in vitro* by treating peripheral blood mononuclear cells of nonpregnant women with estrogen or progesterone³⁶. In this study we only included women with well-controlled mild to moderate asthma and did not select for women with fluctuations in asthma symptoms with their menstrual cycle, which translated into a small range of ACT scores in our cohort. We found small, yet significant, differences in ACT scores and eNO levels between OC and non-OC patients. These results are consistent with previous reports³⁷ suggesting that OC use may improve asthma symptoms (higher ACT scores and lower eNO levels). Of note, ACT scores have been shown to negatively correlate with eNO levels in patients with asthma³⁸ and changes in ACT scores of as little as 1.88 have been considered a better asthma control rating³⁹. All patients in our cohort exhibited FEV1s >80% of predicted and the fluctuations were not significant in either OC or non-OC patients, similar to previous observations³⁷. Our results, though, further support the idea of sex steroid hormones as allergic disease modifiers through a mechanism that involves the blockage of iTreg differentiation. Asthma exacerbations have been observed in the ovulatory phase when the levels of estrogen are high⁴⁰. The effects of estrogen on Treg suppressor activity and proliferation have been conducted with contradictory results, probably due to the complexity of estrogen-mediated T cell effects: while estrogen treatment increased the

proliferation of TCR/CD28-stimulated human Tregs⁴¹ and induced the expression of FoxP3 in TCR/CD28-treated murine T cells⁴², estrogen-treated dendritic cells reduced the proliferation of murine Tregs but potentiated their suppressor capacity⁴³.

The *in vitro* iTreg polarization of CD4⁺ T cells from women with asthma was performed in the absence of sex hormones. Since a negative correlation was found between serum hormone levels present during CD4⁺ T cell isolation and the frequency of iTregs generated 5 days later in culture, we believe that the T cells were already primed towards a phenotype that facilitated (or prevented) their differentiation into iTregs. In agreement with others⁴¹, we did not observe significant differences in the levels of circulating nTregs at different phases of the menstrual cycle, and we also found similar levels of nTregs between patients taking and not taking OCs. However, a limitation of our study is that the effector function of iTregs was not determined, and thus changes in their suppressor activity cannot be ruled out as potential contributors to the observed clinical differences.

The manipulation of peripheral T cell plasticity in pathological systems might represent a novel strategy for future therapeutic approaches. The conversion of memory pathogenic Th2 cells into Tregs was recently achieved through TGF- β stimulation in the presence of retinoic acid and rapamycin, and these “converted” Tregs were able to migrate to the airways and suppress Th2- mediated responses²⁷. Further studies are necessary to evaluate whether treatment with sex steroid hormones can induce the conversion of pathogenic Th2 or Th17 cells into Tregs. For instance, the possible conversion of Th17 cells into Tregs could be of great importance in patients with steroid-resistant asthma. We did not look for the potential mechanisms of action of estrogen and progesterone in the development of iTregs. It is known that T cells express estrogen receptor alpha (ER α)⁴⁴, that estrogen may affect T cell function through non-genomic, ER α -independent mechanisms⁴⁵ and that progesterone promotes the development of Th2 cells and cytokines⁴⁶. Tamoxifen is an ER ligand that can act as a full agonist, partial agonist or even as an antagonist of the ER depending on the target tissue⁴⁷. The fact that Tamoxifen behaved in an agonistic manner in our *in vitro* system (Supp. Fig. 3) further supports the involvement of functional ERs in iTreg differentiation.

Others have reported alterations in the percentages of Tregs present in the bronchoalveolar lavage fluid of patients with asthma^{48,49} and in the peripheral blood of allergic individuals⁵⁰. However, to our knowledge, this is the first report evaluating the generation of iTregs in asthma patients. In spite of the small sample size tested in this study, and the fact that our donor pool of healthy subjects did not exclude males or OC users, our findings suggest a link between patients’ clinical symptoms and serum hormone levels. A better understanding of the impact that sex steroid hormones have on iTreg generation could help us clarify patient-to-patient variability in the efficacy of current treatments and benefit future therapies for a plethora of pathologies, including asthma.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors wish to thank Jennifer Strange and Greg Bauman from the University of Kentucky Flow Cytometry Service Facility for their assistance in the collection of flow cytometry samples. For the statistical analysis, support was obtained from the Biostatistics, Epidemiology and Research Design (BERD) at the Center for Clinical and Translational Science (CCTS) and the Applied Statistics Laboratory (ASL) from the University of Kentucky.

This work was performed at University of Kentucky, College of Medicine, Lexington, KY and supported by the National Center for Research Resources (NCRR), National Institutes of Health through Grant 2P20 RR020171 and by the NCRR and the National Center for Advancing Translational Sciences through Grant UL1TR000117 to F.M. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. J.T was the recipient of the Young Faculty Support Award, from The Foundation of the American College of Allergy, Asthma and Immunology.

References

1. Holgate ST, Polosa R. Treatment strategies for allergy and asthma. *Nat Rev Immunol.* 2008; 8(3): 218–30. [PubMed: 18274559]
2. McCombe PA, Greer JM, Mackay IR. Sexual dimorphism in autoimmune disease. *Curr Mol Med.* 2009; 9(9):1058–79. [PubMed: 19747114]
3. Skobeloff EM, Spivey WH, St Clair SS, et al. The influence of age and sex on asthma admissions. *JAMA.* 1992; 268(24):3437–40. [PubMed: 1460733]
4. Thomsen SF, Ulrik CS, Kyvik KO, et al. The incidence of asthma in young adults. *Chest.* 2005; 127(6):1928–34. [PubMed: 15947304]
5. Venn A, Lewis S, Cooper M, et al. Questionnaire study of effect of sex and age on the prevalence of wheeze and asthma in adolescence. *BMJ.* 1998; 316(7149):1945–6. [PubMed: 9641931]
6. Temprano J, Mannino DM. The effect of sex on asthma control from the National Asthma Survey. *J Allergy Clin Immunol.* 2009; 123(4):854–60. [PubMed: 19181370]
7. Gissler M, Järvelin MR, Louhiala P, et al. Boys have more health problems in childhood than girls: follow-up of the 1987 Finnish birth cohort. *Acta Paediatr.* 1999; 88(3):310–4. [PubMed: 10229043]
8. Skobeloff EM, Spivey WH, Silverman R, et al. The effect of the menstrual cycle on asthma presentations in the emergency department. *Arch Intern Med.* 1996; 156(16):1837–40. [PubMed: 8790078]
9. Shames RS, Heilbron DC, Janson SL, et al. Clinical differences among women with and without self-reported perimenstrual asthma. *Ann Allergy Asthma Immunol.* 1998; 81(1):65–72. [PubMed: 9690575]
10. Salam MT, Wenten M, Gilliland FD. Endogenous and exogenous sex steroid hormones and asthma and wheeze in young women. *J Allergy Clin Immunol.* 2006; 117(5):1001–7. [PubMed: 16675325]
11. Chandler MH, Schuldheisz S, Phillips BA, et al. Premenstrual asthma: the effect of estrogen on symptoms, pulmonary function, and beta 2-receptors. *Pharmacotherapy.* 1997; 17(2):224–34. [PubMed: 9085312]
12. Tan KS, McFarlane LC, Lipworth BJ. Modulation of airway reactivity and peak flow variability in asthmatics receiving the oral contraceptive pill. *Am J Respir Crit Care Med.* 1997; 155(4):1273–7. [PubMed: 9105066]
13. Tam A, Morrish D, Wadsworth S, et al. The role of female hormones on lung function in chronic lung diseases. *BMC Womens Health.* 2011; 11:24. [PubMed: 21639909]
14. Hellings PW, Kasran A, Liu Z, et al. Interleukin-17 orchestrates the granulocyte influx into airways after allergen inhalation in a mouse model of allergic asthma. *Am J Respir Cell Mol Biol.* 2003; 28(1):42–50. [PubMed: 12495931]
15. McKinley L, Alcorn JF, Peterson A, et al. TH17 cells mediate steroid-resistant airway inflammation and airway hyperresponsiveness in mice. *J Immunol.* 2008; 181(6):4089–97. [PubMed: 18768865]
16. Baecher-Allan C, Brown JA, Freeman GJ, et al. CD4+CD25high regulatory cells in human peripheral blood. *J Immunol.* 2001; 167(3):1245–53. [PubMed: 11466340]

17. Ray A, Khare A, Krishnamoorthy N, et al. Regulatory T cells in many flavors control asthma. *Mucosal Immunol.* 2010; 3(3):216–29. [PubMed: 20164832]
18. Cools N, Ponsaerts P, Van Tendeloo VF, et al. Regulatory T cells and human disease. *Clin Dev Immunol.* 2007; 2007:89195. [PubMed: 18317534]
19. Vukmanovic-Stejić M, Agius E, Booth N, et al. The kinetics of CD4⁺Foxp3⁺ T cell accumulation during a human cutaneous antigen-specific memory response in vivo. *J Clin Invest.* 2008; 118(11):3639–50. [PubMed: 18924611]
20. Curotto de Lafaille MA, Kutchukhidze N, Shen S, et al. Adaptive Foxp3⁺ regulatory T cell-dependent and -independent control of allergic inflammation. *Immunity.* 2008; 29(1):114–26. [PubMed: 18617425]
21. Murai M, Krause P, Cheroutre H, et al. Regulatory T-cell stability and plasticity in mucosal and systemic immune systems. *Mucosal Immunol.* 2010; 3(5):443–9. [PubMed: 20505662]
22. Lloyd CM, Hessel EM. Functions of T cells in asthma: more than just T(H)2 cells. *Nat Rev Immunol.* 2010; 10(12):838–848. [PubMed: 21060320]
23. Lloyd CM, Hawrylowicz CM. Regulatory T cells in asthma. *Immunity.* 2009; 31(3):438–49. [PubMed: 19766086]
24. Provoost S, Maes T, van Durme YM, et al. Decreased FOXP3 protein expression in patients with asthma. *Allergy.* 2009; 64(10):1539–46. [PubMed: 19392991]
25. Venuprasad K, Kong YC, Farrar MA. Control of Th2-mediated inflammation by regulatory T cells. *Am J Pathol.* 2010; 177(2):525–31. [PubMed: 20566752]
26. Kim BS, Kim IK, Park YJ, et al. Conversion of Th2 memory cells into Foxp3⁺ regulatory T cells suppressing Th2-mediated allergic asthma. *Proc Natl Acad Sci U S A.* 2010; 107(19):8742–7. [PubMed: 20421479]
27. National Asthma Education and Prevention Program. Expert Panel Report 3 (EPR-3): Guidelines for the Diagnosis and Management of Asthma-Summary Report 2007. *J Allergy Clin Immunol.* 2007; 120(5 Suppl):S94–138. [PubMed: 17983880]
28. American Thoracic Society. Standardization of Spirometry, 1994 Update. *Am J Respir Crit Care Med.* 1995; 152(3):1107–36. [PubMed: 7663792]
29. Nathan RA, Sorkness CA, Kosinski M, et al. Development of the asthma control test: a survey for assessing asthma control. *J Allergy Clin Immunol.* 2004; 113(1):59–65. [PubMed: 14713908]
30. Khalili B, Boggs PB, Bahna SL. Reliability of a new hand-held device for the measurement of exhaled nitric oxide. *Allergy.* 2007; 62(10):1171–4. [PubMed: 17845587]
31. Recommendations for standardized procedures for the on-line and off-line measurement of exhaled lower respiratory nitric oxide and nasal nitric oxide in adults and children-1999. This official statement of the American Thoracic Society was adopted by the ATS Board of Directors, July 1999. *Am J Respir Crit Care Med.* 1999; 160(6):2104–17. [PubMed: 10588636]
32. Reneer MC, Estes DJ, Vélez-Ortega AC, et al. Peripherally induced human regulatory T cells uncouple Kv1.3 activation from TCR-associated signaling. *Eur J Immunol.* 2011; 41(11):3170–5. [PubMed: 21834013]
33. Ellis GI, Reneer MC, Vélez-Ortega AC, et al. Generation of induced regulatory T cells from primary human naïve and memory T cells. *J Vis Exp.* 2012; (62):e3738.
34. Estes DJ, Memarsadeghi S, Lundy SK, et al. High-throughput profiling of ion channel activity in primary human lymphocytes. *Anal Chem.* 2008; 80(10):3728–35. [PubMed: 18433147]
35. Tang AL, Teijaro JR, Njau MN, et al. CTLA4 expression is an indicator and regulator of steady-state CD4⁺ FoxP3⁺ T cell homeostasis. *J Immunol.* 2008; 181(3):1806–13. [PubMed: 18641318]
36. Mjösberg J, Svensson J, Johansson E, et al. Systemic reduction of functionally suppressive CD4^{dim}CD25^{high}Foxp3⁺ Tregs in human second trimester pregnancy is induced by progesterone and 17beta-estradiol. *J Immunol.* 2009; 183(1):759–69. [PubMed: 19535629]
37. Mandhane PJ, Hanna SE, Inman MD, et al. Changes in exhaled nitric oxide related to estrogen and progesterone during the menstrual cycle. *Chest.* 2009; 136(5):1301–7. [PubMed: 19617403]
38. Senna G, Passalacqua G, Schiappoli M, et al. Correlation among FEV₁, nitric oxide and asthma control test in newly diagnosed asthma. *Allergy.* 2007; 62(2):207–8. [PubMed: 17298431]

39. Schatz M, Sorkness CA, Li JT, et al. Asthma Control Test: reliability, validity, and responsiveness in patients not previously followed by asthma specialists. *J Allergy Clin Immunol.* 2006; 117(3): 549–56. [PubMed: 16522452]
40. Eliasson O, Scherzer HH, DeGraff AC Jr. Morbidity in asthma in relation to the menstrual cycle. *J Allergy Clin Immunol.* 1986; 77(1 Pt 1):87–94. [PubMed: 3944377]
41. Prieto GA, Rosenstein Y. Oestradiol potentiates the suppressive function of human CD4 CD25 regulatory T cells by promoting their proliferation. *Immunology.* 2006; 118(1):58–65. [PubMed: 16630023]
42. Polanczyk MJ, Carson BD, Subramanian S, et al. Estrogen drives expansion of the CD4+CD25+ regulatory T cell compartment. *J Immunol.* 2004; 173(4):2227–30. [PubMed: 15294932]
43. Polanczyk MJ, Hopke C, Vandenbark AA, et al. Estrogen-mediated immunomodulation involves reduced activation of effector T cells, potentiation of Treg cells, and enhanced expression of the PD-1 costimulatory pathway. *J Neurosci Res.* 2006; 84(2):370–8. [PubMed: 16676326]
44. Törnwall J, Carey AB, Fox RI, et al. Estrogen in autoimmunity: expression of estrogen receptors in thymic and autoimmune T cells. *J Gend Specif Med.* 1999; 2(5):33–40. [PubMed: 11252833]
45. Suzuki T, Yu HP, Hsieh YC, et al. Mitogen activated protein kinase (MAPK) mediates non-genomic pathway of estrogen on T cell cytokine production following trauma-hemorrhage. *Cytokine.* 2008; 42(1):32–8. [PubMed: 18343154]
46. Piccinni MP, Giudizi MG, Biagiotti R, et al. Progesterone favors the development of human T helper cells producing Th2-type cytokines and promotes both IL-4 production and membrane CD30 expression in established Th1 cell clones. *J Immunol.* 1995; 155(1):128–33. [PubMed: 7541410]
47. Smith CL, O'Malley BW. Coregulator function: a key to understanding tissue specificity of selective receptor modulators. *Endocr Rev.* 2004; 25(1):45–71. [PubMed: 14769827]
48. Smyth LJ, Eustace A, Kolsum U, et al. Increased airway T regulatory cells in asthmatic subjects. *Chest.* 2010; 138(4):905–12. [PubMed: 20453071]
49. Hartl D, Koller B, Mehlhorn AT, et al. Quantitative and functional impairment of pulmonary CD4+CD25hi regulatory T cells in pediatric asthma. *J Allergy Clin Immunol.* 2007; 119(5):1258–66. [PubMed: 17412402]
50. Akdis M, Verhagen J, Taylor A, et al. Immune responses in healthy and allergic individuals are characterized by a fine balance between allergen-specific T regulatory 1 and T helper 2 cells. *J Exp Med.* 2004; 199(11):1567–75. [PubMed: 15173208]

Figure 1. Women with asthma taking OCs have reduced levels of sex hormones in serum
 Serum levels of the sex hormones estradiol (**A**) and progesterone (**B**) in asthmatic patients taking (+) (n = 6) and not taking (-) OCs (n = 7) measured by ELISA at three different phases of the menstrual cycle: Menses, midcycle and luteal. The asterisk indicates a statistically significant difference between the two groups of patients measured by a two-way ANOVA with repeated measurements. Statistically significant differences between cycle phases were calculated with a paired Student's *t* test and are indicated by the symbol #. NS: not significant.

Figure 2. CD4⁺ T cells from women with asthma taking OCs can generate greater levels of iTregs

(A) Representative flow cytometry plots showing the gates on CD45⁻CD25^{hi} cells that were used to measure the frequency of nTregs in peripheral blood of patients taking (+) and not taking (-) OCs. CD4⁺ gate was applied. (B) Mean frequencies of nTregs for OC (n = 6) and non- OC (n = 7) patients during the three menstrual cycle phases: menses, midcycle and luteal. (C) Representative flow cytometry plots showing the gating strategy used to measure the percentages of CD45⁻CD25^{hi}FoxP3⁺ iTregs generated *ex vivo* for OC (+) and non-OC (-) patients. Only cells that belonged to both gates (using Boolean logic) were considered as iTregs. (D) Mean frequencies for both groups of patients during the three menstrual cycle phases: menses, midcycle and luteal. Data from the two groups of patients were analyzed with a two-way ANOVA with repeated measurements, excluding patients with missing data (1 patient per group). One asterisks (*) indicates a value of P < 0.05. NS: not significant.

Figure 3. Luteal sex hormone levels negatively correlate with the levels of iTregs generated *ex vivo* in patients not taking OCs

The serum levels of estradiol (A) and progesterone (B) at the luteal phase were plotted against the frequency of CD45⁻CD25^{hi} iTregs generated *ex vivo* for each patient with asthma not taking OCs (n = 7).

Figure 4. Exogenous estradiol and progesterone reduce the frequency of iTregs generated *in vitro* Representative flow cytometry charts showing the percentage of iTregs (CD45RA⁺CD25^{hi} gate) generated *in vitro* in the absence of hormonal treatment (**A**) or in the presence of different concentrations of estrogen (**B**) or progesterone (**C**). The histograms underneath the dot plots show the expression of FoxP3 in the iTreg (black) and non-iTreg cells (gray). (**D**) Mean frequencies of iTregs generated in each condition (n = 3–4). Each culture condition was compared against the control without hormonal treatment and statistically significant differences were calculated with a paired Student's *t* test. One asterisk (*) indicates a value of *P* < 0.05 and two asterisks (**) a value of *P* < 0.01.

Figure 5. Women with asthma taking OCs report better ACT scores

Mean ACT scores (A, D, G), eNO levels (B, E, H) and FEV1s (C, F, I) during the menses (A–C), midcycle (D–F) and luteal (G–I) phases of the menstrual cycle for patients with asthma taking (+) (n = 6) and not taking (–) (n = 7) OCs. The asterisk indicates a statistically significant difference between both groups of patients using the Mann Whitney test.

Table 1Demographic data and clinical parameters of the patients^a.

Characteristic	Patients taking OCs	Non-OC Patients
N	6	7
Sex, Female (Male)	6 (0)	7 (0)
Age ^b , years	28 (21 – 34)	30 (23 – 35)
ACT score ^b	22.4 (20.5 – 25)	21.4 (17 – 24.5)
FEV1s ^b , % of predicted	99.3 (89.5 – 112.5)	96.8 (75 – 116)
eNO ^b , ppb ^c	16.5 (7 – 32)	25.9 (11.5 – 74.5)

^a Averaged clinical parameters obtained in the first visits of each menstrual cycle (*i.e.* averaged data from visits 1 and 4)^b Data presented as mean (range)^c Parts per billion