CORRESPONDENCE

Treatment Resistant Hypertension—Investigation and Conservative Management

by Prof. Dr. med. Franz Weber, Prof. Dr. med. Manfred Anlauf
in issue 25/2014

Better Risk Assessment for Individual Substances

The colleagues Weber and Anlauf list substances with blood pressure-increasing effects to be considered in differential diagnosis (Box, p. 427).

However, for psychiatric medications it is not correct to link entire drug classes, such as antipsychotics, monoamine oxidase (MAO) inhibitors and tricyclic antidepressants, with arterial hypertension in the way presented in the article.

In contrast to the statements in the article, antipsychotics (or neuroleptics) are rather associated with orthostatic dysregulation and hypotension (e.g. quetiapine, risperidone, clozapine) than with hypertension, especially when patients are newly started on the medication; here, α_1 -antagonistic effects play a major role. Likewise, MAO inhibitors have mostly blood pressure-lowering effects. Only if dietary restrictions (low-tyramine diet) are not followed, patients are at risk for hypertensive crisis, with the risk of blood pressure increases being much higher for tranylcypromine, an irreversible, unselective MAO inhibitor, than for moclobemide or rasagiline. The same applies to tricyclic antidepressants where patients on the medication rather develop arterial hypotension and only in exceptional cases hypertension (e.g. under desipramine).

Not listed are some psychiatric medications frequently associated with marked increases in blood pressure. These include antidepressants with noradrenergic effects, such as venlafaxine (1), duloxetine (2), bupropion (3), reboxetine (reimbursed by statutory health insurances until 2011), and (levo-)miltacipran, and also prescription stimulants, such as modafinil/adranafil, atomoxetine, and (lis-)dexamfetamine.

Overall, we would prefer if the risk assessment for blood pressure-elevating effects was performed for individual compounds instead of drug classes.

DOI: 10.3238/arztebl.2015.0120a

REFERENCES

1. Diaper A, Rich AS, Wilson SJ, et al.: Changes in cardiovascular function after venlafaxine but not pregabalin in healthy volunteers: a double-blind, placebo-controlled study of orthostatic challenge, blood pressure and heart rate. *Hum Psychopharmacol* 2013; 28: 562–75.
2. Pigott TA, Prakash A, Arnold LM, Aaronson ST, Mallinckrodt CH, Wohlreich MM: Duloxetine versus escitalopram and placebo: an 8-month, double-blind trial in patients with major depressive disorder. *Current medical research and opinion* 2007; 23: 1303–18.
3. Wilens TE, Hammerness PG, Biederman J, et al.: Blood pressure changes associated with medication treatment of adults with attention-deficit/hyperactivity disorder. *The Journal of clinical psychiatry* 2005; 66: 253–9.

4. Weber F, Anlauf M: Treatment resistant hypertension—investigation and conservative management. *Dtsch Arztebl Int* 2014; 111: 425–31.

Prof. Dr. med. Roland W. Freudenmann
Dr. med. Ninja Freudenmann
Dr. med. Bartosz Zurowski
Prof. Dr. med. Carlos Schönfeldt-Lecuona
Prof. Dr. med. Bernhard J. Connemann
Dr. med. Maximilian Gahr

On behalf of the authors:

Prof. Dr. med. Roland Freudenmann

Universitätsklinikum Ulm, Klinik für Psychiatrie und Psychotherapie III,
roland.freudenmann@uni-ulm.de

Conflict of interest statement

Prof. Connemann has received reimbursement of meeting participation expenses for congresses from Pfizer und Astra Zeneca. He has received reimbursement for travel and accommodation expenses from Synthelabo, Astra Zeneca and Pfizer. He has received financial support from Janssen-Cilag, Altana Pharma and Novartis for a research project he initiated.

The remaining authors declare that no conflict of interest exists.

In Reply:

We like to thank our colleagues Freudenmann et al. for their constructive and differentiated criticism. However, we like to point out that in the summaries of product characteristics (SmPCs), especially of those antipsychotics named by you, hypertension, apart from orthostatic dysregulation and hypotension, is listed as very common adverse event (quetiapine), common adverse event (clozapine) or after the use of an injectable formulation (risperidone). This also applies to amitriptyline as a tricyclic antidepressant. Our article's perspective is that of a physician investigating possible causes of treatment-resistant hypertension. Here, the patient's medication history should also take rare adverse reactions to psychiatric medications into consideration.

The new antidepressants highlighted by you represent an important addition to the substances we listed in our table in a very summarizing fashion due to lack of space.

DOI: 10.3238/arztebl.2015.0120b

REFERENCES

1. Weber F, Anlauf M: Treatment resistant hypertension—investigation and conservative management. *Dtsch Arztebl Int* 2014; 111: 425–31.

Prof. Dr. med. Franz Weber
St. Walburga-Krankenhaus, Meschede
fc.weber@t-online.de

Prof. Dr. med. Manfred Anlauf
Privatpraxis im Medizinischen Versorgungszentrum, Dialyse-Zentrum,
Cuxhaven
manfred.anlauf@t-online.de

Conflict of interest statement

The authors declare that no conflict of interest exists.