Published in final edited form as:

Antiviral Res. 2014 September ; 109: 125–131. doi:10.1016/j.antiviral.2014.06.019.

A NEW ANTIVIRAL: CHIMERIC 3TC-AZT PHOSPHONATE EFFICIENTLY INHIBITS HIV-1 IN HUMAN TISSUES *EX VIVO*

Christophe Vanpouille^{*,1}, Anastasia Khandzhinskaya^{*,2}, Inna Karpenko^{*,2}, Sonia Zicari¹, Victor Barreto-de-Souza¹, Svetlana Frolova², Leonid Margolis^{1,+}, and Sergey Kochetkov^{2,+}

¹Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, MD

²Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Moscow, Russian Federation

Abstract

Although more-recently developed antivirals target different molecules in the HIV-1 replication cycle, nucleoside reverse transcriptase inhibitors (NRTIs) remain central for HIV-1 therapy. Here, we test the anti-HIV activity of a phosphonate chimera of two well-known NRTIs, namely AZT and 3TC. We show that this newly synthesized compound suppressed HIV-1 infection in lymphoid tissue *ex vivo* more efficiently than did phosphonates of other NRTIs. Moreover, the new compound was not toxic for tissue cells, thus making the chimeric phosphonate strategy a valid approach for the development of anti HIV-1 compound heterodimers.

INTRODUCTION

Despite intensive research to find new drugs to fight HIV-1/AIDS, nucleoside reverse transcriptase inhibitors (NRTIs) remain at the core of HIV-1 treatment and are an important component of HAART. Among eight NRTIs that have been used before now or are currently being used for treatment of HIV-infected patients, the most extensively studied is 3'-azido-3'-deoxythymidine (AZT, zidovudine, Retrovir[®]) (De Clercq, 2002). Although AZT has been widely used since the beginning of the antiretroviral era, this drug has significant side effects. In particular, zidovudine induces mitochondrial disorder with massive liver steatosis, myopathy, lactic acidosis, and mitochondrial DNA depletion (Chariot et al., 1999). Also, upon AZT monotherapy, resistant virions are quickly selected. In particular, five mutations in HIV reverse transcriptase (RT) contributing to the development of high-level resistance to zidovudine have been described (Kellam et al., 1992; Ren et al., 1998).

*Correspondence to: Leonid Margolis, Program in Physical Biology, Eunice Kennedy Shriver, National Institute of Child Health and Human Development, Building 10, Room 9D58, 10 Center Drive, MSC 1855, Bethesda, MD 20892, Phone: 301-594-2476, Fax: 301-480-0857, margolis@helix.nih.gov; or Sergey Kochetkov, Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, 32 Vavilov Street 119991, Moscow, Russian Federation, Phone: +7 4991350590, Fax: +7 4991351405, kochet@eimb.ru.

⁺These authors contributed equally to this work

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Phosphonate derivatives of AZT have shown a significant decrease in cellular toxicity and improvement of AZT therapeutic properties (Khandazhinskaya et al., 2010; Wainberg and Cameron, 1998). In particular, we recently showed that AZT 5'-aminocarbonylphosphonate (**2**) shares similar pharmacokinetic parameters with AZT 5'-H-phosphonate (**1**), i.e., a slow release of the drug following oral administration, efficient penetration into cells, and decreased toxicity (Khandazhinskaya et al., 2009). Similar studies have been done on 2',3'-dideoxy-3'-thiacytidine (3TC, lamivudine, Epivir®), a cytidine analogue originally developed against HBV that was approved for HIV-1 treatment by the FDA in 1995. Like AZT phosphonate derivatives, 3TC 5'-H-phosphonate (**3**) and 3TC 5'-aminocarbonylphosphonate (**4**) were found to be much less toxic than the parent 3TC in cell cultures. Also, in laboratory animals, prodrug transformation into the active nucleoside 3TC was slower (Khandazhinskaya et al., 2011), thus making phosphonate derivatives of NRTIs promising candidates for extended-release forms of the parent NRTI.

Here, using a similar strategy, we report on the development and the antiviral activity in *ex vivo* human tissues of a phosphonate heterodimer of 3TC and AZT, (3TC-AZT heterodimer), O-(L-2',3'-dideoxy-3'-thiacytidine-5'-yl)-O'-(3'-azido-3'-deoxythymidine-5'-yl)aminocarbonyl phosphonate (**5**). At this, the initial stage of development of this new antiviral, two major questions that need to be addressed are whether this new compound significantly inhibits HIV-1, and if the answer to this question is “yes” is this inhibition due to potential cell toxicity? We answer these two questions below.

The heterodimer strategy has been successfully used before now. For example, Imbach and his associates synthesized the first bis-nucleoside phosphates in 1990 (Puech et al., 1990). However, that heterodimer was relatively resistant to enzymatic hydrolysis by phosphodiesterases and thus too stable to release the active compounds. Other described compounds (Laduree et al., 2003; Pontikis et al., 2000; Velazquez et al., 1995) were designed not as depot forms but rather as non-hydrolysable molecules that supposedly have double activity. In contrast, our strategy was intended to design a molecule that is quickly hydrolyzed into active fragments, two of which (AZT and 3TC) might act synergistically.

Since *in vivo* the critical pathogenic events of HIV infection occur in tissues which are not faithfully reflected by single-cell cultures, we utilized a system of human lymphoid tissue *ex vivo* developed in our laboratory. Human tissues *ex vivo* retain tissue cytoarchitecture, support HIV-1 replication, and have been shown to complement pre-clinical drug testing against different pathogens (Grivel and Margolis, 2009; Rohan et al., 2010; Vanpouille et al., 2012).

We found that the 3TC-AZT heterodimer (**5**) significantly suppressed HIV-1 replication at a level that surpasses some of the clinically used antivirals and exhibited low toxicity towards MT-4 cell cultures and various tissue lymphocytes.

MATERIALS AND METHODS

For details of the *synthesis of (5) and the evaluation of its stability* see the Supplement data file.

Human tissue culture *ex vivo*

Tonsillar tissues from routine surgery were obtained from the Children's National Medical Center (Washington, DC). Tissues were obtained according to an IRB approved protocol. Tissues were dissected into 2-mm³ blocks and cultured as described earlier (Grivel and Margolis, 2009). Human tissue blocks were inoculated with HIV-1 X4_{LAI.04} as described elsewhere (Grivel and Margolis, 2009). Human tonsillar tissues (27 blocks of tissue from each of *n* donors for each experimental condition) were treated with phosphonate derivatives overnight and then infected with a prototypical X4 variant of HIV-1 (HIV-1 X4_{LAI.04}) (Rush University Virology Quality Assurance Laboratory, Chicago, IL).

MT-4 cell cultures

The MT-4 T-cell line was obtained from ATCC and maintained in RPMI with 10% FCS. MT-4 cells (10⁵ cells) were inoculated with 10 µL of viral stock of X4_{LAI.04} containing 0.7-ng/mL p24gag antigen. Cells and HIV-1 were incubated in a vial for 1h30 at 37°C. Infected cell suspensions were then transferred to 12-well plates, mixed with 1 mL of medium containing the test compound at an appropriate dilution, and further incubated at 37° C and cultured for 3 days.

Antiviral assay

We evaluated the antiviral activity of each compound by measuring inhibition of human HIV-1 replication in MT-4 cell cultures and in human lymphoid tissues. For each compound, in MT-4 cell culture or in lymphoid tissue *ex vivo*, HIV-1 inhibition, at each single concentration, was defined by the following formula: inhibition = $(1 - R_{\text{compounds}}/R_{\text{Ctl}}) * 100$, where $R_{\text{compounds}}$ and R_{Ctl} are the amounts of p24 accumulated in the medium in compound-treated cultures and in untreated cultures, respectively. We calculated the EC₅₀ values (with 95% confidence interval (CI)) by fitting the data points to a sigmoidal dose-response curve, using Prism software, (version 4.0; GraphPad). The EC₅₀ (50% effective concentration) is defined as the compound concentration required for inhibition of viral replication by 50%.

Flow cytometry

To assess the cytotoxicity of (1), (2), and (5) in human tonsillar tissues after 12 days of culture, cells isolated from untreated tissue blocks and from those treated with compounds were stained with combinations of the following fluorescence- labeled monoclonal antibodies: anti-CD3-QD605, anti-CD4-QD655, anti-CD8-QD705, anti-CD25-APC, anti-CD38-PE, anti-HLA-DR-APC-Cy7, anti-CXCR4-Brilliant violet 421, anti-CCR5-PR-Cy5 anti-CD45RA-FITC, and anti-CCR7-PE-Cy7 (Caltag Laboratories; Biolegend). Detection and enumeration of HIV-1- infected cells were performed with intracellular staining by means of anti-p24-PE (Beckman Coulter). Data were acquired and analyzed as described elsewhere (Grivel and Margolis, 2009). We quantified cell depletion using Trucount beads (Becton Dickinson) for volumetric control and normalized cell numbers by tissue-block weights.

RESULTS

3TC-AZT heterodimer synthesis

First, we compared two potential synthetic schemes for the 3TC-AZT heterodimer (**5**) by assessing the yields of the final product and the separation conditions of the reaction mixtures. The alternative was the coupling of (**2**) with 3TC or (**4**) with AZT (see structures of the compounds in Fig. 1). The TLC on silica gel 60 F₂₅₄ plates permitted the conclusion that the yields of the heterodimer (**5**) under each of these couplings were close to one another and achieved 40–50%. However, in the reaction of (**2**) with 3TC the isolation and purification of (**5**) was complicated because of the remaining 3TC, which was taken in excess and whose mobility was similar to that of the target product. In contrast, after the coupling of (**4**) with AZT, AZT was easily removed because of a large difference in the chromatographic mobility. Therefore, the following synthesis of the heterodimer phosphonate was proposed (Fig 2): We prepared ethoxycarbonylphosphonic acid by treating ethyl diethoxyphosphonoformate with trimethylbromosilane at room temperature. The interaction of the resulting acid with 3TC in the presence of dicyclohexylcarbodiimide in pyridine led to 3TC 5'-ethoxycarbonyl phosphonate. The latter compound was then treated with 25% aqueous ammonia at room temperature for 18 h to result in the formation of (**4**). We obtained 3TC-AZT heterodimer (**5**) in a yield of 45% from the reaction of (**4**) with AZT in the presence of triisopropylbenzenesulfonyl chloride in pyridine for 72 h. We confirmed its structure using ¹H, ¹³C and ³¹P-NMR spectroscopy and HRMS.

The stock solution of (**5**) was prepared in DMSO, in which the compound is stable with a half-life greater than 72 h (Fig. 3A). The half-life of the heterodimer at pH 2.2 exceeded 24 h; at pH 5.15 it was about 12 h; at pH 6.6, 1 h; and at pH 8.6, less than 5 min. In all cases the hydrolysis products were the active compounds AZT, 3TC, (**2**), and (**4**) at a ratio of 2:2:3:3 (Fig. 3B). Thus, the heterodimer under physiological conditions is quickly hydrolyzed into active RT inhibitors.

Phosphonate derivatives inhibit HIV-1 replication

We compared the anti-HIV activity of AZT 5'-H-phosphonate (**1**), AZT 5'-aminocarbonylphosphonate (**2**), and 3TC-AZT heterodimer (**5**). (**1**) (Nikavir[®]), (**2**), and (**5**) efficiently inhibited HIV-1 X4_{Lai.04} in human lymphoid tissues *ex vivo* (Fig 4). The absolute HIV replication level in tissues from different donors varied from 8 to 70 ng/mL with a median value of 16 ng/mL. The suppression of HIV-1 replication by (**5**), measured as p24gag accumulation in culture medium, was dose dependent (Fig. 4) with an EC₅₀ in the low nanomol range (Table 1). In the same system the EC₅₀s of AZT and 3TC *ex vivo* were respectively 3 nM (95% CI: 1–16; *n*=2–13 for each tested concentration) and 17 nM (95% CI: 3–102; *n*=2–13 for each tested concentration) (data not shown).

The ability of the heterodimer to suppress HIV-1 replication was confirmed in single-cell culture (MT-4 cells) in which we compared the antiviral activities of the newly developed heterodimer (**5**) with those of Nikavir[®] (**1**) and of (**2**) (Table 1). All compounds efficiently and dose-dependently inhibited HIV-1 replication in MT-4 cells, although the EC₅₀s were different from those found in tissues (Table 1). Nevertheless, in this culture system made of

identical cells, (5) also proved to be the most active of all the phosphonate compounds (Table 1).

Phosphonate derivatives are not toxic to tonsillar tissues

We used human tissues *ex vivo* that contain various cells to evaluate the effects of the phosphonate compounds on the viability of various lymphocyte subsets. After 12 days in culture, we isolated cells from tissue blocks treated with 10- μ M heterodimer (5) (a concentration that completely suppresses HIV replication in this system) and from donor-matched untreated tissue and stained them for CD3, CD4, CD8, HIV-1 coreceptors (CXCR4, CCR5), activation markers (CD25, CD38, HLADR), and memory naïve markers (CD45RA, and CCR7). We evaluated cellular depletion in all three main subsets of memory CD4⁺ T cells: central (TCM; CD45RA–CCR7⁺), effector (TEM; CD45RA–CCR7[–]), and terminally differentiated effector (TEMRA; CD45RA+CCR7[–]), as well as in naïve (CD45RA+CCR7⁺) CD4⁺ T cells and activated cells. We evaluated the cytopathicity of the three compounds by comparing the numbers of cells in infected tissues with those in matched uninfected tissues. To account for size differences in tissue blocks, we normalized the data by the weight of the tissue (Fig. 6).

We compared the numbers of various subsets of lymphocytes. On day 12, the numbers of lymphocytes in corresponding subsets of CD3⁺, CD3+CD8⁺, CD3+CD4⁺, CD3+CD4+CXCR4⁺, CD3+CD4+CCR5⁺, as well as in activated (CD25⁺, CD38⁺ or HLA-DR⁺), Tnaïve, T_{CM}, T_{EM}, T_{EMRA}, and T cells were statistically not different ($p>0.05$) in untreated tissues and in tissues treated with (1), (2), and (5) (Fig. 6). Thus, none of the compounds seems to deplete tissues of T lymphocytes of any of the subsets tested. The low toxicity of the compounds for tissue cells was confirmed in MT-4 cell cultures, since the compound concentrations that kill 50% of MT-4 cells (CC₅₀) were 500 μ M, 180 μ M, and 2.7 mM respectively for compounds (5), (1), and (2) (data not shown).

DISCUSSION

Although more than 30 drugs, including entry inhibitors as well as inhibitors of crucial viral enzymes, have been approved for the treatment of HIV/AIDS, nucleoside analogs targeting HIV-1 RT remain important components in drug development (De Clercq, 2009). Inside the cells, NRTIs must be phosphorylated into NRTI triphosphates. As the efficacy of the triphosphorylation is low, drug doses must therefore be high, generally leading to significant toxicity and clearance of the compound. An alternative is to design the corresponding depot form, i.e., a prodrug-like derivative capable of delivering the active compound in the organism at a controlled rate (Stanczak and Ferrá, 2006). Many depot forms of anti-HIV drugs have been developed (Beaumont et al., 2003; Calogeropoulou et al., 2003). For example, the two depot forms of AZT that we developed, despite being inferior to AZT in their antiviral efficacy in cell cultures, have the advantage of a slow release of AZT in a variety of laboratory animals, thus exhibiting low toxicity (Khandazhinskaya et al., 2010). A similar phenomenon has recently been described for two phosphonate forms of 3TC, 3TC 5'-H-phosphonate (3) and 3TC 5'-aminocarbonylphosphonate (4) (Khandazhinskaya et al., 2011).

Here, we describe a new phosphonate derivative, namely (3TC-AZT heterodimer: (O-L-2', 3'-dydeoxy-3'-thiacytidine-5'-yl)-O'-(3'-azido-3'-deoxythymidine-5'-yl)aminocarbonylphosphonate, or 3TC-AZT heterodimer (**5**). This compound is a chimera of AZT-5'- and 3TC-5'-aminocarbonylphosphonate. In earlier studies, both of the latter compounds showed anti-HIV-1 activity *in vitro* and lower toxicity in experimental animals than did their respective parental forms (Khandazhinskaya et al., 2010; Khandazhinskaya et al., 2009). We developed the aminocarbonylphosphonate derivative and not the *H*-phosphonate derivative, as bis-substituted aminocarbonylphosphonates have been shown to be more stable than *H*-phosphonates (Khandazhinskaya et al., 2000). In this work, we compared the antiviral activity of the newly developed chimera (**5**) with that of Nikavir[®], another AZT phosphate derivative (**1**).

In contrast to the testing of the majority of the newly developed antivirals, we tested the anti-HIV activity of our heterodimer in human lymphoid tissue *ex vivo*, an experimental system that faithfully reflects important aspects of lymphoid tissue *in vivo*, where critical events of HIV pathogenesis occur in HIV-infected individuals. Human lymphoid tissue supports productive HIV infection without exogenous activation and stimulation, and it retains tissue cytoarchitecture as well as the pattern of expression of key cell surface molecules relevant to HIV infection (Grivel and Margolis, 2009). Also, such a system reflects *in vivo* donor-to-donor variability and allows testing of various drugs as a preliminary step before engaging in costly and lengthy clinical trials. Importantly, *ex vivo* tissue cultures may reveal anti-HIV-1 activities of compounds, such as acyclovir, that were not revealed through use of conventional cell line cultures (Lisco et al., 2008).

We found that 3TC-AZT heterodimer (**5**) efficiently suppressed HIV-1 replication in tonsil histocultures with an EC₅₀ in the range of clinically approved drugs. For example, we found that it has an EC₅₀ lower than that of 3TC or Nikavir[®], but higher than that of AZT. We also evaluated this activity for some of these compounds in single-cell cultures of the MT-4 cell line, where HIV-1 replication is typically much greater than in histocultures. In MT-4 cell cultures, the 3TC-AZT heterodimer (**5**) was also superior in suppressing HIV-1 compared with other tested phosphonate derivatives. However, the EC₅₀s of all the tested phosphonate compounds were closer to each other in MT-4 than in human tissues *ex vivo*, probably because all the cells in MT-4 cell cultures are equally accessible to the compounds and they do not reflect the diversity of cells present in tissues. In human tissues *ex vivo*, the EC₅₀ of (**5**) was in the range of two FDA-approved NRTIs, namely AZT and 3TC (in the nM range). However, the depot form technology allows slow and continuous release of 3TC and AZT as active compounds, thus allowing slower general clearance of the two NRTIs. In tissues *ex vivo* and in MT-4 cell cultures, (**5**) was more efficient in inhibiting HIV-1 than was Nikavir[®], a depot form of AZT.

It is very important for any compound that suppresses viral replication to test its toxicity, since what appears to be an antiviral effect can be the result of killing viral cell targets. To test whether it may be the case for the heterodimer, we enumerated cells in tissues treated with 3TC-AZT heterodimer (**5**) and in untreated control. There was no significant cell loss in total lymphocytes or in CD4+ or CD8+ lymphocyte subsets. However, we further investigated whether the compound might be toxic for some specific subpopulations of T

cells essential for HIV-1 infection, in particular for memory and/or activated CD4⁺ T cells, which preferentially support HIV replication (Douek et al., 2002). We focused on the possible toxicity of the heterodimer for naïve, TCM, TEM, and TEMRA CD4⁺ T cells and for CD4⁺ T cells expressing any one of the three following activation markers: CD25, CD38, and HLA-DR. The compound did not trigger significant loss of any of the studied subsets in human tissue *ex vivo*.

The 3TC-AZT heterodimer (**5**) tissue metabolism is relatively complex: it is eventually broken down into AZT, AZT -5' aminocarbonyl phosphonate (**2**), 3TC, and 3TC-5' aminocarbonyl phosphonate (**4**). For the first hydrolysis, which releases two active compounds, to occur quickly the heterodimer should not be too stable in the cells. It seems that this hydrolysis is indeed quick and spontaneous and does not seem to be mediated by any enzymes. After an 18-h incubation, phosphonate derivatives (**2**) and (**4**) are in significant excess in blood plasma, as revealed by the HPLC analysis. These two released phosphonate derivatives are further slowly transformed into their unmodified NRTI form, i.e., AZT or 3TC, with a kinetic constant of reaction difficult to measure. The advantage of using a heterodimer is not that it is significantly more active than 3TC and AZT, but rather that some of the heterodimer will release 3TC and AZT immediately upon hydrolysis while the other two products of the heterodimer hydrolysis, namely the phosphonates of 3TC and AZT, will continue to be further hydrolyzed into 3TC and AZT, therefore smoothing the pharmacokinetic profile by prolonging their presence, creating a depot effect.

Detailed study of the heterodimer metabolism, together with evaluation of the relative presence of 3TC and AZT in *ex vivo* and *in vivo* systems at each time-point, is to be addressed in *in vivo* experiments that are beyond the scope of the current paper, in which we have demonstrated that the development of bis phosphonate derivatives is feasible and that such a bis phosphonate of 3TC and AZT inhibits HIV-1 in human lymphoid tissue *ex vivo* more efficiently than does Nikavir[®]. Its low toxicity and its complex metabolism, which is associated with a slow release of active compounds, make it a candidate for future development and demonstrate that the phosphonate strategy may be useful for the development of heterodimers of various anti-HIV-1 compounds.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by joint NIH-RFBR grant (RFBR 12-04-91450-NIH). VBS was partially supported by fellowships from the Brazilian Ministry of Education/CAPES and the Brazilian Ministry of Science and Technology/CNPq. We thank the entire staff of the Department of Pathology of Children's National Medical Center for their generous assistance in obtaining human tonsillar tissues. MT-4 cells were obtained through the AIDS Reagent Program, Division of AIDS, NIAID, NIH.

References

- Beaumont K, Webster R, Gardner I, Dack K. Design of ester prodrugs to enhance oral absorption of poorly permeable compounds: challenges to the discovery scientist. *Current drug metabolism*. 2003; 4:461–485. [PubMed: 14683475]

- Calogeropoulou T, Detsi A, Lekkas E, Koufaki M. Strategies in the design of prodrugs of anti-HIV agents. *Current topics in medicinal chemistry*. 2003; 3:1467–1495. [PubMed: 14529522]
- Chariot P, Drogou I, de Lacroix-Szmania I, Eliezer-Vanerot MC, Chazaud B, Lombes A, Schaeffer A, Zafrani ES. Zidovudine-induced mitochondrial disorder with massive liver steatosis, myopathy, lactic acidosis, and mitochondrial DNA depletion. *Journal of hepatology*. 1999; 30:156–160. [PubMed: 9927163]
- De Clercq E. New developments in anti-HIV chemotherapy. *Biochimica et biophysica acta*. 2002; 1587:258–275. [PubMed: 12084468]
- De Clercq E. Anti-HIV drugs: 25 compounds approved within 25 years after the discovery of HIV. *International journal of antimicrobial agents*. 2009; 33:307–320. [PubMed: 19108994]
- Douek DC, Brenchley JM, Betts MR, Ambrozak DR, Hill BJ, Okamoto Y, Casazza JP, Kuruppu J, Kunstman K, Wolinsky S, Grossman Z, Dybul M, Oxenius A, Price DA, Connors M, Koup RA. HIV preferentially infects HIV-specific CD4+ T cells. *Nature*. 2002; 417:95–98. [PubMed: 11986671]
- Grivel JC, Margolis L. Use of human tissue explants to study human infectious agents. *Nature protocols*. 2009; 4:256–269.
- Kellam P, Boucher CA, Larder BA. Fifth mutation in human immunodeficiency virus type 1 reverse transcriptase contributes to the development of high-level resistance to zidovudine. *Proceedings of the National Academy of Sciences of the United States of America*. 1992; 89:1934–1938. [PubMed: 1371886]
- Khandazhinskaya A, Matyugina E, Shirokova E. Anti-HIV therapy with AZT prodrugs: AZT phosphonate derivatives, current state and prospects. *Expert opinion on drug metabolism & toxicology*. 2010; 6:701–714. [PubMed: 20380483]
- Khandazhinskaya AL, Jasko MV, Karpenko IL, Solyev PN, Golubeva NA, Kukhanova MK. 5'-phosphonate derivatives of 2',3'-dideoxy-3'-thiacytidine as new anti-HIV prodrugs. *Chemical biology & drug design*. 2011; 78:50–56. [PubMed: 21518262]
- Khandazhinskaya AL, Shirokova EA, Karpenko IL, Zakirova NF, Tarusova NB, Krayevsky AA. P-(alkyl)-nucleoside 5'-hydrogenphosphonates as depot forms of antiviral nucleotide analogues. *Nucleosides, nucleotides & nucleic acids*. 2000; 19:1795–1804.
- Khandazhinskaya AL, Yanvarev DV, Jasko MV, Shipitsin AV, Khalizev VA, Shram SI, Skoblov YS, Shirokova EA, Kukhanova MK. 5'-aminocarbonyl phosphonates as new zidovudine depot forms: antiviral properties, intracellular transformations, and pharmacokinetic parameters. *Drug metabolism and disposition: the biological fate of chemicals*. 2009; 37:494–501. [PubMed: 19106116]
- Laduree D, Sugeac E, Fossey C, Schmidt S, Laumond G, Aubertin AM. Synthesis of certain heterodimers expected as HIV-1 reverse transcriptase inhibitors. *Nucleosides, nucleotides & nucleic acids*. 2003; 22:873–875.
- Lisco A, Vanpouille C, Tchesnokov EP, Grivel JC, Biancotto A, Brichacek B, Elliott J, Fromentin E, Shattock R, Anton P, Gorelick R, Balzarini J, McGuigan C, Derudas M, Gotte M, Schinazi RF, Margolis L. Acyclovir is activated into a HIV-1 reverse transcriptase inhibitor in herpesvirus-infected human tissues. *Cell host & microbe*. 2008; 4:260–270. [PubMed: 18779052]
- Pontikis R, Dolle V, Guillaumel J, Dechaux E, Note R, Nguyen CH, Legraverend M, Bisagni E, Aubertin AM, Grierson DS, Monneret C. Synthesis and evaluation of “AZT-HEPT”, “AZT-pyridinone”, and “ddC-HEPT” conjugates as inhibitors of HIV reverse transcriptase. *Journal of medicinal chemistry*. 2000; 43:1927–1939. [PubMed: 10821705]
- Puech F, Gosselin G, Balzarini J, Good SS, Rideout JL, De Clercq E, Imbach JL. Synthesis and biological evaluation of dinucleoside methylphosphonates of 3'-azido-3'-deoxythymidine and 2', 3'-dideoxycytidine. *Antiviral research*. 1990; 14:11–23. [PubMed: 1964371]
- Ren J, Esnouf RM, Hopkins AL, Jones EY, Kirby I, Keeling J, Ross CK, Larder BA, Stuart DI, Stammers DK. 3'-Azido-3'-deoxythymidine drug resistance mutations in HIV-1 reverse transcriptase can induce long range conformational changes. *Proceedings of the National Academy of Sciences of the United States of America*. 1998; 95:9518–9523. [PubMed: 9689112]
- Rohan LC, Moncla BJ, Kunjara Na Ayudhya RP, Cost M, Huang Y, Gai F, Billitto N, Lynam JD, Pryke K, Graebing P, Hopkins N, Rooney JF, Friend D, Dezzutti CS. In vitro and ex vivo testing

of tenofovir shows it is effective as an HIV-1 microbicide. PloS one. 2010; 5:e9310. [PubMed: 20174579]

Stanczak A, Ferra A. Prodrugs and soft drugs. Pharmacological reports : PR. 2006; 58:599–613. [PubMed: 17085852]

Vanpouille C, Arakelyan A, Margolis L. Microbicides: still a long road to success. Trends in microbiology. 2012; 20:369–375. [PubMed: 22705107]

Velazquez S, Alvarez R, San-Felix A, Jimeno ML, De Clercq E, Balzarini J, Camarasa MJ. Synthesis and anti-HIV activity of [AZT]-[TSAO-T] and [AZT]-[HEPT] dimers as potential multifunctional inhibitors of HIV-1 reverse transcriptase. Journal of medicinal chemistry. 1995; 38:1641–1649. [PubMed: 7538589]

Wainberg MA, Cameron DW. HIV resistance to antiviral drugs: public health implications. Drug resistance updates : reviews and commentaries in antimicrobial and anticancer chemotherapy. 1998; 1:104–108. [PubMed: 16904396]

- Human tissues *ex vivo* complement pre-clinical drug testing against different pathogens.
- Chimeric 3TC-AZT phosphonate strongly inhibits HIV-1.
- The 3TC-AZT chimera is not toxic.
- The phosphonate heterodimer strategy is a promising strategy for the development of new anti HIV-1 compounds.

Figure 1. Structural formulae of AZT, 3TC, and their phosphonate derivatives

Shown are structural formulae of AZT and its H-phosphonate (Nikavir®) (**1**) and 5'-aminocarbonyl phosphonate derivative (**2**), and structural formulae of 3TC and its H-phosphonate (**3**) and 5'-aminocarbonyl phosphonate derivative (**4**). Compound (**5**) is an O-(L-2',3'-dydeoxy-3'-thiacytidine-5'-yl)-O'-(3'-azido-3'-deoxythymidine-5'-yl)aminocarbonyl phosphonate, or 3TC-AZT heterodimer (**5**).

Figure 2. 3TC-AZT heterodimer (5) synthesis

Presented are the synthesis of compound 5 and the yields of each chemical reaction.

Figure 3. The assessment of chemical stability of 3TC-AZT heterodimer 5 was tested at different pH levels and time points

A. Elution profiles of heterodimer (5) (solution in DMSO) under different HPLC conditions (I, reverse phase mode) and (II, pseudo-ion-pair mode).

B. Elution profile after incubation at pH 5.15 for 24 h under conditions (II, pseudo-ion-pair mode) is shown in panel B. For details, see Materials and Methods.

Figure 4. Antiviral activity of NRTI phosphonate derivatives in human lymphoid tissues
 Blocks of human tonsillar tissue were treated with increasing concentrations of AZT H-phosphonate **1** (A), AZT 5'-aminocarbonylphosphonate **2** (B), and heterodimer **5** (C) and then inoculated *ex vivo* with X4_{LAI.04}. The medium was changed and the drugs were replenished every three days. We monitored HIV-1 replication by measuring p24_{gag} accumulated in culture media over 3-day periods. For each donor, each datum represents pooled viral release from 27 tissue blocks. Drugs were added at concentrations ranging from 10 pM to 3 μ M, and their anti-HIV activity was evaluated from the suppression of viral

replication compared with donor-matched HIV-infected tissues not treated with phosphonate derivatives. We estimated the 50% effective concentration (EC_{50}) and 95% confidence interval by fitting the data to four-parameter logistic regression. EC_{50} were estimated from independent experiments performed with tonsillar tissues from seven different donors.

Figure 5. Antiviral activity of NRTI phosphonate derivatives in MT-4 cell cultures
MT-4 cells (10^5 cells) were inoculated with 10 μ L of viral stock of X4_{LAI.04} containing 0.7-ng/mL p24gag antigen. Infected cell suspensions were then transferred to 12-well plates, mixed with 1 mL of medium containing the test compound at an appropriate dilution, further incubated at 37° C, and cultured for 3 days. Cells were treated with increasing concentrations of AZT H-phosphonate (**1**) (A), AZT 5'-aminocarbonylphosphonate (**2**) (B), or heterodimer (**5**) (C). Drugs were added at concentrations ranging from 0.1 pM to 10 μ M, and their anti-HIV activities were evaluated from the suppression of viral replication

compared with untreated MT-4 cell cultures. Depending on the concentration and the compound, HIV-1 suppression was calculated from two to nine independent experiments. We estimated the EC_{50} and 95% confidence interval by fitting the data to four-parameter logistic regression.

Figure 6. Cell depletion of phosphonate derivative-inoculated *ex vivo*-cultured human lymphoid tissues

Tissue blocks (27 from each individual human tonsil donor) were treated with 10 μ M heterodimer (5). At day 12, cells were stained for anti-CD3-QD605, anti-CD4-QD655, anti-CD8-QD705, anti-CD25-APC, anti-CD38-PE, anti-HLA-DR-APC-Cy7, anti-CXCR4-Brilliant violet 421, anti-CCR5-PR-Cy5, anti-CD45RA-FITC, and anti-CCR7-PE-Cy7. We evaluated the cytopathicity of the heterodimer for different cell subsets by comparing the number of cells in infected tissues with that in matched uninfected tissues. To account for size differences in tissue blocks, we normalized the data by the weight of the

tissues. Data are means $\pm$ SEM (error bars) of 10 independent experiments performed with tissues from 10 different donors.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Antiviral activity in MT-4 cell cultures and in human lymphoid tissue *ex vivo* infected with HIV-1 X4_{LAI.04} MT-4 cells (10⁵ cells per condition) or human tonsillar tissue cultured *ex vivo* (27 blocks per condition) were treated with increasing concentrations of AZT H-phosphonate **1** (A), AZT 5'-aminocarbonylphosphonate **2** (B), heterodimer **5** (C), and then inoculated *ex vivo* with X4_{LAI.04}. In MT-4 or human tissues, the anti-HIV activity was evaluated from the suppression of viral replication compared with untreated control cultures. In MT-4, depending on the compound concentration, HIV-1 inhibition was calculated from 2–9 independents experiments. In human lymphoid tissues, HIV-1 inhibition was calculated from 7 independents experiments. We estimated the 50% effective concentration (EC₅₀) and 95% confidence interval by fitting the data to four-parameter logistic regression.

	MT-4 cell cultures		Human tissue ex vivo	
Compound	EC ₅₀ (nM)*	95% confidence interval (nM)	EC ₅₀ (nM)*	95% confidence interval (nM)
Heterodimer 5	3	1–11	11	8–17
AZT 5'-H-phosphonate 1 (Nikavir®)	9	3–26	91	39–212
AZT 5'-aminocarbonyl phosphonate 2	339	4–28,000	191	85–431