Published in final edited form as:

J Stroke Cerebrovasc Dis. 2015 March ; 24(3): 618–621. doi:10.1016/j.jstrokecerebrovasdis.2014.10.008.

Poststroke Fatigue: Hints to a Biological Mechanism

Kyra Becker, MD^{*}, Ruth Kohen, MD[†], Richard Lee, BS^{*}, Patricia Tanzi, BSN, RN^{*}, Dannielle Zierath, BS^{*}, Kevin Cain, PhD[‡], Pamela Mitchell, PhD, RN[§], and Jonathan Weinstein, MD, PhD^{*}

^{*}Department of Neurology, University of Washington School of Medicine, Seattle, Washington

[†]Department of Psychiatry, University of Washington School of Medicine, Seattle, Washington

[‡]Department of Biostatistics, University of Washington School of Medicine, Seattle, Washington

[§]Biobehavioral Nursing and Health Systems, University of Washington School of Nursing, Seattle, Washington.

Abstract

Background—Poststroke fatigue (PSF) is common, but the biological basis of this fatigue is unknown. We explored the possibility that PSF is related to systemic inflammation by investigating polymorphisms in 2 genes that affect the immune response.

Methods—In a substudy of a larger trial that evaluated the role of the immune response on stroke outcome, fatigue was assessed at 30, 90, 180, and 365 days after ischemic stroke using the Fatigue Assessment Scale. Subjects were genotyped for 3 single nucleotide polymorphisms, one in the interleukin-1 receptor antagonist gene (*IL1RN*; rs4251961, a T/C substitution) and two in the toll-like receptor-4 (*TLR4*) gene (1063 A/G [Asp299Gly] rs4986790 and 1363 C/T [Thr399Ile] rs4986791).

Results—Of the 39 participants, 22 (56%) endorsed fatigue during the study. The degree of fatigue was remarkably constant over time and independent of stroke outcome. The C allele of the rs4251961 single nucleotide polymorphism (SNP) in *IL1RN* was associated with self-reported fatigue ($P = .03$), whereas the cosegregating polymorphisms in *TLR4* were associated with lower levels of fatigue ($P = .04$).

Conclusions—SNPs in 2 genes with opposing effects on inflammatory immune responses were significantly, but differentially, associated with PSF. These findings suggest a direct link between immune signaling dysregulation and PSF.

Keywords

Poststroke fatigue; inflammation; polymorphisms; *IL1RN*; *TLR4*

Poststroke fatigue (PSF) is common after stroke and adversely affects quality of life.^{1,2} Despite the high prevalence of PSF and its negative impact on recovery, the biological underpinnings of PSF are unknown. A potential role for inflammation in the genesis of PSF has been considered.³⁻⁵ We hypothesized that if inflammation contributes to PSF, single nucleotide polymorphisms (SNPs) in genes that affect the immune response may be associated with PSF. The SNP rs4251961 is located in the promoter region of *IL1RN*; its C allele is associated with lower circulating concentrations of the gene product (interleukin, IL-1ra) and higher concentrations of proinflammatory cytokines.^{6,7} In the toll-like receptor-4 (*TLR4*) gene, 2 functional polymorphisms, 1063 A/G (Asp299Gly; rs4986790) and 1363 C/T (Thr399Ile; rs4986791), are described. In Caucasians, these 2 SNPs cosegregate such that it is more common for them to occur together rather than independently.⁸ These *TLR4* SNPs result in altered TLR4 proteins with decreased responsiveness to TLR4 ligands.^{8,9} Given their opposing effects on systemic inflammatory responses, we hypothesized that the *IL1RN* SNP and the 2 cosegregating *TLR4* SNPs would be associated with different rates of PSF.

Methods

Research Subjects

The parent-patient population is described elsewhere.¹⁰ Briefly, patients with ischemic stroke admitted to Harborview Medical Center from September 2005 through May 2009 who were at least 18 years of age were enrolled within 72 hours of symptom onset. Individuals with ongoing therapy for malignancy, known history of human immunodeficiency virus, hepatitis B or C, history of brain tumor, anemia (hematocrit <35 on admission), and those taking immunomodulatory drugs were excluded. All study procedures were approved by the University of Washington Institutional Review Board.

Clinical Data

Clinical and demographic data were collected on all subjects. Stroke severity was determined by the National Institutes of Health Stroke Scale score. Outcome was assessed by the modified Rankin Scale (mRS) score. Total infarct volume on initial diffusion-weighted magnetic resonance imaging was calculated by the ABC/2 method.¹¹ Subjects were asked about fatigue by the study nurse using the Fatigue Assessment Scale (FAS), a well characterized scale for assessing PSF.¹² Approval to administer the FAS was obtained approximately 30 months after study onset. This article includes data from the 39 subjects who provided FAS data at one or more time points. Subjects were also asked if they felt sad or blue at these same time points.

Genotyping

DNA was extracted from blood plasma samples using QIAamp DNA Blood Mini Kit (Qiagen, Valencia, CA) per manufacturer's protocols. For all 3 of the SNPs examined, genotyping was carried out using TaqMan SNP Genotyping Assay Sets and Master Mix (Applied Bio-systems, Carlsbad, CA). In brief, 2 ng of sample DNA was genotyped per manufacturer's protocols on StepOne-Plus Real-Time PCR (polymerase chain reaction) System (Applied Biosystems) under the following cycling conditions: 95° C for 10 minutes,

then 40 cycles of 95° C for 15 seconds, and 60° C for 1 minute. An allelic discrimination plot was then generated using StepOne Software, v2.0 (Applied Biosystems). Target SNP reference identification numbers were rs4986790 and rs4986791 for the 2 *TLR4* SNPs and rs4251961 for the *IL1RN* SNP. All samples were processed in triplicate. Reproducibility of the geno-typing method was confirmed as described.¹⁰ In brief, plasma-based PCR genotyping method was confirmed by carrying out identical PCR-based genotyping on DNA extracted from isolated leukocytes in a subset ($n = 42$) of patients. In these 42 patients, there was 100% concordance between the plasma-based and leukocyte-based samples. Genotype distributions for all 3 SNPs did not differ significantly from Hardy–Weinberg equilibrium (not shown).

Statistics

Descriptive data for continuous variables are presented as mean and standard deviation or median and interquartile range and compared using *t* tests for normally distributed data and the Mann–Whitney *U* test for non–normally distributed data. Data for categorical variables are presented as percentages and compared using the linear-by-linear association. Good outcome was defined as mRS less than 2. Patients were categorized based on the highest observed FAS score using previously defined cut points: 10–21 = not fatigued, 22–34 = fatigued, and 35–50 = very fatigued.¹³ Significance was set at *P* less than .05.

Results

Individual FAS scores over time are shown in Figure 1. Median FAS scores did not differ over time and were similar among those with good outcome (mRS <2) and those without. Among our 39 participants, 17 (44%) did not endorse fatigue (FAS, 10–21) at any time point after stroke, 14 (36%) had fatigue (FAS, 22–34) at one or more time points, and 8 (20%) felt extremely fatigued (FAS, 35–50) at one or more time points in the year after stroke. The clinical characteristics of these subjects are shown in Table 1. In this cohort, there was no relationship between fatigue and infarct volume, infarct location, or infarct etiology (as determined by the Trial of Org 10172 in Acute Stroke Treatment criteria¹⁴). Among subjects who endorsed feeling sad at one or more time points during the study period, 12 of 15 (80%) had at least one FAS score of 22 or more compared with 10 of 24 (40%) of patients who consistently denied feeling sad ($P = .016$).

Of our 39 patients, 14 (36%) had one copy and 8 (20%) had 2 copies of the C allele of the rs4251961 SNP in *IL1RN*. Four (10%) of subjects carried 1 variant allele at both of the *TLR4* coding SNPs. *IL1RN* rs4251961 C allele carriers were more prone to fatigue, whereas carriers of *TLR4* variant SNPs appeared to be protected against PSF (Table 1). Figure 2 shows median FAS scores over time by *IL1RN* C allele carrier status and *TLR4* coding variant carrier status. At both 30 and 90 days, patients with at least one rs4251961 C allele had significantly higher scores on the FAS.

Discussion

Fatigue is a common and disabling symptom that affects over half of stroke survivors. Few studies have addressed the biological basis of this symptom, and to our knowledge, this

study is the first to evaluate a genetic contribution to PSF. Although this exploratory study is small, it indicates a potentially substantial role for genetic factors in PSF. Large studies found that individuals with one or more *IL1RN* rs4251961 C alleles (those with more fatigue in our study) had lower circulating concentrations of IL-1ra and higher concentrations of inflammatory biomarkers like IL-1 β and C reactive protein.^{6,7} In contrast, individuals with *TLR4* coding variants, who appeared to be protected from PSF in our study, had decreased responsiveness to TLR4 ligands in previous studies.^{8,9} Our data thus support the hypothesis that PSF might result from a proinflammatory state.

This study adds to data suggesting that fatigue is independent of functional outcome given that FAS scores were similar in those with and without a good outcome. Among patients with mild stroke, fatigue is the most common complaint and is identified as the major issue restricting their life style.^{15,16} It is thus not surprising that fatigue is more predictive of a lower health-related quality of life in young adults than actual neurologic dysfunction.¹ Fatigue also prevents individuals who experience an otherwise good stroke outcome from returning to work, which means that there are economic and psychologic implications to poststroke fatigue.¹⁷ An additional observation in our cohort is that the degree of fatigue, as assessed by the FAS, remains remarkably constant over the year after stroke. Identification of treatments to lessen fatigue might therefore improve the quality of life.

In summary, PSF is a common sequela of stroke. The etiology of PSF is not known, but available data suggest that systemic inflammation may be important. Polymorphisms in genes that affect the immune response, such as *IL1RN* and *TLR4*, appear to influence the occurrence of fatigue and support the idea that inflammation is important in the genesis of fatigue. These findings need to be validated in a larger cohort and could provide valuable insight into possible interventions to alleviate PSF.

Acknowledgments

This study was funded by NINDS R01NS049197.

References

1. Naess H, Lunde L, Brogger J, et al. Fatigue among stroke patients on long-term follow-up. The Bergen Stroke Study. *J Neurol Sci.* 2011; 312:138–141. [PubMed: 21862037]
2. Feigin VL, Barker-Collo S, Parag V, et al. Prevalence and predictors of 6-month fatigue in patients with ischemic stroke: a population-based stroke incidence study in Auckland, New Zealand, 2002–2003. *Stroke.* 2012; 43:2604–2609. [PubMed: 22851549]
3. Ormstad H, Aass HC, Amthor KF, et al. Serum cytokine and glucose levels as predictors of poststroke fatigue in acute ischemic stroke patients. *J Neurol.* 2011; 258:670–676. [PubMed: 21365457]
4. Ormstad H, Aass HC, Amthor KF, et al. Serum levels of cytokines, glucose, and hemoglobin as possible predictors of poststroke depression, and association with post-stroke fatigue. *Int J Neurosci.* 2012; 122:682–690. [PubMed: 22812657]
5. McKechnie F, Lewis S, Mead G. A pilot observational study of the association between fatigue after stroke and C-reactive protein. *J R Coll Physicians Edinb.* 2010; 40:9–12. [PubMed: 21125032]
6. Reiner AP, Wurfel MM, Lange LA, et al. Polymorphisms of the IL1-receptor antagonist gene (*IL1RN*) are associated with multiple markers of systemic inflammation. *Arterioscler Thromb Vasc Biol.* 2008; 28:1407–1412. [PubMed: 18451331]

7. Rafiq S, Stevens K, Hurst AJ, et al. Common genetic variation in the gene encoding interleukin-1-receptor antagonist (IL-1RA) is associated with altered circulating IL-1RA levels. *Genes Immun.* 2007; 8:344–351. [PubMed: 17443229]
8. Ferwerda B, McCall MB, Alonso S, et al. TLR4 polymorphisms, infectious diseases, and evolutionary pressure during migration of modern humans. *Proc Natl Acad Sci U S A.* 2007; 104:16645–16650. [PubMed: 17925445]
9. Arbour NC, Lorenz E, Schutte BC, et al. TLR4 mutations are associated with endotoxin hyporesponsiveness in humans. *Nat Genet.* 2000; 25:187–191. [PubMed: 10835634]
10. Becker KJ, Dankwa D, Lee R, et al. Stroke, IL-1ra, IL1RN, infection and outcome. *Neurocrit Care.* 2014; 21:140–146. [PubMed: 24233813]
11. Sims JR, Gharai LR, Schaefer PW, et al. ABC/2 for rapid clinical estimate of infarct, perfusion, and mismatch volumes. *Neurology.* 2009; 72:2104–2110. [PubMed: 19528517]
12. Michielsen HJ, De Vries J, Van Heck GL. Psychometric qualities of a brief self-rated fatigue measure: the Fatigue Assessment Scale. *J Psychosom Res.* 2003; 54:345–352. [PubMed: 12670612]
13. Michielsen HJ, Drent M, Peros-Golubicic T, et al. Fatigue is associated with quality of life in sarcoidosis patients. *Chest.* 2006; 130:989–994. [PubMed: 17035429]
14. Adams HP Jr, Bendixen BH, Kappelle LJ, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. *Stroke.* 1993; 24:35–41. [PubMed: 7678184]
15. Carlsson GE, Moller A, Blomstrand C. Consequences of mild stroke in persons, <75 years—a 1-year follow-up. *Cerebrovasc Dis.* 2003; 16:383–388. [PubMed: 13130180]
16. Parks NE, Eskes GA, Gubitz GJ, et al. Fatigue impact scale demonstrates greater fatigue in younger stroke survivors. *Can J Neurol Sci.* 2012; 39:619–625. [PubMed: 22931703]
17. Andersen G, Christensen D, Kirkevold M, et al. Post-stroke fatigue and return to work: a 2-year follow-up. *Acta Neurol Scand.* 2012; 125:248–253. [PubMed: 21692753]

Figure 1.

Dot plot of individual patient Fatigue Assessment Scale (FAS) scores at each time point after stroke. Patients with good outcome (modified Rankin Scale; mRS <2) are displayed with open circles, those with mRS are displayed with closed circles. Not every subject contributed data at each time point. The median (interquartile range) FAS for the entire cohort at each time point is displayed along the x axis.

Figure 2.

Fatigue Assessment Scale scores at each time point based on genotype: (A) compares the scores of those with an IL1RN rs4251961 C allele to those without and (B) compares the scores of those with a functional polymorphism in TLR4 to those without. Not every subject contributed data at each time point. Statistics are by Kruskal–Wallis H test. *denotes $P < .05$. Box plots depict the median and interquartile range.

Table 1

Differences in clinical characteristics and genetics in those who were not tired, those who were tired, and those who were extremely tired (based on the highest observed FAS score)

	FAS, 10-21, N = 17	FAS, 22-34, N = 14	FAS, 35-50, N = 8	P value
Clinical characteristics at presentation				
Age (years)	47 (40-61)	52 (48-68)	61 (50-70)	.08
Female gender, n (%)	6 (35)	8 (57)	2 (25)	>.20
Initial NIHSS score	3 (2-8)	5 (4-10)	7 (3-10)	.10
Infarct volume (mm ³)	3.8 (.5-15.0)	4.4 (1.0-32.2)	.8 (.2-8.6)	>.20
Endorsed sadness,* n (%)	3 (18)	7 (50)	5 (62)	<.05
Hypertension, n (%)	8 (47)	8 (57)	4 (50)	>.20
Hyperlipidemia, n (%)	13 (76)	9 (64)	7 (88)	>.20
Coronary heart disease, n (%)	4 (24)	1 (7)	1 (12)	>.20
Atrial fibrillation, n (%)	3 (18)	4 (29)	1 (12)	>.20
Diabetes mellitus, n (%)	3 (18)	3 (21)	2 (25)	>.20
Smoker, n (%)	8 (47)	4 (29)	2 (25)	>.20
Prior stroke on imaging, n (%)	5 (29)	3 (21)	1 (12)	>.20
Genetic polymorphisms, n (%)				
<i>IL1RN</i> rs4251961 (C/T or C/C genotype)	7 (41)	8 (57)	7 (88)	.03
<i>TLR4</i> 299Gly/399Ile variant allele carriers	4 (24)	0	0	.04

Abbreviations: FAS, Fatigue Assessment Scale; *IL1RN*, interleukin-1 receptor antagonist gene; NIHSS, National Institutes of Health Stroke Scale; *TLR4*, toll-like receptor-4 gene.

* Endorsed sadness at any time point during the study likelihood ratio.