Intimate Partner Violence During Pregnancy: Maternal and Neonatal Outcomes

Jeanne L. Alhusen, PhD, CRNP, RN,¹ Ellen Ray, DNP, CNM,¹ Phyllis Sharps, PhD, RN, FAAN,¹
and Linda Bullock, PhD, RN, FAAN²

Abstract

The effects of intimate partner violence (IPV) on maternal and neonatal outcomes are multifaceted and largely preventable. During pregnancy, there are many opportunities within the current health care system for screening and early intervention during routine prenatal care or during episodic care in a hospital setting. This article describes the effects of IPV on maternal health (e.g., insufficient or inconsistent prenatal care, poor nutrition, inadequate weight gain, substance use, increased prevalence of depression), as well as adverse neonatal outcomes (e.g., low birth weight [LBW]), preterm birth [PTB], and small for gestational age [SGA]) and maternal and neonatal death. Discussion of the mechanisms of action are explored and include: maternal engagement in health behaviors that are considered “risky,” including smoking and alcohol and substance use, and new evidence regarding the alteration of the hypothalamic-pituitary-adrenal axis and resulting changes in hormones that may affect LBW and SGA infants and PTB. Clinical recommendations include a commitment for routine screening of IPV in all pregnant women who present for care using validated screening instruments. In addition, the provision of readily accessible prenatal care and the development of a trusting patient–provider relationship are first steps in addressing the problem of IPV in pregnancy. Early trials of targeted interventions such as a nurse-led home visitation program and the Domestic Violence Enhanced Home Visitation Program show promising results. Brief psychobehavioral interventions are also being explored. The approach of universal screening, patient engagement in prenatal care, and targeted individualized interventions has the ability to reduce the adverse effects of IPV and highlight the importance of this complex social disorder as a top priority in maternal and neonatal health.

Introduction

INTIMATE PARTNER VIOLENCE (IPV) during pregnancy is a serious public health issue with significant negative health consequences for women and children.^{1–6} The majority of research has found that between 3% and 9% of women experience abuse during pregnancy,^{7,8} though there are well established risk factors that are associated with higher rates of abuse, including young age, single relationship status, minority race/ethnicity, and poverty.^{9,10} Indeed, studies conducted among samples of low-income, predominantly single women have noted higher prevalence with rates up to 50%.^{11–13} The wide range in prevalence estimates has been attributed to several causes including differences among the populations sampled, measurements used to assess for IPV, inconsistencies in defining IPV, and differences in defining the time frame for abuse during the perinatal period. This review

uses the U.S. Centers for Disease Control and Prevention's uniform definition for IPV;¹⁴ that is, IPV is a pattern of coercive control of one intimate partner by the other that includes physical and sexual violence, threats of physical or sexual violence, and emotional abuse in the context of physical and sexual violence.

Research that has assessed IPV in all three trimesters demonstrates a higher prevalence than those studies that most commonly screen for IPV only once, typically during the first prenatal appointment.¹⁵ Both early clinic studies,¹⁶ and large-scale population-based studies using Pregnancy Risk Assessment Monitoring System (PRAMS) data^{5,7,8} found that the prevalence of IPV during pregnancy was lower than the year prior to pregnancy. An in-depth qualitative study examining abuse patterns during pregnancy suggested that for approximately one-third of battered women, pregnancy was a protective period, while for another group (approximately 15% of

¹Johns Hopkins University, School of Nursing, Baltimore, Maryland.

²The University of Virginia, School of Nursing, Charlottesville, Virginia.

those abused during pregnancy), abuse started or worsened during pregnancy often due to partners doubting the baby was theirs. However, abuse patterns remained consistent for the largest proportion of women abused during pregnancy.¹⁷

Intimate Partner Violence and Maternal Health

Health behaviors

Experiencing IPV during pregnancy is associated with a multitude of pregnancy-specific behaviors. Research has shown that women abused during pregnancy are twice as likely to miss prenatal care appointments or initiate prenatal care later than recommended.^{18–20} Women experiencing IPV are also twice as likely to not initiate prenatal care until the third trimester²¹ and are significantly more likely to miss three or more prenatal visits than their nonabused counterparts (45% vs. 28%).²² Extant research supports an association between insufficient prenatal care and adverse birth outcomes, including preterm delivery and low birth weight (LBW).^{23,24}

Poor nutrition and inadequate gestational weight gain have also been associated with experiencing abuse during pregnancy. Several studies have documented an association between IPV during pregnancy and poor weight gain.^{25–27} Research has demonstrated the significant impact of IPV on women's health behaviors during pregnancy, including higher rates of smoking,^{11,28} alcohol use, and substance use.^{29–31} Inadequate weight gain, smoking, alcohol use, and substance use are well-established risk factors for adverse neonatal outcomes and will be discussed in greater detail below.

A growing body of literature includes studies examining the role of IPV in affecting sexual health. A systematic review of IPV and sexual health found that the majority of studies addressing sexual risk taking, inconsistent condom use, or partner nonmonogamy found that experiencing IPV was associated with sexual risk taking of the woman or her partner. Similarly, IPV was associated with having an unplanned pregnancy or an induced abortion in the majority of studies (13 out of 16) reviewed. Finally, nearly 80% of those studies examining the association between IPV and sexually transmitted infections or urinary tract infections found an association.³²

Maternal mental health

IPV during pregnancy is associated with depression, both during pregnancy^{19,33,34} and in the postpartum period.^{35–37} Indeed, women experiencing abuse during pregnancy are 2.5 times more likely to report depressive symptomatology than their nonabused counterparts.²² Depression has been identified as the most common mental health consequence of IPV, with nearly 40% of abused women reporting depressive symptomatology.^{34,38} Post-traumatic stress disorder (PTSD) is also a common sequelae of IPV with reported rates of PTSD ranging between 19% and 84%.^{39–41} Abused women experience depression and PTSD as comorbidities at significantly higher rates than nonabused women. Research examining depression and PTSD had found that in 49%–75% of the cases, major depression co-occurred with PTSD.^{42,43}

The gravest consequences of IPV during pregnancy include homicide and suicide. Several studies have indicated that maternal injury is a leading cause of maternal mortality.^{44–46} Homicide and suicide are two potentially preventable

causes of maternal injury. A recent study utilized a multistate sample from the National Violent Death Reporting System and found pregnancy-associated suicide and homicide rates of 2.0 and 2.9 deaths per 100,000 live births, respectively. Further, 54.3% of pregnancy-associated suicides involved intimate partner conflict attributable to the suicide, and 45.3% of pregnancy-associated homicides were associated with IPV.⁴⁷ Similar results have been found in several other studies.^{48–51} These findings highlight that pregnancy-associated suicide and homicide each account for more deaths than many of the more “traditional” obstetrical causes of maternal mortality that receive greater attention.

Intimate Partner Violence and Neonatal Outcomes

Low birth weight and preterm birth

The effects of IPV extend to the consequent health of the neonate. A sizeable body of research supports the role of IPV in low birth weight and increased rates of preterm birth (PTB).^{1,3–6} LBW and PTB are well established leading causes of neonatal morbidity and mortality.¹³ An analysis of the U.S. PRAMS data revealed a significant association between IPV during pregnancy and delivering a neonate classified as LBW, after adjusting for smoking maternal age, government assistance, and education.⁵ A study examining maternal, fetal, neonatal, and infant outcomes of women hospitalized for assault during pregnancy found that women who delivered an infant during the assault hospitalization were over three times as likely to deliver a LBW neonate (odds ratio [OR]=3.10, 95% confidence interval [95% CI]=1.25–7.71) as compared with unassaulted women. Further, women discharged after an assault, delivering at a subsequent hospitalization had increased risk of abruption (OR=1.8, 95% CI=1.3–2.5), hemorrhage (OR=1.8, 95% CI=1.4–2.5), and LBW (OR=1.7, 95% CI=1.5–1.9) as compared with unassaulted women. This highlights the need to monitor undelivered assaulted women closely for the duration of their pregnancies.⁶ Similarly, women who had documented police reports of IPV occurring during pregnancy were more likely to have a LBW neonate (OR=1.67, 95% CI=1.12–2.49) or a very LBW infant (e.g., less than 1500 grams) (OR=2.78, 95% CI=1.35–5.74).⁵² When assessing the influence of frequency of abuse during pregnancy, research has demonstrated more frequent abuse (e.g., greater than 10 times during pregnancy) as predictive of delivering a LBW neonate (OR=2.9, 95% CI=1.2–6.6).⁵³ Finally, two recent meta-analyses of IPV-pregnancy studies found women experiencing IPV during pregnancy were at increased risk for PTB and delivering a LBW neonate.^{1,3} It is important to note that several studies have not found a significant relationship between IPV and delivering preterm or having a LBW neonate.^{54–57} A possible explanation for the failure of previous research to find associations between abuse and adverse neonatal outcomes is that the study sample sizes were relatively small, and the reported prevalence of abuse during pregnancy was low. Several of the studies had samples of less than 600 women, and the prevalence of violence in five studies was $\leq 10\%$.^{54,55,57–60} Similarly, it is important to note that generalizations are difficult across studies finding positive associations due to different populations sampled, assessments, methods, and data analysis. For example, studies do not uniformly control for background and other related

factors associated with either IPV or newborn outcomes (i.e., maternal age, socioeconomic status, health behaviors).

Small for gestational age

Less studied is the association between IPV during pregnancy and delivering a neonate classified as small for gestational age (SGA). SGA neonates are those who are smaller in size than normal for their gestational age, most commonly defined as a birth weight below the 10th percentile for the gestational age. SGA neonates are at increased risk of early childhood developmental and behavioral problems.^{61,62} Importantly, existing research has demonstrated an association between being born SGA and an increased rate of coronary heart disease, stroke, noninsulin-dependent diabetes mellitus, adiposity, and metabolic syndrome in adulthood.^{63–65} While limited, research examining the association between IPV during pregnancy and delivering a SGA neonate has yielded mixed results. This may be in part due to a wide variation in the prevalence (e.g., 3%–19%) of disclosed abuse in these studies.^{12,60,66} There is a need for studies to explicate how intimate partner violence was defined, how the assessment was conducted (i.e., who assessed, where the assessment was conducted), and at what point during the perinatal period violence was assessed. A recent study conducted in a low-income, urban, predominantly African American sample found that after adjusting for education, income, marital status, and substance use, experiencing IPV was associated with over five times increased odds for any adverse neonatal outcome (e.g. SGA, LBW, PTB) (adjusted odds ratio [aOR] = 5.34; 95% CI = 1.97–14.46) and specifically with a fourfold increase in having a SGA neonate (aOR = 4.00; 95% CI = 1.58–9.97).¹²

Perinatal death

Several studies have investigated the association between IPV during pregnancy and perinatal death (e.g., fetal loss after 20 weeks gestation up to neonatal death occurring ≤ 28 days after delivery). In a large sample of women attending family practice clinics in South Carolina, abuse during pregnancy was significantly associated with an increased risk of perinatal death (adjusted relative risk = 2.1, 95% CI = 1.3–3.4).⁵³ In a large study that examined maternal and neonatal outcomes for those women hospitalized during pregnancy after sustaining an assault, researchers found that women delivering in the same hospital stay as the assaultive episode had more than an 8-fold increased risk of fetal death (aOR = 8.13, 95% CI = 4.6–14.3) and nearly a 6-fold increased risk of neonatal death (aOR = 5.94; 95% CI = 3.43–10.28).⁶ Finally, in a large sample of low-income women residing in Texas, experiencing physical abuse during pregnancy was associated with an increased neonatal death rate (1.5% versus 0.2%, $p = 0.004$), physical-abuse group versus no-abuse group (i.e., verbal, physical), respectively.⁶⁷

Mechanisms of action

There is a growing recognition of the need to better understand how the experience of IPV may contribute to poor maternal and neonatal outcomes. IPV may contribute to poor outcomes, including an increased risk of spontaneous abortion, fetal loss, PTB, LBW, and neonatal death, as a direct result of blunt physical trauma to the mother. Physical trauma

is an important predictor of mortality in young women, and is the leading cause of death in pregnancy.^{68,69} Furthermore, African American trauma patients have higher mortality rates and worse outcomes than Caucasians, highlighting the need for a more comprehensive assessment of contributions to evident health disparities in neonatal outcomes.⁷⁰ A recent study examining trauma center registries in a level-1 urban trauma center over a period of 2 years found injuries occurring in the context of IPV were reported in 22% of cases. Importantly, an additional 40% of the cases could not be characterized further than being an assault. Further, women injured by stabbing were three times more likely to have a history of psychiatric illness or report IPV.⁷¹

In addition to blunt physical or sexual trauma to the mother, abuse during pregnancy may increase the risk of perinatal death, PTB, LBW, and SGA through one or more of the following mechanisms: negative maternal coping behaviors, inadequate or poor maternal nutrition, isolation and poor or limited access to prenatal care, and elevated physical or psychological stress levels. Indirect mechanisms may impact adverse pregnancy outcomes through varied pathways. For example, psychological stress may exacerbate preexisting conditions such as hypertension and gestational diabetes, or it may lead to pregnancy complications including preeclampsia or preterm labor.⁷²

Women with significant psychosocial stressors, including the experience of IPV, are more likely to engage in risky health behaviors such as smoking, alcohol use, and substance use and are less likely to see health care.^{11,27,28,73} These behaviors are associated with poor maternal and neonatal outcomes, and also place women at greater risk for unintended pregnancies and sexually transmitted infections.^{74–76} Similarly, research supports an association between high levels of depressive symptoms and PTB or LBW in a high risk sample of women.⁷⁷ Depression may act as a mediator for engaging in other negative health behaviors including the use of tobacco, illicit substances, and alcohol during pregnancy.⁷⁸

Finally, the stress of experiencing abuse during pregnancy may alter a woman's hypothalamic-pituitary-adrenal (HPA) axis. Our understanding of this causal pathway is growing, largely due to animal research. Animal studies have demonstrated that exposure to stress in the perinatal period causes heightened sensitivity to induced HPA hormone secretion in both mother and offspring.⁷⁹ As a result, higher levels of HPA hormones, including corticotrophin-releasing hormone (CRH), could initiate labor as well as restrict utero-placental perfusion.⁸⁰ Increasing evidence suggests that CRH could act as a "placental clock" that may be accelerated by exposure to physiological or psychological stress.⁸¹ Several prospective studies have found that maternal levels of CRH were significantly elevated by mid gestation in women who subsequently delivered prematurely and that maternal psychosocial stress levels at mid gestation significantly predicted the magnitude of CRH increase between mid and late gestation.^{82–84} There is an appreciable need for further research in this area, particularly to better understand how timing of abuse and severity of abuse may result in clinically meaningful biological changes in both mother and child.

Clinical Implications

Routine screening for IPV in the health care setting could identify women at risk of or experiencing IPV and lead to

interventions that reduce violence and improve maternal and child outcomes. Proponents of screening offer several key aspects for its benefit including the high prevalence of IPV and documented negative sequelae, the acceptability of screening among women, the availability of feasible screening instruments and techniques, and the opportunity to provide additional resources and referrals in those women screening positive.^{85–88} The U.S. Preventive Services Task Force recently updated its recommendations on screening women for IPV and concluded that screening instruments accurately identify women experiencing IPV and screening could reduce IPV and improve outcomes depending on the population screened and outcome measured.⁸⁹ Furthermore, several professional organizations and health care organizations support screening for violence.^{90,91}

Despite multiple organizations endorsing screening for IPV, evidence suggests actual screening rates in health care settings remain low. There are several reasons for low screening rates, including gaps in provider knowledge and lack of education regarding IPV; provider perception that patients won't be compliant (i.e., patients will not disclose abuse); lack of effective interventions to adequately address IPV; provider self-efficacy; fear of offending patients; providers' personal experience with abuse; fears of being involved with the judicial system; and lack of standardized recommendations on timing and frequency of screening.⁹² Clearly, efforts to improve universal screening for IPV in health care settings are necessary. Further research is needed on validating screening tools in a variety of geographic, cultural and practice settings. Additionally, more research is needed to determine the most accurate, efficient, and consistent way to administer screening instruments.

Prenatal care presents a unique window of opportunity in which health care providers can foster trusting relationships with pregnant women, thereby increasing the likelihood of IPV detection and mitigating its related negative consequences to both mother and child. Importantly, for many women, pregnancy is the only time they maintain regular contact with health care providers further highlighting the need to form a trusting patient–provider relationship.⁹³

There is a growing consensus that screening for IPV is a safe and effective practice and a necessary first step in addressing IPV. Less clear is how to intervene with IPV in the perinatal care setting and which interventions should be adopted. Home visitation (HV) is one such targeted intervention designed to prevent or reduce IPV victimization and perpetration. The Nurse–Family Partnership, developed by David Olds et al., is a well-known HV program that has been tested in several randomized controlled trials with young high-risk pregnant women.⁹⁴ Addressing IPV was not an original goal of the program, yet the intensity of the program presents a unique opportunity for fostering relationships between nurses and mothers, thereby offering an opportunity to intervene with women experiencing IPV. In one trial, nurse-visited women reported significantly less IPV exposure in the previous 6 months at the 4-year follow-up as compared with women in the control group.⁹⁵ Another promising IPV intervention includes the Domestic Violence Enhanced Home Visitation Program (DOVE), delivered within the context of HV. This program is being tested in urban and rural sites within the United States.⁹⁶ An important challenge for HV programs is identifying and retaining abused pregnant wo-

men in their programs, and this is an area in need of additional research.

Other research has examined the efficacy of a psychobehavioral intervention in reducing IPV during pregnancy and in the postpartum period finding women in the intervention group were less likely to have recurrent episodes of IPV victimization (OR=0.48, 95% CI=0.26–0.80), and women reporting severe IPV showed significantly reduced episodes at follow-up in the postpartum period (OR=0.39, 95% CI=0.18–0.82).⁹⁷ These results support the use of a relatively brief intervention delivered during pregnancy. Yet, other studies implementing interventions did not find significant differences in IPV between the intervention and control groups however these were pilot studies.^{98,99} Furthermore, in one of the pilot studies, the majority of the sample reported “severe” abuse, which related to greater impairment. Thus, the impact of a brief intervention may be limited in women with severe forms of abuse.⁹⁸ This highlights the need for additional large-scale, rigorous research to provide evidence about the effects of interventions in mitigating the impact of violence.

Discussion

Taken together, a burgeoning body of literature supports the role of IPV during pregnancy in contributing to adverse maternal and neonatal outcomes. Yet, maternal complications of pregnancy (e.g. preeclampsia, gestational diabetes) that are associated with a lower prevalence of maternal and fetal death than IPV receive greater interest and support for assessment, intervention, and prevention. Several mechanisms for how IPV may influence adverse maternal and neonatal outcomes have been proposed, including direct effects, mental health effects, behavioral effects, and biological effects, all of which offer health care providers opportunities to intervene. Screening for IPV during pregnancy is an important first step, and training of providers in identifying and treating IPV during the perinatal period is critical. In addition, developing and testing rigorous, evidence-based programs that can reduce IPV and improve maternal, neonatal, and early childhood outcomes is needed.

Efforts to effectively prevent the initiation of IPV should focus on healthy relationships across the lifespan, with an emphasis on children and youth. Programs and policies must be developed that are culturally based and responsive to those populations at greatest risk. The causes of IPV are complex and often the result of individual, familial, community, and societal factors. It is important to find ways to include diverse professional perspectives as well as increased community participation to address IPV in diverse communities. These perspectives may include, but are not limited to, law enforcement, nurses, physicians, social workers, advocacy groups, and academic–community collaborations. Effective responses to IPV require comprehensive, well-coordinated policies and procedures that maximize community resources. Collaborations across disciplines are essential if we are going to make progress in reducing the disparities in health outcomes for mothers and children.

Author Disclosure Statement

No competing financial interests exist.

References

- Shah PS, Shah J. Knowledge synthesis group on determinants of preterm/LBW births. Maternal exposure to domestic violence and pregnancy and birth outcomes: A systematic review and meta-analyses. *J Womens Health (Larchmt)* 2010;19:2017–2031.
- Campbell JC. Health consequences of intimate partner violence. *Lancet* 2002;359:1331–1336.
- Murphy CC, Schei B, Myhr TL, Du Mont J. Abuse: A risk factor for low birth weight? A systematic review and meta-analysis. *CMAJ* 2001;164:1567–1572.
- Kiely M, El-Mohandes AA, Gantz MG, Chowdhury D, Thornberry JS, El-Khorazaty MN. Understanding the association of biomedical, psychosocial and behavioral risks with adverse pregnancy outcomes among African Americans in Washington, DC. *Matern Child Health J*. 2011;15: S85–S95.
- Silverman JG, Decker MR, Reed E, Raj A. Intimate partner violence around the time of pregnancy: Association with breastfeeding behavior. *J Womens Health (Larchmt)* 2006;15:934–940.
- El Kady D, Gilbert WM, Xing G, Smith LH. Maternal and neonatal outcomes of assaults during pregnancy. *Obstet Gynecol* 2005;105:357–363.
- Martin SL, Mackie L, Kupper LL, Buescher PA, Moracco KE. Physical abuse of women before, during, and after pregnancy. *JAMA* 2001;285:1581–1584.
- Saltzman LE, Johnson CH, Gilbert BC, Goodwin MM. Physical abuse around the time of pregnancy: An examination of prevalence and risk factors in 16 states. *Matern Child Health J* 2003;7:31–43.
- Tjaden P, Thoennes N. Full report of the prevalence, incidence, and consequences of violence against women: Findings from the National Violence Against Women Survey. Research Report. Washington, DC: National Institute of Justice and the Centers for Disease Control and Prevention, 2000;181867.
- Vest JR, Catlin TK, Chen JJ, Brownson RC. Multistate analysis of factors associated with intimate partner violence. *Am J Prev Med* 2002;22:156–164.
- Bailey BA, Daugherty RA. Intimate partner violence during pregnancy: Incidence and associated health behaviors in a rural population. *Matern Child Health J* 2007;11:495–503.
- Alhusen JL, Lucea MB, Bullock L, Sharps P. Intimate partner violence, substance use, and adverse neonatal outcomes among urban women. *J Pediatr* 2013;163:471–476.
- Bailey BA. Partner violence during pregnancy: Prevalence, effects, screening, and management. *Int J Womens Health* 2010;2:183–197.
- Saltzman LE, Fanslow JL, McMahon PM, Shelley GA. Intimate partner violence surveillance: Uniform definitions and recommended data elements, version 1.0. Atlanta, GA: Centers for Disease Control and Prevention, National Center for Injury Prevention and Control, 2002.
- Campbell JC, Garcia-Moreno C, Sharps P. Abuse during pregnancy in industrialized and developing countries. *Violence Against Women* 2004;10:770–789.
- McFarlane J, Campbell JC, Sharps P, Watson K. Abuse during pregnancy and femicide: Urgent implications for women's health. *Obstet Gynecol* 2002;100:27–36.
- Campbell J, Oliver C, Bullock L. The dynamics of battering during pregnancy: Women's explanations of why. In: Campbell J, ed. *Empowering survivors of abuse: Health care for battered women and their children*. Thousand Oaks, CA: Sage, 1998:81–89.
- Subramanian S, Katz KS, Rodan M, et al. An integrated randomized intervention to reduce behavioral and psychosocial risks: Pregnancy and neonatal outcomes. *Matern Child Health J* 2012;16:545–554.
- Chambliss LR. Intimate partner violence and its implication for pregnancy. *Clin Obstet Gynecol* 2008;51:385–397.
- Cha S, Masho SW. Discussions about intimate partner violence during prenatal care in the United States: The role of race/ethnicity and insurance status. *Matern Child Health J* 2014;18:1413–1422.
- Goodwin MM, Gazmararian JA, Johnson CH, Gilbert BC, Saltzman LE. Pregnancy intendedness and physical abuse around the time of pregnancy: Findings from the pregnancy risk assessment monitoring system, 1996–1997. *Matern Child Health J* 2000;4:85–92.
- Dunn LL, Oths KS. Prenatal predictors of intimate partner abuse. *J Obstet Gynecol Neonatal Nurs* 2004;33:54–63.
- Vintzileos A, Ananth CV, Smulian JC, Scorza WE, Knuppel RA. The impact of prenatal care on postneonatal deaths in the presence and absence of antenatal high-risk conditions. *Am J Obstet Gynecol* 2002;187:1258–1262.
- Kogan MD, Alexander GR, Kotelchuck M, Nagey DA. Relation of the content of prenatal care to the risk of low birth weight: Maternal reports of health behavior advice and initial prenatal care procedures. *J Am Med Assoc* 1994;271:1340–1345.
- Beydoun HA, Tamim H, Lincoln AM, Dooley SD, Beydoun MA. Association of physical violence by an intimate partner around the time of pregnancy with inadequate gestational weight gain. *Soc Sci Med* 2011;72:867–873.
- Kearney MH, Haggerty LA, Munro BH, Hawkins JW. Birth outcomes and maternal morbidity in abused pregnant women with public versus private health insurance. *J Nurs Scholarsh* 2003;35:345–349.
- Kearney MH, Munro BH, Kelly U, Hawkins JW. Health behaviors as mediators for the effect of partner abuse on infant birth weight. *Nurs Res* 2004;53:36–45.
- Anderson BA, Marshak HH, Hebbeler DL. Identifying intimate partner violence at entry to prenatal care: Clustering routine clinical information. *J Midwifery Womens Health* 2002;47:353–359.
- Caetano R, McGrath C, Ramisetty-Mikler S, Field CA. Drinking, alcohol problems and the five-year recurrence and incidence of male to female and female to male partner violence. *Alcohol Clin Exp Res* 2005;29:98–106.
- El-Bassel N, Gilbert L, Wu E, Go H, Hill J. Relationship between drug abuse and intimate partner violence: A longitudinal study among women receiving methadone. *Am J Public Health* 2005;95:465–470.
- Gilbert L, El-Bassel N, Chang M, Wu E, Roy L. Substance use and partner violence among urban women seeking emergency care. *Psychol Addict Behav* 2012;26:226–235.
- Coker AL. Does physical intimate partner violence affect sexual health? A systematic review. *Trauma Violence Abuse* 2007;8:149–177.
- Kiely M, Gantz MG, El-Khorazaty MN, El-Mohandes AA. Sequential screening for psychosocial and behavioural risk during pregnancy in a population of urban African Americans. *BJOG*. 2013;120:1395–1402.
- Connelly CD, Hazen AL, Baker-Ericzen MJ, Landsverk J, Horwitz SM. Is screening for depression in the perinatal

- period enough? the co-occurrence of depression, substance abuse, and intimate partner violence in culturally diverse pregnant women. *J Womens Health (Larchmt)* 2013;22:844–852.
35. Agrawal A, Ickovics J, Lewis JB, Magriples U, Kershaw TS. Postpartum intimate partner violence and health risks among young mothers in the United States: A prospective study. *Matern Child Health J*. 2014 Feb 23. [Epub ahead of print].
 36. Trabold N, Waldrop DP, Nochajski TH, Cerulli C. An exploratory analysis of intimate partner violence and postpartum depression in an impoverished urban population. *Soc Work Health Care* 2013;52:332–350.
 37. Malta LA, McDonald SW, Hegadoren KM, Weller CA, Tough SC. Influence of interpersonal violence on maternal anxiety, depression, stress and parenting morale in the early postpartum: A community based pregnancy cohort study. *BMC Pregnancy Childbirth* 2012;12:153.
 38. Witt WP, Wisk LE, Cheng ER, et al. Poor prepregnancy and antepartum mental health predicts postpartum mental health problems among US women: A nationally representative population-based study. *Womens Health Issues* 2011;21:304–313.
 39. Hellmuth JC, Jaquier V, Swan SC, Sullivan TP. Elucidating posttraumatic stress symptom profiles and their correlates among women experiencing bidirectional intimate partner violence. *J Clin Psychol* 2014 Apr 19. doi: 10.1002/jclp.22100. [Epub ahead of print]
 40. Huth-Bocks AC, Krause K, Ahlfs-Dunn S, Gallagher E, Scott S. Relational trauma and posttraumatic stress symptoms among pregnant women. *Psychodyn Psychiatry* 2013;41:277–301.
 41. Woods SJ, Hall RJ, Campbell JC, Angott DM. Physical health and posttraumatic stress disorder symptoms in women experiencing intimate partner violence. *J Midwifery Womens Health* 2008;53:538–546.
 42. Stein MB, Kenney CJ. Major depressive and post-traumatic stress disorder comorbidity in female victims of intimate partner violence. *J Affect Disord*. 2001;66:133–138.
 43. Nixon RD, Resick PA, Nishith P. An exploration of comorbid depression among female victims of intimate partner violence with posttraumatic stress disorder. *Journal of Affective Disorders*. 2004;82:315–320.
 44. Krulewitch CJ, Pierre-Louis ML, de Leon-Gomez R, Guy R, Green R. Hidden from view: Violent deaths among pregnant women in the district of columbia, 1988–1996. *J Midwifery Womens Health*. 2001;46:4–10.
 45. Nannini A, Weiss J, Goldstein R, Fogerty S. Pregnancy-associated mortality at the end of the twentieth century: Massachusetts, 1990–1999. *J Am Med Womens Assoc* 2002;57:140–143.
 46. Campbell JC, Glass N, Sharps PW, Laughon K, Bloom T. Intimate partner homicide: Review and implications of research and policy. *Trauma Violence Abuse* 2007;8:246–269.
 47. Palladino CL, Singh V, Campbell J, Flynn H, Gold KJ. Homicide and suicide during the perinatal period: Findings from the national violent death reporting system. *Obstet Gynecol* 2011;118:1056–1063.
 48. Shadigian E, Bauer ST. Pregnancy-associated death: A qualitative systematic review of homicide and suicide. *Obstet Gynecol Surv* 2005;60:183–190.
 49. Martin SL, Macy RJ, Sullivan K, Magee ML. Pregnancy-associated violent deaths: The role of intimate partner violence. *Trauma Violence Abuse* 2007;8:135–148.
 50. Chang J, Berg CJ, Saltzman LE, Herndon J. Homicide: A leading cause of injury deaths among pregnant and postpartum women in the United States, 1991–1999. *Am J Public Health* 2005;95:471–477.
 51. Cheng D, Horon IL. Intimate-partner homicide among pregnant and postpartum women. *Obstet Gynecol* 2010;115:1181–1186.
 52. Lipsky S, Holt VL, Easterling TR, Critchlow CW. Police-reported intimate partner violence during pregnancy and the risk of antenatal hospitalization. *Matern Child Health J* 2004;8:55–63.
 53. Coker AL, Sanderson M, Dong B. Partner violence during pregnancy and risk of adverse pregnancy outcomes. *Paediatr Perinat Epidemiol* 2004;18:260–269.
 54. Berenson AB, Wiemann CM, Wilkinson GS, Jones WA, Anderson GD. Perinatal morbidity associated with violence experienced by pregnant women. *Am J Obstet Gynecol* 1994;170:1760–1769.
 55. Bullock LF, McFarlane J. The birth-weight/battering connection. *Am J Nurs* 1989;1153–1155.
 56. O'Campo P, Gielen AC, Faden RR, Kass N. Verbal abuse and physical violence among a cohort of low-income pregnant women. *Womens Health Issues* 1994;4:29–37.
 57. Grimstad H, Schei B, Backe B, Jacobsen G. Physical abuse and low birthweight: A case-control study. *Br J Obstet Gynaecol* 1997;104:1281–1287.
 58. Schei B, Samuelsen SO, Bakketeig LS. Does spousal physical abuse affect the outcome of pregnancy? *Scand J Soc Med* 1991;19:26–31.
 59. Fernandez FM, Krueger PM. Domestic violence: Effect on pregnancy outcome. *J Am Osteopath Assoc* 1999;99:254–256.
 60. Janssen PA, Holt VL, Sugg NK, Emanuel I, Critchlow CM, Henderson AD. Intimate partner violence and adverse pregnancy outcomes: A population-based study. *Am J Obstet Gynecol* 2003;188:1341–1347.
 61. Silva PA, McGee R, Williams S. A longitudinal study of the intelligence and behavior of preterm and small for gestational age children. *J Dev Behav Pediatr* 1984;5:1–5.
 62. Neuwald MF, Agranonik M, Portella AK, et al. Transgenerational effects of maternal care interact with fetal growth and influence attention skills at 18months of age. *Early Hum Dev* 2014;90:241–246.
 63. Barker DJ, Eriksson JG, Forsen T, Osmond C. Fetal origins of adult disease: Strength of effects and biological basis. *Int J Epidemiol* 2002;31:1235–1239.
 64. Barker DJ. Adult consequences of fetal growth restriction. *Clin Obstet Gynecol* 2006;49:270–283.
 65. Bonamy AK, Parikh NI, Cnattingius S, Ludvigsson JF, Ingelsson E. Birth characteristics and subsequent risks of maternal cardiovascular disease: Effects of gestational age and fetal growth. *Circulation* 2011;124:2839–2846.
 66. Urquia ML, O'Campo PJ, Heaman MI, Janssen PA, Thiessen KR. Experiences of violence before and during pregnancy and adverse pregnancy outcomes: An analysis of the canadian maternity experiences survey. *BMC Pregnancy Childbirth* 2011;11:42.
 67. Yost NP, Bloom SL, McIntire DD, Leveno KJ. A prospective observational study of domestic violence during pregnancy. *Obstet Gynecol* 2005;106:61–65.
 68. Horon IL, Cheng D. Enhanced surveillance for pregnancy-associated mortality—Maryland, 1993–1998. *JAMA* 2001;285:1455–1459.
 69. Ikossi DG, Lazar AA, Morabito D, Fildes J, Knudson MM. Profile of mothers at risk: An analysis of injury and

- pregnancy loss in 1,195 trauma patients. *J Am Coll Surg* 2005;200:49–56.
70. Harris AR, Fisher GA, Thomas SH. Homicide as a medical outcome: Racial disparity in deaths from assault in US level I and II trauma centers. *J Trauma Acute Care Surg* 2012;72:773–782.
 71. Jacovides CL, Bruns B, Holena DN, et al. Penetrating trauma in urban women: Patterns of injury and violence. *J Surg Res* 2013;184:592–598.
 72. Coker AL, Sanderson M, Dong B. Partner violence during pregnancy and risk of adverse pregnancy outcomes. *Paediatr Perinat Epidemiol* 2004;18:260–269.
 73. Martin SL, Beaumont JL, Kupper LL. Substance use before and during pregnancy: Links to intimate partner violence. *Am J Drug Alcohol Abuse* 2003;29:599–617.
 74. Bauer HM, Gibson P, Hernandez M, Kent C, Klausner J, Bolan G. Intimate partner violence and high-risk sexual behaviors among female patients with sexually transmitted diseases. *Sex Transm Dis* 2002;29:411–416.
 75. Bonomi AE, Thompson RS, Anderson M, et al. Intimate partner violence and women's physical, mental, and social functioning. *Am J Prev Med* 2006;30:458–466.
 76. Seth P, Raiford JL, Robinson LS, Wingood GM, Diclemente RJ. Intimate partner violence and other partner-related factors: Correlates of sexually transmissible infections and risky sexual behaviours among young adult African American women. *Sex Health* 2010;7:25–30.
 77. Orr ST, James SA, Blackmore Prince C. Maternal prenatal depressive symptoms and spontaneous preterm births among African-American women in Baltimore, Maryland. *Am J Epidemiol* 2002;156:797–802.
 78. Wang J, Patten SB. Prospective study of frequent heavy alcohol use and the risk of major depression in the Canadian general population. *Depress Anxiety* 2002;15:42–45.
 79. Takahashi LK, Turner JG, Kalin NH. Prolonged stress-induced elevation in plasma corticosterone during pregnancy in the rat: Implications for prenatal stress studies. *Psychoneuroendocrinology* 1998;23:571–581.
 80. Kalantaridou SN, Zoumakis E, Makrigiannakis A, Lavasidis LG, Vrekoussis T, Chrousos GP. Corticotrophin-releasing hormone, stress and human reproduction: An update. *J Reprod Immunol* 2010;85:33–39.
 81. Rich-Edwards J, Krieger N, Maljoub J, Zierler S, Lieberman E, Gillman M. Maternal experiences of racism and violence as predictors of preterm birth: Rationale and study design. *Paediatr Perinat Epidemiol* 2001;15:124–135.
 82. Hobel C, Cunkel-Schetter C, Roesch S, Castro L, Arora C. Maternal plasma corticotrophin-releasing hormone associated with stress at 20 weeks' gestation in pregnancies ending in preterm delivery. *Am J Obstet Gynecol* 1999;180:S257–S263.
 83. McLean M, Bisits A, Davies J, et al. Predicting risk of preterm delivery by second-trimester measurement of maternal plasma corticotrophin-releasing hormone and alpha-fetoprotein concentrations. *Am J Obstet Gynecol* 1999;181:207–215.
 84. Leung TN, Chung TK, Madsen G, McLean M, Chang AM, Smith R. Elevated mid-trimester maternal corticotrophin-releasing hormone levels in pregnancies that delivered before 34 weeks. *Br J Obstet Gynaecol* 1999;106:1041–1046.
 85. Rodriguez MA, Bauer HM, McLoughlin E, Grumbach K. Screening and intervention for intimate partner abuse: Practices and attitudes of primary care physicians. *JAMA* 1999;282:468–474.
 86. Phelan MB. Screening for intimate partner violence in medical settings. *Trauma Violence Abuse* 2007;8:199–213.
 87. Coker AL, Flerx VC, Smith PH, Whitaker DJ, Fadden MK, Williams M. Partner violence screening in rural health care clinics. *Am J Public Health* 2007;97:1319–1325.
 88. Rabin RF, Jennings JM, Campbell JC, Bair-Merrett MH. Intimate partner violence screening tools: A systematic review. *Am J Prev Med* 2009;36:439–445.e4.
 89. Nelson HD, Bougatsos C, Blazina I. Screening women for intimate partner violence: A systematic review to update the U.S. preventive services task force recommendation. *Ann Intern Med* 2012;156:796–808, W-279, W-280, W-281, W-282.
 90. ACOG committee opinion no. 518: Intimate partner violence. *Obstet Gynecol* 2012;119(2 Pt 1):412–417.
 91. MacMillan HL, Wathen CN, Jamieson E, et al. Approaches to screening for intimate partner violence in health care settings: A randomized trial. *JAMA* 2006;296:530–536.
 92. Taft A, O'Doherty L, Hegarty K, Ramsay J, Davidson L, Feder G. Screening women for intimate partner violence in healthcare settings. *Cochrane Database Syst Rev* 2013;4:CD007007.
 93. McFarlane JM, Groff JY, O'Brien JA, Watson K. Secondary prevention of intimate partner violence: A randomized controlled trial. *Nurs Res* 2006;55:52–61.
 94. Olds DL, Henderson CR, Jr, Phelps C, Kitzman H, Hanks C. Effect of prenatal and infancy nurse home visitation on government spending. *Med Care* 1993;31:155–174.
 95. Olds DL, Kitzman H, Cole R, et al. Effects of nurse home-visiting on maternal life course and child development: Age 6 follow-up results of a randomized trial. *Pediatrics* 2004;114:1550–1559.
 96. Sharps P, Alhusen JL, Bullock L, et al. Engaging and retaining abused women in perinatal home visitation programs. *Pediatrics* 2013;132:S134–S139.
 97. Kiely M, El-Mohandes AA, El-Khorazaty MN, Gantz MG. An integrated intervention to reduce intimate partner violence in pregnancy: A randomized controlled trial. *Obstet Gynecol* 2010;115:273–283.
 98. Zlotnick C, Capezza NM, Parker D. An interpersonally based intervention for low-income pregnant women with intimate partner violence: A pilot study. *Arch Womens Ment Health* 2011;14:55–65.
 99. Cripe SM, Sanchez SE, Sanchez E, et al. Intimate partner violence during pregnancy: A pilot intervention program in lima, peru. *J Interpers Violence* 2010;25:2054–2076.
 100. Curry MA, Durham L, Bullock L, Bloom T, Davis J. Nurse case management for pregnant women experiencing or at risk for abuse. *J Obstet Gynecol Neonatal Nurs* 2006;35:181–192.

Address correspondence to:

Jeanne L. Alhusen, PhD, CRNP, RN
525 N. Wolfe Street
Suite 528
Baltimore, MD 21205

E-mail: jalhuse1@jhu.edu