Published in final edited form as:

Bone. 2015 April ; 73: 160–166. doi:10.1016/j.bone.2014.12.013.

Parity, Lactation, Bone Strength, and 16-year Fracture Risk in Adult Women: Findings From the Study of Women's Health Across the Nation (SWAN)

Takahiro Mori, MD MSHS¹, Shinya Ishii, MD MS², Gail A. Greendale, MD³, Jane A. Cauley, DrPH⁴, Kristine Ruppert, PhD⁴, Carolyn J. Crandall, MD MS⁵, and Arun S. Karlamangla, PhD MD³

¹Department of Medicine/Division of General Internal Medicine, Kameda Medical Center, 929 Higashi-cho, Kamogawa, Chiba 296-8602, Japan

²Department of Geriatric Medicine, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan

³Department of Medicine/Division of Geriatrics, David Geffen School of Medicine at UCLA, 10945 Le Conte Ave, Ste. 2339, Los Angeles CA 90095, USA

⁴Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, 130 DeSoto Street, A510 Crabtree Hall, Pittsburgh PA 15260, USA

⁵Department of Medicine/Division of General Internal Medicine and Health Services Research, David Geffen School of Medicine at UCLA, 911 Broxton Ave, 1st floor, Los Angeles CA 90024, USA

Abstract

Our objective was to examine the associations of lifetime parity and accumulated length of lactation with bone strength in women prior to the menopause transition and fracture risk during and after the transition. Participants were 2239 pre- or early perimenopausal women from the Study of Women's Health Across the Nation (SWAN), ages 42–53 at baseline, who had no childbirths after age 42. Bone mineral density (BMD) was measured in the femoral neck and the lumbar spine at the baseline SWAN visit using dual-energy x-ray absorptiometry, and composite

© 2014 Elsevier Inc. All rights reserved.

Corresponding Author: Takahiro Mori, MD, MSHS, Department of Medicine/Division of General Internal Medicine, Kameda Medical Center, 929 Higashi-cho, Kamogawa, Chiba 296-8602, Japan, Telephone: 81-4-7099-1103, takahiromori@outlook.com.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Supplemental data

No supplemental data have been included.

Authors' roles

Study concept and design: TM, GAG, JAC, CJC, and ASK. Acquisition of data: GAG, JAC, and ASK. Analysis and interpretation of data: TM, SI, and ASK. Drafting manuscript: TM. Revising manuscript content: TM, SI, GAG, JAC, KR, CJC, and ASK. Approving final version of manuscript: TM, SI, GAG, JAC, KR, CJC, and ASK. TM takes responsibility for the integrity of the data analysis.

Disclosures

All authors state that they have no conflicts of interest.

indices of femoral neck strength relative to load (in three failure modes: compression, bending, and impact) were calculated from femoral neck BMD, femoral neck size, and body size. Data on fractures after age 42 were collected for a median follow-up of 15.7 years (interquartile range, 11.4–18.5 years). In multiple linear regression adjusted for covariates, lifetime parity was associated positively with femoral neck strength relative to load (0.024 standard deviation (SD) increment in impact strength index per childbirth, $p=0.049$), but accumulated length of lactation was associated negatively with lumbar spine BMD (0.018 SD decrement per every additional 6 months of lactation $p=0.040$). In Cox proportional hazards regression adjusted for covariates, neither parity nor lactation was associated with fracture hazard after age 42. In conclusion, parity and lactation have little impact on peak bone strength prior to menopause, and do not affect fracture risk after age 42 over 16-year follow-up.

Keywords

parity; lactation; fracture; bone strength; bone mineral density; composite strength indices

Introduction

During the last trimester of pregnancy and while breast feeding, a woman is at risk of losing bone mass to provide adequate calcium for the child's skeletal development (1). There is a good evidence that, in the short-term, both pregnancy and lactation can cause bone mineral density (BMD) loss of up to 5%, and that there may be a dose-dependent relationship between duration of lactation and amount of bone loss (2, 3). However, the long-term effects of parity and lactation on bone health are not clear. Some studies have even found that parity and lactation are associated with higher BMD later in life, while others have reported lower BMD, or no association with BMD (2).

BMD, however, is not the only bone characteristic that affects bone strength. Bone size relative to body size also plays an important role (4–6), and there are some studies suggesting associations between parity or lactation and bone size later in life (7–9). Both parity and lactation also have long-term consequences on a woman's body weight (10–12). Greater body weight independently enhances bone re-accrual (via greater skeletal loading) (13), but also leads to higher impact forces on bone in a fall (6, 14). The combined effect of these changes in BMD, bone size and body size on bone strength relative to load (i.e., relative to fall impact forces) is not known.

Composite indices of femoral neck strength, which integrate body size with femoral neck size and BMD (both measured from dual-energy x-ray absorptiometry [DXA] scans of the hip), gauge bone strength relative to load during a fall (15). These indices are inversely associated with incident fractures (15, 16), and, unlike BMD, can stratify fracture risk correctly between diabetics and non-diabetics (17), and across race/ethnicity groups (18). In addition, unlike BMD, the composite indices of femoral neck strength relative to load predict fracture risk in middle-aged women without requiring race/ethnicity information (16).

The primary objective of this study was to examine the associations of lifetime parity and cumulative duration of lactation with BMD and composite indices of femoral neck strength relative to load in pre- or early perimenopausal women between the ages of 42 and 53 years who have completed their child-bearing. A woman's peak bone strength prior to entering the menopause transition is a reliable indicator of her fracture risk later in life (19–21). The second objective of this study was to examine the associations of parity and lactation with the risk of fracture after age 42 years. We used longitudinal data from the Study of Women's Health Across the Nation (SWAN) to study these associations.

Materials and Methods

Study Participants

SWAN is a multi-site, prospective cohort study of the menopausal transition in a community-based sample of 3302 women from one of five ethnic/racial backgrounds in the United States: Caucasian, African-American, Japanese, Chinese, and Hispanic. The eligibility criteria, described in detail elsewhere (22), included age 42–52 years, intact uterus and at least one intact ovary, and not using sex-steroid hormones at the time of screening, at least one menses in the three months before screening, and self-identification as a member of one of the five eligible ethnic/racial backgrounds. Participants were enrolled in 1996–1997 at seven clinical sites in the following areas: Boston, Chicago, Detroit, Pittsburgh, Los Angeles, Newark and Oakland. The Chicago and Newark sites did not perform BMD measurement, and did not contribute to the SWAN bone cohort. Each of the other five sites enrolled Caucasians, and also enrolled women from another ethnic group: African American in Boston, Detroit, and Pittsburgh, Japanese in Los Angeles, and Chinese in Oakland. These women were followed annually for 10 years and then biennially twice (visits 11 and 12) by 2010–11.

Of 2413 participants at the five SWAN Bone Study sites, 2335 were enrolled in the bone cohort at baseline. The main reason for the exclusion was excess body weight; 46 women could not undergo DXA scans because their body weights exceeded the scanners' weight limit of 136kg. A SWAN ancillary study, the Hip Strength Across the Menopause Transition study, measured femoral neck size using archived hip DXA scans from the 1960 women in the SWAN bone cohort who had a baseline and two or more follow-up scans by follow-up visit 10. From the SWAN bone cohort, we excluded data from one woman who had initiated sex steroid hormones (a SWAN exclusion criterion) between screening and the baseline visit, 32 women who gave birth after the age of 42 years (29 before the SWAN baseline and 3 after the baseline visit), 2 women who did not report their age at the time of a fracture after age 42 but before SWAN baseline, 36 women who reported use of tamoxifen either prior to SWAN baseline or at any time during the study, 18 women for whom menopausal transition stage information was missing at the baseline visit, and 7 women for whom baseline BMI measurement was missing. The final sample sizes were 2235 for femoral neck BMD analysis, 2022 for lumbar spine BMD analysis, 1881 for analyses of composite indices of femoral neck strength, and 2239 for fracture analysis. The SWAN and sub-study protocols were approved by the Institutional Review Board at each site, and all participants gave written informed consent.

Measurements of bone strength

DXA scans were acquired with Hologic instruments (Hologic, Inc., Waltham, MA, USA). At baseline, two sites (Pittsburgh and Oakland) used QDR 2000, and three sites (Boston, Detroit, and Los Angeles) used QDR 4500. OsteoDyne's Hip Positioner System was used at every site. The DXA quality control protocols in SWAN have been previously described (23). At the baseline visit, the projected (areal) BMD in the femoral neck and the lumbar spine were recorded, and two femoral neck dimensions were measured using the region of interest (ROI) window, which was repositioned and resized by the DXA operator so that a side of the ROI window spanned the geometric measures of interest. Then the pixel locations of relevant window corners were recorded, and used to calculate the relevant distances in millimeters, using pixel dimensions provided by the manufacturer, Hologic, Inc. They were femoral neck axis length (FNAL): the distance along the femoral neck axis from the lateral margin of the base of the greater trochanter to the apex of the femoral head, and femoral neck width (FNW): the smallest thickness of the femoral neck along any line perpendicular to the femoral neck axis (Fig. 1). Composite indices of femoral neck strength relative to load during a fall were created as follows

$$\text{Compression strength index (CSI)} = \text{BMD} * \text{FNW} / \text{weight}$$

$$\text{Bending strength index (BSI)} = \text{BMD} * (\text{FNW})^2 / (\text{FNAL} * \text{weight})$$

$$\text{Impact strength index (ISI)} = \text{BMD} * \text{FNW} * \text{FNAL} / (\text{height} * \text{weight})$$

CSI reflects the ability of the femoral neck to withstand an axial compressive load proportional to body weight, BSI reflects the ability to withstand bending forces proportional to body weight, and ISI reflects the ability of the femoral neck to absorb the potential energy of impact in a fall from standing height, regardless of the failure mode: bending or compression (15). While CSI and BSI assume only that forces on the bone are proportional to body weight, ISI accounts for differences in the forces in a fall that result from differences in a woman's height.

Measurements of total duration of lactation, and parity

Standardized interview and self-reported questionnaires were used to obtain information about parity and lactation at the baseline visit. For each pregnancy, participants were asked to choose one of the outcomes; livebirth(s), stillbirth, miscarriage, abortion or tubal/ectopic, and the total numbers of pregnancies leading to livebirth(s) or stillbirth(s) were counted to obtain lifetime parity. For each pregnancy that led to livebirth(s), participants were asked the duration of breastfeeding, and cumulative length of lactation was calculated. Missing values of parity (n=4) and lactation (n=4) were counted as zero. For analysis as continuous predictors, we top-censored both parity and lactation at their 99th percentiles (6 and 72 months, respectively).

Fracture ascertainment and time to first fracture

At the baseline visit, participants reported prior fractures in adult life, along with their age at the time of the fractures. Because years but not dates of the prior fractures were reported at baseline, we imputed the dates using the midpoints of the year in which the fracture was reported to have occurred. Only fractures after reaching age 42 were included in this

analysis. During each of the follow-up visits, fractures since the previous visit were self-reported using a standardized interviewer-administered questionnaire. In all visits, the number of fractures, body site(s) affected, and how fractures occurred were recorded. SWAN initiated collection of the date of fracture at visit 7. Because dates of fractures were not collected in the first 6 follow-up visits, we imputed the dates using the midpoints between the participants' previous and index visits. Medical records were obtained for self-reported non-digital non-cranio-facial fractures reported at visit 7 and later, and 95% were confirmed. Using the 42nd birthday as the start time, we computed time to first fracture after age 42 years, and censored women who did not report any fracture at their last SWAN visit.

Measurements of covariates

Standardized interview and self-reported questionnaires at baseline were used to obtain the following covariate information: age (continuous; years), race/ethnicity (Caucasian, African-American, Japanese, Chinese), menopause transition stage (premenopausal [regular menses], early peri-menopausal [menses within 3 months but menses less predictable]), smoking status (never smoking, ex-smoker, or current), smoking pack-years (zero, less than or equal to 10 years, greater than 10 years but less than or equal to 30 years, or greater than 30 years), alcohol categories (abstainer, infrequent: greater than zero but less than or equal to 1 drink per week, light-to-moderate: greater than 1 but less than or equal to 7 drinks per week, heavy: greater than 7 drinks per week), employment status (no vs. yes), history of diabetes (no vs. yes), history of hyperthyroidism (no vs. yes), current (i.e., at baseline visit) use of supplementary calcium, current use of supplementary vitamin D, and 6 binary indicator variables (none vs. any) for use of medications: 1) prior (i.e., before SWAN baseline) use of any sex steroid hormone pills, patch, or injection other than birth control pills, 2) prior use of birth control pills, 3) prior use of depo-provera injections, 4) current or prior use of oral corticosteroids, 5) current use of proton pump inhibitors, and 6) use of other bone-adverse medications (including current or prior use of antiepileptic medications, or current uses of chemotherapy, Gonadotropin-releasing hormone agonist, aromatase inhibitors, or thiazolidinediones). At the baseline visit, no one in the bone cohort reported use of osteoporosis medications (bisphosphonates, selective estrogen receptor modulators, calcitonin, parathyroid hormone, prescription vitamin D, or denosumab).

Medication use information was also collected at every follow-up visit. For fracture analysis, self-reported medication uses from visits 1 to 12 were combined with medication variables collected at the baseline visit to create 6 indicator variables for ever (prior to baseline, at baseline, or at follow-up) use of medications (none vs. any) from the following classes: 1) sex steroid hormone pills or patch other than birth control pills, 2) birth control pills, 3) depo-provera injections, 4) oral corticosteroids, 5) proton pump inhibitors, and 6) other bone-adverse medications (defined as described above).

Physical activity was assessed at the baseline visit with an adapted version of the Kaiser Physical Activity Survey, which is based on the Baecke questionnaire (24). This self-report instrument grades physical activity in four domains: sport, home, active daily living (walking or biking to work, not watching television), and work. Home activity consists of five components: child or dependent adult care, meal preparation and cleanup, light chores

such as dusting, moderate chores such as vacuuming, and heavy chores such as home repair. Scores representing the average responses to domain-specific questions range from 1 to 5 for each domain. We calculated a total physical activity score, ranging from 4 to 20, by adding scores across the four domains, with work activity score set at 1 for those who did not work outside the home (25). Height and weight were measured at the baseline visit with a fixed stadiometer and a digital scale with the participants wearing light clothing and no shoes. BMI was calculated as weight in kilo grams divided by the square of height in meters.

A total of 369 women (15.8%) had one or more missing covariates at the baseline. We imputed the missing values from values reported in follow-up visits (for menopausal transition stage, height, and alcohol consumption level), and by using default values of zero/no/never for smoking status, smoking pack-years, history of hyperthyroidism, and medications. Those who still had missing values of menopausal transition stage or BMI were excluded from the analyses. Finally the missing values of physical activity score were imputed using predictive mean matching (n=74), as the missingness pattern was monotone (26, 27).

Statistical Analysis

We performed multiple linear regressions separately to examine the associations of parity and cumulative length of lactation with each of the bone strength measures (femoral neck BMD, lumbar spine BMD, and three composite indices of femoral neck strength relative to load) at baseline, adjusted for the following covariates also measured at baseline: age, race/ethnicity, menopausal transition stage, BMI, smoking status, smoking pack-years, alcohol consumption level, physical activity level, employment status, history of diabetes, history of hyperthyroidism, current use of supplementary calcium, current use of supplementary vitamin D, 6 medication indicator variables: prior use of any sex steroid hormone pills or patch other than birth control pills, prior use of birth control pills, prior use of depo-provera injections, current or prior use of oral corticosteroids, current use of proton pump inhibitors, other bone-adverse medications, and study site. We included BMI as a continuous (linear) term, plus a squared (quadratic) term to allow for possible higher-order associations, plus multiplicative interaction terms between BMI and race/ethnicity because of the large race/ethnicity differences in BMI.

In exploratory analyses, we re-ran the models after excluding the BMI terms, or physical activity level and employment status from the regression models. To test for possible effects of parity and lactation duration on bone size, we also ran parallel models with FNW and femoral neck cross-sectional bone mineral content (given by $FNW \times$ femoral neck BMD) as the dependent variables (outcomes) (15).

Next, we performed Cox proportional hazard regressions to model time to first fracture (after age 42 years) as a function of parity or cumulative length of lactation prior to age 42, after we had verified the proportional hazards assumption. We did not distinguish between traumatic and non-traumatic fractures in the analysis, as information regarding the mechanism of fracture (i.e. trauma vs. minimal trauma) was not available for fractures before the SWAN baseline. We did exclude fractures not typically associated with osteoporosis, in particular fractures of the face, skull, fingers, and toes (28, 29). Women who

initiated osteoporosis medications were censored at the time of the first visit in which the participants reported the use. We adjusted for race/ethnicity, select covariates measured at SWAN baseline (BMI, smoking status, smoking pack-years, alcohol consumption level, physical activity level, employment status, history of diabetes, history of hyperthyroidism, supplementary calcium, and supplementary vitamin D), and the following 6 medication variables as time-invariant covariates: ever use (before baseline or any time during the study till visit 12) of sex steroid hormone pills or patch other than birth control pills, birth control pills, depo-provera injections, oral corticosteroids, proton pump inhibitors, and other bone-adverse medications, and study site.

In sensitivity analyses, we a) included both parity and lactation in the same models to mutually adjust parity for lactation, and lactation for parity, and b) excluded stillbirths from the parity count. All analyses were performed using the STATA Version 13.1 (StataCorp LP, College Station, Texas, U.S.A.). Two-sided $p < 0.05$ was considered significant.

Results

The median age of study participants was 46 years, 49.8% were Caucasian, 28.5% were African American, 11.3% were Japanese, and 10.5% were Chinese. The median and the interquartile range (IQR) of parity were 2, [1, 3] and the mean, the median, and the IQR of cumulative length of lactation were 8.6, 1, and [0, 12] months (table 1). Pearson's correlation between parity and lactation was 0.38.

Associations with Bone Strength Measures

In multiple linear regression, parity was positively associated only with ISI and not with any of the other four measures of bone strength (Table 2): Each additional childbirth before age 42 years was associated with 0.024 standard deviation (SD) (95% confidence interval (CI): 0.0001, 0.048) increment in ISI ($p=0.049$). Additional adjustment for duration of lactation did not change the point estimate of the parity-ISI association, but the CI widened (95% CI: -0.003, 0.051) and made it statistically marginally significant ($p=0.080$). Excluding stillbirths from the parity count also made the association with ISI become marginally significant ($p=0.059$). Otherwise it did not substantially change the associations between parity and bone strength. Parity also had no association with FNW ($p=0.66$) or cross-sectional bone mineral content ($p=0.42$).

Duration of lactation was itself inversely associated only with lumbar spine BMD, and not with any of the other four bone strength measures (Table 2): Lumbar spine BMD was 0.018 SD (95% CI: -0.036, -0.001) lower for every additional 6 months of lactation before age 42 years, ($p=0.040$). After further adjusting for parity, the inverse association with lumbar spine BMD became weaker (standardized effect size = - 0.015 SD) and statistically non-significant ($p=0.13$). Lactation duration also had no association with FNW ($p=0.35$) or cross-sectional bone mineral content ($p=0.52$).

To explore the reasons for the positive association between parity and ISI, and the lack of strong negative associations between parity/lactation and bone strength measures, we examined the associations of parity and lactation with total physical activity level, home

physical activity level, and BMI. We speculated that any negative effects of child bearing and lactation on bone health were at least partly negated by the potentially higher home physical activity (child and home care) of child rearing, and its effects on body weight. In multiple linear regressions, both parity and lactation were associated with more total physical activity level and more home physical activity level: Each additional childbirth was associated with 0.09 SD (95% CI; 0.06, 0.13, $p<0.001$) increment in total physical activity score, and every additional 6 months of lactation was associated with 0.06 SD (95% CI; 0.04, 0.08, $p<0.001$) increment in total physical activity score. In addition, each additional childbirth was associated with 0.20 SD (95% CI; 0.17, 0.23, $p<0.001$) increment in home physical activity score, and every additional 6 months of lactation was associated with 0.08 SD (95% CI; 0.06, 0.10, $p<0.001$) increment in home physical activity score. In multiple linear regressions, parity was associated with higher BMI: Each additional childbirth was associated with 0.32 kg/m² (95% CI; 0.13, 0.52, $p<0.01$) increment in BMI. Lactation was not significantly associated with BMI.

After excluding physical activity level and employment status from the regression models, parity remained positively associated with ISI (effect size before adjusting for lactation: 0.034 SD, 95% CI; 0.010, 0.058, $p<0.01$), but lactation was no longer negatively associated with lumbar spine BMD. After excluding the BMI terms (but retaining physical activity and employment status), parity was no longer positively associated with ISI, but lactation remained negatively associated with lumbar spine BMD: (effect size before adjusting for parity: -0.019 SD, 95% CI: -0.038, -0.0004, $p=0.045$). After excluding physical activity level, employment status, and the BMI terms, parity was no longer associated with ISI, and lactation became marginally significantly associated with lumbar spine BMD (effect size before adjusting for parity; -0.017 SD, 95% CI: -0.036, 0.001, $p=0.064$).

Associations with Fracture

After a median follow-up of 15.7 years (interquartile range [IQR] 11.4, 18.5), which included median 4.1 years between age 42 and the SWAN baseline visit and median 13.2 years of prospective follow-up after the baseline visit, 357 women (15.9%) had at least one fracture, at a rate of 11.0 fractures per 1000 person-years. At visit 12, 1678 (96.8 %) out of 1733 participants had reached post-menopausal status, including those who had a hysterectomy and /or both ovaries removed (175 participants, 10.1%). Median age of natural (non-surgical) final menstrual period was 52 years (IQR 50, 53). In multivariable Cox proportional hazard regression, neither lifetime parity before age 42 years nor accumulated length of lactation before age 42 years was associated with the hazard of fracture after age 42. The adjusted relative hazards (with 95% CI) were 0.97 (0.89, 1.05) per additional childbirth and 0.97 (0.92, 1.02) per every additional 6 months of lactation, respectively. The sensitivity analyses (addition of mutual adjustment for parity and lactation, and exclusion of stillbirths from the parity count) did not substantially alter the conclusions of the fracture analyses.

Discussion

Similar to some previous studies (30–33), this study also found that cumulative length of lactation before age 42 years was associated inversely with BMD in pre- or early perimenopausal women ages 42–53 years, but only with BMD in the lumbar spine, not in the femoral neck. Duration of lactation was also not associated with any of the composite indices of femoral neck strength relative to load. Lifetime parity before age 42 years was associated with only one of three composite indices of femoral neck strength relative to load, and not associated with BMD in either femoral neck or lumbar spine. The two associations that were found (of the ten that were tested) were small: 0.024 SD increment in ISI per childbirth and 0.018 SD decrement in lumbar spine BMD for every 6 months of lactation. In addition, as seen in some previous studies (34–38), neither parity nor lactation was associated with fracture hazard after age 42 years (over median 15.7 years of follow-up). Taken together, these findings suggest that parity and lactation have no (or minimal, if any) long-term implications on bone strength and fracture risk.

Any bone mass that may have been lost during pregnancy and breast feeding appears to be regained before a woman enters the menopause transition. This recovery may be partly attributable to higher levels of physical activity in those with higher parity: we found that both parity and lactation were associated with greater total physical activity level and home physical activity level, which has beneficial effects on bone health (25). We also found that parity, not lactation, was positively associated with BMI in later life, which is consistent with previous studies that have shown that parity may be associated with greater body weight in later life, while the long-term effect of lactation on weight appears to be unclear (10–12). Greater weight enhances bone re-accrual (via greater skeletal loading) (13), which could increase BMD. At the same time greater body weight increases impact forces during a fall (6, 14). It is, therefore, theoretically possible that the recovery in bone mass is not enough to compensate for the increase in impact forces. We found, however, that composite indices of femoral neck strength relative to load were not lower in women with higher parity or longer duration of lactation.

Like ours, one other recent study also reported an association between longer duration of lactation and lower BMD in the lumbar spine, but not in the femoral neck or the total hip (33). Compared with femoral neck BMD, lumbar spine BMD has a higher proportion of trabecular bone, which is more metabolically active (39, 40), and possibly more susceptible to hormonal influences and reduction in calcium reserves than femoral neck BMD. Although others and we have seen lower BMD in the lumbar spine in women with longer duration of lactation, the size of the effect in women approaching the menopause transition is small, and may not impact fracture risk in later life. No study that we are aware of has found that lactation is associated with higher fracture rate in the long-term.

The assumptions and implications of our fracture study design, in particular of not distinguishing between traumatic and non-trauma fracture, deserve mention. Just like low bone strength is a risk factor for non-trauma fracture, it is also a risk factor for traumatic fracture, in that when there is trauma, those with lower bone strength are more likely to fracture (41). However, for traumatic fractures to be useful as indicators of osteoporosis (or

low bone strength), one has to assume that the occurrence of trauma is random and not related independently to the predictors of interest (parity and lactation, in this analysis). This is analogous to the assumption made when one examines non-traumatic fractures exclusively, which is that fall risk is not related to the predictors of interest. It is not clear that either assumption is more defensible than the other. Under these two assumptions (note that both are needed here), our study implies that parity and cumulative lactation by age 42 are not related to the subsequent hazard of fracture over a median follow-up of 15.7 years.

The limitations of our study need to be acknowledged. Firstly, our assessment of duration of lactation could have been affected by recall bias. Previous studies, however, have suggested that long-term recall of lactation duration is reproducible and accurate (42, 43). Secondly, our study was not powered to find small effects on fracture risk. A previous meta-analysis showed that the relative risk of all fractures associated with one SD decrement in lumbar spine BMD is 1.5 (44); thus, the expected relative increase in risk of fracture per 6 additional months of lactation would be only 0.7%. Thirdly, we do not have information about non-clinical vertebral fractures, which might underestimate the incidence of fractures. The above-mentioned meta-analysis reported that the relative risk of spine fracture associated with one SD decrement in lumbar spine BMD is 2.3 (44). Fourthly, fractures were self-reported. However, medical records were obtained for 67% of self-reported non-digital non-cranio-facial fractures and 95% were confirmed. Furthermore, the fracture analyses examined time to first fracture after age 42 years, but covariate data was collected at SWAN baseline, when median age was 46 years. Covariates such as BMI, physical activity level, and alcohol consumption level may have changed from the baseline visit, and introduced some bias in findings. In addition, effects of pregnancy after age 42 were not addressed, and effects of adolescent pregnancy were not distinguished. Finally, the cohort was middle-aged and the rate of fractures was low.

Despite these limitations, our study has several strengths, including the multi-site design and size of the study sample, long length (nearly 16 years) of follow-up, assessment of parity and accumulated duration of lactation up to the same age (42 years) for every woman and assessment of fractures from that time point (age 42 years) forward. In addition, we investigated potential factors that might have contributed to nullify the effects of parity and lactation on bone strength, such as total physical activity level, home physical activity level, or higher BMI. Finally, to our knowledge, this is the first study to examine the associations between parity or lactation and bone strength relative to load. The importance of incorporating bone size and body size into BMD to assess bone strength relative to load has been demonstrated in multiple cohorts (15, 16, 45–47).

In conclusion, lifetime parity and cumulative duration of lactation had few, small associations with bone strength in pre- and early perimenopausal women. Parity and duration of lactation were also not associated with risk of fracture after age 42, over median follow-up of 16 years. This study adds to the accumulating evidence that parity and lactation have no (or minimal, if any) long-term deleterious effects on bone health.

Acknowledgements

The Study of Women's Health Across the Nation (SWAN) has grant support from the National Institutes of Health (NIH), DHHS, through the National Institute on Aging (NIA), the National Institute of Nursing Research (NINR) and the NIH Office of Research on Women's Health (ORWH) (Grants U01NR004061; U01AG012505, U01AG012535, U01AG012531, U01AG012539, U01AG012546, U01AG012553, U01AG012554, U01AG012495). The Hip Strength Through the Menopausal Transition has grant support from the NIA (AG026463). Takahiro Mori was supported by the VA Special Fellowship Program in Advanced Geriatrics and the VA Greater Los Angeles Geriatric Research Education and Clinical Center. The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the NIA, NINR, ORWH, VA or the NIH.

We thank the study staff at each site and all the women who participated in SWAN.

Appendix

Clinical Centers: *University of Michigan, Ann Arbor – Siobán Harlow, PI 2011 – present, MaryFran Sowers, PI 1994–2011; Massachusetts General Hospital, Boston, MA – Joel Finkelstein, PI 1999 – present; Robert Neer, PI 1994 – 1999; Rush University, Rush University Medical Center, Chicago, IL – Howard Kravitz, PI 2009 – present; Lynda Powell, PI 1994 – 2009; University of California, Davis/Kaiser – Ellen Gold, PI; University of California, Los Angeles – Gail Greendale, PI; Albert Einstein College of Medicine, Bronx, NY – Carol Derby, PI 2011 – present, Rachel Wildman, PI 2010 – 2011; Nanette Santoro, PI 2004 – 2010; University of Medicine and Dentistry – New Jersey Medical School, Newark – Gerson Weiss, PI 1994 – 2004; and the University of Pittsburgh, Pittsburgh, PA – Karen Matthews, PI.*

NIH Program Office: *National Institute on Aging, Bethesda, MD – Winifred Rossi 2012 - present; Sherry Sherman 1994 – 2012; Marcia Ory 1994 – 2001; National Institute of Nursing Research, Bethesda, MD – Program Officers.*

Central Laboratory: *University of Michigan, Ann Arbor – Daniel McConnell (Central Ligand Assay Satellite Services).*

Coordinating Center: *University of Pittsburgh, Pittsburgh, PA – Maria Mori Brooks, PI 2012 -present; Kim Sutton-Tyrrell, PI 2001 – 2012; New England Research Institutes, Watertown, MA -Sonja McKinlay, PI 1995 – 2001.*

Steering Committee: Susan Johnson, Current Chair

Chris Gallagher, Former Chair

References

1. Specker BL. Bone mineral changes during pregnancy and lactation. *Endocrine*. 2002; 17:49–53. [PubMed: 12014704]
2. Karlsson MK, Ahlborg HG, Karlsson C. Female reproductive history and the skeleton—a review. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2005; 112:851–856. [PubMed: 15957983]
3. Møller U, við Streym S, Mosekilde L, Rejnmark L. Changes in bone mineral density and body composition during pregnancy and postpartum. A controlled cohort study. *Osteoporosis International*. 2012; 23:1213–1223. [PubMed: 21607805]

4. Alonso CG, Curiel M, Carranza F, Cano R, Perez A. Femoral bone mineral density, neck-shaft angle and mean femoral neck width as predictors of hip fracture in men and women. Multicenter Project for Research in Osteoporosis. Osteoporosis international: a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA. 2000; 11:714.
5. Faulkner KG, Cummings SR, Black D, Palermo L, Glüer CC, Genant HK. Simple measurement of femoral geometry predicts hip fracture: the study of osteoporotic fractures. Journal of bone and mineral research. 2009; 8:1211–1217. [PubMed: 8256658]
6. Beck TJ, Petit MA, Wu G, LeBoff MS, Cauley JA, Chen Z. Does Obesity Really Make the Femur Stronger? BMD, Geometry, and Fracture Incidence in the Women's Health Initiative - Observational Study. Journal of Bone and Mineral Research. 2009; 24:1369–1379. [PubMed: 19292617]
7. Specker B, Binkley T. High parity is associated with increased bone size and strength. Osteoporosis international. 2005; 16:1969–1974. [PubMed: 16091837]
8. Wiklund P, Xu L, Wang Q, Mikkola T, Lyytikäinen A, Völgyi E, Munukka E, Cheng S, Alen M, Keinänen-Kiukaanniemi S. Lactation is associated with greater maternal bone size and bone strength later in life. Osteoporosis International. 2012; 23:1939–1945. [PubMed: 21927916]
9. Laskey M, Price R, Khoo B, Prentice A. Proximal femur structural geometry changes during and following lactation. Bone. 2011; 48:755–759. [PubMed: 21130909]
10. Gunderson EP. Childbearing and obesity in women: weight before, during, and after pregnancy. Obstet. Gynecol. Clin. North Am. 2009; 36:317–332. [PubMed: 19501316]
11. Nehring I, Schmoll S, Beyerlein A, Hauner H, von Kries R. Gestational weight gain and long-term postpartum weight retention: a meta-analysis. The American journal of clinical nutrition. 2011; 94:1225–1231. [PubMed: 21918221]
12. Chung M, Raman G, Chew P, Magula N, Trikalinos T, Lau J. Breastfeeding and maternal and infant health outcomes in developed countries. Evid Technol Asses (Full Rep). 2007; 153:1–186.
13. Felson DT, Zhang Y, Hannan MT, Anderson JJ. Effects of weight and body mass index on bone mineral density in men and women: The framingham study. Journal of Bone and Mineral Research. 1993; 8:567–573. [PubMed: 8511983]
14. Ishii S, Cauley JA, Greendale GA, Nielsen C, Karvonen-Gutierrez C, Ruppert K, Karlamangla AS. Pleiotropic Effects of Obesity on Fracture Risk: The Study of Women's Health Across the Nation. Journal of Bone and Mineral Research. 2014
15. Karlamangla AS, Barrett-Connor E, Young J, Greendale GA. Hip fracture risk assessment using composite indices of femoral neck strength: the Rancho Bernardo study. Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA. 2004; 15:62–70.
16. Ishii S, Greendale GA, Cauley JA, Crandall CJ, Huang M-H, Danielson ME, Karlamangla AS. Fracture risk assessment without race/ethnicity information. Journal of Clinical Endocrinology & Metabolism. 2012; 97:3593–3602. [PubMed: 22865903]
17. Ishii S, Cauley JA, Crandall CJ, Srikanthan P, Greendale GA, Huang M-H, Danielson ME, Karlamangla AS. Diabetes and femoral neck strength: findings from the Hip Strength Across the Menopausal Transition Study. Journal of Clinical Endocrinology & Metabolism. 2012; 97:190–197. [PubMed: 22072739]
18. Ishii S, Cauley J, Greendale G, Danielson M, Nili NS, Karlamangla A. Ethnic differences in composite indices of femoral neck strength. Osteoporosis international. 2012; 23:1381–1390. [PubMed: 21927926]
19. Heaney R, Abrams S, Dawson-Hughes B, Looker A, Looker A, Marcus R, Matkovic V, Weaver C. Peak bone mass. Osteoporosis International. 2000; 11:985–1009. [PubMed: 11256898]
20. Riis BJ, Hansen MA, Jensen AM, Overgaard K, Christiansen C. Low bone mass and fast rate of bone loss at menopause: equal risk factors for future fracture: a 15-year follow-up study. Bone. 1996; 19:9–12. [PubMed: 8830981]
21. Cauley JA, Danielson ME, Greendale GA, Finkelstein JS, Chang Y-F, Lo JC, Crandall CJ, Neer RM, Ruppert K, Meyn L. Bone resorption and fracture across the menopausal transition: the Study of Women's Health Across the Nation. Menopause (New York, NY). 2012; 19:1200.

22. Sowers, MCS.; Sternfeld, B.; Morganstein, D.; Gold, E.; Greendale, G.; Evans, D.; Neer, R.; Matthews, K.; Sherman, S.; Lo, A.; Weiss, G.; Kelsey, J. Menopause: biology and pathobiology. Academic: San Diego; 2000 Design, survey, sampling and recruitment methods of SWAN: a multi-center, multi-ethnic, community based cohort study of women and the menopausal transition. p. 175-188.
23. Finkelstein JS, Brockwell SE, Mehta V, Greendale GA, Sowers MR, Ettinger B, Lo JC, Johnston JM, Cauley JA, Danielson ME, Neer RM. Bone mineral density changes during the menopause transition in a multiethnic cohort of women. The Journal of clinical endocrinology and metabolism. 2008; 93:861–868. [PubMed: 18160467]
24. Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. The American journal of clinical nutrition. 1982; 36:936–942. [PubMed: 7137077]
25. Mori T, Ishii S, Greendale GA, Cauley JA, Sternfeld B, Crandall CJ, Han W, Karlamangla AS. Physical activity as determinant of femoral neck strength relative to load in adult women: findings from the hip strength across the menopause transition study. Osteoporosis International. 2014; 25:265–272. [PubMed: 23812598]
26. Horton NJ, Lipsitz SR. Multiple imputation in practice: comparison of software packages for regression models with missing variables. The American Statistician. 2001; 55:244–254.
27. Heitjan DF, Little RJ. Multiple imputation for the fatal accident reporting system. Applied Statistics. 1991:13–29.
28. Seeley DG, Browner WS, Nevitt MC, Genant HK, Scott JC, Cummings SR. Which fractures are associated with low appendicular bone mass in elderly women? Ann. Intern. Med. 1991; 115:837–842. [PubMed: 1952469]
29. Stone KL, Seeley DG, Lui LY, Cauley JA, Ensrud K, Browner WS, Nevitt MC, Cummings SR. BMD at multiple sites and risk of fracture of multiple types: long - term results from the Study of Osteoporotic Fractures. Journal of Bone and Mineral Research. 2003; 18:1947–1954. [PubMed: 14606506]
30. Lissner L, Bengtsson C, Hansson T. Bone mineral content in relation to lactation history in pre-and postmenopausal women. Calcif. Tissue Int. 1991; 48:319–325. [PubMed: 2054717]
31. Kojima N, Douchi T, Kosha S, Nagata Y. Cross-sectional study of the effects of parturition and lactation on bone mineral density later in life. Maturitas. 2002; 41:203–209. [PubMed: 11886766]
32. Dursun N, Akin S, Dursun E, Sade I, Korkusuz F. Influence of duration of total breast-feeding on bone mineral density in a Turkish population: does the priority of risk factors differ from society to society? Osteoporosis international. 2006; 17:651–655. [PubMed: 16508701]
33. Tsvetov G, Levy S, Benbassat C, Shraga-Slutsky I, Hirsch D. Influence of number of deliveries and total breast-feeding time on bone mineral density in premenopausal and young postmenopausal women. Maturitas. 2013
34. Grisso J, Kelsey J, Gammon M, O'Brien L. Parity, lactation and hip fracture. Osteoporosis International. 1993; 3:171–176. [PubMed: 8338971]
35. Alderman BW, WEISS NS, DALING JR, URE C, BALLARD JH. Reproductive history and postmenopausal risk of hip and forearm fracture. Am. J. Epidemiol. 1986; 124:262–267. [PubMed: 3728442]
36. O'Neill TW, Silman AJ, Diaz MN, Cooper C, Kanis J, Felsenberg D. Influence of hormonal and reproductive factors on the risk of vertebral deformity in European women. Osteoporosis International. 1997; 7:72–78. [PubMed: 9102068]
37. Johnell O, Gullberg B, Kanis JA, Allander E, Elffors L, Dequeker J, Dilsen G, Gennari C, Vaz LA, Lyritis G. Risk factors for hip fracture in European women: the MEDOS study. Journal of Bone and Mineral Research. 1995; 10:1802–1815. [PubMed: 8592959]
38. Bjørnerem Å, Ahmed LA, Jørgensen L, Størmer J, Joakimsen RM. Breastfeeding protects against hip fracture in postmenopausal women: The Tromsø study. Journal of Bone and Mineral Research. 2011; 26:2843–2850. [PubMed: 21898594]
39. Kent GN, Price RI, Gutteridge DH, Allen JR, Barnes MP, Hickling CJ, Retallack RW, Wilson SG, Devlin RD, Smith M. Human lactation: forearm trabecular bone loss, increased bone turnover, and

- renal conservation of calcium and inorganic phosphate with recovery of bone mass following weaning. *Journal of Bone and Mineral Research*. 1990; 5:361–369. [PubMed: 2343775]
40. Greendale GA, Sowers M, Han W, Huang MH, Finkelstein JS, Crandall CJ, Lee JS, Karlamangla AS. Bone mineral density loss in relation to the final menstrual period in a multiethnic cohort: Results from the Study of Women's Health Across the Nation (SWAN). *Journal of Bone and Mineral Research*. 2012; 27:111–118. [PubMed: 21976317]
41. Mackey DC, Lui L-Y, Cawthon PM, Bauer DC, Nevitt MC, Cauley JA, Hillier TA, Lewis CE, Barrett-Connor E, Cummings SR. High-trauma fractures and low bone mineral density in older women and men. *JAMA: the journal of the American Medical Association*. 2007; 298:2381–2388.
42. Li R, Scanlon KS, Serdula MK. The validity and reliability of maternal recall of breastfeeding practice. *Nutr. Rev.* 2005; 63:103–110. [PubMed: 15869124]
43. Tomeo CA, Rich-Edwards JW, Michels KB, Berkey CS, Hunter DJ, Frazier AL, Willett WC, Buka SL. Reproducibility and validity of maternal recall of pregnancy-related events. *Epidemiology*. 1999; 10:774–776. [PubMed: 10535796]
44. Marshall D, Johnell O, Wedel H. Meta-analysis of how well measures of bone mineral density predict occurrence of osteoporotic fractures. *BMJ (Clinical research ed.)*. 1996; 312:1254–1259.
45. Faulkner KG, Wacker W, Barden H, Simonelli C, Burke P, Ragi S, Del Rio L. Femur strength index predicts hip fracture independent of bone density and hip axis length. *Osteoporosis international*. 2006; 17:593–599. [PubMed: 16447009]
46. Dufour A, Roberts B, Broe K, Kiel D, Bouxsein M, Hannan M. The factor-of-risk biomechanical approach predicts hip fracture in men and women: the Framingham Study. *Osteoporosis International*. 2012; 23:513–520. [PubMed: 21344243]
47. Leslie W, Pahlavan P, Tsang J, Lix L. Prediction of hip and other osteoporotic fractures from hip geometry in a large clinical cohort. *Osteoporosis international*. 2009; 20:1767–1774. [PubMed: 19238304]

Highlights

We analyzed longitudinal data from the Study of Women's Health Across the Nation.

Parity and lactation have little impact on peak bone strength prior to menopause.

Parity and lactation do not affect fracture risk after age 42 over 16-year follow-up.

Fig. 1. Femoral neck size measurements. AB is the femoral neck axis length (FNAL): the distance from the base of the greater trochanter to the apex of the femoral head. DE is the femoral neck width (FNW): the smallest thickness of the femoral neck along any line perpendicular to the femoral neck axis. C is where the femoral neck axis meets the inner pelvic rim.

Table 1Characteristics^a of the study participants at baseline^b

Characteristics	Study sample with composite indices of femoral neck strength data (n=1881 ^c)	Study sample with fracture data (n=2239 ^d)
Age (year)	46 [44, 48]	46 [44, 48]
Race/ethnicity		
Caucasian	936 (49.8%)	1115 (49.8%)
African American	499 (26.5%)	637 (28.5%)
Japanese	231 (12.3%)	252 (11.3%)
Chinese	215 (11.4%)	235 (10.5%)
Body mass index (kg/m ²)	25.6 [22.3, 31.1]	26.0 [22.5, 31.6]
Menopausal transition stage		
Premenopausal	1,066 (56.7%)	1210 (54.3%)
Early perimenopausal	815 (43.3%)	1017 (45.7%)
Smoking status		
Current	281 (15.1%)	367 (16.5%)
Ex-smoker	475 (25.4%)	570 (25.7%)
Never smoked	1111 (59.5%)	1285 (57.8%)
Smoking pack-year		
0	1,190 (64.2%)	1382 (62.7%)
10 years	300 (16.2%)	351 (15.9%)
>10–30 years	291 (15.7%)	364 (16.5%)
>30 years	74 (4.0%)	108 (4.9%)
Alcohol consumption level		
Abstainer	899 (51.2%)	1074 (51.3%)
Infrequent	391 (22.3%)	459 (21.9%)
Light to moderate	364 (20.7%)	442 (21.1%)
Heavy	101 (5.8%)	120 (5.7%)
History of diabetes	85 (4.5%)	117 (5.2%)
History of hyperthyroidism	68 (3.6%)	81 (3.6%)
Current use of supplementary calcium	844 (44.9%)	988 (44.2%)
Current use of supplementary vitamin D	723 (38.5%)	851 (38.1%)
Medication use at baseline		
Prior use of sex steroid hormones (pills, patches, or injections) other than birth control pills	119 (6.4%)	159 (7.1%)
Prior use of birth control pills	1382 (73.8%)	1643 (73.7%)
Prior use of depo-provera injections	13 (0.7%)	18 (0.8%)
Current or prior use of oral corticosteroids	109 (5.8%)	142 (6.3%)
Current use of proton pump inhibitors	24 (1.3%)	29 (1.3%)
Current or prior use of bone adverse medications ^e	50 (2.7%)	57 (2.5%)

Characteristics	Study sample with composite indices of femoral neck strength data (n=1881 ^c)	Study sample with fracture data (n=2239 ^d)
Medication use ever (till 12 th follow up) ^f		
Sex steroid hormones (pills, patches, or injections) other than birth control pills	711 (37.8%)	825 (36.8%)
Birth control pills	1423 (75.7%)	1687 (75.3%)
Depo-provera injections	25 (1.3%)	30 (1.3%)
Oral corticosteroids	449 (23.9%)	523 (23.4%)
Proton pump inhibitors	399 (21.2%)	457 (20.4%)
Bone adverse medications ^e	312 (16.6%)	350 (15.6%)
Physical activity score (ranging from 4 to 20) ^g	9.7 [8.4, 11.1]	9.7 [8.5, 11.1]
Home activity score (ranging from 1 to 5)	2.6 [2.2, 3.4]	2.6 [2.2, 3.4]
Employment status	1548 (82.3%)	1914 (82.1%)
Parity and lactation		
Parity (including live births and stillbirth)	2 [1, 3]	2 [1, 3]
Duration of lactation (months)	2 [0, 12]	1 [0, 12]
Bone Strength Measurements		
Femoral neck bone mineral density (g/cm ²)	0.83 [0.74, 0.92]	0.84 [0.75, 0.93]
Lumbar spine bone mineral density (g/cm ²)	1.06 [0.97, 1.15]	1.07 [0.97, 1.15]
Compression strength index (g/kg-m)	3.28 [2.86, 3.70]	-
Bending strength index (g/kg-m)	1.00 [0.86, 1.15]	-
Impact strength index (g/kg-m)	0.18 [0.16, 0.21]	-

^a Median and interquartile range for continuous variables and number of participants and percentage for categorical variables.

^b All characteristics reported were measured at baseline except the 'medication use ever' variables, which were used in the fracture analysis

^c Femoral neck size was measured in a subset of women in the Hip Strength Across the Menopause Transition SubStudy. Sample sizes were smaller than 1889 for femoral neck bone mineral density (n=1880) and lumbar spine bone mineral density (n=1708).

^d Sample sizes were smaller than 2239 for femoral neck bone mineral density (n=2235) and lumbar spine bone mineral density (n=2022).

^e Included antiepileptic medications, chemotherapy, gonadotropin-releasing hormone agonist, aromatase inhibitors, or thiazolidinediones.

^f Any use including either prior or at baseline, or during the follow-up till visit 12.

^g Sum of four domains of physical activity: sport, home, active living, and work.

Table 2

Adjusted¹ associations² (with 95 % confidence interval) of lifetime parity (before age 42) and accumulated length of lactation (before age 42) with bone strength measurements at study baseline.

	Femoral neck BMD (n=2235, mean 0.85, SD 0.14)	Lumbar spine BMD (n=2022, mean 1.07, SD 0.13)	Compression strength index (n=1881, mean 3.3, SD 0.64)	Bending strength index (n=1881, mean 1.02, SD 0.22)	Impact strength index (n=1881, mean 0.18, SD 0.04)
Lifetime Parity (per child/birth)	0.0002 (-0.025, 0.025)	-0.026 (-0.058, 0.006)	0.022 (-0.002, 0.046)	0.024 (-0.004, 0.051)	0.024* (0.0001 0.048)
Lactation Duration (per every 6 months)	0.002 (-0.012, 0.016)	-0.018* (-0.036 0.001)	0.007 (-0.007, 0.020)	0.009 (-0.006, 0.024)	0.006 (-0.007, 0.020)

Abbreviations: BMD=bone mineral density, SD=standard deviation

¹ Multiple linear regressions adjusted for age, race/ethnicity, menopausal transition stage, body mass index, smoking status, smoking pack-years, alcohol consumption level, physical activity level, employment status, diabetes, hyperthyroidism, current use of supplementary calcium, current use of supplementary vitamin D, prior use of sex steroid hormones, prior use of birth control pills, prior use of depo-provera injections, current or prior use of oral corticosteroids, current use of proton pump inhibitors, other bone-adverse medications, and study site.

² Units: BMD or strength index standard deviation.

* p<0.05