In brief

Surgeon admits manslaughter: Consultant surgeon Steven Walker, aged 47, who was accused of killing two patients through gross negligence, changed his plea last week and admitted manslaughter of one of the women. The prosecution decided not to proceed with the charges over the second patient. Mr Walker, from Camberley, Surrey, was due to be sentenced on 23 June after the *BMJ* went to press.

Dutch nurse guilty of murdering seven patients: An appeal court in the Hague has sentenced nurse Lucy de Berk, aged 42, to life imprisonment and compulsory psychiatric care for the murder of seven patients and the attempted murder of another three. Her crimes took place at three hospitals in the Hague where she worked from 1997 to 2001 (*BMJ* 2003;326:680).

Department criticised for not knowing its spending on AIDS: A report from the National Audit Office last week criticised the Department for International Development for not knowing how much it was spending on its AIDS programmes. Estimates for 2002-3 varied from £103m (\$190m; €155m) to £274m, depending on whether reproductive health programmes were included. See www.nao.org.uk

Israeli surgeon imprisoned for manslaughter: Senior Israeli surgeon Professor Vladimir Yakirevich was sentenced to seven years in prison and given a \$44 000 (£24 000; €36 000) fine by the Tel Aviv District Court after having been convicted on two counts of manslaughter, blackmail, theft, and taking bribes while at the Sourasky Medical Centre in the 1990s (*BMJ* 2000;320:1292).

Report calls for reform of private finance initiative: A report from the NHS Confederation says the private finance initiative (PFI) in the NHS must be reformed. It calls for the PFI regime to be more flexible and for bed requirements to be more accurately assessed. *Getting the Best out of Future Capital Investment in Health* is available from Will Little, tel 07880 500 726.

Girls in southern Sudan are more likely to die in childbirth than complete primary school

Peter Moszynski *Geneva*

Girls in southern Sudan are more likely to die in pregnancy or childbirth than complete primary education, says a Unicef report published last week, to mark the International Day of the African Child.

"After 21 years of war, southern Sudan ranks as the worst place in the world for many key indicators of women and children's well-being, including its rates of chronic malnutrition, primary school completion, immunisation and antenatal care," says Unicef, announcing the study.

Describing the state of women and children as "shocking," Unicef calculates that: "A girl born in southern Sudan has a better chance of dying in pregnancy or childbirth than of completing primary school. To put it another way, one in nine women dies in pregnancy or childbirth but only one in a hundred girls completes primary school."

Some 95 000 children aged

under 5 years are estimated to have died last year in southern Sudan (population 7.5 million), mostly from preventable disease. The total death toll of under 5s in the world's 31 industrialised countries (with a combined population of 938 million) was 76 000, the report says.

Only one in 50 children completes primary education, which is thought to be "the lowest rate in the world," and only one in four adults (one in 10 women) is literate.

The study, conducted by the New Sudan Centre for Statistics and Evaluation (the statistical arm of the de facto government of South Sudan) in conjunction with Unicef, reviews surveys conducted in southern Sudan during recent years and compares them with models and figures from neighbouring countries and statistics from the rest of Sudan. There have been virtually no reliable data from southern Sudan since

before the war, and the last reliable census was before independence in 1956.

A new census is due to be conducted during the interim period following the signing of a comprehensive peace agreement, and this would provide a baseline for planning and monitoring social progress, says Unicef. □

Towards a Baseline: Best Estimates of Social Indicators for Southern Sudan is accessible at www.unsudanig.org/Publications/assessments

Only 1 in 100 girls in southern Sudan complete primary school

Oestrogen doesn't protect mental function in older women

Janice Hopkins Tanne *New York*

The "oestrogen only" treatment arm of the women's health initiative memory study shows that oestrogen does not protect against mild cognitive impairment or dementia and slightly increases the risk, say two reports published in *JAMA* (2004;291:2947-58, 2959-68). Mild cognitive impairment strongly predicts dementia.

Previous studies have suggested that oestrogen might prevent cognitive decline, said Lon Schneider, professor of psychiatry, neurology, and gerontology at the University of Southern California in Los Angeles, who commented on the studies in an editorial (*JAMA* 2004;291:3005-7). He told the *BMJ* that the finding which showed oestrogen

might increase the risk of cognitive impairment and dementia was unexpected. "The studies show that giving oestrogen or oestrogen and progestin [progesterone] to women over 65 in hopes of delaying dementia or increasing cognitive function is misplaced. They should not be getting oestrogen."

Two years ago, the women's health initiative—a trial to evaluate oestrogen and progesterone in 27 000 postmenopausal women—was stopped early (*BMJ* 2002;325:61). The oestrogen only part of the trial continued until February 2004 (*BMJ* 2004;328:602).

The two new reports cover cognitive function in the oestrogen only study and pooled results with the oestrogen-progesterone study.

The oestrogen only study included 2808 women aged 65-79 who had a baseline test of global cognitive function and at least one follow up test with the modified mini-mental state examination (3MSE): 1387 women who took 0.625 mg of

combined equine oestrogen daily and 1421 who took a placebo. During a mean follow up of 5.4 years, probable dementia was diagnosed in 28 women receiving oestrogen and in 19 receiving placebo (hazard ratio 1.49; 95% confidence interval 0.83 to 2.66).

Lead author Mark Espeland, professor of public health sciences at Wake Forest University School of Medicine in Winston-Salem, North Carolina, told the *BMJ* that women with relatively low cognitive function at baseline were at particularly high risk of deterioration.

Dr Sally Shumaker, the principal national investigator of the memory study and associate dean for research at Wake Forest, and colleagues pooled data from the oestrogen only trial and the trial of combined 0.625 mg of oestrogen plus 2.5 mg of medroxyprogesterone acetate per day in 4532 women aged 65-79. They found the overall hazard ratio for probable dementia was 1.76 (95% confidence interval 1.19 to 2.60; $P=0.005$). □