Published in final edited form as:

Plant J. 2011 June ; 66(5): 844–851. doi:10.1111/j.1365-3113X.2011.04550.x.

Identification of the MBF1 heat-response regulon of *Arabidopsis thaliana*

Nobuhiro Suzuki¹, Hiroe Sejima², Rachel Tam², Karen Schlauch², and Ron Mittler^{1,3,*}

¹Department of Biological Sciences, College of Arts and Sciences, University of North Texas, 1155 Union Circle #305220, Denton, TX 76203

²Department of Biochemistry and Molecular Biology, University of Nevada, Mail Stop 200, Reno NV 89557, USA

³Department of Plant Sciences, Hebrew University of Jerusalem, Jerusalem 91904, Israel

SUMMARY

Brief periods of heat stress of even a few days can have a detrimental effect on yield production worldwide causing devastating economical and sociological impacts. Here we report on the identification of a new heat-response regulon in plants controlled by the multiprotein bridging factor 1c (MBF1c) protein of *Arabidopsis thaliana*. Members of the highly conserved MBF1 protein family function as non DNA-binding transcriptional co-activators involved in regulating metabolic and development pathways in different organisms from yeast to humans. Nonetheless, our studies suggest that MBF1c from *Arabidopsis* functions as a transcriptional regulator which binds DNA and controls the expression of 36 different transcripts during heat stress, including the important transcriptional regulator DRE-binding protein 2A (DREB2A), two heat shock transcription factors (HSFs), and several zinc finger proteins. We further identify CTAGA as a putative response element for MBF1c, demonstrate that the DNA-binding domain of MBF1c has a dominant-negative effect on heat tolerance when constitutively expressed in plants, and show that constitutive expression of MBF1c in soybean enhances yield production in plants grown under controlled growth conditions without causing adverse effects on growth. Our findings could have a significant impact on improving heat tolerance and yield of different crops subjected to heat stress.

Keywords

Heat stress; transcription; regulon; MBF1; HSF; HSP

INTRODUCTION

The heat stress response is a highly conserved response involving multiple pathways, regulatory networks, and cellular compartments (Kotak *et al.*, 2007). At least two key

*Corresponding Author: Ron Mittler, Department of Biological Sciences, College of Arts and Sciences, University of North Texas, 1155 Union Circle #305220, Denton, TX 76203-5017 ron.mittler@unt.edu, TEL: 940-369-5367, FAX: 940-565-3821.

Microarray accession data: Array Express (E-MEXP-2760).

components coordinating this response in plants have been identified. They include a network of heat shock transcription factors (HSFs), and the recently identified multiprotein bridging factor 1c (MBF1c) protein (von Koskull-Döring *et al.*, 2007; Suzuki *et al.*, 2008). The function of HSFs as DNA-binding transcriptional regulators that control the heat stress response by binding to a defined heat shock response element (HSE) has been established in multiple biological systems (Shamovsky and Nudler, 2008). In contrast, MBF1 was only recently identified as a key regulator of thermotolerance in plants and its mode of action is largely unknown (Suzuki *et al.*, 2008). MBF1 is a highly conserved protein thought to function as a non DNA-binding transcriptional co-activator involved in different developmental and metabolic pathways in different organisms, ranging from yeast to humans (Takemaru *et al.*, 1998; Kabe *et al.*, 1999; Liu *et al.*, 2003). In the model plant *Arabidopsis thaliana* MBF1 is encoded by 3 different genes (*Mbf1a*, *b*, and *c*; Tsuda and Yamazaki, 2004) of which *Mbf1c* (At3g24500) is required for thermotolerance (Suzuki *et al.*, 2008). Genetic analysis of MBF1c in *Arabidopsis* using gain- and loss-of-function mutants demonstrated that MBF1c functions upstream to salicylic acid, ethylene, and trehalose signaling during heat stress (Suzuki *et al.*, 2005, 2008) MBF1c is not however required for the expression of many heat shock proteins (HSP) or ascorbate peroxidase 1 during heat stress (Suzuki *et al.*, 2005, 2008). Nonetheless, its mode of action during heat stress remains unknown. Here we show that MBF1c functions as a transcriptional regulator which most likely binds DNA and controls a regulon of 36 different transcripts during heat stress, including the important transcriptional regulator DRE-binding protein 2A (DREB2A; Schramm *et al.*, 2007), two HSFs, and several zinc finger proteins. We further identify a putative response element for MBF1 and test it using gel-shift and yeast one-hybrid assays, as well as demonstrate that ectopic expression of the DNA-binding domain of MBF1c in transgenic plants has a dominant-negative effect on heat tolerance. Our study identifies a new heat stress-response regulon in plants and opens the way for the development of new strategies to enhance thermotolerance in different crops.

RESULTS

Identification of the MBF1c regulon

To identify transcripts that function downstream to MBF1c during heat stress we subjected wild type and *mbf1c* mutants (Suzuki *et al.*, 2008) to heat stress and analyzed their transcriptome using ATH1® DNA chips at 0, 20 and 60 min following heat stress application. For each time point we used 3 technical repeats (3 different chips each representing 60 different plants), and the entire experiment was conducted with 3 biological repeats. Statistical analysis identified 36 different transcripts significantly less elevated in *mbf1c* compared to wild type during heat stress (Tables 1 and S1, S3 and S4), and several of these transcripts were confirmed by quantitative real-time PCR (qPCR) analysis in wild type and *mbf1c* plants subjected to heat stress (Figure 1), and in wild type and *mbf1c* plants complemented by expression of MBF1c under the control of the CaMV35S promoter (Suzuki *et al.*, 2008) subjected to heat stress (Figure S1).

The majority of transcripts identified by our analysis were also identified as elevated during heat or salinity stresses in transcriptome databases (Table 1; Horan *et al.*, 2008),

demonstrating that they belong to the plant abiotic stress-response network. Interestingly, MBF1c was required for the elevated expression of two HSFs (B2B and B2A), the transcriptional regulator DREB2A (that functions upstream to HSFA3; Schramm *et al.*, 2007), and a component of the RNA polymerase basal apparatus (TBP-Associated Factor 7). Because MBF1 proteins were postulated to associate with the TBP complex (Takemaru *et al.*, 1998; Kabe *et al.*, 1999; Liu *et al.*, 2003), the later could represent a protein involved in the association of MBF1c with the RNA polymerase complex.

Genetic analysis of selected transcripts

To confirm that some of the transcripts identified by our analysis are involved in the heat stress response of plants we chose 3 genes encoding these transcripts (*Dreb2A* – *At5g05410*, *Sen1* - *At4g35770*, and the zinc finger protein - *At3g21890*), and obtained two independent homozygous knockout lines for each gene. We then subjected these lines to heat stress and measured their survival. The three genes encoding the MBF1c-dependent transcripts were required for heat tolerance in Arabidopsis, demonstrating that they are an integral part of the plant thermotolerance pathway (Figure 2).

Identification of a putative MBF1c DNA-binding element

MBF1 proteins are not thought to be DNA binding proteins (Takemaru *et al.*, 1998; Kabe *et al.*, 1999; Liu *et al.*, 2003). Nevertheless, they contain a putative helix-turn-helix domain that may bind DNA (de Koning *et al.*, 2009). Bioinformatics analysis of the promoter regions of the 36 MBF1c-dependent transcripts (Table 1) identified two DNA sequences that appeared at high frequencies in these promoters (Table 1 and Figure 3). SELEX analysis (Bouvet, 2001; Clouaire *et al.*, 2005) using purified MBF1c protein identified two additional DNA sequences that showed binding to MBF1c (Table 1 and Figure S2). All four sequences identified by our analysis (Table 1) showed binding to the purified MBF1c protein in gel mobility shift assays (Figure S3). Nonetheless, only one sequence identified by our transcriptome analysis (CTAGA) showed specific MBF1c-dependent transcriptional activation in yeast (Figure 3), with another sequence identified by our SELEX assay (GGAGG) showing weaker activity in this system (Figure S4).

When the frequency of appearance of the different DNA elements in the 500 bp region of promoters from all heat stress response genes identified in this study (60 min heat stress) was compared to that of all Arabidopsis genes (Table 2), it was found that the CTAGA element was disproportionally more represented in the promoters of heat response genes. Based on this analysis (Table 2), the yeast one-hybrid assays (Figure 3) and the gel mobility shift assays (Figure S3), CTAGA is a good candidate for a putative MBF1c DNA-binding element.

Dominant-negative function of the DNA-binding domain of MBF1c

A dominant-negative effect can occur when the DNA-binding domain of a transcription factor is expressed in cells and competes with the endogenous protein without activating transcription. We therefore fused the putative DNA-binding domain of MBF1c (67 aa at the C-terminal of MBF1c; HTH-GFP), or the rest of the MBF1c protein (90 aa at the N-terminal of MBF1c; N-term-GFP) to GFP and expressed these fusion proteins in wild type

Arabidopsis plants (Figure 4a). The GFP-fusion protein, containing the C-terminal DNA-binding domain of MBF1c, suppressed thermotolerance to levels similar to that of the null *mbf1c* mutant when expressed in transgenic plants (Figure 4b). In contrast, the GFP-fusion protein, containing the N-terminal of MBF1c without the DNA binding domain, did not significantly suppress thermotolerance when expressed in transgenic plants (Figure 4c). Although the two fusion proteins were localized to the nuclei of transgenic plants following heat stress, as is the full-length MBF1c (Suzuki *et al.*, 2008), the HTH-GFP protein appeared to do so more efficiently than the N-term-GFP protein (Figures 4b and 4c).

Ectopic expression of MBF1c in soybean improves yield in plants grown under controlled growth conditions

Our previous studies demonstrated that ectopic expression of MBF1c in Arabidopsis enhances tolerance to heat and osmotic stress without a negative effect on plant growth and yield (Suzuki *et al.*, 2005, 2008). Moreover, ectopic expression of MBF1c was found to improve the growth and yield of transgenic Arabidopsis plants grown under controlled growth conditions (Suzuki *et al.*, 2005). To test whether MBF1c has a similar effect on a crop plant we transformed soybean with Arabidopsis MBF1c. Ectopic expression of MBF1c in soybean resulted in enhanced yield in plants grown under controlled growth conditions in a greenhouse without any adverse effects on growth of plants (Figure 5).

DISCUSSION

Current and predicted climatic conditions, such as prolonged drought and heat episodes, pose a serious challenge to agricultural production worldwide, affecting plant growth and yield, and causing annual losses estimated at billions of dollars (Mittler and Blumwald, 2010). Dissecting the heat stress response of plants could therefore lead to the development of crops with enhanced thermotolerance (Mittler, 2006; Kotak *et al.*, 2007; von Koskull-Döring *et al.*, 2007; Mittler and Blumwald, 2010). Here we report on the identification of a new heat stress-response regulon in plants controlled by MBF1c. This regulon includes important transcriptional regulators such as DREB2A (that controls HSF3; Schramm *et al.*, 2007), two HSFs (B2A, B2B), and at least two other proteins required for thermotolerance in Arabidopsis (Figure 2). We further demonstrate that MBF1c is likely to function as a DNA-binding transcriptional regulator and identify at least one putative DNA-binding element for MBF1c (CTAGA; Table 2; Figure 3). The finding that MBF1c could function as a DNA-binding protein is new because this protein was previously proposed to function as a non DNA-binding transcriptional co-activator in different systems (Takemaru *et al.*, 1998; Kabe *et al.*, 1999; Liu *et al.*, 2003; de Koning *et al.*, 2009; Tsuda *et al.*, 2004). Unlike the MBF1a and MBF1b proteins of Arabidopsis, MBF1c is highly responsive to abiotic stress and shows the least homology to MBF1 proteins from other sources (Tsuda and Yamazaki, 2004; Tsuda *et al.*, 2004). It is therefore possible that MBF1c has undergone significant changes during evolution making it a DNA-binding protein involved in the abiotic stress response of plants. This possibility is in agreement with the unique function of MBF1c as a stress-response protein, which is different from the function proposed for many MBF1 proteins from other sources, generally thought to be involved in regulating development and metabolism (Takemaru *et al.*, 1998; Kabe *et al.*, 1999; Liu *et al.*, 2003). Moreover, MBF1c

shares only 54–55% identity in its putative DNA-binding domain with MBF1a or MBF1b, whereas MBF1a and MBF1b share 93% identity within their putative DNA-binding domains. MBF1c might therefore represent a newer branching of the MBF1 protein family involved in stress regulation.

Our functional characterization of MBF1c adds to our knowledge of the plant thermotolerance network demonstrating that MBF1c functions by regulating a new heat response regulon that is intertwined with the important HSF network. Our demonstration that MBF1c can enhance thermotolerance in *Arabidopsis*, as well as yield of *Arabidopsis* and soybean plants grown under controlled growth conditions (Suzuki *et al.*, 2005; Figure S5), could open the way for the testing of this gene in laboratory and field studies in different crops. These studies could have a significant impact on improving heat tolerance and yield in regions subjected to the detrimental effects of global warming.

EXPERIMENTAL PROCEDURES

Plant material, growth conditions and stress treatment

Arabidopsis thaliana plants (cv. Columbia) were grown on peat pellets (Jiffy-7, Shippagan, Canada), under controlled conditions: 21°C, 14 h light cycle, 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and a relative humidity of 70% (Suzuki *et al.*, 2005). Null mutants for the zinc finger protein At3g21890 (CS27058 and CS25831), DREB2A (Salk_016495 and Salk_084889) and SEN1 (Salk_116080 and Salk_151963) were obtained, backcrossed and homozygous as described in Rizhsky *et al.*, (2004), Davletova *et al.*, (2005a, b). GFP, or GFP fused in-frame to the C- or N-terminal domains of MBF1c were expressed in plants under the control of the 35S CaMV promoter, using a pGreen-based vector, and visualized using an Olympus, IX 81 FV 1000 confocal microscope (Olympus, Tokyo), as described in Luhua *et al.*, (2008), Suzuki *et al.*, (2008). Two T3 homozygous lines obtained from 2 individual transformation events were used in these studies. Survival rate of seedlings subjected to heat stress was performed as described in (Suzuki *et al.*, 2005). Two independent *mbf1c* mutants expressing MBF1c under the control of the CaMV35S promoter were obtained as described in Suzuki *et al.*, (2008). Transgenic soybean (*Glycine max* cv. Thorne) plants constitutively expressing MBF1c under the control of the CaMV35S promoter were generated at the Iowa State Plant Transformation lab (Paz *et al.*, 2006). Six homozygous transgenic lines were tested by RNA blots for MBF1c expression and 3 independent homozygous lines were grown in a greenhouse under controlled growth conditions and assayed for growth and yield as described in Suzuki *et al.*, (2005).

DNA chip analysis

For microarray analysis, 11-day-old WT and *mbf1c* plants grown in peat pellets as described above were heat stressed at 40°C, 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$, and sampled at 0, 20, and 60 min (each time point had 3 technical repeats and the entire experiment had 3 biological repeats). RNA samples were used to perform chip hybridization analyses (*Arabidopsis* ATH1® chips; Affymetrix, Santa Clara, CA) at the Nevada Genomics Center in University of Nevada, Reno. Conditions for RNA isolation, labeling and hybridization are described in Davletova *et al.*, (2005a). All GeneChip® arrays were processed first by RMA (Robust Multi-Array

Average; Irizarry *et al.*, 2003) using the R package affy (Gautier *et al.*, 2004). Expression values were computed from raw CEL files by first applying the RMA model of probe-specific correction of PM (perfect match) probes. These corrected probe values were then normalized via quantile normalization, and a median polish was applied to compute one expression measure from all probe values. Resulting RMA expression values were log₂-transformed (Davletova *et al.*, 2005a). Statistical analysis (ANOVA) was performed as described in Davletova *et al.*, (2005a). Microarray data from this experiment was deposited in Array Express (E-MEXP-2760).

cDNA synthesis and qPCR analysis

cDNA synthesis, qPCR and RT-PCR data analysis were performed as described in Miller *et al.*, (2009). First strand cDNAs were produced from 1µg of total RNA that was used for the microarray analysis. Threshold cycle values for MBF1c (At3g24500), zinc finger protein (At3g21890), DREB2A (At5g05410), MYB7 (At2g16720) and HSP70 (At3g12580) were calculated with the *C_T* of EF1-α (At5g60390) as internal control. Primer pairs used for amplifications are shown in Table S2.

Systematic evolution of ligands by exponential enrichment (SELEX) assay

DNA-binding specificity of the MBF1c was determined by SELEX as described in Bouvet, (2001), Clouaire *et al.*, (2005) with minor modifications (Figure S1).

In Silico analysis

Promoters of the genes identified by the microarray analysis (Table 1) were assembled by ELEMENT (<http://element.cgrb.oregonstate.edu/>) as described in Nemhauser *et al.*, (2004), Koussevitzky *et al.*, (2007). Oligonucleotide sequences recovered by SELEX were analyzed by MEME (<http://meme.sdsc.edu/meme/website/meme.html>). Genes containing the identified consensus sequences in their 1000-bp promoter regions were identified by Patmatch (<http://www.arabidopsis.org/cgi-bin/patmatch/nph-patmatch.pl>). DNA element frequency analysis was performed with Motif Analysis from TAIR (<http://www.arabidopsis.org/tools/bulk/motiffinder/index.jsp>).

Yeast one-hybrid

Yeast one-hybrid assay was performed using Matchmaker™ Gold Yeast One-Hybrid Library Screening System (Clontech; Mountain View, CA) as described by the manufacturer instructions. Thirty-six bp oligonucleotides with 3 or 5 tandem repeats of consensus sequences or two mutated forms were cloned into pAbAi vector, carrying the AUR1-C gene. Plasmid was linearized and integrated into the genome of yeast (Y1H Gold). Yeast cells were then transformed with pGADT7-AD plasmid (Invitrogen; Carlsbad, CA) that contains MBF1c (Suzuki *et al.*, 2008). Interaction was determined based on the ability of transformed yeast to grow on -Leu media in the presence of 200 to 1000ng/ml of aureobasidin A.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Supported by funding from The National Science Foundation (IBN-0420033, NSF-0431327, IOS-0639964 and IOS-0743954), The Nevada Agricultural Experimental Station, NIH Grant Number P20 RR-016464 from the INBRE Program of the National Center for Research Resources, and EU grant FP7 - MARRIE CURIE 447.

References

- Bouvet, P. DNA-protein interactions. In: Moss, T., editor. *Methods in Molecular Biology*. Totowa: Humana Press; 2001. p. 603–610.
- Clouaire T, Roussigne M, Ecochard V, Mathe C, Amalric F, Girard JP. The THAP domain of THAP1 is a large C2CH module with zinc-dependent sequence-specific DNA-binding activity. *Proc Natl Acad Sci USA*. 2005; 102:6907–6912. [PubMed: 15863623]
- Davletova S, Rizhsky L, Liang H, Shengqiang Z, Oliver DJ, Couto J, Shulaev V, Schlauch K, Mittler R. Cytosolic ascorbate peroxidase 1 is a central component of the reactive oxygen gene network of Arabidopsis. *Plant Cell*. 2005a; 17:268–281. [PubMed: 15608336]
- Davletova S, Schlauch K, Couto J, Mittler R. The zinc-finger protein Zat12 plays a central role in reactive oxygen and abiotic stress signaling in Arabidopsis. *Plant Physiol*. 2005b; 139:847–856. [PubMed: 16183833]
- de Koning B, Blombach F, Wu H, Brouns SJ, van der Oost J. Role of multiprotein bridging factor 1 in archaea: bridging the domains? *Biochem Soc Trans*. 2009; 37:52–57. [PubMed: 19143601]
- Gautier L, Cope L, Bolstad B, Irizarry RA. affy: analysis of Affymetrix GeneChip data at the probe level. *Bioinformatics*. 2004; 20:307–315. [PubMed: 14960456]
- Horan K, Jang C, Bailey-Serres J, Mittler R, Shelton C, Harper JF, Zhu JK, Cushman JC, Gollery M, Girke T. Annotating genes of known and unknown function by large-scale coexpression analysis. *Plant Physiol*. 2008; 147:41–57. [PubMed: 18354039]
- Irizarry RA, Hobbs R, Collin R, Beazer-Barclay YD, Antonellis KJ, Scherf U, Speed TP. Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics*. 2003; 4:249–264. [PubMed: 12925520]
- Kabe Y, Goto M, Shima D, Imai T, Wada T, Morohashi K, Shirakawa M, Hirose S, Handa H. The role of human MBF1 as a transcriptional coactivator. *J Biol Chem*. 1999; 274:34196–34202. [PubMed: 10567391]
- Kotak S, Larkindale J, Lee U, von Koskull-Döring P, Vierling E, Scharf KD. Complexity of the heat stress response in plants. *Curr Opin Plant Biol*. 2007; 10:310–316. [PubMed: 17482504]
- Koussevitzky S, Nott A, Mockler TC, Hong F, Sachetto-Martins G, Surpin M, Lim J, Mittler R, Chory J. Signals from chloroplasts converge to regulate nuclear gene expression. *Science*. 2007; 316:715–719. [PubMed: 17395793]
- Liu QX, Jindra M, Ueda H, Hiromi Y, Hirose S. *Drosophila* MBF1 is a co-activator for tracheae defective and contributes to the formation of tracheal and nervous systems. *Development*. 2003; 130:719–728. [PubMed: 12506002]
- Luhua S, Ciftci-Yilmaz S, Harper JF, Cushman JC, Mittler R. Enhanced tolerance to oxidative stress in transgenic Arabidopsis plants expressing proteins of unknown function. *Plant Physiol*. 2008; 148:280–292. [PubMed: 18614705]
- Miller G, Schlauch K, Tam R, Cortes D, Torres MA, Shulaev V, Dangel JL, Mittler R. The plant NADPH oxidase RBOHD mediates rapid systemic signaling in response to diverse stimuli. *Sci Signal*. 2009; 18:ra45. [PubMed: 19690331]
- Mittler R. Abiotic stress, the field environment and stress combination. *Trends Plant Sci*. 2006; 11:15–19. [PubMed: 16359910]
- Mittler R, Blumwald E. Genetic engineering for modern agriculture: Challenges and perspectives. *Ann Rev Plant Biol*. 2010; 61:443–462. [PubMed: 20192746]
- Nemhauser JL, Mockler TC, Chory J. Interdependency of brassinosteroid and auxin signaling in Arabidopsis. *PLOS Biol*. 2004; 2:e258. [PubMed: 15328536]

- Paz MM, Martinez JC, Kalvig AB, Fonger TM, Wang K. Improved cotyledonary node method using an alternative explant derived from mature seed for efficient *Agrobacterium*-mediated soybean transformation. *Plant Cell Rep.* 2006; 25:206–213. [PubMed: 16249869]
- Rizhsky L, Davletova S, Liang H, Mittler R. The zinc finger protein Zat12 is required for cytosolic ascorbate peroxidase 1 expression during oxidative stress in *Arabidopsis*. *J Biol Chem.* 2004; 279:11736–11743. [PubMed: 14722088]
- Schramm F, Larkindale J, Kiehlmann E, Ganguli A, Englich G, Vierling E, von Koskull-Döring P. A cascade of transcription factor DREB2A and heat stress transcription factor HsfA3 regulates the heat stress response of *Arabidopsis*. *Plant J.* 2007; 53:264–274. [PubMed: 17999647]
- Shamovsky I, Nudler E. New insights into the mechanism of heat shock response activation. *Cell Mol Life Sci.* 2008; 65:855–861. [PubMed: 18239856]
- Suzuki N, Bajad S, Shuman J, Shulaev V, Mittler R. The transcriptional co-activator MBF1c is a key regulator of thermotolerance in *Arabidopsis thaliana*. *J Biol Chem.* 2008; 283:9269–9275. [PubMed: 18201973]
- Suzuki N, Rizhsky L, Liang H, Shuman J, Shulaev V, Mittler R. Enhanced tolerance to environmental stresses in transgenic plants expressing the transcriptional co-activator MBF1. *Plant Physiol.* 2005; 139:1313–1322. [PubMed: 16244138]
- Takemaru K, Harashima S, Ueda H, Hirose S. Yeast coactivator MBF1 mediates GCN4-dependent transcriptional activation. *Mol Cell Biol.* 1998; 18:4971–4976. [PubMed: 9710580]
- Tsuda K, Tsuji T, Hirose S, Yamazaki K. Three *Arabidopsis* MBF1 homologs with distinct expression profiles play roles as transcriptional co-activators. *Plant Cell Physiol.* 2004; 45:225–231. [PubMed: 14988493]
- Tsuda K, Yamazaki K. Structure and expression analysis of three subtypes of *Arabidopsis* MBF1 genes. *Biochim Biophys Acta.* 2004; 1680:1–10. [PubMed: 15451167]
- von Koskull-Döring P, Scharf KD, Nover L. The diversity of plant heat stress transcription factors. *Trends Plant Sci.* 2007; 12:452–457. [PubMed: 17826296]

Figure 1. Expression of heat-response transcripts in wild type and *mbf1c* plants

The expression of *MBF1c*, 3 transcripts identified by our microarray analysis as dependent on MBF1c for expression during heat (*DREB2A*, *MYB7*, and *At3g21890*), and one MBF1c-independent transcript (*HSP70*) was determined using qPCR in wild type and *mbf1c* plants subjected to heat stress (40°C) for 0, 20 and 60 min. Results are average and SD of 3 independent experiments, each performed with 5 technical repeats.

Figure 2. Heat stress survival of Arabidopsis seedlings deficient in different genes encoding MBF1c-dependent transcripts

Survival assays were performed with two independent knockout mutants for the zinc finger protein At3g21890 (*z-finger*), *Dreb2A*, and *Sen1*. Results are average and SD of 3 independent experiments, each performed with 5 technical repeats.

(a)

CTAGA			
	Word		Count
20min HS	-----T	CTAGA A-----	4
	-----TT	CTAGA A-----	4
60min HS	-----	CTAGA -----	91
	-----T	CTAGA -----	48
	-----	CTAGA A-----	37
	-----T	CTAGA A-----	22
	-----	CTAGA AG----	17
	----TT	CTAGA A-----	17
	---CTT	CTAGA -----	12
	---AAT	CTAGA -----	8
	-----	CTAGA AGC---	7
	-----	CTAGA TAT---	6

(b)

Figure 3. Identification of a putative DNA-binding element for MBF1c
(a) Alignment of sequences from the 1000 bp promoter region of genes encoding MBF1c-dependent transcripts (Table 1) identifying the CTAGA consensus.
(b) Yeast one-hybrid assay showing specific CTAGA transcriptional activation by MBF1c.

Figure 4. Dominant-negative function of the DNA-binding domain of MBF1c

(a) The different DNA constructs used for ectopic expression of the DNA-binding (HTH), or N-terminal (N-term) domains of MBF1c in transgenic plants.

(b) *Left panel:* Heat stress survival of Arabidopsis seedlings expressing the DNA-binding (HTH) GFP fusion (HTH 1 and 2 are two independent homozygous transgenic lines). *Right panel:* Localization of the HTH-GFP fusion in root cells of seedlings subjected to heat stress (40°C) for 1 hour.

(c) Same as B, but for the N-terminal of MBF1c fused to GFP (Nterm). Results are average and SD of 3 independent experiments, each performed with 5 technical repeats.

Figure 5. Ectopic expression of MBF1c in soybean improves yield in plants grown under controlled growth conditions

(a) RNA gel blot analysis showing expression of MBF1c in three independent transgenic lines of soybean.

(b) Percent germination, plant height and weight of seeds per plant of three independent homozygous lines ectopically expressing MBF1c grown in a greenhouse to maturity at 25–28°C. No differences in individual seed weight were found between the different plants. Results are average and SD of 3 independent experiments, each performed with 12 technical repeats arranged in a mixed plot setting.

Table 1
MBF1c-dependent heat stress-response transcripts in Arabidopsis

Transcripts significantly less elevated in *mbf1c* compared to wild type during heat stress (20 and 60 min; 40°C) are shown. Locus identifier, gene name, and response to drought, heat, salt, or cold as found in microarray databases (Horan *et al.*, 2008) is indicated for each transcript (red-elevated, blue-suppressed, white-unchanged). The frequency of putative MBF1c DNA-binding sites is indicated in the 1000 bp promoter region of each gene (Left two binding elements are from bioinformatics analysis of microarray data; right two binding elements are from a SELEX assay). Fold change (log2) is also shown for each gene in wild type and *mbf1c* plants at 60 min heat stress.

Locus	Gene Name	Response to Abiotic Stress				Putative DNA Element				Fold change (log2)	
		Heat	Drought	Salt	Cold	CTAGA	CTTTA	GGAGG	ACGGAA	WT	<i>mbf1c</i>
20 min											
AT3G24500	MBF1C	<div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div><div></div></div>									

Locus	Gene Name	Response to Abiotic Stress				Putative DNA Element				Fold change (log2)	
		Heat	Drought	Salt	Cold	CTAGA	CTTTA	GGAGG	ACGGAA	WT	<i>mbf1c</i>
AT5G03380	Heavy-metal-associated protein	Heat				2	1	0	0	1.454	0.725
AT5G35320	Hypothetical protein	Heat				2	0	1	0	5.639	4.7
AT1G78080	RAP2.4 (related to AP2.4)	Heat				0	1	0	1	1.422	0.866
AT1G62760	Invertase/pectin methylesterase inhibitor	Heat				0	2	0	0	3.214	2.201
AT4G11660	Heat shock transcription factor B2B	Heat				1	3	0	0	3.7	3.084
AT5G62020	Heat shock transcription factor B2A	Heat				0	1	0	0	4.175	3.275
AT1G55620	CLC-F voltage-gated chloride channel	Heat				0	2	0	0	0.735	-0.131
AT1G59980	ARL2/ATDJC39 heat shock protein	Heat				1	2	0	0	1.834	1.248
AT5G47610	Zinc finger (C3HC4-type RING)	Heat				2	1	3	0	4.638	3.36
AT3G12320	Unknown protein	Heat				2	3	1	0	1.637	1.551
AT1G13600	ATBZIP58 (Basic Leucine Zipper)	Heat				1	1	1	0	1.221	0.447
AT2G34590	Transketolase	Heat				0	2	0	0	0.732	0.308
AT1G26800	Zinc finger (C3HC4-type RING)	Heat				3	0	0	0	4.557	4.054
AT5G37710	Lipase/calmodulin-binding heat-shock	Heat				0	5	1	1	1.482	0.868
AT2G44130	Kelch repeat F-box protein	Heat				3	2	0	0	0.993	-1.14
AT1G56610	Syntaxin	Heat				0	2	0	0	0.621	0.126
AT5G66090	Adenine nucleotide alpha hydrolases-like	Heat				0	0	1	0	1.587	1.195
AT2G27580	Zinc finger (AN1-like)	Heat				0	1	1	0	3.559	2.886
AT5G52570	Beta-carotene hydroxylase 2	Heat				2	5	0	0	2.256	0.893
AT5G06280	Hypothetical protein	Heat				2	2	0	0	1.477	1.013
AT5G25280	Serine-rich protein	Heat				0	3	0	0	1.285	0.584

The frequency of DNA element appearance in the 500 or 1000 bp region of promoters from all heat response genes (60 min heat stress) from this study is compared to that of all Arabidopsis genes.

Table 2
Frequency of DNA element appearance in the promoter region of Arabidopsis genes

In 60 min heat responsive genes (1039 genes)			In all the genes (33518 genes)		
<u>500bp</u>					
Sequence	Number of genes	Ratio (%)	Number of genes	Ratio (%)	p-value
CTAGA	224	21.56	3989	11.90	5.96E-19
CTTTA	270	25.99	7859	23.45	4.56E-03
GGAGG	65	6.26	1870	5.58	3.30E-02
ACGGAA	148	14.23	3980	11.87	2.52E-03

<u>1000bp</u>					
Sequence	Number of genes	Ratio (%)	Number of genes	Ratio (%)	p-value
CTAGA	287	27.68	7506	22.39	1.01E-05
CTTTA	417	40.21	12807	38.21	1.05E-02
GGAGG	120	11.57	3980	11.87	3.70E-02
ACGGAA	264	25.46	7498	22.37	1.80E-03