Published in final edited form as:

Cancer Immunol Immunother. 2015 March ; 64(3): 367–379. doi:10.1007/s00262-014-1640-x.

A Phase I Dose Escalation Trial of MAGE-A3 and HPV16 Specific Peptide Immunomodulatory Vaccines in Patients with Recurrent / Metastatic (RM) Squamous Cell Carcinoma of the Head and Neck (SCCHN)

Dan P Zandberg^{1,2}, Sandra Rollins³, Olga Goloubeva², Robert E Morales⁴, Ming Tan⁵, Rodney Taylor⁶, Jeffrey S Wolf⁶, Lisa M. Schumaker², Kevin J Cullen^{1,2}, Ann Zimrin^{1,2}, Robert Ord^{2,7}, Joshua E Lubek^{2,7}, Mohan Suntharalingam⁸, John C. Papadimitriou³, Dean Mann³, Scott E Strome^{*,6}, and Martin J. Edelman^{*,1,2}

¹Department of Medicine, Division of Medical Oncology, University of Maryland School of Medicine, Baltimore, MD, USA

²University of Maryland Marlene and Stewart Greenebaum Cancer Center, Baltimore, MD, USA

³Department of Pathology, University of Maryland School of Medicine, Baltimore, MD, USA

⁴Department of Radiology, University of Maryland School of Medicine, Baltimore, MD, USA

⁵Department of Biostatistics, Bioinformatics and Biomathematics, Georgetown Lombardi Comprehensive Cancer Center, Washington DC, USA

⁶Department of Otorhinolaryngology-Head and Neck Surgery, University of Maryland, Baltimore, MD, USA

⁷Department of Oral and Maxillofacial Surgery, University of Maryland School of Medicine, Baltimore, MD, USA

⁸Department of Radiation Oncology, University of Maryland School of Medicine, Baltimore MD, USA

Abstract

Background—We conducted a phase I dose escalation study to evaluate the safety and immunologic response to peptide immunomodulatory vaccines GL-0810 (HPV16) and GL-0817 (MAGE-A3) in HPV16 and MAGE-A3 positive RM-SCCHN patients, respectively.

Methods—Three dose levels (500 µg, 1000 µg, 1500 µg) of GL-0810 or GL-0817 with adjuvants Montanide (1.2ml) and GM-CSF (100µg/m2) were administered subcutaneously q2 weeks for a total of 4 vaccinations in HPV16 and MAGE-A3 positive RM-SCCHN patients, respectively.

Corresponding Author: Dan P Zandberg MD, Assistant Professor of Medicine University of Maryland Greenebaum Cancer Center, 22 S. Greene Street Rm N9E29, Baltimore, MD 21201, USA, Phone: 410-328-6373, Fax: 410-328-6896.

*Dr. Edelman and Dr. Strome share senior authorship.

Disclosures: Dr. Strome is a co-founder and stockholder of Gliknik Inc., a biotechnology company and licensee of the Trojan Peptide Vaccines. Dr. Strome also receives royalties from the Mayo Clinic College of Medicine for intellectual property related to B7-H1 (PD-L1). All other authors have no disclosures.

Results—Nine and 7 patients were enrolled in the HPV16 and MAGE-A3 cohorts, respectively. No dose limiting toxicities were observed and toxicity was predominantly local and grade 1 (erythema, pain, and itching at the injection site). In those patients that received all 4 vaccinations, 80% (4/5) of the HPV16 cohort, and 67% (4/6) of the MAGE-A3 cohort developed antigen specific T cell and antibody responses to the vaccine. Significant concordance between T cell and antibody responses was observed for both groups. No clear dose-response correlation was seen. All patients progressed by RECIST at first repeat imaging, except for one patient in the MAGE-A3 500µg cohort who had stable disease for 10.5 months. The median PFS and OS for the MAGE-A3 cohorts were 79 and 183 days respectively and for the HPV16 cohort 80 and 196 days respectively.

Conclusions—GL-0810 and GL-0817 were well tolerated in patients with RM-SCCHN with T cell and antibody responses observed in the majority of patients who received all 4 vaccinations.

Keywords

SCCHN; peptide; vaccine; immune response

Introduction

Squamous cell carcinoma of the head and neck (SCCHN) is the 6th most common malignancy worldwide.[1] Despite aggressive initial definitive therapy, select groups of patients with SCCHN are at high risk for the development of second primary tumors and a significant proportion of patients will recur, commonly within the first 2 years.[2] Unfortunately, a significant proportion of these patients will not have a salvage surgical or radiation option, and the median survival for locoregionally recurrent or metastatic disease treated with palliative chemotherapy alone is a dismal 8-10 months.[3] New therapeutic modalities are desperately needed for this patient population.

In an effort to improve outcomes for recurrent/metastatic SCCHN patients, we developed two separate peptide immunomodulatory vaccines using HLA-I and HLA-II T cell epitopes of the human papillomavirus type 16 (HPV16) and melanoma antigen E (MAGE)-A3 tumor associated antigens. The human papillomavirus (HPV) now has an established role in the etiology of squamous cell carcinoma of the oropharynx.[4] HPV16 was chosen because it accounts for the majority of cases (68 to 87%) of HPV positive head and neck cancer worldwide.[5], [6] Additionally, tumor specificity and the presence of well-defined peptide epitopes make it an ideal target.[7-9]

Melanoma antigen E (MAGE-A3) is a cancer-testis antigen. Cancer-testis antigens are expressed by human tumors of different histological types and have limited expression in normal parenchyma.[10] For example, MAGE-A3 was originally discovered in melanomas and has subsequently been identified in various epithelial cell malignancies including head and neck cancer where it has been found in upwards of 51% of cases.[10, 11] Importantly, MAGE-A3 expression may be associated with a more aggressive course, as several studies suggest that MAGE-A3 may be a negative prognostic indicator in hematologic malignancies.[12, 13] Additionally, expression has been associated with more advanced stage in squamous cell carcinoma of the larynx.[11] Therefore, MAGE-A3 was chosen

based on its tumor specificity, high percent expression, and the existence of previously defined HLA-I and HLA-II immunostimulatory epitopes.[14, 15]

In order to improve the efficacy of these vaccines, we incorporated several unique features into their design. First, we included a “penetrin” sequence derived from HIV-TAT (RKKRRQRRR) which facilitates peptide translocation through the cell membrane directly to the endoplasmic reticulum and golgi apparatus to form peptide-HLA complexes.[16] Additionally, the junction of multiple HLA-I and HLA-II peptide-epitopes via furin-cleavable linkers, allows the release of individual epitopes when cleaved in the Golgi.[17, 18] Furthermore, the inclusion of both HLA-I and HLA-II epitopes facilitates both cytotoxic T cell activation and stimulation of the CD4 T helper cell response, important for development of tumor-specific memory T cell and humoral immunity.[19-22] Finally, the combination of Montanide ISA 51 and granulocyte macrophage colony stimulating factor (GM-CSF) is postulated to enhance antigen presentation and promote dendritic cell migration to the site of vaccination.[23]

Based on promising pre clinical data with these immunomodulatory peptide vaccines[17] we initiated a pilot study in 5 patients with recurrent or metastatic SCCHN (RM-SCCHN). This pilot study demonstrated that these immunomodulatory peptide vaccines in combination with GM-CSF and montanide adjuvant were well tolerated and able to stimulate antigen-specific cellular and humoral immunity. Although there were no documented responses, one patient was still alive 31 months post therapy.[18] In order to evaluate the safety of these vaccines in a larger patient cohort and investigate how peptide dose correlates with development of antigen specific immunity, we performed the current phase I dose escalation trial. Our data from this study reaffirm the feasibility and safety of this vaccine based approach and document a striking correlation between the development of antigen specific T cell and humoral immunity which will facilitate monitoring of subsequent pivotal studies. Furthermore, the lack of clinical efficacy in the face of a clear immunological response suggests that these vaccines may be most clinically relevant in resected SCCHN patients at high risk for disease recurrence and/or in combination with manipulation of specific co signaling pathways.

Materials and Methods

We conducted a single center phase I dose escalation study at the University of Maryland Greenebaum Cancer Center (UMGCC), to evaluate the effect and safety of three dose levels (500µg, 1000µg, and 1500µg) of HPV16 (GL-0810) or MAGE-A3 (GL-0817) immunomodulatory peptide vaccine in combination with adjuvant montanide and GM-CSF in HPV16 or MAGE-A3 expressing RM-SCCHN patients respectively. The primary objectives of the study were to establish safety and the adequate biologic dose for this vaccine defined by immune response (see below).

Patient Selection

Eligibility criteria included age >18 yo, PS 0-2, biopsy proven progressive, recurrent (post-surgical, radiation therapy, chemotherapy, combination therapy) or metastatic SCCHN which by the judgment of the treating physician was incurable by standard treatment

modalities or the patient was unwilling to be treated with surgery, chemotherapy or radiation therapy. A patient's tumor must have tested positive for either MAGE-A3 or HPV16. Main exclusion criteria included known HIV infection, patients with any malignant disease to the CNS or in a location that would put the patient at significant risk should an inflammatory response occur, patients undergoing treatment with systemic immunosuppressants and/or had an immunocompromising condition, or other active cancer requiring therapy or history of malignancy that was expected to effect life expectancy. Patients could not be treated with other cancer therapy concurrent with vaccine and could not have received chemotherapy, immunotherapy, biologic therapy, or radiation therapy within 4 weeks of the date of first vaccination.

Importantly, an HLA-A*02 genotype was not included as an entry criteria for this trial. The decision to eliminate HLA typing from the inclusion criteria was based on the fact that observed immunologic responses in the pilot study were predominantly directed against the HLA II vaccine epitopes. Based on the knowledge that class II molecules have an open binding groove and, as such, have less stringent requirements for recognition (add reference), we anticipated that patients bearing diverse HLA II genotypes could respond to the class II vaccine epitopes.[18]

Study Design

Patients meeting initial inclusion/exclusion criteria underwent a biopsy of accessible tumor for determination of HPV16 and MAGE-A3 status. Patients with HPV16 positive tumors were assigned to cohort 1 and were treated with HPV16 immunomodulatory peptide vaccine GL-0810 plus adjuvant and those with MAGE-A3 positive tumors were assigned to cohort 2 and treated with MAGE-A3 immunomodulatory peptide vaccine GL-0817 plus adjuvant. If a patient's tumor was both MAGE-A3 and HPV16 positive they were enrolled in the cohort with the fewest patients.

Both cohorts 1 and 2 ran independently of each other. Three dose levels (500 µg, 1000 µg, 1500 µg) of HPV16 or MAGE-A3 immunomodulatory vaccines were tested in sequential groups of patients. Each peptide was admixed with fixed doses of the adjuvants Montanide (1.2ml) and GM-CSF (100µg/m2) and given together as one injection. A total of 4 vaccinations were administered subcutaneously in the inguinal region at 2-week intervals. Dose escalation occurred in each cohort once three subjects from the preceding dose level received at least two of the planned four vaccinations and no patient experienced a dose limiting toxicity (DLT). Doses were not escalated in any individual patient. If two or more patients experienced a DLT at a given dose level, accrual at that dose would be halted and the preceding dose level would be expanded to six patients. Data was to be analyzed after at least 3 vaccinated patients had received a dose that was tolerated. The dose that provided an adequate immunologic response was to be expanded by 9 patients for a total of 12 patients at that particular dose. Therefore, the maximum number of subjects that could be enrolled in each of the two cohorts was 24 (6+6+12).

An adequate biological dose was to be defined by a five-fold increase in interferon (IFN)-γ expression level in the ELISPOT assay after the first 2 weeks of therapy. If an adequate

biological response was not generated in a majority of patients at any dosage, than the dose at which the greatest increase in IFN- γ would be expanded.

Screening of patients took place within 30 days prior and baseline evaluations within 7 days prior to the first vaccination. During the screening phase, patients underwent PET/CT for baseline radiographic evaluation (up to 6 weeks prior to first vaccination) as well as tumor biopsy to determine MAGE-A3 and HPV16 expression. After each vaccination the patient was observed for 4 hours with vital signs and evaluation of the injection site pre dose, immediately post injection, 0.5, 1, 2 and 4 hours post dosage. Subjects were evaluated at each vaccination visit, 2 weeks and 6 weeks after the fourth vaccination, and thereafter every 3 months through month 24. At these visits history and physical examination, adverse event recording, ECOG performance status, physical tumor measurements (if applicable) and blood draws for immune response evaluations were taken. A repeat PET/CT was done 6 weeks (+/- 7 days) after the fourth vaccination (3 months after first vaccination).

Patients who, prior to completing the allotted number of peptide vaccinations, had disease progression, intolerable toxicity, or refused further treatment, had treatment discontinued and were followed until the end of the study.

Adverse events monitoring

Toxicity was defined as per the NCI Common Toxicity criteria for adverse events (CTCAE) version 3.0. Dose-limiting toxicity criteria are shown in supplemental table 1.

Discontinuation of an individual patient from the study was based on adverse events deemed related to the study medication as detailed in supplemental table 2.

Disease monitoring

A maximum of 3 measurable lesions were identified as target lesions in each patient and were measured and recorded at baseline by physical examination and/or intravenous contrast enhanced PET/CT. Definition of complete response (CR), partial response (PR), stable disease (SD), or progressive disease (PD) was based on response evaluation criteria in solid tumors (RECIST 1.1).

Laboratory Methods

Descriptions of MAGE-A3 and HPV16 peptides, isolation of peripheral blood mononuclear cells, MAGE-A3 and HPV16 analysis by PCR, Trojan-specific T-cell response monitoring, Interferon- γ restimulation elispot assay, MAGE-Trojan and HPV16-Trojan ELISA used in this study are found in our previously published pilot study.[18] Importantly, all Elispot assay results reported in this study included in vitro stimulation, to enhance response detection.

Statistical Analysis

This phase I study was designed to evaluate the safety of the vaccines and establish an adequate biological dose of the HPV16 and MAGE-A3 peptide immunomodulatory vaccines. Based on the results of the pilot study,[18] we did not anticipate serious toxicity and therefore a true MTD was not anticipated, but an adequate biologic dose was sought.

Descriptive statistics were utilized to describe the incidence of adverse effects as defined by CTCAE version 3.0.

A biological (immunological) response was defined as a binary (positive versus negative) response measured in each patient after treatment with either HPV16 or MAGE-A3 immunomodulatory peptide vaccine. Positive response was defined as an at least fivefold increase in interferon (IFN)- γ expression level in the ELISPOT assay after the first 2 weeks of therapy. An adequate dose was defined as a dose that provided a high response rate. The goal of the design was to find a biologically 'adequate' dose while using a very small number of patients, because optimal biological dose may not exist or require a large number of patients.

Additional endpoints included progression free survival (PFS), response and overall survival (OS). PFS and OS functions were estimated using the Kaplan-Meier approach. PFS time was defined as the time from first treatment to documentation of progression or death. If a patient died without a progression documented, the patient would be considered to have progressed on their death date. OS time was defined as the time from first treatment until death or censored at the last date of patient contact. Response evaluations including CR, PR, SD, and PD were defined as detailed above.

The exact McNemar test at the 0.05 level of significance was used to assess whether there was a concordance between T cell and antibody responses for patients treated with the HPV16 and MAGE-A3 vaccines. Statistical analysis was conducted using R (v.3.0.2).

Results

Patient Demographics

Between 2008 and 2012, 56 patients were screened for enrollment on the clinical trial. Forty-four patients underwent biopsy for evaluation and 9 patients tested positive for MAGE-A3, 12 for HPV16, and 2 were positive for both HPV16 and MAGE-A3. A total of 17 patients were enrolled on the protocol. One patient was subsequently discovered to be ineligible and was taken off of the study. Therefore, there were 16 evaluable patients including 9 patients in the HPV16 cohort and 7 patients in the MAGE-A3 cohort (Figure 1). The study was closed prematurely on 7/11/13 due to poor accrual. All patients enrolled had completed treatment by this time and one patient remained in survival follow up.

Baseline patient characteristics are shown for the HPV16 and MAGE-A3 positive cohorts in tables 1A and 1B, respectively. The median age of the entire study population was 60, and all patients were male and predominantly white. In the HPV16 cohort, 89% of patients had an oropharyngeal primary, whereas in the MAGE-A3 cohort, the primary site varied between oropharynx, hypopharynx, and larynx. The HPV16 cohort contained younger patients with disease that was predominantly metastatic whereas MAGE-A3 positive patients had predominantly locoregional recurrence. HPV16 positive patients were more heavily pretreated with palliative chemotherapy compared to the MAGE-A3 positive patients.

Feasibility and Toxicity

In the HPV16 cohort, 3 patients were vaccinated at each dose level (500µg, 1000µg, and 1500µg). In the MAGE-A3 cohort, 3 patients were vaccinated at the 500µg and 1000µg dose levels while only one patient was treated at the 1500µg dose level. Eleven out of the 16 treated patients received all four vaccinations. In the HPV16 cohort, 5 out of 9 patients completed all 4 vaccinations and in the MAGE-A3 cohort, 6 out of 7 patients received all 4 vaccines. Five patients progressed during vaccine therapy and therefore did not receive all four vaccinations.

Toxicity data is shown in tables 2A and 2B for HPV16 and MAGE-A3 cohorts, respectively. Overall, the vaccine was well tolerated with predominantly local grade 1 toxicity including erythema, pain, and itching at the injection site. The patient receiving the MAGE-A3 vaccine at the highest dosage did experience grade 1 systemic symptoms including headache, nausea and chills. One patient receiving the MAGE-A3 vaccine at 500 µg, developed a pharyngocutaneous fistula 7 days after his first vaccination. This was treated conservatively and he received all remaining doses. No patients discontinued treatment early because of toxicity.

Immunologic Outcomes

Interferon gamma (IFN γ) Elispot and ELISA assays were used to evaluate post vaccination peptide vaccine-specific T cell and antibody responses, respectively. Elispot and ELISA results for individual responding patients at the three vaccine dosages in the HPV16 cohort and MAGE-A3 cohort are shown in Figures 2 and 3, respectively and a summary of immunologic responses is shown in Table 3. In the HPV16 cohort, all patients that had a T cell response also had an antibody response, with 44% (4/9) of patients having both a T cell and antibody response to the Trojan peptide vaccine. One patient (043) had a T cell response and antibody to HPV16 at baseline, which increased after the fourth vaccination. One patient (003) had an antibody response but not a T cell response and four patients had no detectable immunologic response to the vaccine. Amongst patients that received all four vaccinations, 80% (4/5) had both T cell and antibody responses.

In the MAGE-A3 cohort, all patients that had a T cell response also had an antibody response, with 57% (4/7) of patients having both a T cell and antibody response to the peptide immunomodulatory vaccine. One patient (004) did not have a T cell response but had antibody to MAGE-A3 at baseline. Two patients had no response to the peptide antigens by either Elispot or ELISA. Amongst patients that received all four vaccinations, 67% (4/6) of patients had both a T cell and antibody response.

For both groups, there was significant concordance between T cell and antibody responses ($p < 0.001$), with T cell responses preceding antibody detection. Antibody titers were sustained in responding patients at the last time point of evaluation, with some patients having rising titers even at 3 months after the first vaccination. Higher dosages of vaccine did not appear to increase response rate but may lead to a more rapid T cell response. Due to the lower accrual, we were unable to determine the adequate biologic dosage based on our pre planned analysis parameters, however given that all dosages were well tolerated, we

anticipate that the highest dosage of 1500µg will be selected for further study in both cohorts.

Clinical Outcomes

There were no complete or partial responses by RECIST. All patients progressed at first repeat imaging except for one patient in the MAGE 500µg cohort (patient 017) who had stable disease. This patient withdrew consent but did continue to follow up at UMGCC and his disease was stable for 10.5 months off any therapy before progressing. The median PFS and OS for the MAGE-A3 cohort were 79 and 183 days respectively and for the HPV16 cohort 80 days and 196 days respectively.

Discussion

This novel phase I dose escalation trial, evaluated the safety and immune response at three different dose levels of MAGE-A3 (GL-0817) or HPV16 (GL-0810) immunomodulatory peptide vaccines with adjuvants montanide and GM-CSF in patients with locally recurrent or metastatic MAGE-A3 or HPV16 positive SCCHN, respectively. The vaccines were well tolerated and no dose limiting toxicity was reached. In patients who received all four vaccinations, 80% of HPV16 positive and 67% of MAGE-A3 positive patients developed both a T cell and antibody response. In both groups there was a significant correlation between T cell response and subsequent antibody response. However, these immune responses did not translate into any clinical responses as measured by RECIST.

Consistent with our findings from our pilot study [18], the preponderance of T cell responses were directed against the whole Trojan vaccine or the class II epitopes. The increased recognition of the whole Trojan vaccine may reflect an intrinsic bias in our assay design, necessitated by limited sample availability, in which PBMC were initially stimulated by the whole Trojan vaccine in vitro, with secondary stimulation using the intrinsic Class I and II peptide epitopes. Alternatively, it is possible that the increased responses to the whole vaccine are a result of “neo epitopes” generated by artificial linking of multiple epitopes. Finally, since this trial was conducted in patients with existing disease, if peptide T cells trafficked to the tumor site, their actual numbers could be artificially reduced in the peripheral blood. Unfortunately, we were unable to distinguish these possibilities in our study secondary to limited sample availability.

There was a significant correlation between T cell and antibody responses. As anticipated, in the majority of responding patients, the T cell response preceded the antibody response. While some T cell responses began to wane, most antibody responses continued to increase after the last vaccination. Higher doses in both the MAGE-A3 and HPV16 cohorts appeared to generate earlier but not more robust responses. Although lower than expected accrual prevented statistical analysis to determine an adequate biologic dosage, given that all dosages were well tolerated and higher dosages may be associated with an earlier response, we anticipate that the 1500 µg dosage will be used in subsequent trials.

Interestingly, 2 patients (1 in HPV16 cohort, and 1 in MAGE-A3 cohort) had baseline antibodies against their respective peptide vaccine. While antibodies to HPV16 E6 have

been reported in up to 35% of patients with squamous cell carcinoma of the oropharynx, even upwards of ten years before a diagnosis of cancer[24], baseline antibodies to MAGE-A3 are rarely found.[21, 25]

Despite a T cell and antibody response in the majority of patients that completed all four vaccinations, we observed no clinical responses in either cohort. One patient in the MAGE-A3 500µg cohort did have stable disease for 10.5 months. While appropriate endpoints for trials continue to be debated, despite the efficacy and recent approval of Sipuleucel-T for the treatment of metastatic castrate resistant prostate cancer, a pooled analysis of therapeutic vaccine trials published since 2004, involving multiple tumor types with a total of 973 patients, found an overall response rate of 3.6%, and a response rate of 3.7% in trials that included only peptide vaccines.[26] Similar to our findings, analysis of published therapeutic cancer vaccine trials have shown that evidence of activation of the immune system by laboratory correlates has not translated into clinical responses.[26]

To our knowledge this trial as well as our previously published pilot study [18], represent the first published experience using therapeutic immunomodulatory peptide vaccines to treat recurrent or metastatic SCCHN. However, MAGE-A3 and HPV16 peptide vaccinations have been evaluated in the treatment of other cancers. For example, vaccination with MAGE-A3 peptide vaccines can stimulate cellular and humoral immune responses in both melanoma and NSCLC. [27, 28] While in NSCLC the use of this vaccine has only been tested in the adjuvant setting[28], data evaluating the clinical efficacy of a MAGE-A3 peptide vaccine against active macroscopic disease exists in metastatic melanoma, demonstrating some activity in cutaneous or lymph node metastasis but not visceral metastasis.[25, 29, 30] For example, Marchand et al analyzed clinical response in 33 patients with stage III or IV metastatic melanoma treated with recombinant MAGE-A3 protein with immunological adjuvant SBAS-2. There were 2 partial responses and 2 mixed responses; however there were no clinical responses in any patients that had visceral metastasis.[29] Therapeutic vaccines targeted against HPV have shown some clinical efficacy for pre cancerous lesions.[31]

Potential reasons for lack of clinical efficacy in the face of an active immune response must be understood in order to effectively inform subsequent studies. Two of the major impediments to clinical efficacy are immunologic tolerance and patient selection. The causes of immunologic tolerance in SCCHN are the result of both tumor expression of molecules which dampen the immune response and active cellular suppression. For example, SCCHN cells can secrete immunosuppressive cytokines such as TGF- β , IL-10, and prostaglandin E2 (PGE2).[32, 33] Additionally, tumor expression of B7-H1 (PD-L1) which inhibits the activity of PD-1 expressing CD4 and CD8 T cells, has been shown to be expressed in 40-70% of SCCHN patients with higher expression in HPV positive patients. [34-36] Furthermore, in HPV positive SCCHN, early viral oncogenes E5, E6 and E7 can down regulate antigen processing and presentation.[37, 38] E5 and E7 can decrease cell surface expression of HLA-I and E7 can down regulate expression of transporter associated with antigen processing (TAP).[37, 38]. Importantly, direct down regulation of MAGE-A3 expression may prove unlikely, as in melanoma patients treated in the therapeutic trials

discussed above, MAGE-A3 expression was still present in the combined 25 patients that underwent subsequent resection of cutaneous metastasis.[23, 29]

Vaccine induced anti-tumor immunity also competes with and may actually augment regulatory immune responses. For example, regulatory T cells (Tregs), which suppress effector T cells and mediate peripheral tolerance are increased in the blood and tumor microenvironment in SCCHN patients.[39, 40] Myeloid derived suppressor cells (MDSC), which also have been found to be increased in the SCCHN tumor microenvironment, are known to suppress the immune response through secretion of IL-10, increasing Tregs and decreasing T cell activation and migration.[32, 41] Our understanding of the role that each of these elements plays, alone and collectively in the regulation of antitumor vaccine based immune responses, is hindered by the lack of post treatment tissue analysis, as findings in peripheral blood may not correlate with responses in the tumor microenvironment.

The second factor that likely contributes to the lack of observed efficacy was the patient cohort selected for study. The majority of the studied patient population had progressed through multiple lines of therapy, and represents a population where advancement of disease is typically rapid. For example progression free survival with second line chemotherapy in recurrent/metastatic SCCHN is 1-3 months.[42, 43] In this heavily pre-treated population, the immune response to vaccination did not develop until weeks and in some cases months after the start of vaccine therapy – mitigating its potential efficacy in the face of actively progressing disease. For example, 5 patients (4 in HPV 16 and 1 in MAGE-A3 cohort) progressed before they could even receive all four vaccinations.

The potential means to surmount these hurdles include combining the peptide based vaccine with other immunotherapy agents known to inhibit death of anti-tumor effector T cells e.g. anti-B7-H1 or to mitigate the effects of cellular suppression e.g. Cytosine.[44, 45] In addition, the delay between vaccination and development of the immune response, suggests that the efficacy/utility of this regimen may best be determined in the adjuvant setting.

In conclusion, peptide immunomodulatory vaccines against HPV16 (GL-0810) and MAGE-A3 (GL-0817) in recurrent/metastatic SCCHN are well tolerated and stimulated potentially meaningful T cell and antibody responses in the majority of patients who completed all four vaccinations. In responding patients, T cell responses significantly correlated with antibody responses in both the HPV16 and MAGE-A3 cohorts – a finding with important biologic implications, which may also simplify monitoring of subsequent studies. While clinical efficacy was lacking in this refractory patient population, protocol development is underway to evaluate the efficacy of the MAGE-A3 immunomodulatory peptide vaccine in the adjuvant setting after definitive therapy in high-risk squamous cell carcinoma of the oral cavity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

None

References

1. Jemal A, Bray F, Center MM, et al. Global cancer statistics. *CA Cancer J Clin.* 2011; 61:69–90.10.3322/caac.20107 [PubMed: 21296855]
2. McDonald MW, Lawson J, Garg MK, et al. ACR appropriateness criteria retreatment of recurrent head and neck cancer after prior definitive radiation expert panel on radiation oncology-head and neck cancer. *Int J Radiat Oncol Biol Phys.* 2011; 80:1292–1298.10.1016/j.ijrobp.2011.02.014 [PubMed: 21530100]
3. Vermorken JB, Specenier P. Optimal treatment for recurrent/metastatic head and neck cancer. *Ann Oncol.* 2010; 21(7):vii252–261.10.1093/annonc/mdq453 [PubMed: 20943624]
4. Gillison ML, Koch WM, Capone RB, et al. Evidence for a causal association between human papillomavirus and a subset of head and neck cancers. *J Natl Cancer Inst.* 2000; 92:709–720.10.1093/jnci/92.9.709 [PubMed: 10793107]
5. Kreimer AR, Clifford GM, Boyle P, Franceschi S. Human papillomavirus types in head and neck squamous cell carcinomas worldwide: a systematic review. *Cancer Epidemiol Biomarkers Prev.* 2005; 14:467–475.10.1007/s12105-010-0171-9 [PubMed: 15734974]
6. Munoz N, Bosch FX, de Sanjose S, et al. Epidemiologic classification of human papillomavirus types associated with cervical cancer. *N Engl J Med.* 2003; 348:518–527.10.1056/NEJMoa021641 [PubMed: 12571259]
7. Kenter GG, Welters MJ, Valentijn AR, et al. Phase I immunotherapeutic trial with long peptides spanning the E6 and E7 sequences of high-risk human papillomavirus 16 in end-stage cervical cancer patients shows low toxicity and robust immunogenicity. *Clin Cancer Res.* 2008; 14:169–177.10.1158/1078-0432.CCR-07-1881 [PubMed: 18172268]
8. Rensing ME, Sette A, Brandt RM, et al. Human CTL epitopes encoded by human papillomavirus type 16 E6 and E7 identified through in vivo and in vitro immunogenicity studies of HLA-A*0201-binding peptides. *J Immunol.* 1995; 154:5934–5943. [PubMed: 7538538]
9. van der Burg SH, Rensing ME, Kwappenberg KM, et al. Natural T-helper immunity against human papillomavirus type 16 (HPV16) E7-derived peptide epitopes in patients with HPV16-positive cervical lesions: identification of 3 human leukocyte antigen class II-restricted epitopes. *Int J Cancer.* 2001; 91:612–618.10.1002/1097-0215 [PubMed: 11267969]
10. Cuffel C, Rivals JP, Zaugg Y, et al. Pattern and clinical significance of cancer-testis gene expression in head and neck squamous cell carcinoma. *Int J Cancer.* 2011; 128:2625–2634.10.1002/ijc.25607 [PubMed: 20715104]
11. Figueiredo DL, Mamede RC, Spagnoli GC, et al. High expression of cancer testis antigens MAGE-A, MAGE-C1/CT7, MAGE-C2/CT10, NY-ESO-1, and gage in advanced squamous cell carcinoma of the larynx. *Head Neck.* 2011; 33:702–707.10.1002/hed.21522 [PubMed: 20886663]
12. Jungbluth AA, Ely S, DiLiberto M, et al. The cancer-testis antigens CT7 (MAGE-C1) and MAGE-A3/6 are commonly expressed in multiple myeloma and correlate with plasma-cell proliferation. *Blood.* 2005; 106:167–174. doi: <http://dx.doi.org/10.1182/blood-2004-12-4931>. [PubMed: 15761016]
13. Condomines M, Hose D, Raynaud P, et al. Cancer/Testis Genes in Multiple Myeloma: Expression Patterns and Prognosis Value Determined by Microarray Analysis. *The Journal of Immunology.* 2007; 178:3307–3315.10.4049/jimmunol.178.5.3307 [PubMed: 17312182]
14. van der Bruggen P, Traversari C, Chomez P, et al. A gene encoding an antigen recognized by cytolytic T lymphocytes on a human melanoma. *Science.* 1991; 254:1643–1647.10.1126/science.1840703 [PubMed: 1840703]
15. Kobayashi H, Song Y, Hoon DS, et al. Tumor-reactive T helper lymphocytes recognize a promiscuous MAGE-A3 epitope presented by various major histocompatibility complex class II alleles. *Cancer Res.* 2001; 61:4773–4778. [PubMed: 11406551]

16. Lu J, Wettstein PJ, Higashimoto Y, et al. TAP-independent presentation of CTL epitopes by Trojan antigens. *J Immunol.* 2001; 166:7063–7071.10.4049/jimmunol.166.12.7063 [PubMed: 11390450]
17. Lu J, Higashimoto Y, Appella E, Celis E. Multiepitope Trojan antigen peptide vaccines for the induction of antitumor CTL and Th immune responses. *J Immunol.* 2004; 172:4575–4582.10.4049/jimmunol.172.7.4575 [PubMed: 15034075]
18. Voskens CJ, Sewell D, Hertzano R, et al. Induction of MAGE-A3 and HPV-16 immunity by Trojan vaccines in patients with head and neck carcinoma. *Head Neck.* 2012; 34:1734–1746.10.1002/hed.22004 [PubMed: 22287423]
19. Fruh K, Yang Y. Antigen presentation by MHC class I and its regulation by interferon gamma. *Curr Opin Immunol.* 1999; 11:76–81.10.1016/S0952-7915(99)80014-4 [PubMed: 10047537]
20. Cheever MA, Chen W. Therapy with cultured T cells: principles revisited. *Immunol Rev.* 1997; 157:177–194.10.1111/j.1600-065X.1997.tb00982.x [PubMed: 9255630]
21. Gao FG, Khammanivong V, Liu WJ, et al. Antigen-specific CD4+ T-cell help is required to activate a memory CD8+ T cell to a fully functional tumor killer cell. *Cancer Res.* 2002; 62:6438–6441. [PubMed: 12438231]
22. Voskens CJ, Strome SE, Sewell DA. Synthetic peptide-based cancer vaccines: lessons learned and hurdles to overcome. *Curr Mol Med.* 2009; 9:683–693. [PubMed: 19689295]
23. Pichichero ME. Improving vaccine delivery using novel adjuvant systems. *Hum Vaccin.* 2008; 4:262–270.10.4161/hv.4.4.5742 [PubMed: 18398303]
24. Kreimer AR, Johansson M, Waterboer T, et al. Evaluation of human papillomavirus antibodies and risk of subsequent head and neck cancer. *J Clin Oncol.* 2013; 31:2708–2715.10.1200/JCO.2012.47.2738 [PubMed: 23775966]
25. Kruit WH, van Ojik HH, Brichard VG, et al. Phase 1/2 study of subcutaneous and intradermal immunization with a recombinant MAGE-3 protein in patients with detectable metastatic melanoma. *Int J Cancer.* 2005; 117:596–604.10.1002/ijc.21264 [PubMed: 15945101]
26. Klebanoff CA, Acquavella N, Yu Z, Restifo NP. Therapeutic cancer vaccines: are we there yet? *Immunol Rev.* 2011; 239:27–44.10.1111/j.1600-065X.2010.00979.x [PubMed: 21198663]
27. Atanackovic D, Altorki NK, Stockert E, et al. Vaccine-induced CD4+ T cell responses to MAGE-3 protein in lung cancer patients. *J Immunol.* 2004; 172:3289–3296.10.4049/jimmunol.172.5.3289 [PubMed: 14978137]
28. Vansteenkiste J, Zielinski M, Linder A, et al. Adjuvant MAGE-A3 immunotherapy in resected non-small-cell lung cancer: phase II randomized study results. *J Clin Oncol.* 2013; 31:2396–2403.10.1200/JCO.2012.43.7103 [PubMed: 23715567]
29. Marchand M, Punt CJ, Aamdal S, et al. Immunisation of metastatic cancer patients with MAGE-3 protein combined with adjuvant SBAS-2: a clinical report. *Eur J Cancer.* 2003; 39:70–77. doi: [http://dx.doi.org/10.1016/S0959-8049\(02\)00479-3](http://dx.doi.org/10.1016/S0959-8049(02)00479-3). [PubMed: 12504661]
30. Kruit WH, Suciu S, Dreno B, et al. Selection of immunostimulant AS15 for active immunization with MAGE-A3 protein: results of a randomized phase II study of the European Organisation for Research and Treatment of Cancer Melanoma Group in Metastatic Melanoma. *J Clin Oncol.* 2013; 31:2413–2420.10.1200/JCO.2012.43.7111 [PubMed: 23715572]
31. Brinkman JA, Hughes SH, Stone P, et al. Therapeutic vaccination for HPV induced cervical cancers. *Dis Markers.* 2007; 23:337–352.10.1155/2007/245146 [PubMed: 17627067]
32. Allen CT, Judd NP, Bui JD, Uppaluri R. The clinical implications of antitumor immunity in head and neck cancer. *Laryngoscope.* 2012; 122:144–157.10.1002/lary.21913 [PubMed: 22183634]
33. Young MR. Protective mechanisms of head and neck squamous cell carcinomas from immune assault. *Head Neck.* 2006; 28:462–470.10.1002/hed.20331 [PubMed: 16284974]
34. Lyford-Pike S, Peng S, Young GD, et al. Evidence for a role of the PD-1:PD-L1 pathway in immune resistance of HPV-associated head and neck squamous cell carcinoma. *Cancer Res.* 2013; 73:1733–41.10.1158/0008-5472.CAN-12-2384 [PubMed: 23288508]
35. Strome SE, Dong H, Tamura H, et al. B7-H1 blockade augments adoptive T-cell immunotherapy for squamous cell carcinoma. *Cancer Res.* 2003; 63:6501–6505. [PubMed: 14559843]
36. Ukpo OC, Thorstad WL, Lewis JS Jr. B7-H1 Expression Model for Immune Evasion in Human Papillomavirus-Related Oropharyngeal Squamous Cell Carcinoma. *Head Neck Pathol.* 2012; 7:113–21.10.1007/s12105-012-0406-z [PubMed: 23179191]

37. Li W, Deng XM, Wang CX, et al. Down-regulation of HLA class I antigen in human papillomavirus type 16 E7 expressing HaCaT cells: correlate with TAP-1 expression. *Int J Gynecol Cancer*. 2010; 20:227–232.10.1111/IGC.0b013e3181ccec5 [PubMed: 20134267]
38. Campo MS, Graham SV, Cortese MS, et al. HPV-16 E5 down-regulates expression of surface HLA class I and reduces recognition by CD8 T cells. *Virology*. 2010; 407:137–142.10.1016/j.virol.2010.07.044 [PubMed: 20813390]
39. Schott AK, Pries R, Wollenberg B. Permanent up-regulation of regulatory T-lymphocytes in patients with head and neck cancer. *Int J Mol Med*. 2010; 26:67–75.10.3892/ijmm_00000436 [PubMed: 20514424]
40. Strauss L, Bergmann C, Szczepanski M, et al. A unique subset of CD4+CD25 high Foxp3+ T cells secreting interleukin-10 and transforming growth factor-beta1 mediates suppression in the tumor microenvironment. *Clin Cancer Res*. 2007; 13:4345–4354.10.1158/1078-0432.CCR-07-0472 [PubMed: 17671115]
41. Ostrand-Rosenberg S, Sinha P, Beury DW, Clements VK. Cross-talk between myeloid-derived suppressor cells (MDSC), macrophages, and dendritic cells enhances tumor-induced immune suppression. *Semin Cancer Biol*. 2012; 22:275–281.10.1016/j.semcancer.2012.01.011 [PubMed: 22313874]
42. Herbst RS, Arquette M, Shin DM, et al. Phase II multicenter study of the epidermal growth factor receptor antibody cetuximab and cisplatin for recurrent and refractory squamous cell carcinoma of the head and neck. *J Clin Oncol*. 2005; 23:5578–5587.10.1200/JCO.2005.07.120 [PubMed: 16009949]
43. Zenda S, Onozawa Y, Boku N, et al. Single-agent docetaxel in patients with platinum-refractory metastatic or recurrent squamous cell carcinoma of the head and neck (SCCHN). *Jpn J Clin Oncol*. 2007; 37:477–481.10.1093/jjco/hym059 [PubMed: 17720737]
44. Le DT, Jaffee EM. Regulatory T-cell modulation using cyclophosphamide in vaccine approaches: a current perspective. *Cancer Res*. 2012; 72:3439–3444.10.1158/0008-5472.CAN-11-3912 [PubMed: 22761338]
45. Lutsiak ME, Semnani RT, De Pascalis R, et al. Inhibition of CD4(+)25+ T regulatory cell function implicated in enhanced immune response by low-dose cyclophosphamide. *Blood*. 2005; 105:2862–2868. doi: <http://dx.doi.org/10.1182/blood-2004-06-2410>. [PubMed: 15591121]

Abbreviations

B7H1	B7 homolog 1
CNS	Central Nervous System
CR	Complete Response
CTCAE	Common toxicity criteria for adverse events
DLT	Dose Limiting Toxicity
ECOG	Eastern Cooperative Oncology Group
ELISA	enzyme-linked immunosorbent assay
ELISPOT	Enzyme-Linked ImmunoSpot
GM-CSF	Granulocyte colony stimulating factor
HIV	Human Immunodeficiency Virus
HLA	Human Leukocyte Antigen
HPV16	Human Papillomavirus 16
IFN-γ	Interferon Gamma

IL-10	Interleukin 10
MAGE	Melanoma antigen E
MDSC	Myeloid derived suppressor cells
ml	Milliliter
m²	Meter squared
MTD	Maximum Tolerated Dose
NSCLC	Non-small cell lung cancer
OS	Overall Survival
PD	Progressive Disease
PD-1	Programmed death 1
PD-L1	Programmed death-ligand 1
PET/CT	Positron emission tomography-computed tomography
PGE₂	Prostaglandin E ₂
PR	Partial Response
RECIST	Response Evaluation Criteria in Solid Tumors
R/M	Recurrent/Metastatic
SCCHN	Squamous cell carcinoma of the head and neck
SD	Stable Disease
TAP	Transporter associated with antigen processing
TGF-β	Transforming growth factor beta
Tregs	Regulatory T cells
UMGCC	University of Maryland Greenebaum Cancer Center
μg	Microgram

Figure 1. Consort Diagram

Figure 2.

*. Elispot and ELISA results from HPV16 positive patients that had a response to vaccine.

ELISPOT Assay: Collection of peripheral blood mononuclear cells (PBMCs) occurred before and after vaccination at time points shown in Figure 2. T cell reactivity measurements were carried out against human papillomavirus (HPV16) – Trojan and individual Trojan specific HLA class I (HPV-CTL) and Class II (HPV-HTL) using interferon- γ (IFN- γ) ELISPOT assays. Control values are shown without subtraction from experimental values. ELISPOT assays were performed with in vitro stimulation. Number of spots per 100,000 PBMCs is shown on the Y-axis. Antigen concentration was 10 μ g/ml. Due to variation in response some figures contain different increments for the Y axis.

ELISA Assay: HPV16-Trojan specific immunoglobulin G (IgG) antibody response was evaluated before and after vaccination at time points shown in Figure 2. Absorbance data on the Y-axis represents mean optical density (OD) results from triplicate wells from three different plasma dilutions

A. HPV16 500 μ g cohort

B. HPV16 1000 μ g cohort

C. HPV16 1500 μ g cohort

* = Patient 003 did not have a T cell response

Figure 3.

*, Elispot and ELISA results from MAGE-A3 positive patients that had a response to vaccine.

ELISPOT Assay: Collection of peripheral blood mononuclear cells (PBMCs) occurred before and after vaccination at time points shown in Figure 3. T cell reactivity measurements were carried out against MAGE-A3 - Trojan and individual Trojan specific HLA class I (MAGE-CTL) and Class II (MAGE-HTL) using interferon- γ (IFN- γ) ELISPOT assays. Control values are shown without subtraction from experimental values. ELISPOT assays were performed with in vitro stimulation. Number of spots per 100,000 PBMCs is shown on the Y-axis. Antigen concentration was 10 μ g/ml. Due to variation in response some figures contain different increments for the Y axis.

ELISA Assay: MAGE-A3-Trojan specific immunoglobulin G (IgG) antibody response was evaluated before and after vaccination at time points shown in Figure 3. Absorbance data on the Y-axis represents mean optical density (OD) results from triplicate wells from three different plasma dilutions

A. MAGE-A3 500 μ g cohort

B. MAGE-A3 1000 μ g cohort

C. MAGE-A3 1500 μ g cohort

* = Patient 004 did not have a T cell response

Table 1
A. Baseline characteristics of patients in HPV16 positive cohort

Characteristic	Value
Total number of patients	9
Age, yrs Median(range)	53 (43-68)
Gender	
Male	9
Female	0
Race:	
Black	1 (11%)
White	8 (89%)
Other	0
Performance status	
Median	1
Range	0-1
Primary Site	
Oral	0
Oropharynx	8 (89%)
Hypopharynx	0
Larynx	1 (11%)
Nasopharynx	0
Nasal/Paranasal	0
Disease at enrollment	
Locoregional recurrence only	2 (22%)
Metastatic Disease	7 (78%)
Sites of Metastatic Disease	
Lung	5
Bone	2
Liver	1
Other	1 (retroperitoneum)
Number of prior chemotherapy regimens	
0	2 (22%)
1	3 (33%)
2	2 (22%)
3+	2 (22%)

B. Baseline Characteristics of patients in MAGE-A3 positive cohort	
Characteristic	Value
Total number of patients	7
Age,yrs Median(range)	63 (48-71)
Gender	
Male	7
Female	0
Race	
Black	2 (29%)
White	5 (71%)
Other	0
Performance status	
Median	1
Range	0-1
Primary Site	
Oral	0
Oropharynx	2 (29%)
Hypopharynx	3 (42%)
Larynx	2 (29%)
Nasopharynx	0
Nasal/Paranasal	0
Disease at enrollment	
Locoregional recurrence only	5 (71%)
Metastatic Disease	2 (29%)
Sites of Metastatic Disease	
Lung	2
Bone	0
Liver	0
Other	1 (axilla)
Number of prior chemotherapy regimens	
0	4 (57%)
1	3 (43%)
2	0
3+	0

Table 2
A. Toxicity data for patients receiving the HPV16 Peptide Immunomodulatory Vaccine

Dosage level	Number of patients	Toxicity	Grade 1 *	Grade 2 *	Grade 3 *	Grade 4 *
500µg	3	Induration at injection site	1			
1000µg	3	Pruritus at injection site	3			
		Erythema at injection site	2			
		Edema at injection site		3		
1500µg	3	Pruritus at injection site	3			
		Edema at injection site	1			
* number of events						

B. Toxicity Data for patients receiving the MAGE-A3 Peptide Immunomodulatory Vaccine						
Dosage level	Number of patients	Toxicity	Grade 1 *	Grade 2 *	Grade 3 *	Grade 4 *
500µg	3	Pharyngocutaneous Fistula			1	
		Erythema at injection site	3			
		Pain at injection site	3			
1000µg	3	Pruritus at injection site	1			
		Pain at injection site	1			
		Edema at injection site		1		
1500µg	1	Headache	2			
		Nausea	2			
		Pruritus at injection site	2			
		Chills	2			
		Flu like symptoms	3			
		Joint pain	1			
* number of events						

Table 3
Summary of T Cell and Antibody Response to Vaccines

Cohort	T Cell and Antibody Response*	T cell and Antibody Response in patients that received all four vaccinations
All Patients	50% (8/16)	73% (8/11)
MAGE-A3	57% (4/7)	67% (4/6)
HPV16	44% (4/9)	80% (4/5)

* All patients that had a T cell response also had an antibody response. One patient in the HPV16 500µg cohort had an antibody response but not a T cell response.