CORRESPONDENCE

Non-Alcoholic Fatty Liver Disease Epidemiology, Clinical Course, Investigation, and Treatment

by Dr. med. Johannes Weiß, Dr. med. Monika Rau, Prof. Dr. med. Andreas Geier
in issue 26/2014

Difficult-to-Understand Point

This very well researched review article lists “states of hunger” as one of the possible causes of non-alcoholic fatty liver disease. In this point, I do not fully understand the complex relationships involved. This phenomenon is also incomprehensible to me from a pathophysiological perspective. The opposite would be quite plausible. I think that many colleagues find it difficult to understand this point; therefore, I would like to ask the authors to briefly explain it.

DOI: 10.3238/arztebl.2015.0143a

REFERENCES

1. Weiß J, Rau M, Geier A: Non-alcoholic fatty liver disease: epidemiology, clinical course, investigation, and treatment. *Dtsch Arztebl Int* 2014; 111: 447–52.

Walter Hofmann
Unter der Kastanien 22
65779 Kelkheim
WaHo@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Mediterranean Diet

Certainly, the information about non-alcoholic fatty liver disease (NAFLD) provided by the authors can only be agreed with (1). Nevertheless, I think that the “Fatty Liver Index” (FLI), a validated non-invasive diagnostic tool, should also be mentioned as a relevant aid to decision-making in everyday clinical practice. Developed by the working group around Giorgio Bedogni in 2006, the FLI is calculated using an algorithm based on body-mass index, waist circumference, triglycerides, and gamma-glutamyl transferase. An FLI higher than 60 indicates a more than 80% likelihood of hepatic steatosis (2, 3). Implementing the easy-to-use FLI as part of a tight patient monitoring regimen could offer clinical advantages as patients with an FLI above 60 also have an increased incidence of diabetes, develop early atherosclerosis (thicker intima/media, higher plaque burden) and show an increased mortality compared with patients with low FLIs.

As a nutritional scientist, I like to point out with regard to the alimentary management of the disease—which plays a critical role in NAFLD patients—that only the Mediterranean diet is supported

by some potential evidence (4): plenty of vegetables, fruits, legumes, mainly complex carbohydrates with low glycemic index, fish, olive oil, dairy products, etc. and little processed food, “red meat”, sweets and sugary soft drinks would thus be the primary long-term diet for NAFLD patients.

DOI: 10.3238/arztebl.2015.0143b

REFERENCES

1. Weiß J, Rau M, Geier A: Non-alcoholic fatty liver disease: epidemiology, clinical course, investigation and treatment. *Dtsch Arztebl Int* 2014; 111: 447–52.
2. Bedogni G, Bellentani S, Miglioli L, et al.: The Fatty Liver Index: a simple and accurate predictor of hepatic steatosis in the general population. *BMC Gastroenterol* 2006; 6: 33.
3. Koehler EM, Schouten JN, Hansen BE, et al.: External validation of the fatty liver index for identifying nonalcoholic fatty liver disease in a population-based study. *Clin Gastroenterol Hepatol* 2013; 11: 1201–4.
4. Sofi F, Casini A: Mediterranean diet and non-alcoholic fatty liver disease: New therapeutic option around the corner? *World J Gastroenterol* 2014; 20: 7339–46.

Dr. oec. troph. Martin Hofmeister

Verbraucherzentrale Bayern e. V., Referat Lebensmittel und Ernährung,
München
hofmeister@vzbayern.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Benefit: “Therapeutic Indication”

Liver biopsy as the gold standard for diagnosis should only be performed if a therapeutic indication can be derived from it. Ultrasonography as a non-invasive screening method loses accuracy in early stages of the disease and depends very much on the skills and experiences of the examiner. Therefore, we prefer the “Fatty Liver Index” (FLI) for screening. In their publications (1, 2), Bedogni et al. described a simple and accurate indicator which calculates the FLI using an algorithm based on triglycerides (mg/dL), BMI (kg/m^2), gamma-GT (U/L), and waist circumference (cm). The FLI can vary between 0 and 100. An FLI <30 (negative likelihood ratio = 0.2) rules out fatty liver with a high degree of certainty, while an FLI >60 (positive likelihood ratio = 4.3) rules in fatty liver.

The significance of the FLI was confirmed in subsequent studies. An FLI >60 is indicative of fatty liver with 78% probability, while an FLI of 20 or less rules out fatty liver with 91% probability.

Not only can the FLI be used for screening purposes, it is also an appropriate method for patient monitoring after dietary interventions. A multivariate adjusted analysis of data from more than 3000 patients undergoing coronary angiography (3) demonstrated for the group of patients with an FLI above 75.6 a highly statistically significant increase in cardiovascular mortality, non-cardiovascular mortality (infections, cancer,

liver disease), and a significantly increased overall mortality.

Since “doctors in private practice have an important steering function“, the FLI is ideally suited for both screening and follow-up, as it is easy to use and cost-effective.

DOI: 10.3238/arztebl.2015.0143c

REFERENCES

1. Bedogni G, Bellentani S, Miglioli L, et al.: The fatty liver index: A simple and accurate predictor of hepatic steatosis in the general population. *BMC Gastroenterol* 2006; 6: 33.
2. Bedogni G, Kahn HS, Bellentani S, Tiribelli C: A simple index of lipid overaccumulation is a good marker of liver steatosis. *BMC Gastroenterology* 2010; 10: 98.
3. Lerchbaum E, Pilz S, Grammer TB, et al.: The fatty liver index is associated with increased mortality in subjects referred to coronary angiography; *Nutrition, Metabolism & Cardiovascular Diseases* 2013; 23: 1231–8.
4. Weiß J, Rau M, Geier A: Non-alcoholic fatty liver disease: epidemiology, clinical course, investigation and treatment. *Dtsch Arztebl Int* 2014; 111: 447–52.

Dr. med. Hardy Walle
Bodymed AG, Kitzingen
h.walle@bodymed.com

Conflict of interest statement

Dr. Walle is the CEO of the Bodymed AG and managing director of the Thanaka GmbH.

In Reply

W. Hofmann raises the question of the links between non-alcoholic fatty liver disease (NAFLD) and states of hunger, mentioned in the review as a possible cause of the disease. Here, we would like to point out that in the corresponding table states of hunger are not listed as a cause of NAFLD, but as a cause of secondary hepatic steatosis—this difference is important. Two exemplary conditions are kwashiorkor and marasmus which may be associated with fatty liver, besides muscle wasting. The most likely underlying pathomechanisms are proteolysis of the muscles as well as lipolysis which may lead to fatty liver as the result of an increased production of free fatty acids and a lack of lipoproteins; this was experimentally demonstrated for states of hunger (1). Both M. Hofmeister and H. Walle mention in their letters the “Fatty Liver Index“ (2) which can identify patients with fatty liver with an accuracy of 0.84 based on the parameters body mass index, waist circumference, triglycerides, and gamma-glutamyl transpeptidase. Ultimately, various non-invasive scores are available which can be used to identify high-risk patients. We think that the “NAFLD Fibrosis Score“

mentioned in our article, which was developed by the North American Clinical Research Network (CRN) working group (3), is perfectly suited for this purpose. This score is also based on few routine parameters, viz. age, body mass index, AST (GOT), ALT (GPT), platelet count, and albumin, and can be calculated online; at the same time, the interpretation of the result is provided. This score can rule out advanced fibrosis with a negative predictive value between 0.88 and 0.93 and diagnose advanced fibrosis with a positive predictive value between 0.82 and 0.90; with this, 75% of all liver biopsies could be avoided in the underlying study. The “Fatty Liver Index“ has a negative likelihood ratio of 0.2 for ruling out hepatic steatosis for values <30 and a positive likelihood ratio of 4.3 for values ≥60 (2). However, there is the important difference that the “NAFLD Fibrosis Score“ provides information about advanced fibrosis, while the “Fatty Liver Index“ provides information about hepatic steatosis. A recently published study has demonstrated the superiority of the “NAFLD Fibrosis Score“ over other non-invasive scores, such as the ratio of AST (GOT) and platelet count, the FIB-4 score, or the BARD score (4).

DOI: 10.3238/arztebl.2015.0144

REFERENCES

1. Møller L, Stodkilde-Jørgensen H, Jensen FT, Jørgensen JO: Fasting in healthy subjects is associated with intrahepatic accumulation of lipids as assessed by 1H-magnetic resonance spectroscopy. *Clin Sci (Lond)* 2008; 114: 547–52.
2. Bedogni G, Bellentani S, Miglioli L, et al.: The Fatty Liver Index: a simple and accurate predictor of hepatic steatosis in the general population. *BMC Gastroenterol* 2006; 6: 33.
3. Angulo P, Hui JM, Marchesini G, et al.: The NAFLD fibrosis score: a noninvasive system that identifies liver fibrosis in patients with NAFLD. *Hepatology* 2007; 45: 846–54.
4. Angulo P, Bugianesi E, Björnsson ES, et al.: Simple noninvasive systems predict long-term outcomes of patients with nonalcoholic fatty liver disease. *Gastroenterology* 2013; 145: 782–9.
5. Weiß J, Rau M, Geier A: Non-alcoholic fatty liver disease: epidemiology, clinical course, investigation and treatment. *Dtsch Arztebl Int* 2014; 111: 447–52.

Dr. med. Johannes Weiß

Dr. med. Monika Rau

Prof. Dr. med. Andreas Geier

Universitätsklinik Würzburg, Medizinische Klinik und Poliklinik II, Schwerpunkt Hepatologie, Würzburg
weiss_j1@ukw.de

Conflict of interest statement

Prof. Geier received material resources from Burgerstein Vitamine (study medication SASL34) and from the Velux Foundation (external funds for the SASL34 study).

The remaining authors declare that no conflict of interest exists.