Published in final edited form as:

Exp Gerontol. 2015 August ; 68: 3–7. doi:10.1016/j.exger.2014.09.018.

Cellular senescence and the aging brain

Shankar J. Chinta¹, Georgia Woods¹, Anand Rane¹, Marco Demaria¹, Judith Campisi^{1,2}, and Julie K Andersen^{1,*}

¹Buck Institute for Research on Aging, Novato, CA, 94945, USA

²Lawrence Berkeley National Laboratory, Berkeley, CA, 94720, USA

Abstract

Cellular senescence is a potent anti-cancer mechanism that arrests the proliferation of mitotically competent cells to prevent malignant transformation. Senescent cells accumulate with age in a variety of human and mouse tissues where they express a complex ‘senescence-associated secretory phenotype’ (SASP). The SASP includes many pro-inflammatory cytokines, chemokines, growth factors and proteases that have the potential to cause or exacerbate age-related pathology, both degenerative and hyperplastic. While cellular senescence in peripheral tissues has recently been linked to a number of age-related pathologies, its involvement in brain aging is just beginning to be explored. Recent data generated by several laboratories suggest both aging and age-related neurodegenerative diseases are accompanied by an increase in SASP-expressing senescent cells of non-neuronal origin in the brain. Moreover, this increase correlates with neurodegeneration. Senescent cells in the brain could therefore constitute novel therapeutic targets for treating age-related neuropathologies.

Keywords

Aging; Brain; Cellular Senescence; Senescence Associated Secretory Phenotype; Neurodegeneration

1. Introduction

The development of therapies aimed at mitigating or delaying age-related neurodegenerative diseases is a major priority for the biomedical community due to the enormous social, emotional and economic burden associated with them. The disappointing outcomes of dozens of phase III clinical trials of treatments for Alzheimer’s disease (AD) and Parkinson’s disease (PD) indicate a need for fresh approaches to identify novel targets that drive processes that cause age-related neuropathology.

© 2014 Elsevier Inc. All rights reserved.

*Corresponding author: jandersen@buckinstitute.org, Phone: (415) 209-2070.

Conflict of interest

The authors have no conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A new view has recently emerged suggesting that aging itself may not merely be a major risk factor for these disorders, but may actually be the underlying driving force. This view begs the question as to whether interventions that prevent the occurrence of basic aging processes can prevent or alleviate age-related conditions, including neurodegenerative diseases. One such mechanism currently under investigation by several laboratories is a process known as cellular senescence.

2. Cellular senescence and the SASP

Cellular senescence is a potent anti-cancer mechanism that can occur in virtually all cell types that are capable of cell division. Thus far, replication-competent cell types that undergo senescence include fibroblasts, epithelial cells, melanocytes, endothelial cells, astrocytes and many others (Bitto et al., 2010; Coppe et al., 2010; Coppe et al., 2008; Voghel et al., 2007; Wajapeyee et al., 2008). The senescence response arrests cell proliferation, stably and essentially irreversibly, in response to stresses that puts cells at risk for malignant transformation (Campisi, 2001; Collado and Serrano, 2010; Prieri et al., 2008). These stresses include repeated cell division that erodes telomeres (perceived by cells as severely damaged DNA), DNA damage anywhere in the genome, and disrupted chromatin (epigenomic damage) (Campisi, 2007; Guney et al., 2006; Rodier et al., 2005; Shay and Wright, 2005). Cellular senescence can also be induced by activated oncogenes, strong or persistent mitogenic signals, and several forms of oxidative stress (Adams, 2009; Ben-Porath and Weinberg, 2005; Braig and Schmitt, 2006; Campisi and d'Adda di Fagagna, 2007; Herbig and Sedivy, 2006; Ohtani et al., 2004; Passos and Von Zglinicki, 2006; Toussaint et al., 2000). Many senescence inducers directly or indirectly cause genomic or epigenomic damage. The damage response ultimately activates the p53/p21 and p16^{INK4a}/pRB tumor suppressor pathways, which establish and maintain the senescence growth arrest (Adams, 2009; Campisi and d'Adda di Fagagna, 2007; Herbig and Sedivy, 2006; Ohtani et al., 2004).

Cellular senescence may be an example of evolutionary antagonistic pleiotropy (Campisi, 2003). This evolutionary theory posits that, because the force of natural selection declines with age, processes that were selected to promote fitness in young organisms can have unselected deleterious effects in older organisms (Rose, 1991). Hence, the senescence response protects organisms from cancer early in life; late life in life, however, it may promote phenotypes and pathologies associated with aging. Senescent cells have indeed been demonstrated to accumulate with age in a variety of tissues (Dimri et al., 1995; Erusalimsky and Kurz, 2005; Herbig and Sedivy, 2006; Jeyapalan et al., 2007; Kishi, 2004; Melk et al., 2003; Paradis et al., 2001). A seminal publication recently demonstrated that the elimination of senescent cells that accumulate in a progeroid mouse model prevents the onset of three major aging phenotypes (cataracts, sarcopenia, loss of subcutaneous fat), providing the first evidence that senescent cells play a causal role in at least some age-related pathologies *in vivo* (Baker et al., 2011).

In addition to arresting growth, senescent cells express a senescence-associated secretory phenotype (SASP): the robust secretion of many inflammatory cytokines, growth factors and proteases (Coppe et al., 2010; Coppe et al., 2008). SASP factors include several interleukins

(ILs), monocyte chemotactic proteins (MCPs; aka CCLs), growth-related oncogenes (GROs; aka CXCLs), and inflammatory cytokines such as granulocyte-macrophage colony stimulating factor (GM-CSF) and macrophage inflammatory proteins (MIPs; aka CCLs), amongst others (Coppe et al., 2010; Davalos et al., 2010; Freund et al., 2010). SASP factors can have potent effects on neighboring cells and thus can alter local and systemic tissue milieus.

There are potentially beneficial effects of the SASP. For example, chemokines or cytokines secreted by senescent cells can recruit natural killer cells, thus facilitating the removal of senescent cells and neighbouring tumor cells; this process is termed 'senescence surveillance'. Other SASP factors can communicate cellular damage to the surrounding tissue and stimulate repair or limit damage-induced fibrosis (Krizhanovsky et al., 2008). However, many SASP factors have been shown, or are suspected, to cause or contribute to the loss of tissue structure and function that occurs with age by creating a pro-inflammatory milieu. For example, the SASP has been shown to: (1) disrupt normal tissue structure and function – e.g., the ability of mammary epithelial cells to form alveoli and ducts and express milk proteins (Parrinello et al., 2005; Tsai et al., 2005); (2) induce epithelial-to-mesenchyme transitions in normal and premalignant epithelial cells (Coppe et al., 2008; Coppe et al., 2011); and (3) stimulate premalignant and non-aggressive cancer cells to migrate and invade a basement membrane (Coppe et al., 2010; Rodier et al., 2009). In some cases, the use of blocking antibodies and recombinant proteins allowed assignment of these activities to one or a few SASP factors. *In vivo*, senescent cells can promote the conversion of premalignant cells to full blown malignancy (Krtolica et al., 2001) and stimulate the growth and vascularization of tumors initiated from established tumor cell lines (Coppe et al., 2010; Krtolica et al., 2001).

Most SASP factors are up-regulated at the level of mRNA, in part due to increased activities of nuclear factor kappa light chain enhancer of activated B cells (NF- κ B) and CCAAT/enhancer binding protein (C/EBP) transcription factors (Coppe et al., 2008; Freund et al., 2010). The SASP is dependent on the activation of specific signaling pathways, including the DNA damage response (DDR), p38 mitogen-activated protein kinase (p38MAPK), and mechanistic target of rapamycin (mTOR) pathways (Coppe et al., 2008; Freund et al., 2011).

3. The aging brain

The brain is arguably the most multifaceted tissue in complex organisms, controlling processes that are not only vital to life but also at the heart of cognition and personality. Loss of brain function, whether through trauma or – much more commonly – aging, exacts an enormous human and economic toll, especially among people in developed nations where average life spans are at record highs (Vaupel, 2010; Yankner et al., 2008). As with virtually all aging tissues in the body, the aging brain is characterized by low level, chronic inflammation (Chung et al., 2009; Franceschi et al., 2007). This phenomenon has been termed 'inflammaging' (or 'neuro-inflammaging' in the brain) (Frank-Cannon et al., 2009). Inflammaging is thought to cause or contribute to most, if not all, major pathologies associated with aging. Inflammation evolved to remove foreign bodies, pathogens and damaged cells produced by acute cellular stress or injury (Ferrucci et al., 2005). The acute

inflammatory response causes robust local oxidative and nitrosative damage, but is designed to be short-acting and self-limiting. Chronic inflammation, by contrast, is a more feeble response, but is long-standing and often self-perpetuating (Nathan and Ding, 2010). Inflammaging is considered ‘sterile’ inflammation because it occurs in the absence of an obvious pathogen or foreign body.

In the aging brain, pathological changes associated with chronic inflammation include significant decreases in certain neuronal populations, dendritic and axonal arborization, post-synaptic densities, dendritic spines, presynaptic markers, synapses and cortical volume (Yankner et al., 2008). These cellular and tissue changes result in cognitive and motor impairment, memory loss and other phenotypes characteristic of aged mammals. The etiology of neuro-inflammaging is a crucial unresolved question. An important source of neuroinflammation in the aging brain is the proliferative glial cells (astrocytes, oligodendrocytes and microglia). These cells normally provide structural, metabolic and trophic support to neurons (Allen and Barres, 2009; Chung et al., 2009; Lucin and Wyss-Coray, 2009; McGeer and McGeer, 1998; Ransom et al., 2003). However, they can also have detrimental effects on neighboring neurons due to the chronic production of pro-inflammatory factors, including reactive oxygen species (ROS) and leukocyte-attracting cytokines, which occurs with increasing frequency during aging.

4. Brain cell senescence

A potential contributor to age-related inflammation in the brain is cellular senescence, likely occurring in replication-competent glial cells. Recent studies from several laboratories suggest that senescent cells are detectable in the mammalian brain, where they could contribute to neurodegenerative processes by secreting pro-inflammatory SASP factors and/or disrupting cell-cell contacts needed for the structural and functional neuron-glial interaction that maintains neuronal ion and metabolic homeostasis (Benarroch, 2005; Magistretti, 2006). Senescent cells and their SASPs may therefore constitute a novel, understudied, and potentially important contributor to neuroinflammation and subsequent neurodegeneration. Characterization of cellular senescence in the brain could uncover novel therapeutic targets for the prevention and treatment of chronic age-related neurodegenerative diseases.

4.1 Evidence for cellular senescence in proliferation-competent brain cell types

Astrocytes are involved in a variety of important physiological and pathological processes, including modulation of synaptic neuronal function and plasticity (Finch, 1993; Nichols et al., 1993). They are the most abundant cell type in the brain and the primary responders to central nervous system (CNS) insults, including infection, trauma and neurodegeneration, in response to which they exert important tissue defense mechanisms. Dysfunctional astrocytes are implicated in neuropathology associated with both normal brain aging and various age-related neurodegenerative diseases (Chen and Swanson, 2003). In response to exogenously added H₂O₂, cultured astrocytes have been reported to display numerous characteristics of senescent cells: arrested growth, an enlarged morphology, increased senescence-associated beta-galactosidase (SA-Bgal) activity, and increased expression of the senescent cell markers p21 and p16^{INK4a} (Bitto et al., 2010). Cultured human astrocytes exposed to DNA-

damaging ionizing irradiation (IR) also undergo senescence and develop a SASP, similar to the behavior of cultured human fibroblasts (Zou et al., 2012). Astrocytes cultured from the brains of aging rats stain positively for SA- β gal, in conjunction with a reduced ability to maintain the survival of co-cultured neurons (Pertusa et al., 2007). *In vivo*, astrocytes, as determined by glial acidic fibrillary protein (GFAP)-positivity, demonstrated a flat morphology, a characteristic of senescent cells, as well as age-related synaptic impairment (Nichols et al., 1993). These findings suggest that loss of neuroprotection during brain aging coincides with increased astrocytic senescence.

Microglia, another important replication-competent cell type in the brain, function as resident macrophages in the CNS (Streit, 2002a, b). Microglia provide immune surveillance and mediate innate immune responses to invading pathogens or injury. These responses include the secretion of cytokines, prostaglandins and growth factors, production of external ROS and stimulation of phagocytosis (Doorn et al., 2012). Microglia are normally found in a quiescent (resting) state, characterized by small soma and highly ramified processes. In response to infection or CNS injury, microglia become activated and undergo morphological changes, including shortening of ramified branches and enlargement of the soma. Activated microglia also up-regulate cell surface activation antigens and secrete a variety of pro-inflammatory mediators and other potentially neurotoxic factors (Kreutzberg, 1996). Chronic microglial activation has been implicated in the neuronal death associated with neurodegenerative diseases, including AD and PD (Frank-Cannon et al., 2009; Sugama, 2009).

There is strong evidence to suggest that, with advanced age, functional abnormalities occur in the microglia that impair their ability to respond efficiently to stimuli (Conde and Streit, 2006; Sawada et al., 2008). A comparative study examining both young and old autopsied human brains demonstrated that, with age, microglia transform morphologically from ramified to hypertrophic and dystrophic forms, characterized by loss of ne branches (deramification), formation of cytoplasmic spheroids, beading and fragmentation (Flanary, 2005; Streit et al., 2004).

It has been reported that telomere shortening occurs in rat microglia, both in culture with repeated cell division and *in vivo* with advancing age. In both cases, this telomere shortening can lead to cellular senescence (Flanary and Streit, 2003, 2004). In response to repeated lipopolysaccharide administration, cultured microglial cells also undergo a senescence response as determined by arrested growth, enhanced SA- β gal activity, and senescence-associated heterochromatic foci (Yu et al., 2012). Both normal brain aging and chronic age-related neurodegenerative disease are associated with microglial-mediated increases in components that are associated with the SASP, including increases in pro-inflammatory cytokines such as IL-1 β and IL-6 (Bachstetter et al., 2011).

Other proliferative cell types in the brain that could conceivably undergo senescence include oligodendrocytes, endothelial cells and neural stem cells (NSCs). The senescence of these cell types could have both intrinsic and extrinsic effects on neuronal function. The senescence of oligodendrocytes, for example, could reduce myelination of neuronal axons, thereby decreasing their interneuronal signaling ability. The senescence of endothelial cells

could contribute to the age-related disruption of the blood-brain barrier (BBB), resulting in and influx of peripheral inflammatory factors that can contribute to subsequent neuronal cell loss (Zlokovic, 2008). The BBB deterioration that occurs during aging has also been linked to age-related cognitive decline (Abazov et al., 2009). Finally, the senescence of NSCs could blunt adult neurogenesis. Exposure of cultured NSCs to ionizing radiation was recently reported to result in senescence without a SASP in the small fraction of the population that failed to die by apoptosis, and p16-positive NSCs have been identified in the aging brain (Molofsky et al., 2006; Zou et al., 2012). Neurons are terminally differentiated and would not be predicted to mount a 'classic' senescent response, although a recent study reported senescence markers in non-dividing neurons in the aging mouse brain (Jurk et al., 2012). Given the mounting evidence that senescent cells, largely through the SASP, can disrupt tissue structure and function, cellular senescence may be an important factor to consider in age-related neurodegeneration, including which cell types are involved and how (Figure 1).

4.2 Brain cell senescence in human aging and neurodegenerative diseases

Senescent markers were recently reported in astrocytes in autopsied human brain tissue; both p16^{INK4a} and the SASP factor matrix metalloproteinase (MMP) 3 increased significantly with age and were even higher in affected cortical brain tissues from AD patients relative to age-matched controls (Bhat et al., 2012). Our laboratory similarly found an increased burden of senescent astrocytes in autopsied substantia nigra pars compacta (SNpc) from PD patients compared to age-matched controls based on elevated p16^{INK4a} levels (unpublished data). Likewise, we found an increase in DNA damage (γ H2AX) foci in astrocytes in PD SN; these foci are characteristic of senescent cells and are required for the SASP that develops in response to genomic stress (Rodier et al., 2011). This finding suggests that at least astrocytes, the major cell division-competent cell type in the brain, undergo senescence *in vivo* in humans and are more prominent in both neurodegenerative disease and aging.

5. Future issues to be addressed

Several important biological questions remain to be addressed with regard to cellular senescence in the brain. Do other proliferative cell types undergo senescence in the context of aging and age-related neurodegenerative diseases? What stressors are responsible for eliciting brain cell senescence and the SASP? How do the SASPs compare between various brain cells and other cell types, including fibroblasts, the cell type most commonly used for exploring this phenomenon? Most importantly, does brain cell senescence contribute to neurodegeneration? If yes, is the neurodegeneration due to direct effects of the SASP on neighboring neurons or is it due to activation of glial cells, resulting in the amplification of neuroinflammatory processes?

6. Conclusion

Senescent cells accumulate with age in a variety of human and mouse tissues and their elimination was recently demonstrated to prevent certain age-related pathologies in peripheral tissues in a progeroid mouse model (Baker et al., 2011). Recent studies from various laboratories suggest that senescent cells are also present in the aging brain and in

conjunction with age-related neurodegenerative diseases. Neurodegeneration associated with these conditions is closely tied to neuroinflammation. A potential source of this neuroinflammation is the pro-inflammatory SASP from senescent brain cells. Although there are still many unanswered questions involving brain cell senescence, the ability to critically test the idea that senescent cells can cause or contribute to age-related neuropathology will allow the identification of an important and novel target (senescent cells) for pharmacological interventions aimed at amelioration of age-related neurodegeneration.

Acknowledgments

Financial support was provided by an Ellison Senior Scholar fellowship (JKA), the 2014 Buck Impact Circle (JKA/JC), NIA PPG AG025901 (JKA/JC), CIRM TG2-01155 (SJC), American Italian Cancer Foundation (MD), T32 AG000266 (GW), and NIA R37 AG009909 (JC).

References

- Abazov VM, Abbott B, Abolins M, Acharya BS, Adams M, Adams T, Aguilo E, Ahn SH, Ahsan M, Alexeev GD, et al. Measurement of the semileptonic branching ratio of $B_{\{s\};\{0\}}$ to an orbitally excited $D_{\{s\};\{**\}}$ state: $Br(B_{\{s\};\{0\}} \rightarrow D_{\{s1\};\{-\}}(2536)\mu; \{+\} \nu_X)$. *Physical review letters*. 2009; 102:051801. [PubMed: 19257502]
- Adams PD. Healing and hurting: molecular mechanisms, functions, and pathologies of cellular senescence. *Molecular cell*. 2009; 36:2–14. [PubMed: 19818705]
- Allen NJ, Barres BA. Neuroscience: Glia - more than just brain glue. *Nature*. 2009; 457:675–677. [PubMed: 19194443]
- Bachstetter AD, Xing B, de Almeida L, Dimayuga ER, Watterson DM, Van Eldik LJ. Microglial p38alpha MAPK is a key regulator of proinflammatory cytokine up-regulation induced by toll-like receptor (TLR) ligands or beta-amyloid (Abeta). *Journal of neuroinflammation*. 2011; 8:79. [PubMed: 21733175]
- Baker DJ, Wijshake T, Tchkonja T, LeBrasseur NK, Childs BG, van de Sluis B, Kirkland JL, van Deursen JM. Clearance of p16Ink4a-positive senescent cells delays ageing-associated disorders. *Nature*. 2011; 479:232–236. [PubMed: 22048312]
- Ben-Porath I, Weinberg RA. The signals and pathways activating cellular senescence. *The international journal of biochemistry & cell biology*. 2005; 37:961–976. [PubMed: 15743671]
- Benarroch EE. Neuron-astrocyte interactions: partnership for normal function and disease in the central nervous system. *Mayo Clinic proceedings*. 2005; 80:1326–1338. [PubMed: 16212146]
- Bhat R, Crowe EP, Bitto A, Moh M, Katsetos CD, Garcia FU, Johnson FB, Trojanowski JQ, Sell C, Torres C. Astrocyte senescence as a component of Alzheimer's disease. *PloS one*. 2012; 7:e45069. [PubMed: 22984612]
- Bitto A, Sell C, Crowe E, Lorenzini A, Malaguti M, Hrelia S, Torres C. Stress-induced senescence in human and rodent astrocytes. *Experimental cell research*. 2010; 316:2961–2968. [PubMed: 20620137]
- Braig M, Schmitt CA. Oncogene-induced senescence: putting the brakes on tumor development. *Cancer research*. 2006; 66:2881–2884. [PubMed: 16540631]
- Campisi J. From cells to organisms: can we learn about aging from cells in culture? *Experimental gerontology*. 2001; 36:607–618. [PubMed: 11295503]
- Campisi J. Cellular senescence and apoptosis: how cellular responses might influence aging phenotypes. *Experimental gerontology*. 2003; 38:5–11. [PubMed: 12543256]
- Campisi J. Aging and cancer cell biology, 2007. *Aging cell*. 2007; 6:261–263. [PubMed: 17517036]
- Campisi J, d'Adda di Fagagna F. Cellular senescence: when bad things happen to good cells. *Nature reviews Molecular cell biology*. 2007; 8:729–740.

- Chen Y, Swanson RA. Astrocytes and brain injury. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. 2003; 23:137–149.
- Chung HY, Cesari M, Anton S, Marzetti E, Giovannini S, Seo AY, Carter C, Yu BP, Leeuwenburgh C. Molecular inflammation: underpinnings of aging and age-related diseases. *Ageing research reviews*. 2009; 8:18–30. [PubMed: 18692159]
- Collado M, Serrano M. Senescence in tumours: evidence from mice and humans. *Nature reviews Cancer*. 2010; 10:51–57.
- Conde JR, Streit WJ. Effect of aging on the microglial response to peripheral nerve injury. *Neurobiology of aging*. 2006; 27:1451–1461. [PubMed: 16159684]
- Coppe JP, Patil CK, Rodier F, Krtolica A, Beausejour CM, Parrinello S, Hodgson JG, Chin K, Desprez PY, Campisi J. A human-like senescence-associated secretory phenotype is conserved in mouse cells dependent on physiological oxygen. *PloS one*. 2010; 5:e9188. [PubMed: 20169192]
- Coppe JP, Patil CK, Rodier F, Sun Y, Munoz DP, Goldstein J, Nelson PS, Desprez PY, Campisi J. Senescence-associated secretory phenotypes reveal cell-nonautonomous functions of oncogenic RAS and the p53 tumor suppressor. *PLoS biology*. 2008; 6:2853–2868. [PubMed: 19053174]
- Coppe JP, Rodier F, Patil CK, Freund A, Desprez PY, Campisi J. Tumor suppressor and aging biomarker p16(INK4a) induces cellular senescence without the associated inflammatory secretory phenotype. *The Journal of biological chemistry*. 2011; 286:36396–36403. [PubMed: 21880712]
- Davalos AR, Coppe JP, Campisi J, Desprez PY. Senescent cells as a source of inflammatory factors for tumor progression. *Cancer metastasis reviews*. 2010; 29:273–283. [PubMed: 20390322]
- Dimri GP, Lee X, Basile G, Acosta M, Scott G, Roskelley C, Medrano EE, Linskens M, Rubelj I, Pereira-Smith O, et al. A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proceedings of the National Academy of Sciences of the United States of America*. 1995; 92:9363–9367. [PubMed: 7568133]
- Doorn KJ, Lucassen PJ, Boddeke HW, Prins M, Berendse HW, Drukarch B, van Eruslimsky JD, Kurz DJ. Cellular senescence in vivo: its relevance in ageing and cardiovascular disease. *Experimental gerontology*. 2005; 40:634–642. [PubMed: 15970413]
- Ferrucci L, Corsi A, Lauretani F, Bandinelli S, Bartali B, Taub DD, Guralnik JM, Longo DL. The origins of age-related proinflammatory state. *Blood*. 2005; 105:2294–2299. [PubMed: 15572589]
- Finch C. Neuron atrophy during aging: programmed or sporadic? *TINS*. 1993; 16:104–110. [PubMed: 7681233]
- Flanary B. The role of microglial cellular senescence in the aging and Alzheimer diseased brain. *Rejuvenation research*. 2005; 8:82–85. [PubMed: 15929715]
- Flanary BE, Streit WJ. Telomeres shorten with age in rat cerebellum and cortex in vivo. *Journal of anti-aging medicine*. 2003; 6:299–308. [PubMed: 15142431]
- Flanary BE, Streit WJ. Progressive telomere shortening occurs in cultured rat microglia, but not astrocytes. *Glia*. 2004; 45:75–88. [PubMed: 14648548]
- Franceschi C, Capri M, Monti D, Giunta S, Olivieri F, Sevini F, Panourgia MP, Invidia L, Celani L, Scurti M, et al. Inflammaging and anti-inflammaging: a systemic perspective on aging and longevity emerged from studies in humans. *Mechanisms of ageing and development*. 2007; 128:92–105. [PubMed: 17116321]
- Frank-Cannon TC, Alto LT, McAlpine FE, Tansey MG. Does neuroinflammation fan the flame in neurodegenerative diseases? *Molecular neurodegeneration*. 2009; 4:47. [PubMed: 19917131]
- Freund A, Orjalo AV, Desprez PY, Campisi J. Inflammatory networks during cellular senescence: causes and consequences. *Trends Mol Med*. 2010; 16:238–246. [PubMed: 20444648]
- Freund A, Patil CK, Campisi J. p38MAPK is a novel DNA damage response-independent regulator of the senescence-associated secretory phenotype. *The EMBO journal*. 2011; 30:1536–1548. [PubMed: 21399611]
- Guney I, Wu S, Sedivy JM. Reduced c-Myc signaling triggers telomere-independent senescence by regulating Bmi-1 and p16(INK4a). *Proceedings of the National Academy of Sciences of the United States of America*. 2006; 103:3645–3650. [PubMed: 16537449]
- Herbig U, Sedivy JM. Regulation of growth arrest in senescence: telomere damage is not the end of the story. *Mechanisms of ageing and development*. 2006; 127:16–24. [PubMed: 16229875]

- Jeyapalan JC, Ferreira M, Sedivy JM, Herbig U. Accumulation of senescent cells in mitotic tissue of aging primates. *Mechanisms of ageing and development*. 2007; 128:36–44. [PubMed: 17116315]
- Jurk D, Wang C, Miwa S, Maddick M, Korolchuk V, Tzolou A, Gonos ES, Thrasivoulou C, Saffrey MJ, Cameron K, et al. Postmitotic neurons develop a p21-dependent senescence-like phenotype driven by a DNA damage response. *Aging cell*. 2012; 11:996–1004. [PubMed: 22882466]
- Kishi S. Functional aging and gradual senescence in zebrafish. *Annals of the New York Academy of Sciences*. 2004; 1019:521–526. [PubMed: 15247079]
- Kreutzberg GW. Microglia: a sensor for pathological events in the CNS. *Trends in neurosciences*. 1996; 19:312–318. [PubMed: 8843599]
- Krizhanovsky V, Xue W, Zender L, Yon M, Hernando E, Lowe SW. Implications of cellular senescence in tissue damage response, tumor suppression, and stem cell biology. *Cold Spring Harbor symposia on quantitative biology*. 2008; 73:513–522.
- Krtolica A, Parrinello S, Lockett S, Desprez PY, Campisi J. Senescent fibroblasts promote epithelial cell growth and tumorigenesis: a link between cancer and aging. *Proceedings of the National Academy of Sciences of the United States of America*. 2001; 98:12072–12077. [PubMed: 11593017]
- Lucin KM, Wyss-Coray T. Immune activation in brain aging and neurodegeneration: too much or too little? *Neuron*. 2009; 64:110–122. [PubMed: 19840553]
- Magistretti PJ. Neuron-glia metabolic coupling and plasticity. *The Journal of experimental biology*. 2006; 209:2304–2311. [PubMed: 16731806]
- McGeer PL, McGeer EG. Glial cell reactions in neurodegenerative diseases: pathophysiology and therapeutic interventions. *Alzheimer disease and associated disorders*. 1998; 12(Suppl 2):S1–6. [PubMed: 9769023]
- Melk A, Kittikowit W, Sandhu I, Halloran KM, Grimm P, Schmidt BM, Halloran PF. Cell senescence in rat kidneys in vivo increases with growth and age despite lack of telomere shortening. *Kidney international*. 2003; 63:2134–2143. [PubMed: 12753300]
- Molofsky AV, Slutsky SG, Joseph NM, He S, Pardal R, Krishnamurthy J, Sharpless NE, Morrison SJ. Increasing p16INK4a expression decreases forebrain progenitors and neurogenesis during ageing. *Nature*. 2006; 443:448–452. [PubMed: 16957738]
- Nathan C, Ding A. Nonresolving inflammation. *Cell*. 2010; 140:871–882. [PubMed: 20303877]
- Nichols NR, Day JR, Laping NJ, Johnson SA, Finch CE. GFAP mRNA increases with age in rat and human brain. *Neurobiology of aging*. 1993; 14:421–429. [PubMed: 8247224]
- Ohtani N, Yamakoshi K, Takahashi A, Hara E. The p16INK4a-RB pathway: molecular link between cellular senescence and tumor suppression. *The journal of medical investigation : JMI*. 2004; 51:146–153. [PubMed: 15460900]
- Paradis V, Youssef N, Dargere D, Ba N, Bonvoust F, Deschatrette J, Bedossa P. Replicative senescence in normal liver, chronic hepatitis C, and hepatocellular carcinomas. *Human pathology*. 2001; 32:327–332. [PubMed: 11274643]
- Parrinello S, Coppe JP, Krtolica A, Campisi J. Stromal-epithelial interactions in aging and cancer: senescent fibroblasts alter epithelial cell differentiation. *Journal of cell science*. 2005; 118:485–496. [PubMed: 15657080]
- Passos JF, Von Zglinicki T. Oxygen free radicals in cell senescence: are they signal transducers? *Free radical research*. 2006; 40:1277–1283. [PubMed: 17090417]
- Pertusa M, Garcia-Matas S, Rodriguez-Farre E, Sanfeliu C, Cristofol R. Astrocytes aged in vitro show a decreased neuroprotective capacity. *Journal of neurochemistry*. 2007; 101:794–805. [PubMed: 17250685]
- Prieri F, Gresser E, Le Dreau Y, Obiols J, Kister J. New method of simulation to evaluate the sensitivity to oxidation of lubricating oils: an aging cell coupled with fourier transform infrared spectroscopy. *Applied spectroscopy*. 2008; 62:810–816. [PubMed: 18935833]
- Ransom B, Behar T, Nedergaard M. New roles for astrocytes (stars at last). *Trends in neurosciences*. 2003; 26:520–522. [PubMed: 14522143]
- Rodier F, Coppe JP, Patil CK, Hoeijmakers WA, Munoz DP, Raza SR, Freund A, Campeau E, Davalos AR, Campisi J. Persistent DNA damage signalling triggers senescence-associated inflammatory cytokine secretion. *Nature cell biology*. 2009; 11:973–979.

- Rodier F, Kim SH, Nijjar T, Yaswen P, Campisi J. Cancer and aging: the importance of telomeres in genome maintenance. *The international journal of biochemistry & cell biology*. 2005; 37:977–990. [PubMed: 15743672]
- Rodier F, Munoz DP, Teachenor R, Chu V, Le O, Bhaumik D, Coppe JP, Campeau E, Beausejour CM, Kim SH, et al. DNA-SCARS: distinct nuclear structures that sustain damage-induced senescence growth arrest and inflammatory cytokine secretion. *Journal of cell science*. 2011; 124:68–81. [PubMed: 21118958]
- Rose MR. Evolution and physiology. *Science*. 1991; 254:448. [PubMed: 17742234]
- Sawada M, Sawada H, Nagatsu T. Effects of aging on neuroprotective and neurotoxic properties of microglia in neurodegenerative diseases. *Neuro-degenerative diseases*. 2008; 5:254–256. [PubMed: 18322405]
- Shay JW, Wright WE. Senescence and immortalization: role of telomeres and telomerase. *Carcinogenesis*. 2005; 26:867–874. [PubMed: 15471900]
- Streit WJ. Microglia and the response to brain injury. Ernst Schering Research Foundation workshop. 2002a:11–24. [PubMed: 12066409]
- Streit WJ. Microglia as neuroprotective, immunocompetent cells of the CNS. *Glia*. 2002b; 40:133–139. [PubMed: 12379901]
- Streit WJ, Sammons NW, Kuhns AJ, Sparks DL. Dystrophic microglia in the aging human brain. *Glia*. 2004; 45:208–212. [PubMed: 14730714]
- Sugama S. Stress-induced microglial activation may facilitate the progression of neurodegenerative disorders. *Medical hypotheses*. 2009; 73:1031–1034. [PubMed: 19556067]
- Toussaint O, Medrano EE, von Zglinicki T. Cellular and molecular mechanisms of stress-induced premature senescence (SIPS) of human diploid fibroblasts and melanocytes. *Experimental gerontology*. 2000; 35:927–945. [PubMed: 11121681]
- Tsai KK, Chuang EY, Little JB, Yuan ZM. Cellular mechanisms for low-dose ionizing radiation-induced perturbation of the breast tissue microenvironment. *Cancer research*. 2005; 65:6734–6744. [PubMed: 16061655]
- Vaupel JW. Biodemography of human ageing. *Nature*. 2010; 464:536–542. [PubMed: 20336136]
- Voghel G, Thorin-Trescases N, Farhat N, Nguyen A, Villeneuve L, Mamarbachi AM, Fortier A, Perrault LP, Carrier M, Thorin E. Cellular senescence in endothelial cells from atherosclerotic patients is accelerated by oxidative stress associated with cardiovascular risk factors. *Mechanisms of ageing and development*. 2007; 128:662–671. [PubMed: 18022214]
- Wajapeyee N, Serra RW, Zhu X, Mahalingam M, Green MR. Oncogenic BRAF induces senescence and apoptosis through pathways mediated by the secreted protein IGFBP7. *Cell*. 2008; 132:363–374. [PubMed: 18267069]
- Yankner BA, Lu T, Loerch P. The aging brain. *Annual review of pathology*. 2008; 3:41–66.
- Yu HM, Zhao YM, Luo XG, Feng Y, Ren Y, Shang H, He ZY, Luo XM, Chen SD, Wang XY. Repeated lipopolysaccharide stimulation induces cellular senescence in BV2 cells. *Neuroimmunomodulation*. 2012; 19:131–136. [PubMed: 22248729]
- Zlokovic BV. The blood-brain barrier in health and chronic neurodegenerative disorders. *Neuron*. 2008; 57:178–201. [PubMed: 18215617]
- Zou Y, Zhang N, Ellerby LM, Davalos AR, Zeng X, Campisi J, Desprez PY. Responses of human embryonic stem cells and their differentiated progeny to ionizing radiation. *Biochemical and biophysical research communications*. 2012; 426:100–105. [PubMed: 22917535]

Highlights

- The role of cellular senescence in brain aging is just beginning to be explored.
- Cellular senescence appears to involve proliferative cell types in the brain.
- These cells are characterized by a SASP that can contribute to brain pathology.
- Senescent cell elimination could constitute a novel therapeutic for brain aging.

Fig. 1.