CORRESPONDENCE

The Prognostic Significance of Respiratory Rate in Patients With Pneumonia: A Retrospective Analysis of Data From 705 928 Hospitalized Patients in Germany From 2010–2012

by PD Dr. med. Richard Strauß, Prof. Dr. med. Santiago Ewig, Dr. med. Klaus Richter, Dr. rer. nat. Thomas König, PD Dr. med. Günther Heller, Prof. Dr. med. Torsten T. Bauer in issue 29–30/2014

Far Removed From Patients

Pointing out that measuring the respiratory rate in severely ill patients—which includes patients with pneumonia—is a useful approach to treatment does undoubtedly have its merits. The fact that this parameter is also of prognostic significance would be more obvious to readers, however, if any mention at all had been made of the pathophysiology of this phenomenon. It is concerning that this article may have been written only to point out a self evident clinical diagnosis.

Changes in the respiratory rate may have different causes even in pneumonia, a fact that doctors have been familiar with for some 120 years (1, 2). It is scientifically correct to present a biological parameter in a statistical manner only, but it is clinically pointless, because this does not really get to the bottom of the problem. The estimated prognosis is of little help for patients. What is important is that a doctor draws diagnostic or therapeutic conclusions from what he or she sees. The article does not make any mention of this. It is an example of a way of thinking that is removed from patients and merely focused on organizational economic principles.

DOI: 10.3238/arztebl.2015.0173a

REFERENCES

1. Leube W: Diagnose der inneren Krankheiten. Leipzig: F.C.W. Vogel 1895: 127.
2. Wyngaarden JB, Smith LH: Textbook of medicine. Philadelphia: W. B. Saunders 1985: 1501.
3. Strauß R, Ewig S, Richter K, König T, Heller G, Bauer TT: The prognostic significance of respiratory rate in patients with pneumonia—a retrospective analysis of data from 705 928 hospitalized patients in Germany from 2010–2012. Dtsch Arztebl Int 2014; 111: 503–8.

Prof. Dr. med. Peter von Wichert, Dir. emer
Zentrum Innere Medizin
Philipps Universität Marburg
vonwichert@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Significance of the Respiratory Rate in Emergency Medicine

The authors of the study show that the respiratory rate is an independent risk marker for hospital mortality in community acquired pneumonia (CAP). We wish to add that the respiratory rate is an important risk marker for the

purposes of sensitively identifying emergency patients at vital risk (1). This idea is only insufficiently implemented in clinical practice in Germany and Switzerland: for this reason, we introduced a five step triage system (the emergency severity index, ESI) in the emergency departments at Nuremberg and Basel university medical centers, so as to identify early on critically ill patients by using the obligatory measuring of the respiratory rate, among others.

Furthermore, an interprofessional CAP treatment concept was introduced in Nuremberg, in order to facilitate the recommended treatment within the recommended intervals (local adaptation of CAP guidelines, introduction of checklists, further medical educational events, interactive workshops, process analyses, feedback sessions). The implementation of management bundles has significantly improved the quality of care for affected patients and has reduced in-hospital mortality of CAP patients by 21% (1). On the basis of our experiences we suggest further standardization in medical emergencies that require time critical action.

In the future it would be helpful to establish diagnostic instruments to facilitate the measuring of the respiratory rate in the initial diagnostic evaluation. Appropriate equipment is available in the form of devices that measure oxygen saturation or—by using a video analysis system—the respiratory rate (3). These devices will, however, still have to be validated in the setting of emergency and acute medicine, and their diagnostic importance will need to be evaluated in relevant studies. Measuring the respiratory rate is absolutely crucial in any emergency patient.

DOI: 10.3238/arztebl.2015.0173b

REFERENCES

1. Hortmann M, Heppner HJ, Popp S, Lad T, Christ M: Reduction of mortality in community-acquired pneumonia after implementing standardized care bundles in the emergency department. Eur J Emerg Med 2014; 21: 429–35.
2. Grossmann FF, Nickel CH, Christ M, Schneider K, Spirig R, Bingisser R: Transporting clinical tools to new settings: cultural adaptation and validation of the emergency severity index in German. Ann Emerg Med 2011; 57: 257–64.
3. Addison PS, Watson JN, Mestek ML, Ochs JP, Uribe AA, Bergese SD: Pulse oximetry-derived respiratory rate in general care floor patients. J Clin Monit Comput 2014; epub ahead of print.
4. Strauß R, Ewig S, Richter K, König T, Heller G, Bauer TT: The prognostic significance of respiratory rate in patients with pneumonia—a retrospective analysis of data from 705 928 hospitalized patients in Germany from 2010–2012. Dtsch Arztebl Int 2014; 111: 503–8.

Dr. med. Marcus Hortmann
Universitäts-Herzzentrum Freiburg,
Klinik für Kardiologie und Angiologie I
PD Dr. med. Christian H. Nickel
Prof. Dr. med. Roland Bingisser
Interdisziplinäre Notfallstation, Universitätsspital Basel

Prof. Dr. med. Michael Christ
Klinik für Notfallmedizin und Internistische Intensivmedizin
Paracelsus Medizinische Privatuniversität Nürnberg
michael.christ@klinikum-nuernberg.de

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

We thank Hortmann et al. for their supplementary and positive assessment of the respiratory rate as a risk marker for emergency patients at risk. They have obviously also experienced that knowledge/ awareness of this fact is still insufficiently put into clinical practice. Since the respiratory rate is a component in different established prognostic instruments used in acutely ill patients (among others, of the emergency severity index (ESI) or confusion, respiratory rate, blood pressure (CRB)-65 index), a valid measurement of the respiratory rate is vital for using these instruments.

The systematic use of validated instruments for the structured support of medical triage decisions, such as the colleagues in Nuremberg's and Basel's emergency department successfully practise, has as its vital prerequisite that the respiratory rate is measured reliably. Hortmann et al. have shown impressively how the successful implementation of such an instrument (the ESI, in this case) in the context of a structured multidimensional healthcare concept can make a crucial contribution to improving the quality of healthcare. We can only congratulate our colleagues on their reported results in CAP (1).

If the risk assessment is complemented by appropriate subsequent clinical actions that are adapted to the individual risk (for example, intensive monitoring, rapid treatment, discharge, admission), patients will benefit. However, as Professor von Wichert points out in his letter to the editor, patients will not benefit from the assessment alone—something we addressed in the discussion section in our article: “Thus, abnormal respiratory rates should be further investigated while these patients require close monitoring (e.g. on display devices).” Current guidelines explicitly recommend the use of the CRB-65 index, and thus the respiratory rate, for the purposes of risk assessment and decision support (for example, inpatient admission, intensive monitoring) and thus is of immediate benefit for patients (2, 3). It would have exceeded the scope and objectives of our original article to explain the pathophysiology of respiratory rate regulation. We therefore confined ourselves to mentioning hypercapnia,

hypoxia, and metabolic acidosis as factors that—among others—can lead to an increase in the respiratory rate.

Easy to use automated methods for measuring the respiratory rate can improve acceptance in everyday clinical practice. There are currently different developments in new measuring techniques that—as long as they yield valid results in a cost effective fashion—could be used in emergency medicine in the future. In addition to the approaches mentioned by Hortmann et al., these could also include procedures for analyzing acoustic phenomena of respiration (4).

DOI: 10.3238/arztebl.2015.0174

REFERENCES

1. Hortmann M, Heppner HJ, Popp S, et al.: Reduction of mortality in community-acquired pneumonia after implementing standardized care bundles in the emergency department. *Eur J Emerg Med* 2014; 21: 429–35.
2. Hoffken G, Lorenz J, Kern W, et al.: Epidemiology, diagnosis, antimicrobial therapy and management of community-acquired pneumonia and lower respiratory tract infections in adults. Guidelines of the Paul-Ehrlich-Society for Chemotherapy, the German Respiratory Society, the German Society for Infectiology and the Competence Network CAPNETZ Germany. *Pneumologie* 2009; 63: e1–68.
3. Woodhead M, Blasi F, Ewig S, et al.: Guidelines for the management of adult lower respiratory tract infections—full version. *Clin Microbiol Infect* 2011; 17: e1–59.
4. Sahgal N: Monitoring and analysis of lung sounds remotely. *Int J Chron Obstruct Pulmon Dis* 2011; 6: 407–12.
5. Strauß R, Ewig S, Richter K, König T, Heller G, Bauer TT: The prognostic significance of respiratory rate in patients with pneumonia—a retrospective analysis of data from 705 928 hospitalized patients in Germany from 2010–2012. *Dtsch Arztebl Int* 2014; 111: 503–8.

On behalf of the authors:

PD Dr. med. Richard Strauß
Universitätsklinikum Erlangen
Medizinische Klinik 1
richard.strauss@uk-erlangen.de

Conflict of interest statement

Dr Strauß has received fees for consultancy from Swedish Orphan Biovitrum, Biotest and Pfizer. He has received reimbursements for congress participation fees from Bayer Vital, Pfizer and InfetcoPharm. He has received reimbursement for travel and accommodation expenses and fees for the preparation of scientific meetings from Bayer Vital, Pfizer, InfetcoPharm, and MSD.

The remaining authors declare that no conflict of interest exists.