Published in final edited form as:

Clin Sports Med. 2015 April ; 34(2): 213–231. doi:10.1016/j.csm.2014.12.005.

Current and emerging rehabilitation for concussion: A review of the evidence

Steven P. Broglio, PhD, ATC^{1,2}, Michael W. Collins, PhD³, Richelle M. Williams, MS, ATC¹, Anne Mucha, DPT^{3,4}, and Anthony Kontos, PhD³

¹School of Kinesiology, University of Michigan, Ann Arbor, MI

²University of Michigan Injury Center

³Department of Orthopaedic Surgery/UPMC Concussion Program- University of Pittsburgh, Pittsburgh, PA

⁴UPMC Centers for Rehab Services- UPMC, Pittsburgh, PA

Abstract

Concussion is one of the most hotly debated topics in sports medicine today. Research surrounding concussion has experienced significant growth recently especially in the areas of incidence, assessment, and recovery. However, there is limited research on the most effective rehabilitation approaches for this injury. This review evaluates the current literature for evidence for and against physical and cognitive rest and the emerging areas targeting vestibular, oculomotor, and pharmacological interventions for the rehabilitation of sport-related concussion.

Keywords

Concussion; physical rest; cognitive rest; vestibular rehabilitation; pharmacological interventions

Introduction

The clinical signs and symptoms of sport concussion have long been recognized as (1, 2), brought about by an extrinsic force applied directly or indirectly to the head or body (3). Much of the scientific literature surrounding this injury has focused on injury incidence(4), assessment tools(5, 6), and recovery patterns among athletes(7). Absent from the literature are reviews of empirical studies assessing the effectiveness of different rehabilitation approaches for concussed patients. Therefore, this paper will review and evaluate the evidence supporting consensus-based standard of care (e.g., physical and cognitive rest) and

© 2014 Elsevier Inc. All rights reserved.

Corresponding Author Steven P. Broglio PhD, ATC, 401 Washtenaw Ave, Ann Arbor, MI 48109, Phone: 734.764.9669, Fax: 734.936.1925, broglio@umich.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

emerging, targeted (e.g., vestibular, oculomotor, exertional, pharmacological) rehabilitation approaches for concussion based on an evolving model of clinical concussion care(8).

The concept of physical and cognitive rest as the cornerstone of concussion management was developed and by the International Concussion in Sport Group, and currently states “The cornerstone of concussion management is physical and cognitive rest until the acute symptoms resolve and then a graded program of exertion prior to medical clearance and return to play(3).” The rationale for rest asserts that during the acute (1-7 days- possibly longer in youth) post-injury period of increased metabolic demand and limited adenosine triphosphate (ATP) reserves, non-essential activity draws oxygen and glycogen away from injured neurons. The Concussion in Sport Group recommendation has been interpreted by many clinicians to mean that all concussed athletes should be restricted from all physical and cognitive activity until symptoms resolve. At which point, the athlete could be cleared to begin a return to play progression. This “shut down” or “dark closet” approach following concussion is wrought with potential pitfalls for patients including hyperawareness of symptoms, somatization, social isolation, and other potential co-morbid concerns. Citing the risk for prolonged and exacerbated symptoms that may not be directly related to the concussive injury, other medical organizations have recommended that athletes be permitted to engage in limited physical and cognitive activity so long as it does not worsen symptoms(9).

These two perspectives regarding strict rest versus physical and cognitive activity as tolerated are seemingly at odds with each other, in part because there is no agreed upon definition of what constitutes rest following a concussion in the literature. Such recommendations are also limited as they do not take into account the individualized nature of the injury, potential risk factors that may influence outcomes, and differential responses to recovery. Moreover, and most importantly, there are no known prospective randomized control trials evaluating rest in concussed athletes immediately following a concussion(10). In fact, the evidence for physical and cognitive rest is limited, relying on observational studies and studies of patients from sports medicine clinics during the sub-acute stage (11, 12). With a dearth of literature to support clinical guidelines, expert consensus has been used in its place.

The premise that rest is the most effective management strategy for all concussed patients assumes that all concussions are alike, yet concussion recovery is known to be influenced by several modifying factors including sex(13), concussion history(14), and age(15). Even for injuries occurring within these populations, concussions manifest in varied symptoms (e.g., headache, dizziness, foggiess), cognitive (e.g., memory, reaction time, processing speed) (16), psychological (e.g., depression, anxiety)(16), and vestibular (e.g. dizziness, imbalance, gait, vestibulo-ocular) (17) impairments. As such, this highly individualized injury results in a varied injury presentation, indicating no single rehabilitation strategy will be effective for all patients following concussion necessitating distinct treatment(8).

Physical Rest

Declines in neurocognitive function and motor control and increases in self-report symptoms following concussion are well documented(5, 18). Among the most commonly reported symptoms are headache, dizziness, and confusion immediately following a concussion (19-25). Other research has reported increased rates of depression and fatigue among the same cohort (26). Between 80-90% of concussed individuals will return to pre-injury levels of functioning within two weeks without intervention, but a small percentage (2.5%) will remain symptomatic 45 days post-injury, despite resolution of other objective measures (e.g. neurocognitive and balance assessments)(27). Therefore, management of athletes falling within and outside the range of normal recovery may require different approaches.

Evidence for Physical Rest

Consensus supporting physical rest recommendations are partly predicated on risk management and animal studies demonstrating impaired recovery with the early onset of physical activity. Although secondary to restricting activity to facilitate recovery, restricting physical activity to reduce risk for a second injury has broad support as the sole means to eliminate Second Impact Syndrome(28). Even in absence of a catastrophic outcome, animal models show metabolic dysfunction in the days immediately following injury (1, 29) with an increased energy demand within the cerebral tissue as ion imbalances are returned to pre-injury levels. During this time, the risk for second injury appears to be highest (23) and an additional injury sustained shortly after the first increases recovery time and impairs the ability to learn among rodents(30). Similar metabolic dysfunction findings were reported by Vagnozzi et al among concussed athletes. That is, altered cerebral metabolism lasting up to 30 days was documented using magnetic resonance spectroscopy imaging in concussed athletes(31) and up to 45 days in those sustaining a second injury prior to resolution of the first (32).

In addition subsequent injury risk and prolonged recovery brought about by sport participation, unrestricted physical activity in a controlled and safe environment during the acute recovery stage may be detrimental. For example, rats exposed to a fluid percussion injury and provided unrestricted running wheel access within the first 6 days of injury showed poorer performance on a cognitive task (i.e. Morris water maze) compared to similarly injured rats that were restricted from activity until day 14 post-injury(33). Additionally, a review of medical records from 159 concussed patients found that those returning to play prior to concussion symptom resolution reported worsening concussion related symptoms (34). The mechanistic underpinnings explaining these findings are not entirely clear, but it is possible that early exercise may draw energy (ie glycogen) away from the brain and inhibit the recovery process.

Evidence for Physical Activity

The evidence supporting physical activity following concussion is sparse, but other medical literature suggests that withholding injured, but non-concussed, athletes from exercise increases reports of depression, anxiety, and lower self-esteem when evaluated both at the time of injury and eight weeks following (35). Injured (non-concussed) high school athletes

missing a minimum of three weeks of athletic participation also showed higher rates of depression than non-injured athletes. The authors indicated that high levels of athletic identity partly explained the finding(36). Moreover, onset of migraine has been shown to occur in patients with limited or minimal physical activity(37). Ultimately, removing an athlete from sport may increase the risk for depression and other concussion-like symptoms to develop, yet the point at which an athlete can begin physical activity following concussion is unclear.

In the single human study evaluating exercise shortly after concussion, 95 concussed student-athletes retrospectively self-reported physical and cognitive activity in the 30 days following injury and compared the findings to a neurocognitive assessment. Each athlete was categorized into one of five groups ranging from “no school or exercise activity” to “school activity and participation in sports games” and completed a computerized neurocognitive assessment for both cognitive functioning and self-report symptoms. The results indicated that athletes engaging in a medium level of physical and cognitive activity (i.e. school activity and light activity at home (e.g. slow jogging, mowing the lawn, etc.)) performed better on the neurocognitive test than those with no physical and cognitive activity and those with the reporting the highest levels of physical and cognitive activity. (38). These findings should be cautiously interpreted however, as the physical and cognitive activity was self-report recall by the injured athlete. In addition, it is not known at what point post-injury the athletes elected to begin physical activity. Coupled with the previously discussed animal work, this investigation set the groundwork to suggest that unrestricted exercise in the immediate acute phase of concussion recovery may increase the risk of subsequent injury and/or delay recovery, yet some level of exercise may be beneficial to the recovery process once the athlete has moved beyond the acute injury stage.

When dealing with athletes continuing to experience concussion like symptoms beyond the acute injury stage, stronger literature is available showing the benefit of physical activity as a means to mitigate symptom reports. Leddy et al (39) implemented a graded return to activity protocol on six concussed athletes that had been symptomatic for a minimum of six weeks (mean 19 weeks) following a concussive event. Once enrolled, the athletes were monitored for an additional two weeks, where there was no change in their symptom reports. They then began an exercise protocol five to six days a week with intensity monitored by heart rate. Following six weeks of the intervention, the athletes had a significant decrease in their symptom reports and were able to return to sport. Interestingly, a concussed non-athlete group completed an identical protocol, but did not show the same decline in symptom reports as the athlete cohort. A follow-up investigation enrolled 91 participants that had been experiencing symptoms following a concussion for a minimum of three weeks. Each participant completed a baseline graded exercise test, whereby 26 were able to reach maximum exertion. These individuals were determined to be experiencing symptoms related to something other than concussion, while 35 of the remaining 65 continued with the same heart rate based exercise protocol described above. A return to full functioning by means of the exercise protocol was achieved in 77% of the subset (n=27) (40).

Exercise has been proven to be a powerful modality for cognitive health(41), but the implementation of post-concussion exercise should be carefully considered relative to the

time from injury. The limited literature available to date suggests that athletes experiencing symptoms in the acute stage of injury should avoid full sport participation to avoid secondary injury, as well as exercise in a controlled environment as it may increase recovery time. Animal and retrospective human studies however, suggest that athletes continuing to report symptoms beyond the acute stage of injury may benefit from moderate levels of exercise. Lastly, those athletes that continue to report concussion related symptoms well beyond the acute stage of injury may benefit from a progressively intensive exercise protocol to return them to their sport.

Cognitive Rest and Academic Accommodations

Cognitive impairment following concussion is common among student-athletes and cognitive rest has been suggested to enhance recovery. The cognitive rest theory is based on the premise that increasing cognitive activities following concussion will increase symptom recovery time and prolong recovery. Cognitive rest includes the reduction of brain stimulating activities (e.g. television, videogames, school work, reading, and writing) and despite the limited data to support the use of cognitive rest, it is widely recommended in consensus statements and concussion guidelines(3, 9, 42-45).

To date, few studies have evaluated cognitive rest, however, these studies have found that increased cognitive activity does delay symptom recovery. Moser et al(11) studied 49 high school and collegiate athletes prescribing a minimum of a week of cognitive and physical rest. Both before and after rest periods, individuals performed the ImPACT and cognitive testing measures. The study concluded with a period of cognitive and physical rest with individuals showing increased performance on the ImPACT and cognitive testing, as well as decreased symptom reporting. Similarly, Brown et al (12) studied 335 patients (mean 15 years old) on level of cognitive activities between clinical visits, finding that longer concussion recovery time was related to higher cognitive activity levels. Indeed, those participating in lowest 50% of cognitive activity were completely asymptomatic within 100 to 150 days of injury, while those engaging in the 3rd and 4th quartiles of cognitive active took up to 300 and 500 days to recovery, respectively.

School is a major component to a student's life, requiring the attainment of new knowledge, development of academic skills, and diligent work to complete assignments and prepare for exams. To be successful in academic endeavors, students must engage in classroom learning requiring attention, material memory recall, critical thinking and problem solving. Students who sustain a sports-related concussion may also experience physical, mental, behavioral and social changes that impact their daily life and threaten a their ability to learn and succeed academically(46). Limiting to school activity is one mechanism that affords the injured athlete time to mitigate concussion related symptoms.

Despite limited research on cognitive load following concussion, it has been suggested that some concussed students may benefit from excused or reduced academics (e.g classroom attendance, homework, exams) immediately following injury(47-49). Returning to academic work while symptomatic may cause symptoms to worsen, resulting in a decline in academic performance(48, 50). While there are more formal accommodations available for long-term

or prolonged cases, temporary and targeted accommodations during the acute recovery time is an easy tool to assist a student's return to the classroom(47). Temporary accommodations may include, but are not limited to, excused absences, lighter homework, breaks during the day, starting later or ending the school day earlier and extended exam or homework dates(49). Once the athlete no longer reports concussion related symptoms, a transition period to partial and then full days is recommended (51).

Requiring a student-athlete to attend school immediately following a concussion may place him/her in a compromised academic position. Concussed student-athletes engaging in moderate to intense cognitive activity may have exacerbated symptoms (12), resulting in incomplete school work, making excused absences important to the recovery process. During this period, a time extension to complete academic assignments including homework and exams allows the student-athlete to make up missed schoolwork and take their time with completion of assignments. Implementing delayed testing or project due dates can help the student maintain good academic standing without the penalty of decreased scores. Temporary accommodations such as these are commonly implemented quickly and without burdensome paperwork. To ensure these tools are available, the concussion management teams should pre-arrange their use with school personnel as part of the concussion management plan prior to the athletic season.

Those individuals experiencing a prolonged concussion recovery or those with recurrent injuries may need additional testing to better accommodate or intervene with school as needed(51). In some cases, implementing an individualized academic plan can help with the management of accommodations. If a student-athlete is having symptoms or displays challenges greater than three weeks, a 504 plan may be implemented(47). A 504 Plan refers to the proper section of the Rehabilitation Act that provides medical need accommodations. In order for formal accommodations to be implemented the student would have to display mental impairments that limits greater than one major life activity(52). Clinicians may also consider an Individualized Education Plan (IEP) for those with prolonged concussion recovery. An IEP allows for the school personnel to collaborate with the physician, student-athlete and parents to create a plan that will best help that student receive special education(53). Both IEP's and 504 plans require extensive medical documentation and are a more permanent measure that are embedded into the school system documents, but allow for changes to the student-athlete's academic plan for classroom success.

Despite consensus for cognitive rest, it is important to note that prolonged cognitive rest and reduction of school events have the potential to exacerbate symptoms or cause negative mental health issues. Depression, behavioral and social issues have shown to increase following a concussion as well as many other injuries when the student-athlete is eliminated from team activities, school events and social outings (54). Decreasing school attendance and other social activities can negatively impact some student-athletes and prevent them from going through proper injury coping mechanisms. Decreased school attendance can also add an increased burden and sense of anxiety to the student-athlete because they are not attending school nor completing school assignments. The mental image of being behind in academics can create a highly anxious environment for student-athletes, especially those who already place high priority on increased academic achievement. Ultimately, the medical

team, in conjunction with a trained professional, should balance the neurocognitive and behavioral accommodations of the concussed student-athlete in a way that restricts cognitive activities that trigger or introduce symptom exacerbation, but allow for him/her to become involved in school activities again.

Vestibular and Oculomotor Rehabilitation

In a new clinical model of SRC (SRC) care, researchers have suggested that oculomotor and vestibular symptoms and impairment may constitute unique clinical subtypes of SRC- along with cognitive-fatigue, anxiety-mood, cervical, and post-traumatic migraine (PTM)(8). These clinical subtypes of concussion, which can occur concurrently or independently, require targeted therapies and treatments in order to be managed most effectively (8). For example, an athlete with an oculomotor concussion subtype will benefit most from vision and oculomotor-specific therapies. Without such targeted intervention strategies, an athlete may experience an unnecessarily prolonged recovery from SRC. In a prospective study of recovery times following SRC, researchers reported that 17% of athletes experience prolonged recovery lasting greater than 3 weeks (55). The identification of specific clinical subtypes of concussion together with the application of targeted treatments and rehabilitation strategies will yield the best clinical outcomes for athletes with SRC. Two clinical subtypes that have been associated with poor clinical outcomes, but that may be amenable to rehabilitation and treatment interventions are vestibular and oculomotor.

Vestibular and Oculomotor Impairment and Symptoms

Vestibular and oculomotor impairment and symptoms occur in approximately 60% of athletes following SRC (17). The vestibular system plays an integral role in balance function and in maintaining visual and spatial orientation. Sensory information from each inner ear is used to inform the adjustment of eye movements for clear, stable vision and to adjust muscle reactions of the head and body for balance and gait. Vestibular impairment and dysfunction may involve either the peripheral or central structures of the vestibulo-spinal system and may result in disequilibrium and impaired balance(56). In contrast, dizziness, vertigo, blurred/unstable vision, discomfort in busy environments and nausea often occur with disruption to the vestibulo-ocular system(56). Vestibular symptoms at the time of injury may predict prolonged recovery following SRC. In fact, Lau and colleagues(25) reported that on-field dizziness was the only significant predictor of a prolonged recovery (>21 days) following SRC. This expression of post-SRC dizziness acutely may be the result of disruption to the vestibulo-ocular and gaze stability systems at the time of the injury.

Similar to vestibular dysfunction, impairment in oculomotor control and visual dysfunction are observed frequently following SRC(57-61). Ciuffreda and colleagues(62) indicated that visual dysfunction involving accommodation, version and vergence, strabismus, and cranial nerve (CN) palsy occurred following mTBI. Symptoms attributed to poor oculomotor function may include blurred vision, diplopia, difficulty reading, eyestrain, headache, reading difficulties, and problems with visual scanning. Vestibular and oculomotor impairment and symptoms are prevalent following SRC and may play a role in prolonged recovery and related clinical outcomes. Therefore, we will present and discuss vestibular

rehabilitation and vision therapy interventions that can be used with athletes experiencing vestibular and oculomotor impairment and symptoms following SRC.

Vestibular Rehabilitation Interventions

There are many different types of vestibular rehabilitation interventions than be implemented to mitigate vestibular symptoms and dysfunction following SRC. Among the most common vestibular issues following SRC are benign paroxysmal positional vertigo (BPPV), vestibulo-ocular reflex (VOR) impairment, visual motion sensitivity, balance dysfunction, cervicogenic dizziness, and exercise-induced dizziness. Table 1 summarizes these and other vestibular problems along with targeted therapeutic interventions. It is important to note that these interventions should be performed by licensed physical therapists specializing in vestibular rehabilitation. Detailed descriptions of each

Benign paroxysmal positional vertigo is the most common disorder of the vestibular system and can occur post-traumatically after SRC. In BPPV, small calcium carbonate crystals (otoconia), which are normally housed in the otolith organs of the inner ear, become dislodged and relocate to one or more of the adjacent semicircular canals. With head movement in the plane of the affected semicircular canal, the otoconia shift position and create a false excitatory stimulus and resultant vertigo. Reproduction of vertigo and a characteristic nystagmus pattern during positional testing (Dix-Hallpike and Roll Test) is necessary to diagnose BPPV. Canalith repositioning maneuvers, designed to shift the displaced otoconia out of the affected semicircular canal is the treatment of choice for BPPV(63).

Gaze stability refers to the ability to maintain visual focus while the head is moving. While gaze stability is mediated by different vestibular and ocular motor systems depending on the velocity and context of the task, the VOR is the primary mechanism for maintaining eye position during head movement. The VOR is a fast-acting reflex which keeps the eyes stable by generating ocular movements precisely in proportion, but opposite in direction, from the head motion. In sport, where rapid acceleration and high velocity movement necessitates quick visual responses, intact VOR functioning is particularly important. When the VOR is impaired, visual blurring, dizziness poor visual focus and oscillopsia may occur with head motion. The responses of the VOR can be adapted through exercise designed to induce movement of a visual image on the retina. This motion – inducing retinal slip—is the primary error signal that drives adaptation of the VOR. Thus, vestibular physical therapy exercises for VOR adaptation require patients to maintain visual focus on a target while moving their head. VOR adaptation exercises are manipulated in multiple ways in order to gain maximal benefit, including varying target size and complexity, postures, duration, direction, amplitude and velocity.

Visual motion sensitivity refers to an increased sense of disorientation, dizziness or postural instability in situations with visual and vestibular conflict. It is thought to arise from inability of the central nervous system to effectively integrate sensory information, particularly vestibular information, creating over-reliance on vision. Patients with visual motion sensitivity become particularly symptomatic when exposed to visually disorienting stimuli or environments, such as malls, grocery stores or even busy patterns. Visual motion

sensitivity has also been described as “visual vertigo”, “space and motion discomfort” and “visual vestibular mismatch” in the literature. Visual motion sensitivity has been reported in patients following peripheral vestibular disorders (Pavlou 2013) and in those with migraine (Bronstein 2013) and anxiety (Furman 2005). It has also recently been recognized in patients following SRC(17). Treatment of visual motion sensitivity involves gradual and systematic exposure to provocative stimuli in order to habituate the abnormal responses. Because treatment for visual motion sensitivity has the potential to exacerbate symptoms from concussion, intervention should be introduced in a step-by-step progression that is carefully monitored by a trained vestibular therapist.

Restoring postural control, or balance, is an area of focus for vestibular rehabilitation following SRC(64-66). Because sensory organization is often impaired(67, 68) early after concussion, training the ability to effectively alternate between using visual, somatosensory and vestibular information for postural control is a key component of balance retraining. Graded exercises for sensory organization deficits involve manipulation of these three sensory systems. Examples of sensory organization training activities include performing tasks with eyes closed, while turning the head, with narrowed base of support, on an uneven or soft surface. In addition to sensory organization issues, several studies have shown patients following mTBI have greater difficulty maintaining balance under conditions of divided attention(69-71). Therefore, dual task condition practice, and dynamic balance activities may also be incorporated into vestibular rehabilitation. Although dizziness is most often attributed to vestibular system dysfunction, it may also arise from other impairment following SRC, which may be responsive to intervention. In cervicogenic dizziness, pathology in the cervical spine creates abnormal muscle activity in the deep layers of the upper cervical spine, responsible for providing proprioceptive input to the CNS. Dizziness is thought to occur due to the mismatch between aberrant cervical proprioceptive information in relation to vestibular and visual inputs. Because this cervical afferent information also participates in reflex activity for postural control and eye movements, imbalance and impaired eye movements may occur in addition to dizziness. Management of cervicogenic dizziness is directed toward therapies that treat the underlying cervical spine injury to normalize proprioceptive input with visual and vestibular information; along with treatment of any additional balance or oculomotor impairment through targeted exercises (Treleaven 2011).

Lastly, following concussion, dizziness may arise with exertional activity(72). In a study of soldiers following blast-related concussion, exercise- induced dizziness was categorized as one type of dizziness typically seen by physical therapists in vestibular rehabilitation. While there are no studies that confirm the etiology of exercise-induced dizziness, we postulate that inadequate response of the CNS to combined cardiovascular and vestibular/visual demand may be responsible. Anecdotally, we find in our clinic that stationary cardiovascular activities at high levels of exertion (e.g.: stationary cycling) rarely cause dizziness, whereas cardiovascular activity maintaining similar levels of exertion, when combined with motion (e.g.: forward/backward line drills) often produces significant levels of dizziness. Clearly, more research is needed to validate this hypothesis. Treatment of individuals with exercise-induced symptoms is controversial; however preliminary evidence suggests that graded exercise may be useful in modifying these post-concussive symptoms when chronic(39, 73).

The value of vestibular rehabilitation in managing individual vestibular conditions is well documented. A Cochrane review(74) concluded that there is moderate to strong evidence for efficacy of vestibular rehabilitation in improving VOR impairment and balance deficits due to peripheral vestibular dysfunction; and for the use of canalith repositioning maneuvers performed by vestibular therapists in the management of BPPV. Dizziness due to migraine, as well as patients with central vestibular dysfunction has been shown in studies to improve with vestibular physical therapy intervention (Whitney et al., 2000; Brown et al., 2006). Therapies for visual motion sensitivity, such as optokinetic stimulus exposure, have been shown to be effective with peripheral vestibular disorders. (Pavlou 2013) Several studies have investigated the efficacy of physical therapy treatment of the cervical spine for cervicogenic dizziness (Malmstrom 2007, Heidenreich 2008, Reid 2008), including a recent RCT (Reid 2014) demonstrating significant reduction in intensity and frequency of cervicogenic dizziness with two different manual therapy techniques over placebo. Although vestibular therapies have been shown to be beneficial in the treatment of various vestibular-related impairment, the evidence for utilizing vestibular physical therapy for impairment attributed to SRC is limited, and consists primarily of retrospective, cross sectional, and small cohort studies.

A recent study by Schneider and colleagues(75) conducted a RCT with a sample of 12-30 year olds with dizziness, neck pain and/or headache following SRC. After 8 weekly physical therapy sessions consisting of vestibular and cervical spine rehabilitation, subjects in the treatment group were nearly 4 times more likely to be medically cleared when compared to a control group. In a retrospective chart review, Alsalaheen and colleagues(75) examined the response of a population of concussed patients to vestibular physical therapy. Data from 114 patients referred for vestibular rehabilitation following concussion demonstrated a significant treatment effect for fifteen different measures of dizziness severity, balance confidence, gait and static/dynamic balance. Gottshall and Hoffer (72) assessed computerized VOR and gaze stability measures in eighty-two military individuals who experienced blast-related mild TBI. Impairment was significant at the time of initial evaluation, but returned to normative levels after 4-12 weeks of vestibular physical therapy. Hoffer and colleagues(76)examined the effect of vestibular rehabilitation in a population of fifty-eight active duty military individuals with post-concussive dizziness. They found that after a 6-8 week vestibular rehabilitation program, patients had improved with respect to symptoms of dizziness, perception of balance function, and measures of VOR function. However, the effectiveness of vestibular rehabilitation differed based on type of post-traumatic dizziness. Specifically, patients with post-traumatic migraine-associated dizziness were most responsive to treatment (84%), in contrast to the spatial disorientation group (27%).

Vision Therapy

Most oculomotor problems following SRC such as convergence insufficiency, accommodative insufficiency, impaired version movements and minor ocular misalignments may be managed conservatively with vision therapy(57). However, it is important to note that in rare instances surgical/medical intervention by an ophthalmologist or neuro-ophthalmologist may be warranted for complex diplopia, strabismus that is due to muscle

paralysis or nerve palsy, or other concurrent ocular-health issues. Typically, vision therapy involves vision exercises using eye patches, penlights, mirrors, lenses, prisms, and other non-surgical interventions to improve the function of the ocular muscles.

In spite of anecdotal evidence for the effectiveness of vision therapy following SRC, there is limited empirical support for vision therapy in the literature. However, a 2011 Cochrane review of RCT's for nonsurgical intervention for convergence insufficiency, and another RCT by Scheiman and colleagues for treatment of accommodative insufficiency pointed to the effectiveness of vision therapy in children in managing these two conditions (Scheiman 2011a; Scheiman 2011b). Although empirical support for oculomotor and vision-related therapies following SRC is limited and does not include any RCT's, emerging evidence supports the effectiveness of visual exercises for specific oculomotor problems. A retrospective study by Ciuffreda et al(62) examined patients with mTBI who were enrolled in a vision therapy program consisting of combined vergence, version and accommodative exercises. They reported that 90% of patients improved markedly or completely in symptoms and subjective reports of enhanced reading at a 2-3 month follow-up. A recent study involving 12 subjects following mTBI, Thiagarajan and colleagues(77) demonstrated that an oculomotor training program targeting the version, vergence, and accommodation components of the ocular motor system significantly improved the amplitudes of vergence and accommodation, accuracy of saccadic eye movements and overall reading.

Pharmacological Interventions

It has been reported that as much as 89% of clinicians manage symptoms of athletes with SRC using over the counter (OTC) or prescription medications(78). The most common interventions involve OTC medications such as NSAIDS and acetaminophen. However, many other prescription pharmacological interventions are employed with athletes who are not following a normal recovery trajectory (i.e., recovered within 10-14 days) following SRC. Research indicates that pharmacological treatments usually begin at approximately 10 days post-injury(42). As with vestibular and oculomotor therapies, pharmacological interventions are most effective when they target specific clinical subtypes of SRC. For example, an athlete with a primary cognitive-fatigue subtype following SRC may be prescribed a neurostimulant such as amantadine. In addition to cognitive-fatigue, other clinical subtypes that are amenable to pharmacological treatment include PTM and anxiety/mood (3, 8). Additionally, sleep-related issues are often treated using pharmacological interventions. It is important to acknowledge that there is still no FDA-approved pharmacological treatment for SRC. As such, all pharmacological interventions discussed below involve off-label use of medications that were approved for other primary purposes. Moreover, each medication discussed below may involve side effects that warrant close monitoring from prescribing clinicians. In addition, the use of certain medications, such as neurostimulants, may be in violation of the medication and performance enhancement policies of specific sport governing bodies, thus proper documentation is very important.

Targeted Pharmacological Interventions: Matching Treatments to Clinical Subtypes

Cognitive-fatigue is a commonly targeted clinical subtype for pharmacological intervention. Athletes with this subtype experience difficulty concentrating, memory problems, attentional

issues, decreased vigor, and headaches that worsen throughout the day. These symptoms are often treated effectively with the use of a neurostimulant. The most commonly used neurostimulant is amantadine, with 10% of clinicians reporting that they prescribe amantadine to athletes following SRC(78). There is some empirical evidence that amantadine, a dopaminergic neurostimulant primarily purposed as an anti-viral medication, can improve cognitive-fatigue symptoms and memory in athletes experiencing prolonged recovery following SRC(79). Other neurostimulants that can be used to treat athletes with cognitive-fatigue include methylphenidate, Adderall, and atomoxetine. There is some evidence of the effectiveness of methylphenidate on improving processing speed in moderate TBI (e.g., (80)), but not in athletes with SRC. Of note, some athletes may already be taking these medications for ADHD and related conditions, thereby necessitating close monitoring from clinicians and may need to be adjusted during their recovery period.

Some athletes may develop anxiety/mood issues as a direct result of a SRC or secondary to the injury recovery process with its concomitant frustrations and feelings of isolation and loss of control(16). Other athletes may have pre-existing anxiety/mood issues prior to injury that may be exacerbated following a SRC. Regardless of its underlying cause, anxiety/mood issues following SRC can be treated with tricyclic antidepressants (e.g., amitriptyline). In fact, tricyclic antidepressants are used by up to 23% of clinicians treating young athletes with SRC(78). It is likely that this relatively high percentage of clinicians prescribing tricyclic antidepressants is due in part to its use across multiple clinical subtypes including anxiety/mood, sleep, and PTM. Other common medications used for athletes in the anxiety/mood clinical subtype include selective serotonin reuptake inhibitors (SSRI) and selective norepinephrine reuptake inhibitors (SNRI). There is some anecdotal evidence that short-term, low dosage use of certain benzodiazepines such as Klonopin may be effective in athletes with vestibular-related anxiety. Klonopin and other benzodiazepines are also believed to act on neurons in the vestibular nuclei of the brain to decrease vestibular related symptoms and in turn decrease anxiety. Klonopin can be effective for vestibular related migraines. However, Klonopin can result in elevated anxiety and sleep disruption in some athletes and its use should be monitored closely.

The symptoms of PTM include headache, nausea, photo- or phono-sensitivity and dizziness. These symptoms have been associated with prolonged recovery and impairment following SRC(81). Clinicians may use a variety of pharmacological interventions to treat the symptoms of PTM including tricyclic and SSRI antidepressants, anticonvulsants (e.g., topiramate, gabapentin, valproic acid), or beta-blockers. In addition, triptans (e.g., Imitrex, Maxalt) are often prescribed as abortive medications for PTM. In spite of anecdotal evidence regarding the effectiveness of these treatments, there are no empirical studies of the effectiveness of these medications in athletes with SRC.

There is often a sleep overlay that permeates across each clinical subtype of SRC. Consequently, clinicians often prescribe OTC and prescription sleep medications for athletes with persistent sleep disruptions following SRC. After all, if an athlete is not sleeping well following a SRC, it will be difficult for that athlete to recover. The most commonly used sleep medication is melatonin with 1/5 of clinicians reporting that they prescribe melatonin to athletes with sleep disruptions following SRC(78). Melatonin together with basic sleep

hygiene can help regulate circadian rhythms and promote better sleep-wake cycling(82, 83). Other medications used to improve sleep disruption in athletes following SRC include antidepressants (e.g., amitriptyline, trazodone) and non-benzodiazepine hypnotics (e.g., Ambien, Lunesta)(83).

Summary and Recommendations

In spite of limited empirical support, physical and cognitive rest have been deemed essential components of initial concussion management and treatment. Such recommendations have been developed by consensus and introduced by the International Concussion in Sport Group in 2008. Since that time, there has been limited empirical work evaluating the efficacy of physical and cognitive rest protocols. Some research suggests that prescribed physical and cognitive rest in the acute stage of concussion may be of benefit to some athletes. However, other studies have indicated that an early return to light to moderate physical activity may be effective for other athletes following concussion. The heterogeneous nature of concussion renders a universal prescription of strict rest for all concussed athletes an ineffective strategy. As such strict rest extending beyond the acute injury stage may result in the athlete developing concussion-like symptoms that are unrelated to the injury (e.g. anxiety, migraine, sleep disorders, etc) and may complicate injury management, which may in turn lead to psychological and other concurrent problems. Student-athletes who are unable to attend or participate in academics to the fullest may benefit from a reduced cognitive load following injury, with a graduated return to academics that does not exacerbate symptoms. Reduction of cognitive load requires a coordinated effort between the medical and school academic support staffs with short or long term academic accommodations. Any accommodations may be lifted once a complete academic schedule can be completed without symptom exacerbation, at which time a return to play protocol be undertaken.

Sport-related concussions can involve several different clinical subtypes that warrant a comprehensive clinical assessment and subsequent targeted treatment and rehabilitation strategies. Recent advances in screening for vestibular and oculomotor impairment and symptoms (e.g., Mucha et al (17)) have revealed that many athletes experience these issues following SRC. Research also suggests that athletes with these issues often have longer recovery times and more pronounced impairment and symptoms following SRC(25). In response to these findings, vestibular and vision therapists have begun to apply specific rehabilitation interventions to enhance the recovery process for those athletes with vestibular or oculomotor impairment and symptoms following SRC. Initial empirical evidence indicates that these vestibular and oculomotor interventions may be useful in mitigating these issues and enhancing the recovery of athletes with SRC. However, additional research regarding which interventions are most effective for each type of impairment and symptoms as well as the optimal number and length of therapeutic sessions needed to obtain the desired effect is warranted.

Most clinicians use some sort of OTC or prescription pharmacological intervention to help manage lingering symptoms and impairment following SRC(78). It is clear from clinical experience that when pharmacological treatments are matched appropriately with patients'

clinical subtypes and symptoms they can be an effective intervention. However, there is some overlap for the effectiveness of certain pharmacological interventions across more than one clinical subtype. Most pharmacological treatments are implemented in patients with lingering (10-14+ days) or chronic (3+ months) symptoms and impairments. It is atypical for a patient to be prescribed a medication in the acute and sub-acute phases following a concussion. This “wait and see” approach may result in missed opportunities for effective early pharmacological intervention following SRC. However, researchers have yet to determine how soon after injury the preceding pharmacological treatments should be implemented to have the greatest therapeutic effect. In fact, it has been suggested that clinicians could accelerate recovery for some patients if pharmacological treatments were implemented earlier in the injury process. In addition, researchers need to explore the effectiveness of various dosage levels, treatment regimens, and administration methods in patients following concussion.

This review was conceived to evaluate the evidence supporting current and emerging rehabilitation approaches for sport concussion. Consensus opinion for prescribed physical and cognitive rest is the most common rehabilitation approach for patients with concussion.. However, more active and targeted rehabilitation strategies including vestibular and oculomotor rehabilitation and pharmacological interventions have emerging evidence supporting their use. Ultimately, there is limited empirical support for the rehabilitation strategies discussed in this paper, necessitating additional research on their effectiveness following concussion. This research should employ multi-site, randomized controlled trial research designs to better elucidate the specific effects of individual interventions. In addition, future research should employ comprehensive outcome assessments and targeted rehabilitation strategies that account for the heterogeneous nature of this injury.

REFERENCES

1. Giza CC, Hovda DA. The neurometabolic cascade of concussion. *J Athl Train.* 2001; 36(3):228–35. [PubMed: 12937489]
2. Giza CC, Hovda DA. The new neurometabolic cascade of concussion. *Neurosurgery.* 2014; 75(Suppl 4):S24–33. [PubMed: 25232881]
3. McCrory P, Meeuwisse WH, Aubry M, Cantu RC, Dvorak J, Echemendia RJ, et al. Consensus statement on concussion in sport: the 4th International Conference on Concussion in Sport held in Zurich, November 2012. *British Journal of Sports Medicine.* 2013; 47(5):250–8. [PubMed: 23479479]
4. National Collegiate Athletic A. NCAA Injury Surveillance Summary for 15 sports: 1988-1999 through 2003-2004. *J Athl Train.* 2007; 42(2):165–320.
5. Broglio SP, Puetz TW. The effect of sport concussion on neurocognitive function, self-report symptoms, and postural control: A meta-analysis. *Sports Med.* 2008; 38(1):53–67. [PubMed: 18081367]
6. McCrea M, Barr WB, Guskiewicz KM, Randolph C, Marshall SW, Cantu R, et al. Standard regression-based methods for measuring recovery after sport-related concussion. *J Int Neuropsychol Soc.* 2005; 11:58–69. [PubMed: 15686609]
7. McCrea M, Guskiewicz KM, Marshall SW, Barr W, Randolph C, Cantu RC, et al. Acute effects and recovery time following concussion in collegiate football players: the NCAA Concussion Study. *JAMA.* 2003; 290(19):2556–63. [PubMed: 14625332]

8. Collins MW, Kontos AP, Reynolds E, Murawski CD, Fu FH. A comprehensive, targeted approach to the clinical care of athletes following sport-related concussion. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA*. 2014; 22(2):235–46.
9. Broglia SP, Cantu RC, Gioia GA, Guskiewicz KM, Kutcher JS, Palm M, et al. National athletic trainers' association position statement: management of sport concussion. *J Athl Train*. 2014; 49(2): 245–65. [PubMed: 24601910]
10. Committee on Sports-Related Concussions in Youth, Board on Children Y, Families, Institute of M, National Research C. Graham R, Rivara FP, Ford MA, Spicer CM. The National Academies Collection: Reports funded by National Institutes of Health. *Sports-Related Concussions in Youth: Improving the Science, Changing the Culture*. National Academies Press (US) Washington (DC) Copyright 2014 by the National Academy of Sciences. All rights reserved.; 2014.
11. Moser RS, Glatts C, Schatz P. Efficacy of immediate and delayed cognitive and physical rest for treatment of sports-related concussion. *J Pediatr*. 2012; 161(5):922–6. [PubMed: 22622050]
12. Brown NJ, Mannix RC, O'Brien MJ, Gostine D, Collins MW, Meehan WP 3rd. Effect of cognitive activity level on duration of post-concussion symptoms. *Pediatrics*. 2014; 133(2):e299–304. [PubMed: 24394679]
13. Covassin T, Swanik CB, Sachs M, Kendrick Z, Schatz P, Zillmer E, et al. Sex differences in baseline neuropsychological function and concussion symptoms of collegiate athletes. *Br J Sports Med*. 2006; 40(11):923–7. discussion 7. [PubMed: 16990442]
14. Iverson GL, Gaetz M, Lovell MR, Collins MW. Cumulative effects of concussion in amateur athletes. *Brain Inj*. 2004; 18(5):433–43. [PubMed: 15195792]
15. Field M, Collins MW, Lovell MR, Maroon JC. Does age play a role in recovery from sports-related concussion? A comparison of high school and collegiate athletes. *J Pediatr*. 2003; 142(5): 546–53. [PubMed: 12756388]
16. Kontos APCM, Russo S. An introduction to sport concussion for the sport psychology consultant. *Appl Sport Psych*. 2004; 16(3):220–35.
17. Mucha A, Collins MW, Elbin RJ, Furman JM, Troutman-Enseki C, DeWolf RM, et al. A Brief Vestibular/Ocular Motor Screening (VOMS) Assessment to Evaluate Concussions: Preliminary Findings. *The American journal of sports medicine*. 2014
18. Belanger HG, Vanderploeg RD. The neuropsychological impact of sports-related concussion: A meta-analysis. *J Int Neuropsychol Soc*. 2005; 11:345–57. [PubMed: 16209414]
19. Collins MW, Field M, Lovell MR, Iverson GL, Johnston KM, Maroon JC, et al. Relationship between postconcussion headache and neuropsychological test performance in high school athletes. *American Journal of Sports Medicine*. 2003; 31(2):168–73. [PubMed: 12642248]
20. Collins MW, Iverson GL, Lovell MR, McKeag DB, Norwig J, Maroon JC. On-field predictors of neuropsychological and symptom deficit following sports-related concussion. *Clin J Sport Med*. 2003; 13(4):222–9. [PubMed: 12855924]
21. Delaney JS, Lacroix VJ, Leclerc S, Johnston KM. Concussions among university football and soccer players. *Clin J Sport Med*. 2002; 12(6):331–8. [PubMed: 12466687]
22. Guskiewicz KM, Weaver NL, Padua DA, Garrett WE. Epidemiology of concussion in collegiate and high school football players. *American Journal of Sports Medicine*. 2000; 28(5):643–50. [PubMed: 11032218]
23. Guskiewicz KM, McCrea M, Marshall SW, Cantu RC, Randolph C, Barr W, et al. Cumulative effects associated with recurrent concussion in collegiate football players: The NCAA concussion study. *JAMA*. 2003; 290(19):2549–55. [PubMed: 14625331]
24. McCrory PR, Ariens M, Berkovic SF. The nature and duration of acute concussive symptoms in Australian football. *Clin J Sport Med*. 2000; 10:235–8. [PubMed: 11086747]
25. Lau BC, Kontos AP, Collins MW, Mucha A, Lovell MR. Which on-field signs/symptoms predict protracted recovery from sport-related concussion among high school football players? *Am J Sport Med*. 2011; 39(11):2311–8.
26. Eisenberg MA, Meehan WP 3rd, Mannix R. Duration and course of post-concussive symptoms. *Pediatrics*. 2014; 133(6):999–1006. [PubMed: 24819569]
27. McCrea M, Guskiewicz KM, Randolph C, Barr WB, Hammeke TA, Marshall SW, et al. Incidence, clinical course, and predictors of prolonged recovery time following sport-related concussion in

- high school and college athletes. *J Int Neuropsychol Soc.* 2013; 19(1):22–33. [PubMed: 23058235]
28. Bruce DA, Alavi A, Bilaniuk L, Dolinskas C, Obrist W, Uzzell B. Diffuse cerebral swelling following head injuries in children: the syndrome of “malignant brain edema”. *J Neurosurg.* 1981; 54(2):170–8. [PubMed: 7452330]
 29. Giza CC, Difiori JP. Pathophysiology of sports-related concussion: an update on basic science and translational research. *Sports Health.* 2011; 3(1):46–51. [PubMed: 23015990]
 30. Prins ML, Hales A, Reger M, Giza CC, Hovda DA. Repeat traumatic brain injury in the juvenile rat is associated with increased axonal injury and cognitive impairments. *Developmental neuroscience.* 2010; 32(5-6):510–8. [PubMed: 20829578]
 31. Vagnozzi R, Signoretti S, Cristofori L, Alessandrini F, Floris R, Isgro E, et al. Assessment of metabolic brain damage and recovery following mild traumatic brain injury: a multicentre, proton magnetic resonance spectroscopic study in concussed patients. *Brain.* 2010; 133(11):3232–42. [PubMed: 20736189]
 32. Vagnozzi R, Signoretti S, Tavazzi B, Floris R, Ludovici A, Marziali S, et al. Temporal window of metabolic brain vulnerability to concussion: A pilot H-magnetic resonance spectroscopy study in concussed athletes - Part III. *Neurosurgery.* 2008; 62(6):1286–95. [PubMed: 18824995]
 33. Griesbach GS, Hovda DA, Molteni R, Wu A, Gomez-Pinilla F. Voluntary exercise following traumatic brain injury: brain-derived neurotrophic factor upregulation and recovery of function. *Neuroscience.* 2004; 125(1):129–39. [PubMed: 15051152]
 34. Carson JD, Lawrence DW, Kraft SA, Garel A, Snow CL, Chatterjee A, et al. Premature return to play and return to learn after a sport-related concussion: physician’s chart review. *Canadian family physician Medecin de famille canadien.* 2014; 60(6):e310, e2–5. [PubMed: 24925965]
 35. Leddy MH, Lambert MJ, Ogles BM. Psychological consequences of athletic injury among high-level competitors. *Res Q Exerc Sport.* 1994; 65(4):347–54. [PubMed: 7886284]
 36. Manuel JC, Shilt JS, Curl WW, Smith JA, Durant RH, Lester L, et al. Coping with sports injuries: an examination of the adolescent athlete. *J Adolesc Health.* 2002; 31(5):391–3. [PubMed: 12401424]
 37. Milde-Busch A, Blaschek A, Borggrafe I, Heinen F, Straube A, von Kries R. Associations of diet and lifestyle with headache in high-school students: results from a cross-sectional study. *Headache.* 2010; 50(7):1104–14. [PubMed: 20533961]
 38. Majerske CW, Mihalik JP, Ren D, Collins MW, Reddy CC, Lovell MR, et al. Concussion in sports: postconcussive activity levels, symptoms, and neurocognitive performance. *J Athl Train.* 2008; 43(3):265–74. [PubMed: 18523563]
 39. Leddy JJ, Kozlowski K, Donnelly JP, Pendergast DR, Epstein LH, Willer B. A preliminary study of subsymptom threshold exercise training for refractory post-concussion syndrome. *Clin J Sport Med.* 2010; 20(1):21–7. [PubMed: 20051730]
 40. Baker JG, Freitas MS, Leddy JJ, Kozlowski KF, Willer BS. Return to full functioning after graded exercise assessment and progressive exercise treatment of postconcussion syndrome. *Rehabilitation research and practice.* 2012; 2012:705309. [PubMed: 22292122]
 41. Vancampfort D, Probst M, Adriaens A, Pieters G, De Hert M, Stubbs B, et al. Changes in physical activity, physical fitness, self-perception and quality of life following a 6-month physical activity counseling and cognitive behavioral therapy program in outpatients with binge eating disorder. *Psychiatry Res.* 2014; 219(2):361–6. [PubMed: 24929440]
 42. Giza CC, Kutcher JS, Ashwal S, Barth J, Getchius TS, Gioia GA, et al. Summary of evidence-based guideline update: Evaluation and management of concussion in sports: Report of the Guideline Development Subcommittee of the American Academy of Neurology. *Neurology.* 2013; 80(24):2250–7. [PubMed: 23508730]
 43. Halstead ME, Walter KD, Council on Sports M, Fitness. American Academy of Pediatrics. Clinical report--sport-related concussion in children and adolescents. *Pediatrics.* 2010; 126(3):597–615.
 44. Herring SA, Cantu RC, Guskiewicz KM, Putukian M, Kibler WB, Bergfeld JA, et al. Concussion (mild traumatic brain injury) and the team physician: a consensus statement--2011 update. *Medicine and Science in Sports and Exercise.* 2011; 43(12):2412–22. [PubMed: 22089299]

45. Harmon KG, Drezner J, Gammons M, Guskiewicz KM, Halstead M, Herring SA, et al. American Medical Society for Sports Medicine position statement: concussion in sport. *Clin J Sport Med*. 2013; 23(1):1–18. [PubMed: 23269325]
46. Mealings M, Douglas J, Olver J. Considering the student perspective in returning to school after TBI: a literature review. *Brain Inj*. 2012; 26(10):1165–76. [PubMed: 22571252]
47. Halstead ME, McAvoy K, Devore CD, Carl R, Lee M, Logan K. Returning to learning following a concussion. *Pediatrics*. 2013; 132(5):948–57. [PubMed: 24163302]
48. Sady MD, Vaughan CG, Gioia GA. School and the concussed youth: recommendations for concussion education and management. *Phys Med Rehabil Clin N Am*. 2011; 22(4):701–19. [PubMed: 22050944]
49. McGrath N. Supporting the student-athlete's return to the classroom after a sport-related concussion. *J Athl Train*. 2010; 45(5):492–8. [PubMed: 20831397]
50. Howell D, Osternig L, Van Donkelaar P, Mayr U, Chou LS. Effects of concussion on attention and executive function in adolescents. *Med Sci Sports Exerc*. 2013; 45(6):1030–7. [PubMed: 23274602]
51. McGrath N, Dinn WM, Collins MW, Lovell MR, Elbin RJ, Kontos AP. Post-exertion neurocognitive test failure among student-athletes following concussion. *Brain Inj*. 2013; 27(1): 103–13. [PubMed: 23252441]
52. Rehabilitation Act of 1973 amendment 29 U.S.C. U.S Department of Education; 2012. § 794 (Section 504)
53. Piebes SK, Gourley M, Valovich McLeod TC. Caring for student-athletes following a concussion. *The Journal of school nursing : the official publication of the National Association of School Nurses*. 2009; 25(4):270–81. [PubMed: 19535497]
54. Schneider KJ, Iverson GL, Emery CA, McCrory P, Herring SA, Meeuwisse WH. The effects of rest and treatment following sport-related concussion: a systematic review of the literature. *Br J Sports Med*. 2013; 47(5):304–7. [PubMed: 23479489]
55. Collins MW, Lovell MR, Iverson GL, Ide T, Maroon JC. Examining concussion rates and return to play in high school football players wearing newer helmet technology: a three-year prospective cohort study. *Neurosurgery*. 2006; 58(2):275–86. [PubMed: 16462481]
56. Furman JM, Raz Y, Whitney SL. Geriatric vestibulopathy assessment and management. *Curr Opin Otolaryngol Head Neck Surg*. 2010; 18(5):386–91. [PubMed: 20613528]
57. Kapoor N, Ciuffreda KJ. Vision Disturbances Following Traumatic Brain Injury. *Curr Treat Options Neurol*. 2002; 4(4):271–80. [PubMed: 12036500]
58. Rutner D, Kapoor N, Ciuffreda KJ, Craig S, Han ME, Suchoff IB. Occurrence of ocular disease in traumatic brain injury in a selected sample: a retrospective analysis. *Brain Inj*. 2006; 20(10):1079–86. [PubMed: 17060141]
59. Stiller-Ostrowski JL. Fourth cranial nerve palsy in a collegiate lacrosse player: a case report. *J Athl Train*. 2010; 45(4):407–10. [PubMed: 20617917]
60. Chan RV, Trobe JD. Spasm of accommodation associated with closed head trauma. *Journal of neuro-ophthalmology : the official journal of the North American Neuro-Ophthalmology Society*. 2002; 22(1):15–7. [PubMed: 11937900]
61. Capo-Aponte JE, Tarbett AK, Uroevich TG, Temme LA, Sanghera NK, Kalich ME. Effectiveness of computerized oculomotor vision screening in a military population: pilot study. *Journal of rehabilitation research and development*. 2012; 49(9):1377–98. [PubMed: 23408219]
62. Ciuffreda KJ, Rutner D, Kapoor N, Suchoff IB, Craig S, Han ME. Vision therapy for oculomotor dysfunctions in acquired brain injury: a retrospective analysis. *Optometry (St Louis, Mo)*. 2008; 79(1):18–22.
63. Bhattacharyya N, Baugh RF, Orvidas L, Barrs D, Bronston LJ, Cass S, et al. Clinical practice guideline: benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg*. 2008; 139(5 Suppl 4):S47–81. [PubMed: 18973840]
64. Geurts AC, Ribbers GM, Knoop JA, van Limbeek J. Identification of static and dynamic postural instability following traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*. 1996; 77:639–44. [PubMed: 8669988]

65. Guskiewicz, KM. Regaining balance and postural equilibrium. In: Prentice, WE., editor. *Rehabilitation Techniques in Sports Medicine*. Vol. 3. WCB McGraw-Hill; Boston: 1999. p. 107-33.
66. Guskiewicz KM, Riemann BL, Perrin DH, Nashner LM. Alternative approaches to the assessment of mild head injury in athletes. *Medicine and Science in Sports and Exercise*. 1997; 29(7 Suppl):S213–S21. [PubMed: 9247918]
67. Peterson CL, Ferrara MS, Mrazik M, Piland SG, Elliot R. Evaluation of neuropsychological domain scores and postural stability following cerebral concussion in sports. *Clin J Sport Med*. 2003; 13(4):230–7. [PubMed: 12855925]
68. Guskiewicz KM. Postural stability assessment following concussion: One piece of the puzzle. *Clin J Sport Med*. 2001; 11(3):182–9. [PubMed: 11495323]
69. Parker TM, Osternig LR, Lee H, Donkelaar P, Chou LS. The effect of divided attention on gait stability following concussion. *Clin Biomech*. 2005; 20(4):389–95.
70. Catena RD, von Donkelaar P, Chou LS. Cognitive task effects on gait stability following concussion. *Experimental Brain Research*. 2007; 176(1):23–31. [PubMed: 16826411]
71. Catena RD, von Donkelaar P, Chou LS. Altered balance control following concussion is better detected with an attention test during gait. *Gait Posture*. 2007; 25(3):406–11. [PubMed: 16787746]
72. Gottshall KR, Hoffer ME. Tracking recovery of vestibular function in individuals with blast-induced head trauma using vestibular-visual-cognitive interaction tests. *Journal of neurologic physical therapy : JNPT*. 2010; 34(2):94–7. [PubMed: 20588095]
73. Kozlowski KF, Graham J, Leddy JJ, Devinney-Boymel L, Willer BS. Exercise intolerance in individuals with postconcussion syndrome. *J Athl Train*. 2013; 48(5):627–35. [PubMed: 23952041]
74. Hillier SL, Hollohan V, Cd005397. Vestibular rehabilitation for unilateral peripheral vestibular dysfunction. *Cochrane Database Syst Rev*. 2007; 4
75. Schneider KJ, Meeuwisse WH, Nettel-Aguirre A, Barlow K, Boyd L, Kang J, et al. Cervicovestibular rehabilitation in sport-related concussion: a randomised controlled trial. *Br J Sports Med*. 2014; 48(17):1294–8. [PubMed: 24855132]
76. Hoffer ME, Gottshall KR, Moore R, Balough BJ, Wester D. Characterizing and treating dizziness after mild head trauma. *Otol Neurotol*. 2004; 25(2):135–8. [PubMed: 15021772]
77. Thiagarajan P, Ciuffreda KJ. Versional eye tracking in mild traumatic brain injury (mTBI): effects of oculomotor training (OMT). *Brain Inj*. 2014; 28(7):930–43. [PubMed: 24826956]
78. Kinnaman KA, Mannix RC, Comstock RD, Meehan WP 3rd. Management strategies and medication use for treating paediatric patients with concussions. *Acta Paediatr*. 2013; 102(9):e424–8. [PubMed: 23750873]
79. Reddy CC, Collins M, Lovell M, Kontos AP. Efficacy of amantadine treatment on symptoms and neurocognitive performance among adolescents following sports-related concussion. *J Head Trauma Rehabil*. 2013; 28(4):260–5. [PubMed: 22613947]
80. Whyte J, Hart T, Vaccaro M, Grieb-Neff P, Risser A, Polansky M, et al. Effects of methylphenidate on attention deficits after traumatic brain injury: a multidimensional, randomized, controlled trial. *Am J Phys Med Rehabil*. 2004; 83(6):401–20. [PubMed: 15166683]
81. Kontos AP, Elbin RJ, Lau B, Simensky S, Freund B, French J, et al. Posttraumatic migraine as a predictor of recovery and cognitive impairment after sport-related concussion. *The American journal of sports medicine*. 2013; 41(7):1497–504. [PubMed: 23698389]
82. Petraglia AL, Maroon JC, Bailes JE. From the field of play to the field of combat: a review of the pharmacological management of concussion. *Neurosurgery*. 2012; 70(6):1520–33. discussion 33. [PubMed: 22289786]
83. Meehan WP. Medical therapies for concussion. *Clinics in Sports Medicine*. 2011; 30(1):115–24. [PubMed: 21074086]
84. Pavlou M, Davies RA, Bronstein AM. The assessment of increased sensitivity to visual stimuli in patients with chronic dizziness. *J Vestib Res*. 2006; 16(4-5):223–31. [PubMed: 17538212]
85. Guskiewicz KM, Ross SE, Marshall SW. Postural stability and neuropsychological deficits after concussion in collegiate athletes. *J Athl Train*. 2001; 36(3):263–73. [PubMed: 12937495]

86. Norre ME. Cervical vertigo. Diagnostic and semiological problem with special emphasis upon “cervical nystagmus”. *Acta oto-rhino-laryngologica Belgica*. 1987; 41(3):436–52. [PubMed: 3673567]
87. Kristjansson E, Treleaven J. Sensorimotor function and dizziness in neck pain: implications for assessment and management. *J Orthop Sports Phys Ther*. 2009; 39(5):364–77. [PubMed: 19411769]
88. Leddy JJ, Kozlowski K, Fung M, Pendergast DR, Willer B. Regulatory and autoregulatory physiological dysfunction as a primary characteristic of post concussion syndrome: implications for treatment. *NeuroRehabilitation*. 2007; 22(3):199–205. [PubMed: 17917170]

Key Points

- 1) Concussion rehabilitation policies are largely consensus based
- 2) Emerging evidence is suggesting that exercise and cognitive activity in a controlled and prescriptive manner may benefit recovery.
- 3) Additional rehabilitation strategies (eg vestibular, oculomotor, and pharamocological) also have mounting evidence and should be incorporated by an appropriately trained professional when appropriate.

Table 1

Common Interventions for Vestibular Impairment following Sport-related Concussion.

Impairment	Etiology	Symptoms	Associated Problems/Risk Factors	Physical Therapy Treatment
Benign Paroxysmal Positional Vertigo(63)	Mechanical disruption in the vestibular labyrinth (end organ). Otoconia from otoliths become dislodged and displace in semicircular canal(s)	Vertigo with changes in head position	Older age High impact forces	Canalith repositioning maneuvers
VOR Impairment(76)	Disrupted function in the VOR pathways, peripherally or centrally	Dizziness	Labyrinthine Concussion	Gaze Stability Training
Visual Motion Sensitivity(84)	Impaired central processing/integration of vestibular information with visual and other sensory information	Dizziness	Post-Traumatic Migraine Anxiety	Graded exposure to visually-stimulating environments Virtual reality Optokinetic stimulation
Impaired Postural Control(85)	Disruption/damage to vestibular-spinal reflex pathways, peripherally or centrally	Impaired balance, particularly with: - vision and/or somatosensation reduced - cognitive dual/task demand	Common early finding after concussion; typically resolves before other vestibular deficits	Balance rehabilitation strategies Sensory organization training Divided attention training Dynamic balance training
Cervicogenic Dizziness(86, 87)	Cervical injury results in abnormal afferent input to CNS; mismatch with other sensory information	Dizziness – related to cervical movement/posture Imbalance Impaired oculomotor control	Cervical Pathology Cervicogenic Headaches	Manual therapy for cervical spine Balance training Oculomotor training
Exercise-Induced Dizziness(88)	Inadequate central response to cardiovascular and vestibular/ocular demands of exercise	Dizziness with movement-related cardiovascular exercise	-VOR/Gaze Stability Impairment -Visual Motion Sensitivity -Autonomic dysregulation	Progressive dynamic exertion exercise program