The biology of IQGAP proteins: beyond the cytoskeleton

Andrew C Hedman[†], Jessica M Smith[†] & David B Sacks^{*}

Abstract

IQGAP scaffold proteins are evolutionarily conserved in eukaryotes and facilitate the formation of complexes that regulate cytoskeletal dynamics, intracellular signaling, and intercellular interactions. Fungal and mammalian IQGAPs are implicated in cytokinesis. IQGAP1, IQGAP2, and IQGAP3 have diverse roles in vertebrate physiology, operating in the kidney, nervous system, cardiovascular system, pancreas, and lung. The functions of IQGAPs can be corrupted during oncogenesis and are usurped by microbial pathogens. Therefore, IQGAPs represent intriguing candidates for novel therapeutic agents. While modulation of the cytoskeletal architecture was initially thought to be the primary function of IQGAPs, it is now clear that they have roles beyond the cytoskeleton. This review describes contributions of IQGAPs to physiology at the organism level.

Keywords biology; IQGAP1; IQGAP2; IQGAP3; therapeutics

DOI 10.15252/embr.201439834 | Received 6 November 2014 | Revised 22

December 2014 | Accepted 7 January 2015 | Published online 26 February 2015

EMBO Reports (2015) 16: 427–446

See the Glossary for abbreviations used in this article.

Introduction

IQGAPs are an evolutionarily conserved family of proteins that interact with many partners to regulate diverse cellular processes, including cytokinesis [1,2], cell migration [3], cell proliferation [4], intracellular signaling [4,5], vesicle trafficking [5,6], and cytoskeletal dynamics [7,8]. IQGAP proteins are present in a wide variety of fungi, protist, and animal cells. The majority of vertebrates, including humans, express three related isoforms IQGAP1, IQGAP2, and IQGAP3 (Fig 1). IQGAPs contain several domains that mediate protein–protein interactions (Table 1). While prior reviews have focused on the cellular processes regulated by these interactions [3–5,7,8], attention to the roles of IQGAPs at the organism level has been limited. This review summarizes functions of fungal and vertebrate IQGAP proteins in physiology.

IQGAPs scaffold diverse pathways

The multidomain composition of IQGAPs mediates the formation of protein complexes required for cellular processes. For example, interactions of the IQGAP1 calponin homology domain (CHD) with F-actin and the GAP-related domain (GRD) with small GTPases regulate the cytoskeleton to promote actin binding or polymerization that regulates cytokinesis [1,2], cell migration [9], and stability of cell–cell contacts [10,11]. IQGAPs also scaffold molecules to form signaling complexes, such as components of the mitogen-activated protein kinase (MAPK) pathway [12,13]. The MAPK signaling cascade is activated in response to stimuli, which leads to sequential phosphorylation from Raf to MAPK-ERK kinase (MEK) to extracellular signal-regulated kinase (ERK) [14]. IQGAP1 regulates MAPK signaling by scaffolding several MAPK components, including K-Ras [15], B-Raf [16,17], MEK [13], and ERK [12,13]. These interactions promote ERK activation, which influences myriad cellular processes, ultimately impacting physiology in a variety of tissues. IQGAPs also form complexes with numerous other proteins. These include Ca^{2+} /calmodulin [18–20], Cdc42 [18,21–23], Rac1 [21], and actin [19,20,24,25] to control the actin cytoskeleton, as well as mTor and Akt kinases [26], to modulate Akt activation in processes such as cell growth and survival.

Cytokinesis

Cytokinesis is the culminating event in cell division and is essential for development and tissue maintenance/homeostasis. Defects in cytokinesis can result in aneuploidy, which can lead to developmental defects and has been implicated in cancer [27]. IQGAP proteins have an evolutionarily conserved role in cytokinesis from fungi to mammals. Fungi express a single IQGAP isoform that participates in cytokinesis. A contractile ring, which forms between parent and daughter cells, utilizes myosin motor proteins and the actin cytoskeleton to generate the force necessary to separate cells. Loss-of-function studies for several yeast and fungal IQGAPs, including *Saccharomyces cerevisiae* Iqg1p/Cyk1p [28–30], *Schizosaccharomyces pombe* Rng2p [31–33], and *Candida albicans* Iqg1p [34], result in the formation of multinucleated

Department of Laboratory Medicine, National Institutes of Health, Bethesda, MD, USA

^{*}Corresponding author. Tel: +1 301 496 3386; E-mail: sacksdb@mail.nih.gov

[†]These authors contributed equally to this work

Glossary

AMPK	AMP-activated protein kinase
Arp2/3	actin-related proteins 2/3
[Ca²⁺]_i	intracellular free calcium concentration
CHD	calponin homology domain
CSFV	classical swine fever virus
EB1	microtubule plus end binding protein 1
ERK	extracellular signal-regulated kinase
FAK	focal adhesion kinase
GAP	GTPase-activating protein
GEF	guanine nucleotide exchange factor
GRD	GAP-related domain
IQ	protein sequences containing Iso/Leu and Gln residues
Lis1	lissencephaly 1
M-MuLV	Moloney murine leukemia virus
MAPK	mitogen-activated protein kinase
MEK	MAPK/ERK kinase
MLC	myosin light chain
MLCK	myosin light chain kinase
MLCP	myosin light chain phosphatase
NGF	nerve growth factor
N-WASP	Neuronal Wiskott–Aldrich syndrome protein
PKCϵ	protein kinase C ϵ
PLCϵ1	phospholipase C ϵ 1
PPI	protein–protein interaction
PTPμ	protein-tyrosine phosphatase μ
RGCT	RasGAP_C-terminus domain
RTK	receptor tyrosine kinase
VEGF	vascular endothelial growth factor
VEGFR2	vascular endothelial growth factor receptor 2
WW	tryptophan-containing protein domain

cells, demonstrating a role for IQGAPs in the assembly of the contractile ring and cytokinesis.

Unlike fungi, the amoeba *Dictyostelium discoideum* has four IQGAP-like proteins: DGAP1/ddIQGAP1, GAPA/ddIQGAP2, DDB0233055/ddIQGAP3 (Fig 1), and the hypothetical/putative DDB0232202/ddIQGAP4 [35]. Both DGAP1 and GAPA function in cleavage furrow formation in *D. discoideum* cytokinesis [36–38]. Additionally, GAPA promotes cleavage furrow formation in response to mechanical stress, while DGAP1 inhibits this response [39]. This suggests distinct roles for each protein in response to specific stimuli, that is, DGAP1/biochemical signals and GAPA/mechanosensory inputs.

Less is known about the contribution of IQGAP to cytokinesis in higher eukaryotes. In the nematode *Caenorhabditis elegans*, RNA interference was employed to identify proteins associated with cleavage furrow formation and cytokinesis. Depletion of the *C. elegans* IQGAP PES-7 resulted in the formation of multinucleated germ cells and multinucleated embryos, indicating defects in the completion of meiosis and mitosis [40]. The mid-body assembles microtubules and other proteins necessary for completion of cell division at the end of cytokinesis. In mammalian cells, IQGAP1 was observed at the mid-body or contractile ring during cytokinesis in mouse oocytes and embryos [41], Chinese hamster ovary, as well as human HeLa cells [40].

Anillin proteins form complexes with actin and other proteins necessary for assembling the actomyosin ring at the cleavage furrow [42]. In *S. pombe*, Rng2p is recruited to the cleavage site by Mid1p, an anillin-like protein [43,44]. Similarly, in mammalian cells, anillin

recruits IQGAP3 to the actomyosin ring [45]. Furthermore, loss-of-function studies for IQGAP1 and IQGAP3 demonstrated roles for both proteins in regulating the localization of machinery required for cytokinesis in HeLa cells [45]. In contrast to prior reports, IQGAP1 was not detected at the mid-body in this study. The reason for the discrepancy is unknown. Nevertheless, depletion of either IQGAP1 or IQGAP3 led to defects in cytokinesis and resulted in the formation of multinucleated cells, with a more pronounced defect upon depletion of both IQGAP1 and IQGAP3, suggesting contributions from both proteins to cytokinesis [45]. Further investigation is required to dissect out the specific roles of IQGAP1 and IQGAP3 in cytokinesis.

Physiological relevance

Evidence derived from knockout mice and cultured cells has identified roles for IQGAP proteins, particularly IQGAP1, in multiple organs (Table 2). These studies are summarized here.

Kidney function

Podocytes are unique renal epithelial cells that form foot processes which wrap around glomerular capillaries. The processes of neighboring cells are connected by slit diaphragms, specialized intercellular junctions that mediate glomerular filtration [46] (Fig 2A). Mutations of critical components of slit diaphragms, such as nephrin or podocin, cause the nephrotic syndrome [47]. To further understand slit diaphragm architecture, interactors of the nephrin cytoplasmic domain were examined by mass spectrometry, and IQGAP1 was among the proteins identified [48]. Immunofluorescence microscopy revealed that IQGAP1 co-distributed with nephrin in the podocyte foot processes. IQGAP1 was also observed in kidney tubules and glomeruli [48]. The participation of IQGAP1 in slit diaphragm function was further suggested by the increased *in vitro* permeability of a podocyte layer when IQGAP1 is knocked down [49]. These findings and the association of IQGAP1 with several slit diaphragm components (Fig 2A), including nephrin, α -actinin, α II spectrin, β II spectrin, α -catenin, and podocin [49], suggest that IQGAP1 is an integral component of slit diaphragm organization to facilitate filtration.

Although slit diaphragm junctions are different to adherens junctions, they share key adherens junction proteins, including cadherins and catenins [46]. Adherens junctions are formed through cadherin complexes, which are linked intracellularly to the actin cytoskeleton via α -catenin and β -catenin [50]. IQGAP1 interacts with several adhesion-associated proteins, including E-cadherin (epithelial cadherin) [10,11], N-cadherin (neuronal cadherin) [51], VE-cadherin (vascular endothelial cadherin) [52], and β -catenin [10,53] (Table 1). The interaction of IQGAP1, nephrin, and adherens junction proteins suggests that this multiprotein complex may modulate cadherin-mediated adhesion and cytoskeletal dynamics in the kidney, consistent with previous reports in cultured epithelial cells [11].

The peptide hormone angiotensin II, which activates smooth muscle contraction thus contributing to hypertension, can induce podocyte apoptosis [54]. This can cause podocyte injury or depletion, resulting in glomerulosclerosis, a stiffening of the renal glomeruli. Angiotensin II stimulates podocyte apoptosis via MAPK [55]. Interestingly, angiotensin II increases IQGAP1

Figure 1. Tree of IQGAP proteins.

IQGAP proteins are present in eukaryotes [221]. All contain a GRD. All mammals have five domains: CHD, WW domain, IQ domain, GRD, and RasGAP_C-terminus (RGCT). Domains adapted from the SMART and Pfam databases, tree made as in [221].

expression in both rat glomeruli *in vivo* and cultured podocytes and promotes the interaction of ERK1/2 with IQGAP1 [56]. IQGAP1 knockdown prevents angiotensin II-induced ERK1/2 activation and apoptosis of podocytes. These findings suggest that IQGAP1 participates in angiotensin II-mediated apoptosis by modulating MAPK signaling.

IQGAP1 also interacts with phospholipase C epsilon (PLC ϵ 1) [57]. Mutations in the *PLCE1* gene have been implicated in early-onset nephrotic syndrome, which leads to end-stage kidney disease [57]. IQGAP1 co-immunoprecipitates with PLC ϵ 1 from cultured podocytes. However, PLC ϵ 1-null mice do not manifest renal pathology and it is not known whether PLC ϵ 1—and its association with IQGAP1—contributes to podocyte function in the development of kidney disease.

Neuronal function

The first documentation of IQGAP1 in neuronal cells was published in 2005 [58]. IQGAP1 was observed throughout the cell, along neurites and the developing axon, as well as at the growth cone. Overexpression of IQGAP1 induced neurite outgrowth in NIE-115 mouse neuroblastoma cells, an effect that was enhanced by phosphorylation of IQGAP1 by protein kinase C ϵ (PKC ϵ) [58] (Fig 2Bi). Later work demonstrated that an interaction between IQGAP1 and protein-tyrosine phosphatase PTP μ is required for neurite outgrowth in E8 chick nasal retinal ganglion cells [59] (Fig 2Bi). PTP μ is a cell surface receptor that interacts with cadherin/catenin complexes to mediate cell–cell adhesion [60]. PTP μ forms a complex with IQGAP1, N-cadherin, E-cadherin, and β -catenin [59]. Active Cdc42 promotes the association of PTP μ with IQGAP1 and disruption of

Table 1. Interactors of IQGAPs.

Interactor	Interaction <i>in vitro</i> ^a	Interaction <i>in vivo</i> ^b	Proposed function(s)	Reference(s)
IQGAP1				
Cytoskeleton-associated proteins				
Actin	Yes	Yes	Cross-links actin filaments	[19,20,24,25]
APC	Yes	Yes	Regulates actin dynamics in migrating cells	[129]
Arp2/3	ND	Yes	Stimulates branched actin filament assembly	[130,131]
CD44	ND	Yes	Links hyaluronan to actin cytoskeleton	[132]
CLASP2	Yes	Yes	Links IQGAP1 to microtubules	[133,134]
CLIP-170	Yes	Yes	Links Rac1 and Cdc42 to microtubules	[135]
Cortactin	ND	Yes	Regulates subcellular localization of cortactin, enhances endothelial barrier	[85,136]
EB1	ND	Yes	Enhances endothelial barrier	[85]
Ezrin	Yes ^c	Yes	Unknown	[137]
IFT-A	ND	Yes	Unknown	[138]
ILK	ND	Yes	Regulates microtubule network	[139,140]
Lis1	ND	Yes	Regulates Cdc42 activity during neuronal migration	[63]
mDia1	Yes	Yes	Regulates phagocytosis and phagocytic cup formation	[140,141]
N-WASP	Yes	Yes	Stimulates branched actin filament assembly	[130,131]
NUMB5	ND	Yes	Unknown	[142]
PLD2	ND	Yes	Regulates IQGAP1 subcellular localization and interaction with Rac1	[136]
Protein 4.1R	Yes	Yes	Localizes IQGAP1 at the leading edge of migrating cells	[143]
Vimentin	ND	Yes	Regulates desmosome-like junctions	[144]
Wave2	ND	Yes	Unknown	[145]
Adhesion-associated proteins				
α-actinin	ND	Yes	Unknown	[49]
α-catenin	ND	Yes	Unknown	[49]
αII spectrin	ND	Yes	Unknown	[49]
βII spectrin	ND	Yes	Unknown	[49]
β-catenin	Yes	Yes	Inhibits cell–cell adhesion; enhances β-catenin mediated transcription	[10,53]
β1-integrin	ND	Yes	Regulates actin during mitosis	[146]
β3-integrin	Yes	Yes	Regulates pulmonary vascular permeability	[84]
CD13	ND	Yes	Unknown	[147]
E-cadherin	Yes	Yes	Regulates E-cadherin-mediated cell–cell adhesion	[10,11]
Filamin-A	ND	Yes	Regulates directional cell migration.	[148]
Melusin	Yes	Yes	Regulates cardiomyocyte hypertrophy and survival	[76]
Menin	Yes	Yes	Links menin to E-cadherin/β-catenin	[149]
N-cadherin	ND	Yes	Links N-cadherin to ERK1/2 signaling during fear memory formation, regulates cell–cell adhesion during spermatogenesis	[51,144]
Nectin-1	ND	Yes	Localizes IQGAP1 to cell–cell junctions	[150]
Nephrin	ND	Yes	Unknown	[48,49]
Podocin	ND	Yes	Unknown	[49]
VASP	ND	Yes	Unknown	[151]
VE-cadherin	ND	Yes	Regulates VE-cadherin localization at adherens junctions	[52]
Ca ²⁺ -binding proteins				
Calmodulin	Yes	Yes	Regulates IQGAP1 function	[11,18,19,20]

Table 1 (continued)

Interactor	Interaction <i>in vitro</i> ^a	Interaction <i>in vivo</i> ^b	Proposed function(s)	Reference(s)
Myosin ELC	Yes	ND	Unknown	[152]
S100B	Yes	Yes	Regulates membrane morphology	[153]
S100P	Yes	Yes	Regulates IQGAP1 function in MAPK signaling	[154]
Receptor tyrosine kinases				
EGFR	Yes	Yes	Regulates EGF-induced phosphorylation of EGFR and IQGAP1	[155,156]
FGFR1	Yes	Yes	Bridges FGFR1 to N-WASP-Arp2/3 complex	[130]
HER2	Yes	Yes	Regulates HER2 expression and signaling; modulates trastuzumab resistance	[157]
NGFR/TrkA	ND	Yes ^d	Unknown	[158]
PDGFβR	ND	Yes	Modulates focal adhesion assembly	[79]
VEGFR2	Yes	Yes	Cell migration and proliferation, vascular repair and maintenance, angiogenesis	[52,77]
Receptor serine/threonine kinases				
TGFβR2	Yes	Yes	Regulates TGFβR2 degradation and signaling	[159]
G protein-coupled receptors				
CXCR2	Yes	Yes	Unknown	[160]
GPR161	ND	Yes	Regulates cell migration and proliferation	[161]
KISS1R	ND	Yes	Connects KISS1R to EGFR activation	[162]
LPA1	ND	Yes	Regulates cell migration and invasion	[163]
Other receptors				
AMPA receptor, GluR4 subunit	ND	Yes	Regulates AMPA signaling and synaptic targeting	[164]
NMDAR	ND	Yes	Regulates NR2A signaling, dendritic spine density and memory	[69]
Lipids and lipid-associated proteins				
DGKζ	ND	Yes	Promotes phagocytosis by macrophages.	[165]
PIPKIγ	Yes	Yes	Recruits IQGAP1 to leading edge membrane	[166]
PLCε1	ND	Yes	Unknown	[57]
PtdIns3,4,5P ₃	Yes ^c	Yes	Unknown	[167,168]
PtdIns4,5P ₂	Yes	Yes	Promotes actin polymerization and branching	[166]
PTEN	ND	Yes	Unknown	[169]
Kinases and phosphatases				
Akt	ND	Yes	Regulates Akt activation, cardiac remodeling in response to pressure overload	[72,170–172]
AMPK	ND	Yes	Unknown	[94]
Aurora A	Yes	Yes	Stabilizes Aurora A	[173]
B-Raf	Yes	Yes	Regulates activation of B-Raf and its kinase activity; integrates Ca ²⁺ /calmodulin and B-Raf signaling	[16,17]
CaMKII	ND	Yes	Unknown	[69,174]
C-Raf	Yes	ND	Regulates MAPK activation	[72]
ERK1	Yes	Yes	Scaffold for MAP kinase signaling	[13]
ERK2	Yes	Yes	Scaffold for MAP kinase signaling	[12]
FAK	ND	Yes	Regulates cardiomyocyte hypertrophy and survival	[76]
MEK1	Yes	Yes	Scaffold for MAP kinase signaling	[13]
MEK2	Yes	Yes	Scaffold for MAP kinase signaling	[13]
MTOR	ND	Yes	Regulates cell proliferation	[26,172]
PAK6	ND	Yes	Regulates adherens junction disassembly	[175,176]

Table 1 (continued)

Interactor	Interaction <i>in vitro</i> ^a	Interaction <i>in vivo</i> ^b	Proposed function(s)	Reference(s)
PKA	ND	Yes	Promotes migration	[133]
PKC ϵ	ND	Yes	Substrate; regulates Cdc42 affinity and neurite outgrowth	[58,178]
PP2A	ND	Yes	Regulates interaction of integrins with cytoskeleton	[146,179]
PTP μ	Yes	Yes	Regulates Cdc42-dependent IQGAP1 function and mediates neurite outgrowth	[59]
Src	ND	Yes	Regulates endothelial cell proliferation and VEGF-induced angiogenesis	[78,136]
Scaffolds				
14-3-3	ND	Yes ^d	Unknown	[180]
AKAP79	Yes	Yes	Unknown	[177]
AKAP220	Yes	Yes	Integrates Ca ²⁺ and cAMP signals at the leading edge of migrating cells	[133]
β -arrestin2	ND	Yes	Forms complex with IQGAP1 and LPA1 or GPR161 to regulate cell migration	[161,163]
p14-MP1	ND	Yes	Regulates focal adhesion maturation	[181]
RACK1	ND	Yes	Unknown	[182]
ShcA	Yes	Yes	May couple RTKs to cytoskeleton	[183]
Small GTPases and their regulators				
Arf6	ND	Yes	Regulates Arf6-induced Rac1 activation and glioma cell migration	[184]
Asef	Yes	Yes	Regulates Rac1 activation to enhance endothelial barrier function	[185]
Cdc42	Yes	Yes	Inhibits intrinsic GTPase activity, increasing Cdc42GTP; promotes cell motility	[9,18,21,23]
FGD6	ND	Yes	Regulates podosome formation	[186]
K-Ras	ND	Yes	Regulates interaction of K-Ras with B-RAF	[15]
LRRK2	ND	Yes	Regulates the association of NFAT1 with IQGAP1	[187]
M-Ras	ND	Yes	Unknown	[188]
p190A-RhoGAP	ND	Yes	Inactivates RhoA to regulate airway smooth muscle contractility	[89]
Rab27a	Yes	Yes	Regulates endocytosis of insulin secretory membranes	[92]
Rac1	Yes	Yes	Inhibits intrinsic GTPase activity, increasing Rac1GTP; promotes cell motility	[21]
Rac2	ND	Yes	Unknown	[182]
RacGAP1	ND	Yes	Regulates cell migration and invasion	[189]
Ran	ND	Yes	Regulates β -catenin transcriptional function	[190]
Rap1	ND	Yes	Regulates activation of Rap1	[191]
RhoA	ND	Yes	Modulates RhoA activation; regulates cell proliferation and migration	[89,192]
RhoC	ND	Yes	Regulates RhoC-induced cell migration	[193,194]
TC10 (RhoQ)	Yes	ND	Unknown	[195]
Tiam1	ND	Yes	Unknown	[136]
Wnt signaling molecules				
Dvl	ND	Yes	Facilitates nuclear import of Dvl/ β -catenin complex and modulates Wnt signaling	[190,196]
LGR4	ND	Yes	Required for potentiation of β -catenin signaling by RSPO	[197]
MCAM	ND	Yes	Required for WRAMP structure assembly; bridges MCAM to cytoskeleton	[198]
Nuclear molecules				
ER α	Yes	Yes	Modulates ER α transcriptional function	[199]
ER β	Yes	Yes	Unknown	[199]
Importin- β 5	ND	Yes	Modulates nuclear import of the IQGAP1/ β -catenin/Dvl complex and transactivation of Wnt target genes	[190]

Table 1 (continued)

Interactor	Interaction <i>in vitro</i> ^a	Interaction <i>in vivo</i> ^b	Proposed function(s)	Reference(s)
Mediator	ND	Yes ^d	Unknown	[200]
Nardilysin	ND	Yes ^d	Unknown	[201]
NFAT	ND	Yes	Regulates nuclear translocation and function of NFAT	[202]
Nrf2	Yes	Yes	Stimulates the nuclear translocation and activation of HO-1 stress response	[203,204]
<i>NRON</i>	ND	Yes	Forms RNA-scaffold complex (with GSK3 β , DYRK, and CK1) to regulate NFAT	[202]
PCNA	ND	Yes	Unknown	[205]
PGC-1 α	ND	Yes	Unknown	[206]
RNase L	ND	Yes	Required for ECyd-induced JNK phosphorylation and apoptosis	[207]
RPA32	ND	Yes	Unknown	[205]
TULP3	ND	Yes	Unknown	[138]
WHSC1	ND	Yes	Unknown	[208]
mRNA regulators and co-chaperones				
Aha1	ND	Yes	Unknown	[209]
SMG-9	Yes	Yes	Unknown	[210]
Staufen	ND	Yes	Unknown	[211]
Microbial and viral interactors				
30-C12-HSL	Yes	Yes	<i>Pseudomonas aeruginosa</i> quorum sensing molecule that targets IQGAP1 to modulate epithelial cell migration	[114]
CSFV core protein	Yes	ND	Regulates growth and virulence of CSFV	[118]
Ebola virus VP40	ND	Yes	Regulates viral egress	[117]
Ibe	Yes	Yes	Unknown	[108]
MMLV MA	Yes	Yes	Regulates MMLV invasion and replication	[119]
SopE	ND	Yes	Regulates <i>S. typhimurium</i> invasion	[110]
SseI	Yes	Yes	Modulates SseI-induced inhibition of cell migration	[111]
Tir	Yes	Yes	Regulates actin pedestal formation by EPEC	[107]
YopM	Yes	ND	Promotes caspase-1 activation in <i>Y. pseudotuberculosis</i> -infected cells	[212]
Trafficking proteins				
Exo70	Yes	Yes	Regulates Exo70 subcellular localization	[91,213]
Sec3	Yes	Yes	Regulates formation and activity of invadopodia	[213]
Sec8	Yes	Yes	Regulates formation and activity of invadopodia	[91,213]
SEPT2	Yes	Yes	Regulates septin localization, filament organization and exocytosis	[91]
Syntaxin 1A	ND	Yes	Unknown	[91]
TSG101	Yes	Yes	Unknown	[120]
IQGAP2				
AKAP220	ND	Yes	Recruits active Rac1 to promote membrane ruffling	[214]
Arp2/3	ND	Yes	Regulates actin assembly downstream of thrombin stimulation	[127]
β -catenin	ND	Yes	Unknown	[215]
Calmodulin	Yes ^c	Yes	Unknown	[125,126]
Cdc42	ND	Yes	Inhibits GTPase activity	[126]
Ezrin	Yes ^c	ND	Unknown	[137]
F-actin	ND	Yes	Regulates actin assembly downstream of thrombin stimulation	[127]
LGR4	ND	Yes	Unknown	[197]
<i>NRON</i>	ND	Yes	Unknown	[202]

Table 1 (continued)

Interactor	Interaction <i>in vitro</i> ^a	Interaction <i>in vivo</i> ^b	Proposed function(s)	Reference(s)
PtdIns3,4,5P ₃	Yes	ND	Unknown	[168]
Rac1	ND	Yes	Inhibits GTPase activity	[126]
RhoG	ND	Yes	Unknown	[216]
IQGAP3				
Anillin	Yes ^c	Yes	Recruits IQGAP3 to the contractile ring during cytokinesis	[45]
Calmodulin	Yes ^c	ND	Unknown	[125]
Cdc42	Yes	Yes	Modulates neurite outgrowth in PC12 cells	[62,217]
DGKζ	ND	Yes	Unknown	[165]
ERK1	ND	Yes	Modulates ERK1 activation	[128]
F-actin	Yes	ND	Unknown	[62]
H-Ras	ND	Yes	Modulates Ras/ERK signaling	[217]
LGR4	ND	Yes	Unknown	[197]
Myosin ELC	Yes ^c	ND	Unknown	[125]
Rac1	Yes	Yes	Modulates neurite outgrowth in PC12 cells	[62,217]

^a*In vitro* interactions were demonstrated using pure proteins. In the absence of an *in vitro* interaction, direct binding between IQGAP and the target cannot be inferred. ND, not determined.
^b*In vivo* interactions were demonstrated by co-immunoprecipitation from cell lysate, pulldown with recombinant fusion protein from cell lysate, and/or co-localization unless otherwise noted. ND, not determined
^cInteraction with full-length IQGAP not reported.
^dInteraction identified via mass spectrometry. Targets in mass spectrometry databases not subject to peer review were not included in this table.

this interaction with a cell-permeable peptide inhibitor abrogates PTPμ-mediated neurite outgrowth. Cdc42 is among the best-characterized IQGAP1 binding partners (reviewed in [3,61]). IQGAP1 binding stabilizes active Cdc42 to regulate crosslinking of actin filaments, microtubule dynamics, and E-cadherin-mediated cell–cell adhesion. The studies described above imply that IQGAP1 facilitates changes in the actin cytoskeleton that are required for neurite outgrowth.

In contrast, decreasing endogenous IQGAP1 with siRNA did not impair nerve growth factor (NGF)-stimulated neurite outgrowth in PC12 rat pheochromocytoma cells [62]. However, reducing IQGAP3 attenuated neurite outgrowth induced by NGF. PC12 cells do not

contain IQGAP2 [62]. Therefore, the effect of knockdown of each IQGAP isoform was examined in hippocampal neurons. Reducing IQGAP2 or IQGAP3, but not IQGAP1, decreased axon elongation [62]. Several factors may account for the different reports of IQGAP1 on neurite outgrowth. These include different cell lines (N1E-115 versus PC12), different experimental strategies (induction with or without NGF), and different manipulations of IQGAP1 levels (over-expression versus knockdown).

IQGAP1 participates in neuronal proliferation and migration, which allows neurons to properly organize into a functional neural network. In cultured cerebellar neurons, IQGAP1 and lissencephaly 1 (Lis1) co-localize in axons and growth cones [63]. Lis1 is required

Figure 2.

Table 2. The biological roles of IQGAP1.

Physiology				
Relevant interactors	Cellular function	Physiological process	Putative role in disease	Citation
Kidney function				
Nephrin, Podocin, PLCε1	Organization of slit diaphragms	Glomerular filtration	Nephrotic syndrome	[48,49,57,218,219]
Neuronal function				
PKCε, PTPμ, Cdc42	Regulation of cytoskeleton for neurite outgrowth	Neurite outgrowth, development and maintenance of neurons	Epilepsy, memory formation/loss	[58,59,62]
Lis1, Cdc42, CLIP170, VEGF	Regulation of cytoskeleton for neural migration	Adult neurogenesis	Lissencephaly	[63,66]
Cardiovascular function				
Erk, Akt, Melusin	Erk and Akt activation following cardiac pressure overload	Cardiac remodeling	Myocardial infarction, cardiac hypertrophy	[72,76]
VEGFR2	Migration, proliferation	Neovascularization, angiogenesis	Cancer	[52,66,77,78,81]
αvβ3, EB1, Cortactin	Maintain cell–cell contacts that are linked to the cytoskeleton	Maintenance of vascular endothelial barrier functions	Acute systemic inflammatory diseases	[84,85]
PDGFR, Paxillin, Vincullin	PDGFR signaling for VSMC migration	Neointimal formation	Atherosclerosis, restenosis	[79]
Lung function				
RhoA, P190A-RhoGAP	Modulate RhoA and MLC activity	Airway smooth muscle cell contraction	Asthma	[89]
Insulin secretion				
Exocyst, Rab27a	Insulin secretion	Glucose homeostasis	Diabetes	[91,92]
Tumorigenesis				
Relevant interactors	Cellular function	Putative role in cancer		Citation
K-Ras, B-Raf, MEK1/2, ERK1/2	Proliferation, migration, invasion	Cell growth and differentiation, tumor invasion and metastasis		[101]
Akt, mTor	Proliferation, survival	Tumor growth, proliferation and survival		[172]
Rac1, Cdc42, Actin	Proliferation, migration, invasion	Cell growth and differentiation, tumor invasion and metastasis		[23,80]
Microbial infection				
Pathogen	Relevant interactor	Putative role in infection		Citation
<i>E. coli</i>	Tir	Actin pedestal formation, bacterial attachment		[107]
<i>E. coli</i>	Ibe	Pedestal recruitment, bacterial attachment		[108]
<i>E. coli</i> K1	β-catenin, actin	Disassembly of adherens junctions, invasion of brain endothelial cells, brain oedema in neonatal meningitis		[109]
<i>S. typhimurium</i>	Actin, Cdc42, Rac1, SopE	Actin polymerization and bacterial invasion		[110,112]
<i>S. typhimurium</i>	SseI	Chronic infection		[111]
<i>C. pneumoniae</i>	Unknown	Upregulation of IQGAP1, VSMC migration, atherosclerosis		[113]
<i>P. aeruginosa</i>	3O-C(12)-HSL	Modulates IQGAP1 expression, enhance host cell migration		[114,220]
Ebola virus	VP40	Viral egress		[117]
Marburg virus	TSG101	Viral egress		[121]
M-MULV	Gag	Viral egress		[119]
CSCV	Core protein	Viral egress		[118]

for neurogenesis, neuronal survival, and neuronal migration [64]. IQGAP1 co-immunoprecipitates with Lis1 and knockdown of IQGAP1 impairs neuronal motility [63]. Further, neuronal cells contain a multiprotein complex containing active Cdc42, Lis1,

IQGAP1, and CLIP-170, which appears necessary for optimal motility of neurons (Fig 2Bii). In migrating epithelial cells, IQGAP1 accumulates at the leading edge and associates with CLIP-170, linking Cdc42 and the cortical actin cytoskeleton to the microtubule

network (reviewed in [3]). In cultured cerebellar neurons, increasing intracellular free Ca^{2+} concentrations ($[\text{Ca}^{2+}]_i$) promoted the interaction of Lis1 with IQGAP1 and active Cdc42, suggesting IQGAP1 is a scaffold through which Lis1 links Ca^{2+} influx to Cdc42 and the cytoskeleton [63]. These results are consistent with previous studies showing Ca^{2+} /calmodulin binding to IQGAP1 regulates its interactions (reviewed in [61]).

Adult neurogenesis is the process by which neurons are generated from neural stem cells and progenitor cells. Neural progenitor cells (NPCs) migrate into niches and differentiate into neuronal precursors. Vascular endothelial growth factor (VEGF) stimulates this process [65]. In the absence of IQGAP1, VEGF was unable to stimulate migration of NPCs [66]. Consistent with these results, IQGAP1-null mice exhibit a delay in NPC differentiation. Cdc42, Rac1, and Lis1 binding to IQGAP1 is enhanced in VEGF-stimulated NPC migration [66]. This study supports a model in which IQGAP1 acts as an effector of a VEGF-dependent migratory signal for neural progenitor cells.

IQGAP1 contributes to the regulation of microtubules and the actin cytoskeleton that determines dendritic shape and morphology. Dendritic spines are actin-rich protrusions from a neuron that are responsible for transmission of signals from presynaptic neurons. The spine head connects to the shaft of the dendrite via a neck. Reduction of IQGAP1 in hippocampal neurons decreases the total number of dendrite tips, without significantly altering total dendrite length [67]. Moreover, in the rat hippocampus, the IQGAP1 CHD promotes spine head formation through interactions with the neural Wiskott–Aldrich syndrome protein (N-WASP)–actin-related protein 2/3 (Arp 2/3) complex, while the IQGAP1 GRD is essential for stalk extension [68] (Fig 2Biii). Disruption of the association between IQGAP1 and N-cadherin removes IQGAP1 from hippocampal dendritic spines heads [51]. Importantly, IQGAP1^{-/-} mice have decreased spine density and number in brain areas involved in cognition, emotion, and motivation [69]. IQGAP1^{-/-} mice also have long-term memory deficits, but anxiety and depression-like behavior are unaffected. Loss of dendritic spines are major contributing factors to psychiatric illness, such as schizophrenia and depression, and neurodegenerative disorders, such as Alzheimer's disease [70], and it is tempting to speculate that IQGAP1 may participate in the pathophysiology of these conditions.

Repeated seizures in temporal lobe epilepsy induce loss of neurons, especially from the CA1 and CA3 areas of the hippocampus. In a mouse model of epilepsy induced by pyramidal cell degeneration in the CA3 region, IQGAP1 expression was upregulated in CA1 pyramidal neurons [71]. Detailed analysis indicated that IQGAP1 is increased in uncommitted neural stem cells, leading the authors to speculate that IQGAP1 may contribute to the etiology of epileptogenesis. While additional studies are required to validate this hypothesis, the evidence implicating IQGAP1 in neurite outgrowth, spine development, synaptic plasticity, memory formation, and dendrite formation strongly supports a fundamental role for IQGAP1 in brain function.

The cardiovascular system

Cardiac functions Excessive pressure on the heart activates intracellular signaling pathways that regulate cardiac morphology. Although IQGAP1-null mice have normal basal heart function, prolonged pressure overload leads to unfavorable cardiac remodeling

with thinning of the ventricular walls, decreased contractility, and increased apoptosis [72]. Cardiac pressure overload activates focal adhesion kinase (FAK), which modulates ERK and Akt signaling that control cardiac remodeling [73]. Deletion of the non-receptor tyrosine kinase FAK from cardiac myocytes induces left ventricle thinning and blocks ERK activation [74]. Analogous to FAK, IQGAP1 modulates ERK and Akt activation in response to cardiac pressure overload [72]. At the molecular level, long-term (4-day) transverse aortic band-induced chronic pressure overload of wild-type mouse cardiomyocytes (heart muscle cells) stimulates activation of MEK and ERK, which promote proliferation, and Akt, a kinase that promotes survival [72]. By contrast, MEK, ERK, and Akt activation were abrogated in mice deficient in IQGAP1 [72]. Pressure overload upregulates melusin, a muscle-specific protein [75]. An IQGAP1–melusin complex mediates ERK activation in response to pressure overload [76] (Fig 2C). Additionally, IQGAP1 contribution to cardiac function was demonstrated with transgenic mice overexpressing melusin in the heart and double-transgenic mice that overexpress melusin, but lack IQGAP1. In the absence of IQGAP1, ERK activity was reduced in response to pressure overload and apoptotic death was increased in response to stress, demonstrating a role for IQGAP1 in cardiomyocyte survival [76]. Taken together, these observations implicate IQGAP1 as a signaling platform in cardiac remodeling and morphology.

Vascular functions IQGAP1 influences blood vessel formation. VEGF affects virtually all aspects of blood vessel formation and function. IQGAP1 binds to the VEGF receptor 2 (VEGFR2) and is necessary for VEGF-stimulated endothelial cell migration and proliferation [77] (Fig 2Di). These observations imply that IQGAP1 scaffolds VEGFR2 signaling in maintenance and repair of blood vessels. Subsequent studies showed that the IQGAP1/VEGFR2 interaction regulates angiogenesis. For example, IQGAP1 knockdown suppresses VEGF-stimulated angiogenesis in an *in vivo* model of chicken chorioallantoic membrane [78]. Additional evidence linking IQGAP1 to angiogenesis is derived from studies in mice. Blood vessel formation in response to injury is impaired in mice lacking IQGAP1 [79]. Further, IQGAP1 expression is increased in angiogenesis following ischemia [52] and overexpression of IQGAP1 significantly increased angiogenesis in an *in vivo* mouse tumor model [80]. Finally, IQGAP1-null mice have reduced recovery of blood flow to the leg after hindlimb ischemia [81], further demonstrating the contribution of IQGAP1 to angiogenesis.

Vascular endothelial cells form the barrier between blood and tissues, and disruption of the barrier can result in acute systemic inflammatory diseases. Reduction of IQGAP1 disrupts vascular endothelial barrier integrity [82]. Integrins are important mediators of endothelial barrier function. Mice lacking integrin $\beta 3$ have increased endothelial blood vessel leak in response to VEGF-stimulation [83]. IQGAP1 binds integrin $\beta 3$, and IQGAP1-null mice have reduced localization of integrin $\alpha v\beta 3$ to the cell–cell junction and increased lung vascular permeability [84] (Fig 2Dii). Multiple cytoskeletal signaling proteins, including microtubule plus end binding protein 1 (EB1) and cortactin, control endothelial permeability. A complex comprising IQGAP1, EB1, and cortactin links the actin and microtubule cytoskeletons to strengthen endothelial barrier [85]. Barrier integrity is also affected by shear stress, the mechanical force exerted on endothelial cells by the flow of blood.

IQGAP1 is essential for maintaining endothelial cell alignment under shear stress [86]. Adhesion and alignment of endothelial cells exposed to shear stress is impaired by IQGAP1 knockdown, suggesting that IQGAP1 stabilizes adherens junctions under blood flow. By controlling blood vessel formation and barrier integrity, IQGAP1 is a critical integrator of multiple vascular processes.

Lung function

Asthma is a chronic inflammatory disease that affects ~235 million people and results from airway smooth muscle contraction. Exercise, allergens, microbes, or other stimuli activate the parasympathetic nervous system, leading to release of acetylcholine and histamine, which activate receptors on airway smooth muscle cells to promote contraction [87] (Fig 2E). These receptors induce Ca^{2+} release from intracellular stores and RhoA activation, resulting in myosin light chain (MLC) phosphorylation, enhancing the interaction of myosin with actin, thereby promoting airway smooth muscle cell contractility [88].

IQGAP1 modulates this process [89] (Fig 2E). IQGAP1 co-immunoprecipitates with RhoA and p190A-RhoGAP, a protein that inactivates RhoA, from airway smooth muscle cells. Knockdown of IQGAP1 decreases the RhoA/p190A-RhoGAP co-localization. Consistent with these results, IQGAP1^{-/-} mice have enhanced airway responsiveness, and increased levels of MLC phosphorylation and active RhoA in the posterior trachea [89]. Moreover, IQGAP1 was significantly lower in airway smooth muscle biopsies from patients with asthma than from healthy controls. Collectively, these data imply that IQGAP1 may contribute to the severity of asthma by controlling airway smooth muscle contractility.

Insulin secretion

Increased blood glucose concentration induces insulin release from pancreatic β -cells. Glucose enters the β -cells where it is metabolized, leading to a rise in $[\text{Ca}^{2+}]_i$, which triggers exocytosis of insulin granules [90]. A complex comprising eight subunits, termed the exocyst, tethers insulin-containing vesicles inducing release of insulin at the plasma membrane. IQGAP1 co-immunoprecipitates with the exocyst complex [91]. Knockdown of IQGAP1 significantly reduced the ability of glucose to stimulate insulin secretion from β -cells (Fig 2F). Another mechanism by which IQGAP1 may contribute to insulin secretion is via Rab27a. IQGAP1 forms a complex with Rab27a [92], a small GTPase that is highly expressed in pancreatic β -cells and regulates endocytosis of insulin secretory membranes. Reducing expression of endogenous IQGAP1 with siRNA prevented glucose-induced redistribution of Rab27a from the cytosol to the plasma membrane [92]. Analysis revealed that an association between IQGAP1 and Rab27a is required for endocytosis of secretory membranes. Thus, IQGAP1 participates in both exocytosis and endocytosis of insulin secretory vesicles in response to glucose stimulation (Fig 2F).

Energy homeostasis and insulin secretion are regulated by AMP-activated protein kinase (AMPK) [93]. IQGAP1 was recently identified as an interactor of AMPK, and the proteins co-immunoprecipitated from pancreatic β -cells [94]. Although there is no evidence that this association contributes to β -cell function, the preponderance of evidence suggests that IQGAP1 participates in insulin secretion.

IQGAP2 is expressed predominantly in the liver, an organ that is central to glucose regulation. Knockout mouse models implicate

IQGAP2 in glucose homeostasis. IQGAP2^{-/-} mice had insulin levels similar to those in wild-type mice, but lower fasting blood glucose levels and enhanced insulin sensitivity during a glucose tolerance test [95]. IQGAP2 deficiency led to loss of facilitated long-chain fatty acid synthesis and protection from diet-induced hepatic steatosis. However, conflicting findings were subsequently reported. Another group observed higher blood glucose and insulin levels in IQGAP2-null mice [96]. The IQGAP2^{-/-} mice exhibited aberrant hepatic regulation of glycogenolysis, gluconeogenesis, and lipid homeostasis, leading the authors to conclude that IQGAP2 deficiency predisposes to non-alcoholic fatty liver disease. These differences require further investigation. One notable distinction between the studies was the different genetic backgrounds of the mice, SV129J versus C57BL/6J. While the molecular mechanism is unknown, the collective data argue for the involvement of IQGAP2 in glucose homeostasis.

IQGAP1 as a therapeutic target

Carcinogenesis

Despite advances in chemotherapy, treatment often kills healthy cells, producing severe side effects. Approximately 30% of human neoplasms have mutations in Ras and B-Raf that overactivate ERK [97], promoting tumor proliferation and migration. Although therapeutics targeting B-Raf (e.g., sorafenib, vemurafenib, and dabrafenib) have been developed, responses are highly variable and resistance is common [98]. Therefore, additional molecularly targeted cancer therapeutics are required. IQGAP1 is potentially a new target (Table 2). IQGAP1 is overexpressed in human cancer (reviewed in [99,100]). Overexpression of IQGAP1 is associated with enhanced tumor proliferation, invasion, and angiogenesis [80]. By interacting with several MAPK components, IQGAP1 mediates optimal ERK activation [4,5]. Initial evidence suggests that targeting the IQGAP1/MAPK pathway associations is feasible. Treatment of mice with cell-permeable peptides (corresponding to the WW domain of IQGAP1) disrupts IQGAP1-ERK1/2 interactions and inhibits Ras-driven tumorigenesis [101]. Importantly, the peptides attenuated proliferation of melanoma cells resistant to the B-Raf inhibitor vemurafenib.

Neoplastic transformation by Ras and other oncoproteins often relies on the Rho GTPases, Cdc42, and Rac1 [102]. Cdc42 and Rac1 are not mutated in cancer, but deregulation of their function leads to carcinogenesis [102]. IQGAP1 inhibits the intrinsic GTPase activity of Cdc42 and Rac1 to stabilize the GTP-bound, active forms [23]. Overexpression of IQGAP1 increases the pool of active Cdc42 and Rac1, while knockdown of endogenous IQGAP1 significantly decreases the amount of active Cdc42 and Rac1 in mammalian cells [23,80]. A dominant-negative IQGAP1 construct, which decreases the amount of GTP-bound Cdc42 in cell lysates [23], reduces neoplastic transformation of malignant MCF-7 human breast epithelial cells [80]. These results suggest that blocking the formation of IQGAP1-Cdc42 and IQGAP1-Rac1 complexes will decrease the amount of active Cdc42 and Rac1 in carcinoma cells, reducing tumorigenesis.

Small-molecule inhibitors that disrupt the binding of IQGAP1 to select interactors may be specific chemotherapeutic agents. Targeting a protein-protein interaction (PPI) with a small molecule was thought to be difficult due the large, flat surface areas involved in binding. However, the dynamic PPI interface provides more opportunities for

small molecule binding than traditional 'druggable' binding pockets [103]. Several small-molecule PPI inhibitors are at various stages of development, including phase III clinical trials [104]. As IQGAP1 is an oncogene, but is not required for viability [105], it is an attractive molecule for the development of targeted chemotherapy (Table 2).

Microbial infection

Antibiotics are essential for treating bacterial infection. Typically, antibiotics target bacterial enzymes to inhibit processes such as cell-wall synthesis and protein translation. However, bacteria frequently develop resistance to antibiotics. Novel strategies to combat infection are needed.

Most microbial pathogens usurp signaling pathways of the host cell, particularly cytoskeletal dynamics [106]. Bacterial pathogens manipulate the cytoskeleton to invade the host cell, move within the cell, form vacuoles, and avoid phagocytosis. The role of IQGAP1 in regulation of the cytoskeleton led to investigation of its participation in microbial infection (Table 2). The best-characterized examples include *Escherichia coli*, which usurps IQGAP1 to promote formation of actin pedestals [107,108] and disassembly of adherens junctions [109], and *Salmonella typhimurium*, which injects proteins that 'hijack' IQGAP1 to modulate the cytoskeleton for invasion into host cells [110–112]. More recently, *Chlamydia pneumonia* [113] and *Pseudomonas aeruginosa* [114] were observed to regulate IQGAP1 expression to alter cell adhesion and migration. Potentially, inhibition of IQGAP1 interactions with bacterial proteins could control bacterial infection. A benefit of targeting a host protein is the reduced likelihood of mutation, which commonly occurs with antibiotics directed at bacterial proteins. Disrupting a host protein may produce systemic side effects. The benefits of treatment versus off-target effects are a fundamental question in the therapy of many diseases. Nevertheless, in light of the increasing problem of antibiotic resistance and the lack of new antibiotics coming to market [115], alternative strategies may yield promising results.

During their life cycle, viruses utilize host-cell proteins to mediate entry, replication and budding of viral particles to establish and maintain infection [116]. IQGAP1 interacts with several viral proteins, including Ebola virus protein VP40 [117], classical swine fever virus (CSFV) core protein [118], and Moloney murine leukemia virus (M-MuLV) matrix protein [119] (Table 2). Mutations of these viral proteins that prevent interaction with IQGAP1 or depletion of IQGAP1 from infected cells interfered with viral life cycle. IQGAP1 also forms a complex with host protein TSG101 [120], which mediates release of the Marburg virus [121]. Depletion of IQGAP1 reduced the release of Marburg virus particles. These findings suggest that IQGAP1 plays a critical role in the life cycle of several viruses and is a potential target for antiviral medication.

Conclusions

Accumulating evidence supports diverse roles for IQGAPs in vertebrates. At the molecular level, IQGAPs scaffold multiprotein complexes that regulate similar processes in different tissues. For example, modulation of cytoskeletal dynamics by the association of IQGAP1 with actin, small GTPases and microtubule binding proteins is critical for controlling tissue integrity and morphology. This role is evident in organizing renal slit diaphragms for

Sidebar A: In need of answers

- (i) Do IQGAP1, IQGAP2, and IQGAP3 have differential roles in specific tissues? Do the three IQGAPs have unique, redundant, or complementary functions in physiology?
- (ii) What regulates the interactions of IQGAPs with specific binding partners? Are these complexes tissue specific and/or IQGAP isoform specific? How do IQGAP protein complexes influence cancer, microbial infection, and other diseases?

glomerular filtration [48,49], controlling neural cell morphology for coordinating neural networks [51,67–69], regulating neural cell migration [58,59,63], and maintaining endothelial integrity and stability for barrier functions of blood vessels [77–79,81,82]. Another conserved role for IQGAPs across tissues is the scaffolding of cell signaling pathways, such as MAPK. IQGAP1 enhances activation of MAPK, but different tissues may have different responses. In the kidney, angiotensin II enhances IQGAP1-regulated MAPK signaling to contribute to apoptosis [56], whereas pressure overload of cardiomyocytes promotes IQGAP1-regulated activation of MAPK that leads to cardiac hypertrophy and survival [72,122]. IQGAP1 association with proteins or receptors that have restricted tissue expression may mediate specific cellular responses. For example, the interaction of the muscle-specific protein melusin with IQGAP1 enhances MAPK signaling in cardiomyocytes in response to pressure overload [76].

Although the functions of IQGAP1 have been evaluated in several tissues, the unique, redundant, or complementary roles for IQGAP1, IQGAP2, and IQGAP3 require further investigation. Unique functions may be conferred by the distinct tissue expression of IQGAP isoforms. IQGAP1 is ubiquitously expressed, IQGAP2 is predominantly expressed in liver, while IQGAP3 expression is mainly in the brain [62]. Variations in IQGAP isoform sequence may also contribute to specialized IQGAP functions. The amino acid sequences of IQGAP2 and IQGAP3 are 62 and 59%, respectively, identical to IQGAP1. Therefore, it is possible that IQGAPs are differentially regulated through specific post-translational modifications at residues that are not conserved among all three proteins. For example, quantitative phosphoproteomics studies have identified phosphorylation of IQGAP1 at Ser-330 [123,124], a residue that is not conserved in IQGAP2 or IQGAP3. Further, while IQGAPs share some binding partners, including calmodulin [18,19,125,126] and F-actin [19,20,62,127], differences have been reported. Although both IQGAP1 and IQGAP3 associate with ERK proteins, IQGAP3 binds only ERK1 [128], while IQGAP1 interacts with both ERK1 [13] and ERK2 [12]. Additionally, IQGAP3 co-immunoprecipitates with anillin, whereas IQGAP1 and IQGAP2 do not [45]. Anillin recruits IQGAP3 for specific roles in cytokinesis, yet IQGAP1 may play a complementary role in this process as loss of either IQGAP1 or IQGAP3 leads to defects in cytokinesis. Isoform-specific knockout studies, including tissue specific knockouts, are needed to elucidate the biological roles of the three IQGAP proteins.

IQGAP1 is overexpressed in a variety of cancers [99,100]. Potentially, inhibitors of IQGAP1 functions could prevent tumor invasion, proliferation, and migration. Preliminary studies targeting IQGAP1 are encouraging [101], but efficacy in humans and

potential side effects need to be established. In the 20 years since their discovery, the identified roles of IQGAP proteins have expanded from cytoskeletal regulators to modulators of diverse functions in several organs. We look forward to future studies that expand upon the distinct roles of IQGAPs in physiology and disease.

Acknowledgements

We apologize to those authors whose primary work was omitted due to space restrictions. This work was supported by the Intramural Research Program of the National Institutes of Health.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Lee IJ, Coffman VC, Wu JQ (2012) Contractile-ring assembly in fission yeast cytokinesis: recent advances and new perspectives. *Cytoskeleton* 69: 751–763
- Shannon KB (2012) IQGAP Family members in yeast, *Dictyostelium*, and mammalian cells. *Int J Cell Biol* 2012: 894817
- Noritake J, Watanabe T, Sato K, Wang S, Kaibuchi K (2005) IQGAP1: a key regulator of adhesion and migration. *J Cell Sci* 118: 2085–2092
- Brown MD, Sacks DB (2006) IQGAP1 in cellular signaling: bridging the GAP. *Trends Cell Biol* 16: 242–249
- White CD, Erdemir HH, Sacks DB (2012) IQGAP1 and its binding proteins control diverse biological functions. *Cell Signal* 24: 826–834
- Osman M (2010) An emerging role for IQGAP1 in regulating protein traffic. *Sci World J* 10: 944–953
- Briggs MW, Sacks DB (2003) IQGAP proteins are integral components of cytoskeletal regulation. *EMBO Rep* 4: 571–574
- Mateer SC, Wang N, Bloom GS (2003) IQGAPs: integrators of the cytoskeleton, cell adhesion machinery, and signaling networks. *Cell Motil Cytoskeleton* 55: 147–155
- Mataraza JM, Briggs MW, Li Z, Entwistle A, Ridley AJ, Sacks DB (2003) IQGAP1 promotes cell motility and invasion. *J Biol Chem* 278: 41237–41245
- Kuroda S, Fukata M, Nakagawa M, Fujii K, Nakamura T, Ookubo T, Izawa I, Nagase T, Nomura N, Tani H et al (1998) Role of IQGAP1, a target of the small GTPases Cdc42 and Rac1, in regulation of E-cadherin-mediated cell-cell adhesion. *Science* 281: 832–835
- Li Z, Kim SH, Higgins JM, Brenner MB, Sacks DB (1999) IQGAP1 and calmodulin modulate E-cadherin function. *J Biol Chem* 274: 37885–37892
- Roy M, Li Z, Sacks DB (2004) IQGAP1 binds ERK2 and modulates its activity. *J Biol Chem* 279: 17329–17337
- Roy M, Li Z, Sacks DB (2005) IQGAP1 is a scaffold for mitogen-activated protein kinase signaling. *Mol Cell Biol* 25: 7940–7952
- Pearson G, Robinson F, Beers Gibson T, Xu BE, Karandikar M, Berman K, Cobb MH (2001) Mitogen-activated protein (MAP) kinase pathways: regulation and physiological functions. *Endocr Rev* 22: 153–183
- Matsunaga H, Kubota K, Inoue T, Isono F, Ando O (2014) IQGAP1 selectively interacts with K-Ras but not with H-Ras and modulates K-Ras function. *Biochem Biophys Res Commun* 444: 360–364
- Ren JG, Li Z, Sacks DB (2007) IQGAP1 modulates activation of B-Raf. *Proc Natl Acad Sci U S A* 104: 10465–10469
- Ren JG, Li Z, Sacks DB (2008) IQGAP1 integrates Ca²⁺/calmodulin and B-Raf signaling. *J Biol Chem* 283: 22972–22982
- Hart MJ, Callow MG, Souza B, Polakis P (1996) IQGAP1, a calmodulin-binding protein with a rasGAP-related domain, is a potential effector for cdc42Hs. *EMBO J* 15: 2997–3005
- Ho YD, Joyal JL, Li Z, Sacks DB (1999) IQGAP1 integrates Ca²⁺/calmodulin and Cdc42 signaling. *J Biol Chem* 274: 464–470
- Mateer SC, McDaniel AE, Nicolas V, Habermacher GM, Lin MJ, Cromer DA, King ME, Bloom GS (2002) The mechanism for regulation of the F-actin binding activity of IQGAP1 by calcium/calmodulin. *J Biol Chem* 277: 12324–12333
- Kuroda S, Fukata M, Kobayashi K, Nakafuku M, Nomura N, Iwamatsu A, Kaibuchi K (1996) Identification of IQGAP as a putative target for the small GTPases, Cdc42 and Rac1. *J Biol Chem* 271: 23363–23367
- Mataraza JM, Briggs MW, Li Z, Frank R, Sacks DB (2003) Identification and characterization of the Cdc42-binding site of IQGAP1. *Biochem Biophys Res Commun* 305: 315–321
- Swart-Mataraza JM, Li Z, Sacks DB (2002) IQGAP1 is a component of Cdc42 signaling to the cytoskeleton. *J Biol Chem* 277: 24753–24763
- Erickson JW, Cerione RA, Hart MJ (1997) Identification of an actin cytoskeletal complex that includes IQGAP and the Cdc42 GTPase. *J Biol Chem* 272: 24443–24447
- Fukata M, Kuroda S, Fujii K, Nakamura T, Shoji I, Matsuura Y, Okawa K, Iwamatsu A, Kikuchi A, Kaibuchi K (1997) Regulation of cross-linking of actin filament by IQGAP1, a target for Cdc42. *J Biol Chem* 272: 29579–29583
- Wang JB, Sonn R, Tekletsadik YK, Samorodnitsky D, Osman MA (2009) IQGAP1 regulates cell proliferation through a novel CDC42-mTOR pathway. *J Cell Sci* 122: 2024–2033
- Li R (2007) Cytokinesis in development and disease: variations on a common theme. *Cell Mol Life Sci* 64: 3044–3058
- Epp JA, Chant J (1997) An IQGAP-related protein controls actin-ring formation and cytokinesis in yeast. *Curr Biol* 7: 921–929
- Osman MA, Cerione RA (1998) Iqg1p, a yeast homologue of the mammalian IQGAPs, mediates cdc42p effects on the actin cytoskeleton. *J Cell Biol* 142: 443–455
- Shannon KB, Li R (1999) The multiple roles of Cyk1p in the assembly and function of the actomyosin ring in budding yeast. *Mol Biol Cell* 10: 283–296
- Chang F, Woollard A, Nurse P (1996) Isolation and characterization of fission yeast mutants defective in the assembly and placement of the contractile actin ring. *J Cell Sci* 109(Pt 1): 131–142
- Eng K, Naqvi NI, Wong KC, Balasubramanian MK (1998) Rng2p, a protein required for cytokinesis in fission yeast, is a component of the actomyosin ring and the spindle pole body. *Curr Biol* 8: 611–621
- Wu JQ, Kuhn JR, Kovar DR, Pollard TD (2003) Spatial and temporal pathway for assembly and constriction of the contractile ring in fission yeast cytokinesis. *Dev Cell* 5: 723–734
- Li CR, Wang YM, Wang Y (2008) The IQGAP Iqg1 is a regulatory target of CDK for cytokinesis in *Candida albicans*. *EMBO J* 27: 2998–3010
- Vlahou G, Rivero F (2006) Rho GTPase signaling in *Dictyostelium discoideum*: insights from the genome. *Eur J Cell Biol* 85: 947–959
- Adachi H, Takahashi Y, Hasebe T, Shirouzu M, Yokoyama S, Sutoh K (1997) Dictyostelium IQGAP-related protein specifically involved in the completion of cytokinesis. *J Cell Biol* 137: 891–898

37. Faix J, Dittrich W (1996) DGAP1, a homologue of rasGTPase activating proteins that controls growth, cytokinesis, and development in *Dictyos-
telium discoideum*. *FEBS Lett* 394: 251–257
38. Faix J, Weber I, Mintert U, Kohler J, Lottspeich F, Marriott G (2001) Recruitment of cortexillin into the cleavage furrow is controlled by Rac1 and IQGAP-related proteins. *EMBO J* 20: 3705–3715
39. Kee YS, Ren Y, Dorfman D, Iijima M, Firtel R, Iglesias PA, Robinson DN (2012) A mechanosensory system governs myosin II accumulation in dividing cells. *Mol Biol Cell* 23: 1510–1523
40. Skop AR, Liu H, Yates J 3rd, Meyer BJ, Heald R (2004) Dissection of the mammalian midbody proteome reveals conserved cytokinesis mechanisms. *Science* 305: 61–66
41. Bielak-Zmijewska A, Kolano A, Szczepanska K, Maleszewski M, Borsuk E (2008) Cdc42 protein acts upstream of IQGAP1 and regulates cytokinesis in mouse oocytes and embryos. *Dev Biol* 322: 21–32
42. Piekny AJ, Maddox AS (2010) The myriad roles of anillin during cytokinesis. *Semin Cell Dev Biol* 21: 881–891
43. Almonacid M, Celton-Morizur S, Jakubowski JL, Dingli F, Loew D, Mayeux A, Chen JS, Gould KL, Clifford DM, Paoletti A (2011) Temporal control of contractile ring assembly by Plc1 regulation of myosin II recruitment by Mid1/anillin. *Curr Biol* 21: 473–479
44. Padmanabhan A, Bakka K, Sevugan M, Naqvi NI, D'Souza V, Tang X, Mishra M, Balasubramanian MK (2011) IQGAP-related Rng2p organizes cortical nodes and ensures position of cell division in fission yeast. *Curr Biol* 21: 467–472
45. Adachi M, Kawasaki A, Nojima H, Nishida E, Tsukita S (2014) Involvement of IQGAP family proteins in the regulation of mammalian cell cytokinesis. *Genes Cells* 19: 803–820
46. Grahmmer F, Schell C, Huber TB (2013) The podocyte slit diaphragm - from a thin grey line to a complex signalling hub. *Nat Rev Nephrol* 9: 587–598
47. Wiggins RC (2007) The spectrum of podocytopathies: a unifying view of glomerular diseases. *Kidney Int* 71: 1205–1214
48. Lehtonen S, Ryan JJ, Kudlicka K, Iino N, Zhou H, Farquhar MG (2005) Cell junction-associated proteins IQGAP1, MAGI-2, CASK, spectrins, and alpha-actinin are components of the nephrin multiprotein complex. *Proc Natl Acad Sci U S A* 102: 9814–9819
49. Rigother C, Auguste P, Welsh GI, Lepreux S, Deminiere C, Mathieson PW, Saleem MA, Ripoche J, Combe C (2012) IQGAP1 interacts with components of the slit diaphragm complex in podocytes and is involved in podocyte migration and permeability in vitro. *PLoS ONE* 7: e37695
50. Harris TJ, Tepass U (2010) Adherens junctions: from molecules to morphogenesis. *Nat Rev Mol Cell Biol* 11: 502–514
51. Schrick C, Fischer A, Srivastava DP, Tronson NC, Penzes P, Radulovic J (2007) N-cadherin regulates cytoskeletally associated IQGAP1/ERK signaling and memory formation. *Neuron* 55: 786–798
52. Yamaoka-Tojo M, Tojo T, Kim HW, Hilenski L, Patrushev NA, Zhang L, Fukai T, Ushio-Fukai M (2006) IQGAP1 mediates VE-cadherin-based cell-cell contacts and VEGF signaling at adherence junctions linked to angiogenesis. *Arterioscler Thromb Vasc Biol* 26: 1991–1997
53. Briggs MW, Li Z, Sacks DB (2002) IQGAP1-mediated stimulation of transcriptional co-activation by β -catenin is modulated by calmodulin. *J Biol Chem* 277: 7453–7465
54. Ding G, Reddy K, Kapasi AA, Franki N, Gibbons N, Kasinath BS, Singhal PC (2002) Angiotensin II induces apoptosis in rat glomerular epithelial cells. *Am J Physiol Renal Physiol* 283: F173–F180
55. Grande MT, Lopez-Novoa JM (2008) Therapeutical relevance of MAP-kinase inhibitors in renal diseases: current knowledge and future clinical perspectives. *Curr Med Chem* 15: 2054–2070
56. Liu Y, Liang W, Yang Q, Ren Z, Chen X, Zha D, Singhal PC, Ding G (2013) IQGAP1 mediates angiotensin II-induced apoptosis of podocytes via the ERK1/2 MAPK signaling pathway. *Am J Nephrol* 38: 430–444
57. Hinkes B, Wiggins RC, Gbadegesin R, Vlangos CN, Seelow D, Nurnberg G, Garg P, Verma R, Chaib H, Hoskins BE et al (2006) Positional cloning uncovers mutations in PLCE1 responsible for a nephrotic syndrome variant that may be reversible. *Nat Genet* 38: 1397–1405
58. Li Z, McNulty DE, Marler KJ, Lim L, Hall C, Annan RS, Sacks DB (2005) IQGAP1 promotes neurite outgrowth in a phosphorylation-dependent manner. *J Biol Chem* 280: 13871–13878
59. Phillips-Mason PJ, Gates TJ, Major DL, Sacks DB, Brady-Kalnay SM (2006) The receptor protein-tyrosine phosphatase PTPmu interacts with IQGAP1. *J Biol Chem* 281: 4903–4910
60. Hellberg CB, Burden-Gulley SM, Pietz GE, Brady-Kalnay SM (2002) Expression of the receptor protein-tyrosine phosphatase, PTPmu, restores E-cadherin-dependent adhesion in human prostate carcinoma cells. *J Biol Chem* 277: 11165–11173
61. Briggs MW, Sacks DB (2003) IQGAP1 as signal integrator: Ca^{2+} , calmodulin, Cdc42 and the cytoskeleton. *FEBS Lett* 542: 7–11
62. Wang S, Watanabe T, Noritake J, Fukata M, Yoshimura T, Itoh N, Harada T, Nakagawa M, Matsuura Y, Arimura N et al (2007) IQGAP3, a novel effector of Rac1 and Cdc42, regulates neurite outgrowth. *J Cell Sci* 120: 567–577
63. Kholmanskikh SS, Koeller HB, Wynshaw-Boris A, Gomez T, Letourneau PC, Ross ME (2006) Calcium-dependent interaction of Lis1 with IQGAP1 and Cdc42 promotes neuronal motility. *Nat Neurosci* 9: 50–57
64. Wynshaw-Boris A, Pramparo T, Youn YH, Hirotsune S (2010) Lissencephaly: mechanistic insights from animal models and potential therapeutic strategies. *Semin Cell Dev Biol* 21: 823–830
65. Jin K, Zhu Y, Sun Y, Mao XO, Xie L, Greenberg DA (2002) Vascular endothelial growth factor (VEGF) stimulates neurogenesis in vitro and in vivo. *Proc Natl Acad Sci U S A* 99: 11946–11950
66. Balenci L, Saoudi Y, Grunwald D, Deloulme JC, Bouron A, Bernards A, Baudier J (2007) IQGAP1 regulates adult neural progenitors in vivo and vascular endothelial growth factor-triggered neural progenitor migration in vitro. *J Neurosci* 27: 4716–4724
67. Swiech L, Blazejczyk M, Urbanska M, Pietruszka P, Dortland BR, Malik AR, Wulf PS, Hoogenraad CC, Jaworski J (2011) CLIP-170 and IQGAP1 cooperatively regulate dendrite morphology. *J Neurosci* 31: 4555–4568
68. Jausoro I, Mestres I, Quassollo G, Masseroni L, Heredia F, Caceres A (2013) Regulation of spine density and morphology by IQGAP1 protein domains. *PLoS ONE* 8: e56574
69. Gao C, Frausto SF, Guedea AL, Tronson NC, Jovasevic V, Leaderbrand K, Corcoran KA, Guzman YF, Swanson GT, Radulovic J (2011) IQGAP1 regulates NR2A signaling, spine density, and cognitive processes. *J Neurosci* 31: 8533–8542
70. Koleske AJ (2013) Molecular mechanisms of dendrite stability. *Nat Rev Neurosci* 14: 536–550
71. Yi MH, Kim S, Zhang E, Kang JW, Park JB, Lee YH, Chung CK, Kim YM, Kim DW (2013) IQGAP1 expression in spared CA1 neurons after an excitotoxic lesion in the mouse hippocampus. *Cell Mol Neurobiol* 33: 1003–1012
72. Sbroggio M, Carnevale D, Bertero A, Cifelli G, De Blasio E, Mascio G, Hirsch E, Bahou WF, Turco E, Silengo L et al (2011) IQGAP1 regulates

- ERK1/2 and AKT signalling in the heart and sustains functional remodelling upon pressure overload. *Cardiovasc Res* 91: 456–464
73. Franchini KG, Torsoni AS, Soares PH, Saad MJ (2000) Early activation of the multicomponent signaling complex associated with focal adhesion kinase induced by pressure overload in the rat heart. *Circ Res* 87: 558–565
 74. Domingos PP, Fonseca PM, Nadruz W Jr, Franchini KG (2002) Load-induced focal adhesion kinase activation in the myocardium: role of stretch and contractile activity. *Am J Physiol Heart Circ Physiol* 282: H556–H564
 75. Sbroglio M, Ferretti R, Percivalle E, Gutkowska M, Zylicz A, Michowski W, Kuznicki J, Accornero F, Pacchioni B, Lanfranchi G et al (2008) The mammalian CHORD-containing protein melusin is a stress response protein interacting with Hsp90 and Sgt1. *FEBS Lett* 582: 1788–1794
 76. Sbroglio M, Bertero A, Velasco S, Fusella F, De Blasio E, Bahou WF, Silengo L, Turco E, Brancaccio M, Tarone G (2011) ERK1/2 activation in heart is controlled by melusin, focal adhesion kinase and the scaffold protein IQGAP1. *J Cell Sci* 124: 3515–3524
 77. Yamaoka-Tojo M, Ushio-Fukai M, Hilenski L, Dikalov SI, Chen YE, Tojo T, Fukai T, Fujimoto M, Patrushev NA, Wang N et al (2004) IQGAP1, a novel vascular endothelial growth factor receptor binding protein, is involved in reactive oxygen species-dependent endothelial migration and proliferation. *Circ Res* 95: 276–283
 78. Meyer RD, Sacks DB, Rahimi N (2008) IQGAP1-dependent signaling pathway regulates endothelial cell proliferation and angiogenesis. *PLoS ONE* 3: e3848
 79. Kohno T, Urao N, Ashino T, Sudhahar V, Inomata H, Yamaoka-Tojo M, McKinney RD, Fukai T, Ushio-Fukai M (2013) IQGAP1 links PDGF receptor-beta signal to focal adhesions involved in vascular smooth muscle cell migration: role in neointimal formation after vascular injury. *Am J Physiol Cell Physiol* 305: C591–C600
 80. Jadeski L, Mataraza JM, Jeong HW, Li Z, Sacks DB (2008) IQGAP1 stimulates proliferation and enhances tumorigenesis of human breast epithelial cells. *J Biol Chem* 283: 1008–1017
 81. Urao N, Razvi M, Oshikawa J, McKinney RD, Chavda R, Bahou WF, Fukai T, Ushio-Fukai M (2010) IQGAP1 is involved in post-ischemic neovascularization by regulating angiogenesis and macrophage infiltration. *PLoS ONE* 5: e13440
 82. David S, Ghosh CC, Mukherjee A, Parikh SM (2011) Angiopoietin-1 requires IQ domain GTPase-activating protein 1 to activate Rac1 and promote endothelial barrier defense. *Arterioscler Thromb Vasc Biol* 31: 2643–2652
 83. Su G, Atakilit A, Li JT, Wu N, Bhattacharya M, Zhu J, Shieh JE, Li E, Chen R, Sun S et al (2012) Absence of integrin $\alpha v \beta 3$ enhances vascular leak in mice by inhibiting endothelial cortical actin formation. *Am J Respir Crit Care Med* 185: 58–66
 84. Bhattacharya M, Su G, Su X, Osés-Prieto JA, Li JT, Huang X, Hernandez H, Atakilit A, Burlingame AL, Matthay MA et al (2012) IQGAP1 is necessary for pulmonary vascular barrier protection in murine acute lung injury and pneumonia. *Am J Physiol Lung Cell Mol Physiol* 303: L12–L19
 85. Tian Y, Tian X, Gawlak G, O'Donnell JJ 3rd, Sacks DB, Birukova AA (2014) IQGAP1 regulates endothelial barrier function via EB1 - cortactin crosstalk. *Mol Cell Biol* 34: 3546–3558
 86. Rami L, Auguste P, Thebaud NB, Bareille R, Daculsi R, Ripoche J, Bordenave L (2013) IQ domain GTPase-activating protein 1 is involved in shear stress-induced progenitor-derived endothelial cell alignment. *PLoS ONE* 8: e79919
 87. Somlyo AP, Somlyo AV (2003) Calcium sensitivity of smooth muscle and nonmuscle myosin II: modulated by G proteins, kinases, and myosin phosphatase. *Physiol Rev* 83: 1325–1358
 88. Puetz S, Lubomirov LT, Pfitzer G (2009) Regulation of smooth muscle contraction by small GTPases. *Physiology* 24: 342–356
 89. Bhattacharya M, Sundaram A, Kudo M, Farmer J, Ganesan P, Khalifeh-Soltani A, Arjomandi M, Atabai K, Huang X, Sheppard D (2014) IQGAP1-dependent scaffold suppresses RhoA and inhibits airway smooth muscle contraction. *J Clin Invest* 124: 4895–4898
 90. Seino S, Shibasaki T, Minami K (2011) Dynamics of insulin secretion and the clinical implications for obesity and diabetes. *J Clin Invest* 121: 2118–2125
 91. Rittmeyer EN, Daniel S, Hsu SC, Osman MA (2008) A dual role for IQGAP1 in regulating exocytosis. *J Cell Sci* 121: 391–403
 92. Kimura T, Yamaoka M, Taniguchi S, Okamoto M, Takei M, Ando T, Iwamatsu A, Watanabe T, Kaibuchi K, Ishizaki T et al (2013) Activated Cdc42-bound IQGAP1 determines the cellular endocytic site. *Mol Cell Biol* 33: 4834–4843
 93. Fu A, Eberhard CE, Screaton RA (2013) Role of AMPK in pancreatic beta cell function. *Mol Cell Endocrinol* 366: 127–134
 94. Moon S, Han D, Kim Y, Jin J, Ho WK (2014) Interactome analysis of AMP-activated protein kinase (AMPK)-alpha1 and -beta1 in INS-1 pancreatic beta-cells by affinity purification-mass spectrometry. *Sci Rep* 4: 4376
 95. Chiariello CS, LaComb JF, Bahou WF, Schmidt VA (2011) Ablation of Iqgap2 protects from diet-induced hepatic steatosis due to impaired fatty acid uptake. *Regul Pept* 173: 36–46
 96. Vaitheeswaran B, Hartil K, Navare A, Zheng C, OBroin P, Golden A, Guha C, Lee W, Kurland IJ, Bruce JE (2014) Role of the tumor suppressor IQGAP2 in metabolic homeostasis: possible link between diabetes and cancer. *Metabolomics* 10: 920–937
 97. Schubert S, Shannon K, Bollag G (2007) Hyperactive Ras in developmental disorders and cancer. *Nat Rev Cancer* 7: 295–308
 98. Lito P, Rosen N, Solit DB (2013) Tumor adaptation and resistance to RAF inhibitors. *Nat Med* 19: 1401–1409
 99. Johnson M, Sharma M, Henderson BR (2009) IQGAP1 regulation and roles in cancer. *Cell Signal* 21: 1471–1478
 100. White CD, Brown MD, Sacks DB (2009) IQGAPs in cancer: a family of scaffold proteins underlying tumorigenesis. *FEBS Lett* 583: 1817–1824
 101. Jameson KL, Mazur PK, Zehnder AM, Zhang J, Zarnegar B, Sage J, Khavari PA (2013) IQGAP1 scaffold-kinase interaction blockade selectively targets RAS-MAP kinase-driven tumors. *Nat Med* 19: 626–630
 102. Sahai E, Marshall CJ (2002) Rho-GTPases and cancer. *Nat Rev Cancer* 2: 133–142
 103. Mullard A (2012) Protein-protein interaction inhibitors get into the groove. *Nat Rev Drug Discov* 11: 173–175
 104. Arkin MR, Tang Y, Wells JA (2014) Small-molecule inhibitors of protein-protein interactions: progressing toward the reality. *Chem Biol* 21: 1102–1114
 105. Li S, Wang Q, Chakladar A, Bronson RT, Bernards A (2000) Gastric hyperplasia in mice lacking the putative Cdc42 effector IQGAP1. *Mol Cell Biol* 20: 697–701
 106. Bhavsar AP, Guttman JA, Finlay BB (2007) Manipulation of host-cell pathways by bacterial pathogens. *Nature* 449: 827–834
 107. Brown MD, Bry L, Li Z, Sacks DB (2008) Actin pedestal formation by enteropathogenic *Escherichia coli* is regulated by IQGAP1, calcium, and calmodulin. *J Biol Chem* 283: 35212–35222

108. Buss C, Muller D, Ruter C, Heussipp G, Schmidt MA (2009) Identification and characterization of Ibe, a novel type III effector protein of A/E pathogens targeting human IQGAP1. *Cell Microbiol* 11: 661–677
109. Krishnan S, Fernandez GE, Sacks DB, Prasadarao NV (2012) IQGAP1 mediates the disruption of adherens junctions to promote *Escherichia coli* K1 invasion of brain endothelial cells. *Cell Microbiol* 14: 1415–1433
110. Brown MD, Bry L, Li Z, Sacks DB (2007) IQGAP1 regulates *Salmonella* invasion through interactions with actin, Rac1 and Cdc42. *J Biol Chem* 282: 30265–30272
111. McLaughlin LM, Govoni GR, Gerke C, Gopinath S, Peng K, Laidlaw G, Chien YH, Jeong HW, Li Z, Brown MD et al (2009) The *Salmonella* SPI2 effector SseI mediates long-term systemic infection by modulating host cell migration. *PLoS Pathog* 5: e1000671
112. Kim H, White CD, Li Z, Sacks DB (2011) *Salmonella enterica* serotype Typhimurium usurps the scaffold protein IQGAP1 to manipulate Rac1 and MAPK signalling. *Biochem J* 440: 309–318
113. Zhang L, Li X, Wang B, Zhang T, Ye J (2012) *Chlamydia* (Chlamydia) pneumoniae infection promotes vascular smooth muscle cell adhesion and migration through IQ domain GTPase-activating protein 1. *Microb Pathog* 53: 207–213
114. Karlsson T, Turkina MV, Yakymenko O, Magnusson KE, Vikstrom E (2012) The *Pseudomonas aeruginosa* N-acylhomoserine lactone quorum sensing molecules target IQGAP1 and modulate epithelial cell migration. *PLoS Pathog* 8: e1002953
115. Lewis K (2013) Platforms for antibiotic discovery. *Nat Rev Drug Discov* 12: 371–387
116. Chazal N, Gerlier D (2003) Virus entry, assembly, budding, and membrane rafts. *Microbiol Mol Biol Rev* 67: 226–237
117. Lu J, Qu Y, Liu Y, Jambusaria R, Han Z, Ruthel G, Freedman BD, Harty RN (2013) Host IQGAP1 and Ebola virus VP40 interactions facilitate virus-like particle egress. *J Virol* 87: 7777–7780
118. Gladue DP, Holinka LG, Fernandez-Sainz IJ, Prarat MV, O'Donnell V, Vepkhvadze NG, Lu Z, Risatti GR, Borca MV (2011) Interaction between Core protein of classical swine fever virus with cellular IQGAP1 protein appears essential for virulence in swine. *Virology* 412: 68–74
119. Leung J, Yueh A, Appah FS Jr, Yuan B, de los Santos K, Goff SP (2006) Interaction of Moloney murine leukemia virus matrix protein with IQGAP. *EMBO J* 25: 2155–2166
120. Morita E, Sandrin V, Chung HY, Morham SG, Gygi SP, Rodesch CK, Sundquist WI (2007) Human ESCRT and ALIX proteins interact with proteins of the midbody and function in cytokinesis. *EMBO J* 26: 4215–4227
121. Dolnik O, Kolesnikova L, Welsch S, Strecker T, Schudt G, Becker S (2014) Interaction with tsG101 is necessary for the efficient transport and release of nucleocapsids in Marburg virus-infected cells. *PLoS Pathog* 10: e1004463
122. Pillai VB, Sundaresan NR, Gupta MP (2014) Regulation of Akt signaling by sirtuins: its implication in cardiac hypertrophy and aging. *Circ Res* 114: 368–378
123. Dephoure N, Zhou C, Villen J, Beausoleil SA, Bakalarski CE, Elledge SJ, Gygi SP (2008) A quantitative atlas of mitotic phosphorylation. *Proc Natl Acad Sci U S A* 105: 10762–10767
124. Olsen JV, Vermeulen M, Santamaria A, Kumar C, Miller ML, Jensen LJ, Gnäd F, Cox J, Jensen TS, Nigg EA et al (2010) Quantitative phosphoproteomics reveals widespread full phosphorylation site occupancy during mitosis. *Sci Signal* 3: ra3
125. Atcheson E, Hamilton E, Pathmanathan S, Greer B, Harriott P, Timson DJ (2011) IQ-motif selectivity in human IQGAP2 and IQGAP3: binding of calmodulin and myosin essential light chain. *Biosci Rep* 31: 371–379
126. Brill S, Li S, Lyman CW, Church DM, Wasmuth JJ, Weissbach L, Bernards A, Snijders AJ (1996) The Ras GTPase-activating-protein-related human protein IQGAP2 harbors a potential actin binding domain and interacts with calmodulin and Rho family GTPases. *Mol Cell Biol* 16: 4869–4878
127. Schmidt VA, Scudder L, Devoe CE, Bernards A, Cupit LD, Bahou WF (2003) IQGAP2 functions as a GTP-dependent effector protein in thrombin-induced platelet cytoskeletal reorganization. *Blood* 101: 3021–3028
128. Yang Y, Zhao W, Xu QW, Wang XS, Zhang Y, Zhang J (2014) IQGAP3 promotes EGFR-ERK signaling and the growth and metastasis of lung cancer cells. *PLoS ONE* 9: e97578
129. Watanabe T, Wang S, Noritake J, Sato K, Fukata M, Takefuji M, Nakagawa M, Izumi N, Akiyama T, Kaibuchi K (2004) Interaction with IQGAP1 links APC to Rac1, Cdc42, and actin filaments during cell polarization and migration. *Dev Cell* 7: 871–883
130. Bensenor LB, Kan HM, Wang N, Wallrabe H, Davidson LA, Cai Y, Schafer DA, Bloom GS (2007) IQGAP1 regulates cell motility by linking growth factor signaling to actin assembly. *J Cell Sci* 120: 658–669
131. Le Clairche C, Schlaepfer D, Ferrari A, Klingauf M, Grohmanova K, Veligodskiy A, Didry D, Le D, Egile C, Carlier MF et al (2007) IQGAP1 stimulates actin assembly through the N-WASP-Arp2/3 pathway. *J Biol Chem* 282: 426–435
132. Bourguignon LY, Gilad E, Rothman K, Peyrollier K (2005) Hyaluronan-CD44 interaction with IQGAP1 promotes Cdc42 and ERK signaling, leading to actin binding, Elk-1/estrogen receptor transcriptional activation, and ovarian cancer progression. *J Biol Chem* 280: 11961–11972
133. Logue JS, Whiting JL, Tunquist B, Sacks DB, Langeberg LK, Wordeman L, Scott JD (2011) AKAP220 protein organizes signaling elements that impact cell migration. *J Biol Chem* 286: 39269–39281
134. Watanabe T, Noritake J, Kakeno M, Matsui T, Harada T, Wang S, Itoh N, Sato K, Matsuzawa K, Iwamatsu A et al (2009) Phosphorylation of CLASP2 by GSK-3 β regulates its interaction with IQGAP1, EB1 and microtubules. *J Cell Sci* 122: 2969–2979
135. Fukata M, Watanabe T, Noritake J, Nakagawa M, Yamaga M, Kuroda S, Matsura Y, Iwamatsu A, Perez F, Kaibuchi K (2002) Rac1 and Cdc42 capture microtubules through IQGAP1 and CLIP-170. *Cell* 109: 873–885
136. Usatyuk PV, Gorshkova IA, He D, Zhao Y, Kalari SK, Garcia JG, Natarajan V (2009) Phospholipase D-mediated activation of IQGAP1 through Rac1 regulates hyperoxia-induced p47phox translocation and reactive oxygen species generation in lung endothelial cells. *J Biol Chem* 284: 15339–15352
137. Liu J, Guidry JJ, Worthylake DK (2013) Conserved sequence repeats of IQGAP1 mediate binding to Ezrin. *J Proteome Res* 13: 1156–1166
138. Mukhopadhyay S, Wen X, Chih B, Nelson CD, Lane WS, Scales SJ, Jackson PK (2010) TULP3 bridges the IFT-A complex and membrane phosphoinositides to promote trafficking of G protein-coupled receptors into primary cilia. *Genes Dev* 24: 2180–2193
139. Dobrev I, Fielding A, Foster LJ, Dedhar S (2008) Mapping the integrin-linked kinase interactome using SILAC. *J Proteome Res* 7: 1740–1749
140. Wickstrom SA, Lange A, Hess MW, Polleux J, Spatz JP, Kruger M, Pfaller K, Lambacher A, Bloch W, Mann M et al (2010) Integrin-linked kinase controls microtubule dynamics required for plasma membrane targeting of caveolae. *Dev Cell* 19: 574–588
141. Brandt DT, Marion S, Griffiths G, Watanabe T, Kaibuchi K, Grosse R (2007) Dia1 and IQGAP1 interact in cell migration and phagocytic cup formation. *J Cell Biol* 178: 193–200

142. Karaczyn A, Bani-Yaghoob M, Tremblay R, Kubu C, Cowling R, Adams TL, Prudovsky I, Spicer D, Friesel R, Vary C *et al* (2010) Two novel human NUMB isoforms provide a potential link between development and cancer. *Neural Dev* 5: 31
143. Ruiz-Saenz A, Kremer L, Alonso MA, Millan J, Correias I (2011) Protein 4.1R regulates cell migration and IQGAP1 recruitment to the leading edge. *J Cell Sci* 124: 2529–2538
144. Lui WY, Mruk DD, Cheng CY (2005) Interactions among IQGAP1, Cdc42, and the cadherin/catenin protein complex regulate Sertoli-germ cell adherens junction dynamics in the testis. *J Cell Physiol* 102: 49–66
145. Takahashi K, Suzuki K (2008) Requirement of kinesin-mediated membrane transport of WAVE2 along microtubules for lamellipodia formation promoted by hepatocyte growth factor. *Exp Cell Res* 314: 2313–2322
146. Nakajima E, Suzuki K, Takahashi K (2005) Mitotic dissociation of IQGAP1 from Rac-bound beta1-integrin is mediated by protein phosphatase 2A. *Biochem Biophys Res Commun* 326: 249–253
147. Subramani J, Ghosh M, Rahman MM, Caromile LA, Gerber C, Rezaul K, Han DK, Shapiro LH (2013) Tyrosine phosphorylation of CD13 regulates inflammatory cell-cell adhesion and monocyte trafficking. *J Immunol* 191: 3905–3912
148. Jacquemet G, Morgan MR, Byron A, Humphries JD, Choi CK, Chen CS, Caswell PT, Humphries MJ (2013) Rac1 is deactivated at integrin activation sites through an IQGAP1-filamin-A-RacGAP1 pathway. *J Cell Sci* 126: 4121–4135
149. Yan J, Yang Y, Zhang H, King C, Kan HM, Cai Y, Yuan CX, Bloom GS, Hua X (2009) Menin interacts with IQGAP1 to enhance intercellular adhesion of beta-cells. *Oncogene* 28: 973–982
150. Katata T, Irie K, Fukuhara A, Kawakatsu T, Yamada A, Shimizu K, Takai Y (2003) Involvement of nectin in the localization of IQGAP1 at the cell-cell adhesion sites through the actin cytoskeleton in Madin-Darby canine kidney cells. *Oncogene* 22: 2097–2109
151. Routray C, Liu C, Yaqoob U, Billadeau DD, Bloch KD, Kaibuchi K, Shah VH, Kang N (2011) Protein kinase G signaling disrupts Rac1-dependent focal adhesion assembly in liver specific pericytes. *Am J Physiol Cell Physiol* 301: C66–C74
152. Weissbach L, Bernards A, Herion DW (1998) Binding of myosin essential light chain to the cytoskeleton-associated protein IQGAP1. *Biochem Biophys Res Commun* 251: 269–276
153. Mbele GO, Deloulme JC, Gentil BJ, Delphin C, Ferro M, Garin J, Takahashi M, Baudier J (2002) The zinc- and calcium-binding S100B interacts and co-localizes with IQGAP1 during dynamic rearrangement of cell membranes. *J Biol Chem* 277: 49998–50007
154. Heil A, Nazmi AR, Koltzsch M, Poeter M, Austermann J, Assard N, Baudier J, Kaibuchi K, Gerke V (2011) S100P is a novel interaction partner and regulator of IQGAP1. *J Biol Chem* 286: 7227–7238
155. Blagoev B, Kratchmarova I, Ong SE, Nielsen M, Foster LJ, Mann M (2003) A proteomics strategy to elucidate functional protein-protein interactions applied to EGF signaling. *Nat Biotechnol* 21: 315–318
156. McNulty DE, Li Z, White CD, Sacks DB, Annan RS (2011) MAPK scaffold IQGAP1 binds the EGF receptor and modulates its activation. *J Biol Chem* 286: 15010–15021
157. White CD, Li Z, Dillon DA, Sacks DB (2011) IQGAP1 protein binds human epidermal growth factor receptor 2 (HER2) and modulates trastuzumab resistance. *J Biol Chem* 286: 29734–29747
158. Com E, Lagadeuc C, Page A, El Yazidi-Belkoura I, Slomianky C, Spencer A, Hammache D, Rudkin BB, Hondermarck H (2007) Nerve growth factor receptor TrkA signaling in breast cancer cells involves Ku70 to prevent apoptosis. *Mol Cell Proteomics* 6: 1842–1854
159. Liu C, Billadeau DD, Abdelhakim H, Leof E, Kaibuchi K, Bernabeu C, Bloom GS, Yang L, Boardman L, Shah VH *et al* (2013) IQGAP1 suppresses TβRII-mediated myofibroblastic activation and metastatic growth in liver. *J Clin Invest* 123: 1138–1156
160. Neel NF, Sai J, Ham AJ, Sobolik-Delmaire T, Mernaugh RL, Richmond A (2011) IQGAP1 is a novel CXCR2-interacting protein and essential component of the “chemosynapse”. *PLoS ONE* 6: e23813
161. Feigin ME, Xue B, Hammell MC, Muthuswamy SK (2014) G-protein-coupled receptor GPR161 is overexpressed in breast cancer and is a promoter of cell proliferation and invasion. *Proc Natl Acad Sci U S A* 111: 4191–4196
162. Cvetkovic D, Dragan M, Leith SJ, Mir ZM, Leong HS, Pampillo M, Lewis JD, Babwah AV, Bhattacharya M (2013) KISS1R induces invasiveness of estrogen receptor-negative human mammary epithelial and breast cancer cells. *Endocrinology* 154: 1999–2014
163. Alemayehu M, Dragan M, Pape C, Siddiqui I, Sacks DB, Di Guglielmo GM, Babwah AV, Bhattacharya M (2013) β-Arrestin2 regulates lysophosphatidic acid-induced human breast tumor cell migration and invasion via Rap1 and IQGAP1. *PLoS ONE* 8: e56174
164. Nuriya M, Oh S, Haganir RL (2005) Phosphorylation-dependent interactions of alpha-Actinin-1/IQGAP1 with the AMPA receptor subunit GluR4. *J Neurochem* 95: 544–552
165. Okada M, Hozumi Y, Iwazaki K, Misaki K, Yanagida M, Araki Y, Watanabe T, Yagisawa H, Topham MK, Kaibuchi K *et al* (2012) DGKzeta is involved in LPS-activated phagocytosis through IQGAP1/Rac1 pathway. *Biochem Biophys Res Commun* 420: 479–484
166. Choi S, Thapa N, Hedman AC, Li Z, Sacks DB, Anderson RA (2013) IQGAP1 is a novel phosphatidylinositol 4,5 bisphosphate effector in regulation of directional cell migration. *EMBO J* 32: 2617–2630
167. Dixon MJ, Gray A, Boisvert FM, Agacan M, Morrice NA, Gourlay R, Leslie NR, Downes CP, Batty IH (2011) A screen for novel phosphoinositide 3-kinase effector proteins. *Mol Cell Proteomics* 10: M110 003178
168. Dixon MJ, Gray A, Schenning M, Agacan M, Tempel W, Tong Y, Nedyalkova L, Park HW, Leslie NR, van Aalten DM *et al* (2012) IQGAP proteins reveal an atypical phosphoinositide (aPI) binding domain with a pseudo C2 domain fold. *J Biol Chem* 287: 22483–22496
169. Gunaratne J, Goh MX, Swa HL, Lee FY, Sanford E, Wong LM, Hogue KA, Blackstock WP, Okumura K (2011) Protein interactions of phosphatase and tensin homologue (PTEN) and its cancer-associated G20E mutant compared by using stable isotope labeling by amino acids in cell culture-based parallel affinity purification. *J Biol Chem* 286: 18093–18103
170. Tekletsadik YK, Sonn R, Osman MA (2012) A conserved role of IQGAP1 in regulating TOR complex 1. *J Cell Sci* 125: 2041–2052
171. Ghosh S, Tewari R, Dixit D, Sen E (2009) TNFα induced oxidative stress dependent Akt signaling affects actin cytoskeletal organization in glioma cells. *Neurochem Int* 56: 194–201
172. Chen F, Zhu HH, Zhou LF, Wu SS, Wang J, Chen Z (2010) IQGAP1 is overexpressed in hepatocellular carcinoma and promotes cell proliferation by Akt activation. *Exp Mol Med* 42: 477–483
173. Yin N, Shi J, Wang D, Tong T, Wang M, Fan F, Zhan Q (2012) IQGAP1 interacts with Aurora-A and enhances its stability and its role in cancer. *Biochem Biophys Res Commun* 421: 64–69
174. Takahashi K, Suzuki K (2006) Regulation of protein phosphatase 2A-mediated recruitment of IQGAP1 to beta1 integrin by EGF through activation of Ca²⁺/calmodulin-dependent protein kinase II. *J Cell Physiol* 208: 213–219

175. Fram S, King H, Sacks DB, Wells CM (2014) A PAK6-IQGAP1 complex promotes disassembly of cell-cell adhesions. *Cell Mol Life Sci* 71: 2759–2773
176. Kaur R, Yuan X, Lu ML, Balk SP (2008) Increased PAK6 expression in prostate cancer and identification of PAK6 associated proteins. *Prostate* 68: 1510–1516
177. Nauert JB, Rigas JD, Lester LB (2003) Identification of an IQGAP1/AKAP79 complex in beta-cells. *J Cell Biochem* 90: 97–108
178. Grohmanova K, Schlaepfer D, Hess D, Gutierrez P, Beck M, Kroschewski R (2004) Phosphorylation of IQGAP1 modulates its binding to Cdc42, revealing a new type of rho-GTPase regulator. *J Biol Chem* 279: 48495–48504
179. Suzuki K, Chikamatsu Y, Takahashi K (2005) Requirement of protein phosphatase 2A for recruitment of IQGAP1 to Rac-bound beta1 integrin. *J Cell Physiol* 203: 487–492
180. Meek SE, Lane WS, Piwnicka-Worms H (2004) Comprehensive proteomic analysis of interphase and mitotic 14-3-3-binding proteins. *J Biol Chem* 279: 32046–32054
181. Schiefermeier N, Scheffler JM, de Araujo ME, Stasyk T, Yordanov T, Ebner HL, Offterdinger M, Munck S, Hess MW, Wickstrom SA et al (2014) The late endosomal p14-MP1 (LAMP2/3) complex regulates focal adhesion dynamics during cell migration. *J Cell Biol* 205: 525–540
182. Meng X, Krokkin O, Cheng K, Ens W, Wilkins JA (2007) Characterization of IQGAP1-containing complexes in NK-like cells: evidence for Rac2 and RACK1 association during homotypic adhesion. *J Proteome Res* 6: 744–750
183. Smith MJ, Hardy WR, Li GY, Goudreault M, Hersch S, Metalnikov P, Starostine A, Pawson T, Ikura M (2010) The PTB domain of ShcA couples receptor activation to the cytoskeletal regulator IQGAP1. *EMBO J* 29: 884–896
184. Hu B, Shi B, Jarzynka MJ, Yiin JJ, D'Souza-Schorey C, Cheng SY (2009) ADP-ribosylation factor 6 regulates glioma cell invasion through the IQ-domain GTPase-activating protein 1-Rac1-mediated pathway. *Cancer Res* 69: 794–801
185. Tian Y, Gawlak G, Shah AS, Higginbotham K, Tian X, Kawasaki Y, Akiyama T, Sacks DB, Birukova AA (2014) HGF-induced Asef-IQGAP1 complex controls cytoskeletal remodeling and endothelial barrier. *J Biol Chem* 290: 4097–4109
186. Steenblock C, Heckel T, Czupalla C, Santo AI, Niehage C, Sztacho M, Hoflack B (2014) The Cdc42 guanine nucleotide exchange factor FGD6 coordinates cell polarity and endosomal membrane recycling in osteoclasts. *J Biol Chem* 289: 18347–18359
187. Liu Z, Lee J, Krummey S, Lu W, Cai H, Lenardo MJ (2011) The kinase LRRK2 is a regulator of the transcription factor NFAT that modulates the severity of inflammatory bowel disease. *Nat Immunol* 12: 1063–1070
188. Vasilescu J, Guo X, Kast J (2004) Identification of protein-protein interactions using in vivo cross-linking and mass spectrometry. *Proteomics* 4: 3845–3854
189. Jacquemet G, Green DM, Bridgewater RE, von Kriegsheim A, Humphries MJ, Norman JC, Caswell PT (2013) RCP-driven alpha5beta1 recycling suppresses Rac and promotes RhoA activity via the RacGAP1-IQGAP1 complex. *J Cell Biol* 202: 917–935
190. Goto T, Sato A, Adachi S, Iemura S, Natsume T, Shibuya H (2013) IQGAP1 protein regulates nuclear localization of beta-catenin via importin-beta5 protein in Wnt signaling. *J Biol Chem* 288: 36351–36360
191. Jeong HW, Li Z, Brown MD, Sacks DB (2007) IQGAP1 binds Rap1 and modulates its activity. *J Biol Chem* 282: 20752–20762
192. Casteel DE, Turner S, Schwappacher R, Rangaswami H, Su-Yuo J, Zhuang S, Boss GR, Pilz RB (2012) Rho isoform-specific interaction with IQGAP1 promotes breast cancer cell proliferation and migration. *J Biol Chem* 287: 38367–38378
193. Clark EA, Golub TR, Lander ES, Hynes RO (2000) Genomic analysis of metastasis reveals an essential role for RhoC. *Nature* 406: 532–535
194. Wu Y, Chen YC, Sang JR, Xu WR (2011) RhoC protein stimulates migration of gastric cancer cells through interaction with scaffold protein IQGAP1. *Mol Med Rep* 4: 697–703
195. Neudauer CL, Joberty G, Tatsis N, Macara IG (1998) Distinct cellular effects and interactions of the Rho-family GTPase TC10. *Curr Biol* 8: 1151–1160
196. Goto T, Sato A, Shimizu M, Adachi S, Satoh K, Iemura S, Natsume T, Shibuya H (2013) IQGAP1 functions as a modulator of dishevelled nuclear localization in Wnt signaling. *PLoS ONE* 8: e60865
197. Carmon KS, Gong X, Yi J, Thomas A, Liu Q (2014) RSPO-LGR4 functions via IQGAP1 to potentiate Wnt signaling. *Proc Natl Acad Sci U S A* 111: E1221–E1229
198. Witze ES, Connacher MK, Houel S, Schwartz MP, Morphew MK, Reid L, Sacks DB, Anseth KS, Ahn NG (2013) Wnt5a directs polarized calcium gradients by recruiting cortical endoplasmic reticulum to the cell trailing edge. *Dev Cell* 26: 645–657
199. Erdemir HH, Li Z, Sacks DB (2014) IQGAP1 binds to estrogen receptor-alpha and modulates its function. *J Biol Chem* 289: 9100–9112
200. Ebmeier CC, Taatjes DJ (2010) Activator-Mediator binding regulates Mediator-cofactor interactions. *Proc Natl Acad Sci U S A* 107: 11283–11288
201. Li J, Chu M, Wang S, Chan D, Qi S, Wu M, Zhou Z, Nishi E, Qin J, Wong J (2012) Identification and characterization of nardilysin as a novel dimethyl H3K4-binding protein involved in transcriptional regulation. *J Biol Chem* 287: 10089–10098
202. Sharma S, Findlay GM, Bandukwala HS, Oberdoerffer S, Baust B, Li Z, Schmidt V, Hogan PG, Sacks DB, Rao A (2011) Dephosphorylation of the nuclear factor of activated T cells (NFAT) transcription factor is regulated by an RNA-protein scaffold complex. *Proc Natl Acad Sci U S A* 108: 11381–11386
203. Cheung KL, Lee JH, Shu L, Kim JH, Sacks DB, Kong AN (2013) The Ras GTPase-activating-like protein IQGAP1 mediates Nrf2 protein activation via the mitogen-activated protein kinase/extracellular signal-regulated kinase (ERK) kinase (MEK)-ERK pathway. *J Biol Chem* 288: 22378–22386
204. Kim JH, Xu EY, Sacks DB, Lee J, Shu L, Xia B, Kong AN (2013) Identification and functional studies of a new Nrf2 partner IQGAP1: a critical role in the stability and transactivation of Nrf2. *Antioxid Redox Signal* 19: 89–101
205. Johnson M, Sharma M, Brocardo MC, Henderson BR (2011) IQGAP1 translocates to the nucleus in early S-phase and contributes to cell cycle progression after DNA replication arrest. *Int J Biochem Cell Biol* 43: 65–73
206. Cooper MP, Qu L, Rohas LM, Lin J, Yang W, Erdjument-Bromage H, Tempst P, Spiegelman BM (2006) Defects in energy homeostasis in Leigh syndrome French Canadian variant through PGC-1alpha/LRP130 complex. *Genes Dev* 20: 2996–3009
207. Sato A, Naito T, Hiramoto A, Goda K, Omi T, Kitade Y, Sasaki T, Matsuda A, Fukushima M, Wataya Y et al (2010) Association of RNase L with a Ras GTPase-activating-like protein IQGAP1 in mediating the apoptosis of a human cancer cell-line. *FEBS J* 277: 4464–4473
208. Toyokawa G, Cho HS, Masuda K, Yamane Y, Yoshimatsu M, Hayami S, Takawa M, Iwai Y, Daigo Y, Tsuchiya E et al (2011) Histone lysine methyltransferase Wolf-Hirschhorn syndrome candidate 1 is involved

- in human carcinogenesis through regulation of the Wnt pathway. *Neoplasia* 13: 887–898
209. Sun L, Prince T, Manjarrez JR, Scroggins BT, Matts RL (2012) Characterization of the interaction of Aha1 with components of the Hsp90 chaperone machine and client proteins. *Biochim Biophys Acta* 1823: 1092–1101
 210. Takeda S, Fujimoto A, Yamauchi E, Hiyoshi M, Kido H, Watanabe T, Kaibuchi K, Ohta T, Konishi H (2011) Role of a tyrosine phosphorylation of SMG-9 in binding of SMG-9 to IQGAP and the NMD complex. *Biochem Biophys Res Commun* 410: 29–33
 211. Villace P, Marion RM, Ortin J (2004) The composition of Staufen-containing RNA granules from human cells indicates their role in the regulated transport and translation of messenger RNAs. *Nucleic Acids Res* 32: 2411–2420
 212. Chung LK, Philip NH, Schmidt VA, Koller A, Strowig T, Flavell RA, Brodsky IE, Bliska JB (2014) IQGAP1 is important for activation of caspase-1 in macrophages and is targeted by *Yersinia pestis* type III effector YopM. *MBio* 5: e01402–14
 213. Sakurai-Yageta M, Recchi C, Le Dez G, Sibarita JB, Daviet L, Camonis J, D'Souza-Schorey C, Chavrier P (2008) The interaction of IQGAP1 with the exocyst complex is required for tumor cell invasion downstream of Cdc42 and RhoA. *J Cell Biol* 181: 985–998
 214. Logue JS, Whiting JL, Tunquist B, Langeberg LK, Scott JD (2011) Anchored protein kinase A recruitment of active Rac GTPase. *J Biol Chem* 286: 22113–22121
 215. Schmidt VA, Chiariello CS, Capilla E, Miller F, Bahou WF (2008) Development of hepatocellular carcinoma in *Iqgap2*-deficient mice is IQGAP1 dependent. *Mol Cell Biol* 28: 1489–1502
 216. Wennerberg K, Ellerbroek SM, Liu RY, Karnoub AE, Burridge K, Der CJ (2002) RhoG signals in parallel with Rac1 and Cdc42. *J Biol Chem* 277: 47810–47817
 217. Nojima H, Adachi M, Matsui T, Okawa K, Tsukita S (2008) IQGAP3 regulates cell proliferation through the Ras/ERK signalling cascade. *Nat Cell Biol* 10: 971–978
 218. Boyer O, Benoit G, Gribouval O, Nevo F, Pawtowski A, Bilge I, Bircan Z, Deschenes G, Guay-Woodford LM, Hall M *et al* (2010) Mutational analysis of the PLCE1 gene in steroid resistant nephrotic syndrome. *J Med Genet* 47: 445–452
 219. Liu XL, Kilpelainen P, Hellman U, Sun Y, Wartiovaara J, Morgunova E, Pikkariainen T, Yan K, Jonsson AP, Tryggvason K (2005) Characterization of the interactions of the nephrin intracellular domain. *FEBS J* 272: 228–243
 220. Bridge DR, Novotny MJ, Moore ER, Olson JC (2009) Role of host cell polarity and leading edge properties in *Pseudomonas* type III secretion. *Microbiology* 156: 356–373
 221. Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard JF, Guindon S, Lefort V, Lescot M *et al* (2008) Phylogeny.fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Res* 36: W465–W469