Published in final edited form as:

J Hosp Med. 2015 April ; 10(4): 228–235. doi:10.1002/jhm.2321.

Association of inpatient and outpatient glucose management with inpatient mortality among patients with and without diabetes at a major academic medical center

Neel M. Butala, MD, MBA¹, Benjamin K. Johnson, MA², James D. Dziura, PhD², Jesse S. Reynolds, MS², Janis E. Bozzo, MSN, RN³, Thomas J. Balczak, MD, MPH⁴, Silvio E. Inzucchi, MD⁵, and Leora I. Horwitz, MD, MHS^{6,7,8}

¹ Department of Medicine, Massachusetts General Hospital, Boston, MA

² Yale Center for Analytical Sciences, Yale School of Medicine, New Haven, CT

³ Yale-New Haven Health System, New Haven, CT

⁴ Yale-New Haven Hospital, New Haven, CT

⁵ Section of Endocrinology, Department of Medicine, Yale School of Medicine, New Haven, CT

⁶ Division of Healthcare Delivery Science, Department of Population Health, New York University School of Medicine, New York, NY

⁷ Center for Healthcare Innovation and Delivery Science, New York University Langone Medical Center, New York, NY

⁸ Division of General Internal Medicine & Clinical Innovation, Department of Medicine, New York University School of Medicine, New York, NY

Abstract

Background—Hospitalized patients with diabetes have experienced a disproportionate reduction in mortality over the past decade.

Objective—To examine whether this differential decrease affected all patients with diabetes, and to identify explanatory factors.

Design—Serial, cross-sectional observational study

Setting—Academic medical center

Corresponding Author: Leora Horwitz, MD Department of Population Health 550 First Ave, TRB Room 607 New York, NY 10016
Ph: 646-501-2685 Leora.horwitz@nyumc.org.

Conflicts of Interest:

Dr. Inzucchi serves on a Data Safety Monitoring Board for Novo Nordisk, a manufacturer of insulin products used in the hospital setting. The remaining authors declare no conflicts of interest.

Author contributions:

Design and conduct of the study: NB, JD, SI, TB, LH

Collection, management, analysis, and interpretation of the data: NB, BJ, JD, JR, JB, SI, LH

Preparation, review, or approval of the manuscript: NB, BJ, JD, JR, SI, TB, LH

Access to data:

Leora Horwitz had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Patients—All adult, non-obstetric patients with an inpatient discharge between Jan 1, 2000 and Dec 31, 2010

Measurement—We assessed in-hospital mortality; inpatient glycemic control (percentage of hospital days with glucose below 70, above 299, and between 70-179 mg/dl, and standard deviation of glucose measurements); and outpatient glycemic control (hemoglobin A1c).

Results—We analyzed 322,938 admissions, including 76,758 (23.8%) with diabetes. Among 54,645 ICU admissions, there was a 7.8% relative reduction in the odds of mortality in each successive year for patients with diabetes, adjusted for age, race, payer, length of stay, discharge diagnosis, comorbidities, and service (OR=0.923, 95% CI: 0.906, 0.940). This was significantly greater than the 2.6% yearly reduction for those without diabetes (OR=0.974, CI: 0.963, 0.985; $p<0.001$ for interaction). In contrast, the greater decrease in mortality among non-ICU patients with diabetes did not reach significance. Results were similar among medical and surgical patients. Among ICU patients with diabetes, the significant decline in mortality persisted after adjustment for inpatient and outpatient glucose control (OR=0.953, 95% CI: 0.914, 0.994).

Conclusions—Patients with diabetes in the ICU have experienced a disproportionate reduction in mortality that is not explained by glucose control. Potential explanations include improved cardiovascular risk management or advances in therapies for diseases commonly affecting patients with diabetes.

Keywords

Diabetes; Glucose Management; Mortality; Inpatient Mortality; Outcomes; Hospital Medicine

Background

Patients with diabetes currently comprise over 8 percent of the US population (over 25 million people) and more than 20 percent of hospitalized patients.^{1,2} Hospitalizations of patients with diabetes account for 23 percent of total hospital costs in the US,² and patients with diabetes have worse outcomes after hospitalization for a variety of common medical conditions,³⁻⁶ as well as in intensive care unit (ICU) settings.^{7,8} Individuals with diabetes have historically experienced higher inpatient mortality than individuals without diabetes.⁹ However, we recently reported that patients with diabetes at our large academic medical center have experienced a disproportionate reduction in in-hospital mortality relative to patients without diabetes over the past decade.¹⁰ This surprising trend begs further inquiry.

Improvement in in-hospital mortality among patients with diabetes may stem from improved inpatient glycemic management. The landmark 2001 study by van den Berghe *et al.* demonstrating that intensive insulin therapy reduced post-surgical mortality among ICU patients ushered in an era of intensive inpatient glucose control.¹¹ However, follow-up multicenter studies have not been able to replicate these results.¹²⁻¹⁵ In non-ICU and non-surgical settings, intensive glucose control has not yet been shown to have any mortality benefit, although it may impact other morbidities, such as post-operative infections.¹⁶ Consequently, less stringent glycemic targets are now recommended.¹⁷ Nonetheless, hospitals are being held accountable for certain aspects of inpatient glucose control. For example, the Centers for Medicare & Medicaid Services (CMS) began asking hospitals to

report inpatient glucose control in cardiac surgery patients in 2004.¹⁸ This measure is now publicly reported and, as of 2013, is included in the CMS Value-Based Purchasing Program, which financially penalizes hospitals that do not meet targets.

Outpatient diabetes standards have also evolved in the past decade. The Diabetes Control and Complications Trial in 1993 and the United Kingdom Prospective Diabetes Study in 1997 demonstrated that better glycemic control in Type 1 and newly diagnosed Type 2 diabetes patients, respectively, improved clinical outcomes, and prompted guidelines for pharmacologic treatment of diabetic patients.^{19,20} However, subsequent randomized clinical trials have failed to establish a clear beneficial effect of intensive glucose control on primary cardiovascular endpoints among higher risk patients with longstanding type 2 diabetes,²¹⁻²³ and clinical practice recommendations now accept a more individualized approach to glycemic control.²⁴ Nonetheless, clinicians are also being held accountable for outpatient glucose control.²⁵

To better understand the disproportionate reduction in mortality among hospitalized patients with diabetes that we observed, we first examined whether it was limited to surgical patients or patients in the ICU: the populations that have been demonstrated to benefit from intensive inpatient glucose control. Furthermore, given recent improvements in inpatient and outpatient glycemic control,^{26,27} we examined whether inpatient or outpatient glucose control explained the mortality trends. Results from this study contribute empirical evidence on real-world effects of efforts to improve inpatient and outpatient glycemic control.

Methods

Setting

During the study period, Yale-New Haven Hospital (YNHH) was an urban academic medical center in New Haven, CT with over 950 beds and an average of approximately 32,000 annual adult non-obstetric admissions. YNHH conducted a variety of inpatient glucose control initiatives during the study period. The surgical ICU began an informal medical team-directed insulin infusion protocol in 2000-2001. In 2002, the medical ICU instituted a formal insulin infusion protocol with a target of 100-140 mg/dl, which spread to remaining hospital ICUs by the end of 2003. In 2005, YNHH launched a consultative inpatient diabetes management team to assist clinicians in controlling glucose in non-ICU patients with diabetes. This team covered approximately 10-15 patients at a time and consisted of an advanced-practice nurse practitioner, a supervising endocrinologist and endocrinology fellow, and a nurse educator to provide diabetic teaching. Additionally, in 2005, basal-bolus-correction insulin order sets became available. The surgical ICU implemented a stringent insulin infusion protocol with target glucose of 80-110 mg/dl in 2006, but relaxed it (goal 80-150 mg/dl) in 2007. Similarly, in 2006, YNHH made ICU insulin infusion recommendations more stringent in remaining ICUs (goal 90-130 mg/dl), but relaxed them in 2010 (goal 120-160 mg/dl), based on emerging data from clinical trials and prevailing national guidelines.

Participants and data sources

We included all adult, non-obstetric discharges from Yale-New Haven Hospital between Jan 1, 2000 and Dec 31, 2010. Repeat visits by the same patient were linked by medical record number. We obtained data from YNHH administrative billing, laboratory and point of care capillary blood glucose databases. The Yale Human Investigation Committee approved our study design and granted a Health Insurance Portability and Accountability Act waiver and a waiver of patient consent.

Variables

Our primary endpoint was in-hospital mortality. The primary exposure of interest was whether a patient had diabetes mellitus, defined as the presence of ICD-9 codes 249.x, 250.x, V4585, V5391 or V6546 in any of the primary or secondary diagnosis codes in the index admission, or in any hospital encounter in the year prior to the index admission.

We assessed two effect-modifying variables: ICU status (as measured by a charge for at least 1 night in the ICU) and service assignment to surgery (including neurosurgery and orthopedics), compared to medicine (including neurology). Independent explanatory variables included time between the start of the study and patient admission (measured as days/365), diabetes status, inpatient glucose control, and long-term glucose control (as measured by hemoglobin A1c at any time in the 180 days prior to hospital admission in order to have adequate sample size). We assessed inpatient blood glucose control through point-of-care blood glucose meters (OneTouch® SureStep®, LifeScan, Inc., Milipitas, CA) at YNHH. We used 4 validated measures of inpatient glucose control: the proportion of days in each hospitalization in which there was any hypoglycemic episode (blood glucose value <70 mg/dl), the proportion of days in which there was any severely hyperglycemic episode (blood glucose value >299 mg/dl), the proportion of days in which mean blood glucose was considered to be within adequate control (all blood glucose values between 70 and 179 mg/dl), and the standard deviation of mean glucose during hospitalization as a measure of glycemic variability.²⁸

Covariates included gender, age at time of admission, length of stay in days, race (defined by hospital registration), payer, Elixhauser comorbidity dummy variables (revised to exclude diabetes and to use only secondary diagnosis codes),²⁹ and primary discharge diagnosis grouped using the Clinical Classifications Software (CCS),³⁰ based on established associations with in-hospital mortality.

Statistical analysis

We summarized demographic characteristics numerically and graphically for patients with and without diabetes and compared them using chi-square and t-tests. We summarized changes in inpatient and outpatient measures of glucose control over time numerically and graphically and compared across years using Wilcoxon rank-sum test, adjusted for multiple hypothesis testing.

We stratified all analyses first by ICU status and then by service assignment (medicine versus surgery). Statistical analyses within each stratum paralleled our previous approach to

the full study cohort.¹⁰ Taking each stratum separately (i.e. only ICU patients, or only medicine patients), we used a difference-in-differences approach comparing changes over time in in-hospital mortality among patients with diabetes compared to those without diabetes. This approach enabled us to determine whether patients with diabetes had a different time trend in risk of in-hospital mortality than those without diabetes. That is, for each stratum, we constructed multivariate logistic regression models including time in years, diabetes status, and the interaction between time and diabetes status as well as the aforementioned covariates. We calculated odds of death and confidence intervals for each additional year for patients with diabetes by exponentiating the sum of parameter estimates for time and the diabetes-time interaction term. We evaluated all two-way interactions between year or diabetes status and the covariates in a multiple degree of freedom likelihood ratio test. We investigated non-linearity of the relation between mortality and time by evaluating first and second-order polynomials.

Because we found a significant decline in mortality risk for patients with versus without diabetes among ICU patients but not among non-ICU patients and because service assignment was not found to be an effect modifier, we then limited our sample to ICU patients with diabetes to better understand the role of inpatient and outpatient glucose control in accounting for observed mortality trends. First, we determined the relation between the measures of inpatient glucose control and changes in mortality over time using logistic regression. Then, we repeated this analysis in the subsets of patients that had inpatient glucose data and both inpatient and outpatient glycemic control data, adding inpatient and outpatient measures sequentially. Given the high level of missing outpatient glycemic control data, we compared demographic characteristics for diabetic ICU patients with and without such data using chi-square and t-tests and found that patients with data were younger and less likely to be white and had longer mean length of stay, slightly worse performance on several measures of inpatient glucose control, and lower mortality (see eTable 1).

To explore the effects of dependence among observations from patients with multiple encounters, we compared parameter estimates derived from a model with all patient encounters (including repeated admissions for the same patient) with those from a model with a randomly sampled single visit per patient, and observed that there was no difference in parameter estimates between the two classes of models. For all analyses, we used a type I error of 5% (two-sided) to test for statistical significance using SAS v9.3 (SAS Institute, Cary, NC) or R software (<http://CRAN.R-project.org>).

Results

We included 322,938 patient admissions. Of this sample, 54,645 (16.9%) had spent at least 1 night in the ICU. Overall, 76,758 patients (23.8%) had diabetes, representing 26.3% of ICU patients, 23.2% of non-ICU patients, 28.2% of medical patients and 16.8% of surgical patients (see Table 1 for demographic characteristics).

Mortality trends within strata

Among ICU patients, the overall mortality rate was 9.9%: 10.5% of patients with diabetes and 9.8% of patients without diabetes. Among non-ICU patients, the overall mortality rate was 0.8%: 0.9% of patients with diabetes and 0.7% of patients without diabetes.

Among medical patients, the overall mortality rate was 2.9%: 3.1% of patients with diabetes and 2.8% of patients without diabetes. Among surgical patients, the overall mortality rate was 1.4%: 1.8% of patients with diabetes and 1.4% of patients without diabetes. Figure 1 shows quarterly in-hospital mortality for patients with and without diabetes from 2000 to 2010 stratified by ICU status and by service assignment.

Table 2 describes the difference-in-differences regression analyses, stratified by ICU status and service assignment. Among ICU patients (Table 2, Model 1), each successive year was associated with a 2.6% relative reduction in the adjusted odds of mortality (OR=0.974, 95% CI: 0.963, 0.985) for patients without diabetes compared to a 7.8% relative reduction for those with diabetes (OR=0.923, 95% CI: 0.906, 0.940). In other words, patients with diabetes compared to patients without diabetes had a significantly greater decline in odds of adjusted mortality of 5.3% per year (OR=0.947, 95% CI: 0.927, 0.967). As a result, the adjusted odds of mortality among patients with versus without diabetes decreased from 1.352 in 2000 to 0.772 in 2010.

Among non-ICU patients (Table 2, Model 2), each successive year was associated with a 7.5% relative reduction in the adjusted odds of mortality (OR=0.925, 95% CI: 0.909, 0.940) for patients without diabetes compared to a 9.6% relative reduction for those with diabetes (OR=0.904, 95% CI: 0.879, 0.929); this greater decline in odds of adjusted mortality of 2.3% per year (OR=0.977, 95% CI: 0.946, 1.008; $p=0.148$) was not statistically significant.

We found greater decline in odds of mortality among patients with diabetes than among patients without diabetes over time in both medical patients (3.9% greater decline per year; OR=0.961, 95% CI: 0.942-0.980) and surgical patients (4.5% greater decline per year; OR=0.955, 95% CI: 0.918-0.994), without a difference between the two. Detailed results are shown in Table 2, Models 3 and 4.

Glycemic control

Among ICU patients with diabetes (N=14,364), at least 2 inpatient point-of-care glucose readings were available for 13,136 (91.5%), with a mean of 4.67 readings per day, while hemoglobin A1c data was available for only 5,321 (37.0%). Both inpatient glucose data and hemoglobin A1c were available for 4,989 (34.7%) patients. Figure 2 shows trends in inpatient and outpatient glycemic control measures among ICU patients with diabetes over the study period. Mean hemoglobin A1c decreased from 7.7 in 2000 to 7.3 in 2010. Mean hospitalization glucose began at 187.2, reached a nadir of 162.4 in Q3 2007 and rose subsequently to 174.4 with loosened glucose control targets. Standard deviation of mean glucose and percentage of patient-days with a severe hyperglycemic episode followed a similar pattern, though with nadirs in Q4 2007 and Q2 2008, respectively, while percentage of patient-days with a hypoglycemic episode rose from 1.46% in 2000, peaked at 3.00% in

Q3 2005 and returned to 2.15% in 2010. All changes in glucose control are significant with $p < 0.001$.

Mortality trends and glycemic control

To determine whether glucose control explained the excess decline in odds of mortality among patients with diabetes in the ICU, we restricted our sample to ICU patients with diabetes and examined the association of diabetes with mortality after including measures of glucose control.

We first verified that the overall adjusted mortality trend among ICU patients with diabetes for whom we had measures of inpatient glucose control was similar to that of the full sample of ICU patients with diabetes. Similar to the full sample, we found that the adjusted excess odds of death significantly declined by a relative 7.3% each successive year (OR=0.927, 95% CI: 0.907, 0.947; Table 3, Model 1). We then included measures of inpatient glucose control in the model and found, as expected, that a higher percentage of days with severe hyperglycemia and with hypoglycemia was associated with an increased odds of death ($p < 0.001$ for both; Table 3, Model 2). Nonetheless, after including measures of inpatient glucose control, we found that the rate of change of excess odds of death for patients with diabetes was unchanged (OR=0.926, 95% CI: 0.905, 0.947).

We then restricted our sample to patients with diabetes with both inpatient and outpatient glycemic control data and found that, in this subpopulation, the adjusted excess odds of death among patients with diabetes relative to those without significantly declined by a relative 4.2% each progressive year (OR=0.958, 95% CI: 0.918, 0.998; Table 3, Model 3). Including measures of inpatient glucose control in the model did not significantly change the rate of change of excess odds of death (OR=0.956, 95% CI: 0.916, 0.997; Table 3, Model 4), nor did including both measures of inpatient and outpatient glycemic control (OR=0.953, 95% CI: 0.914, 0.994; Table 3, Model 5).

Discussion

We conducted a difference-in-difference analysis of in-hospital mortality rates among adult patients with diabetes compared to patients without diabetes over 10 years, stratifying by ICU status and service assignment. For patients with any ICU stay, we found that the reduction in odds of mortality for patients with diabetes has been 3 times larger than the reduction in odds of mortality for patients without diabetes. For those without an ICU stay, we found no significant difference between patients with and without diabetes in the rate at which in-hospital mortality declined. We did not find stratification by assignment to a medical or surgical service to be an effect modifier. Finally, despite the fact that our institution achieved better aggregate inpatient glucose control, less severe hyperglycemia and better long-term glucose control over the course of the decade, we did not find that either inpatient or outpatient glucose control explained the trend in mortality for patients with diabetes in the ICU. Our study is unique in its inclusion of all hospitalized patients and its ability to simultaneously assess whether both inpatient and outpatient glucose control are explanatory factors in the observed mortality trends.

The fact that improved inpatient glucose control did not explain the trend in mortality for patients with diabetes in the ICU is consistent with the majority of the literature on intensive inpatient glucose control. In randomized trials, intensive glucose control appears to be of greater benefit for patients without diabetes than for patients with diabetes.³¹ In fact, in one study, patients with diabetes were the only group that did not benefit from intensive glucose control.³² In our study, it is possible that the rise in hypoglycemia nullified some of the benefits of glucose control. Nationally, hospital admissions for hypoglycemia among Medicare beneficiaries now outnumber admissions for hyperglycemia.²⁷

We also do not find that the decline in hemoglobin A1c attenuated the reduction in mortality in the minority of patients for whom these data were available. This is concordant with evidence from three randomized clinical trials that have failed to establish a clear beneficial effect of intensive outpatient glucose control on primary cardiovascular endpoints among older, high-risk patients with type 2 diabetes using glucose-lowering agents.²¹⁻²³ It is notable, however, that the population for whom we had available hemoglobin A1c results was not representative of the overall population of ICU patients with diabetes. Consequently, there may be an association of outpatient glucose control with inpatient mortality in the overall population of ICU patients with diabetes that we were not able to detect.

The decline in mortality among ICU patients with diabetes in our study may stem from factors other than glycemic control. It is possible that patients were diagnosed earlier in their course of disease in later years of the study period, making the population of patients with diabetes younger or healthier. Of note, however, our risk adjustment models were very robust, with c statistics from 0.82-0.92, suggesting that we were able to account for much of the mortality risk attributable to patient clinical and demographic factors. More intensive glucose management may have non-glycemic benefits, such as closer patient observation, which may themselves affect mortality. Alternatively, improved cardiovascular management for patients with diabetes may have decreased the incidence of cardiovascular events. During the study period, evidence from large clinical trials demonstrated the importance of tight blood pressure and lipid management in improving outcomes for patients with diabetes,³³⁻³⁶ guidelines for lipid management for patients with diabetes changed,³⁷ and fewer patients developed cardiovascular complications.³⁸ Finally, it is possible that our findings can be explained by an improvement in treatment of complications for which patients with diabetes previously have had disproportionately worse outcomes, such as percutaneous coronary intervention.³⁹

Our findings may have important implications for both clinicians and policymakers. Changes in inpatient glucose management have required substantial additional resources on the part of hospitals. Our evidence regarding the questionable impact of inpatient glucose control on in-hospital mortality trends for patients with diabetes is disappointing and highlights the need for multifaceted evaluation of the impact of such quality initiatives. There may, for instance, be benefits from tighter blood glucose control in the hospital beyond mortality, such as reduced infections, costs or length of stay. On the outpatient side, our more limited data are consistent with recent studies that have not been able to show a mortality benefit in older diabetic patients from more stringent glycemic control. A

reassessment of prevailing diabetes-related quality measures, as recently called for by some,^{40,41} seems reasonable.

Our study must be interpreted in light of its limitations. It is possible that the improvements in glucose management were too small to result in a mortality benefit. The overall reduction of 25 mg/dl achieved at our institution is less than the 33-50 mg/dl difference between intensive and conventional groups in those randomized clinical trials that have found reductions in mortality.^{11,42} In addition, an increase in mean glucose during the last 1-2 years of the observation period (in response to prevailing guidelines) could potentially have attenuated any benefit on mortality. The study does not include other important clinical endpoints, such as infections, complications, length of stay and hospital costs. Additionally, we did not examine post-discharge mortality, which might have shown a different pattern. The small proportion of patients with A1c results may have hampered our ability to detect an effect of outpatient glucose control. Consequently, our findings regarding outpatient glucose control are only suggestive. Finally, our findings represent the experience of a single, large academic medical center and may not be generalizable to all settings.

Overall, we found that patients with diabetes in the ICU have experienced a disproportionate reduction in in-hospital mortality over time that does not appear to be explained by improvements in either inpatient or outpatient glucose control. While improved glycemic control may have other benefits, it does not appear to impact in-hospital mortality. Our real-world empirical results contribute to the discourse among clinicians and policymakers with regards to refocusing the approach to managing glucose in-hospital and re-adjudication of diabetes-related quality measures.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors would like to acknowledge the Yale-New Haven Hospital diabetes management team: Gael Ulisse APRN, Helen Psarakis APRN, Anne Kaisen APRN, and the Yale Endocrine Fellows.

Funding sources

Dr. Horwitz is supported by the National Institute on Aging (K08 AG038336) and by the American Federation for Aging Research through the Paul B. Beeson Career Development Award Program. This publication was also made possible by CTSA Grant Number UL1 RR024139 from the National Center for Research Resources (NCRR) and the National Center for Advancing Translational Science (NCATS), components of the National Institutes of Health (NIH), and NIH roadmap for Medical Research. Its contents are solely the responsibility of the authors and do not necessarily represent the official view of NIH.

No funding source had any role in design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

References

1. National Diabetes Information Clearinghouse. [November 12, 2013] National Diabetes Statistics. 2011. <http://diabetes.niddk.nih.gov/dm/pubs/statistics/#fast>.

2. Healthcare Cost and Utilization Project. [November 12, 2013] Statistical Brief #93. 2010. <http://www.hcup.us.ahrq.gov/reports/statbriefs/sb93.pdf>.
3. Sarma S, Mentz RJ, Kwasny MJ, et al. Association between diabetes mellitus and post-discharge outcomes in patients hospitalized with heart failure: findings from the EVEREST trial. *Eur J Heart Fail*. 2012; 11:11.
4. Mak KH, Moliterno DJ, Granger CB, et al. Influence of diabetes mellitus on clinical outcome in the thrombolytic era of acute myocardial infarction. GUSTO-I Investigators. Global Utilization of Streptokinase and Tissue Plasminogen Activator for Occluded Coronary Arteries. *J Am Coll Cardiol*. 1997; 30(1):171–179. [PubMed: 9207639]
5. Kornum JB, Thomsen RW, Riis A, Lervang HH, Schonheyder HC, Sorensen HT. Type 2 diabetes and pneumonia outcomes: a population-based cohort study. *Diabetes Care*. 2007; 30(9):2251–2257. [PubMed: 17595354]
6. Mannino DM, Thorn D, Swensen A, Holguin F. Prevalence and outcomes of diabetes, hypertension and cardiovascular disease in COPD. *Eur Respir J*. 2008; 32(4):962–969. [PubMed: 18579551]
7. Slynkova K, Mannino DM, Martin GS, Morehead RS, Doherty DE. The role of body mass index and diabetes in the development of acute organ failure and subsequent mortality in an observational cohort. *Crit Care*. 2006; 10(5):R137. [PubMed: 16999863]
8. Christiansen CF, Johansen MB, Christensen S, O'Brien JM, Tonnesen E, Sorensen HT. Type 2 diabetes and 1-year mortality in intensive care unit patients. *Eur J Clin Invest*. Mar; 2013 43(3): 238–247. [PubMed: 23240763]
9. Holman N, Hillson R, Young RJ. Excess mortality during hospital stays among patients with recorded diabetes compared with those without diabetes. *Diabet Med*. Jul 23.2013
10. Butala NM, Johnson BK, Dziura JD, et al. Decade-long trends in mortality among patients with and without diabetes mellitus at a major academic medical center. *JAMA internal medicine*. Jul 1; 2014 174(7):1187–1188. [PubMed: 24841330]
11. van den Berghe G, Wouters P, Weekers F, et al. Intensive insulin therapy in critically ill patients. *N Engl J Med*. Nov 8; 2001 345(19):1359–1367. [PubMed: 11794168]
12. Finfer S, Chittock DR, Su SY, et al. Intensive versus conventional glucose control in critically ill patients. *N Engl J Med*. Mar 26; 2009 360(13):1283–1297. [PubMed: 19318384]
13. Preiser JC, Devos P, Ruiz-Santana S, et al. A prospective randomised multi-centre controlled trial on tight glucose control by intensive insulin therapy in adult intensive care units: the Glucontrol study. *Intensive care medicine*. Oct; 2009 35(10):1738–1748. [PubMed: 19636533]
14. Arabi YM, Dabbagh OC, Tamim HM, et al. Intensive versus conventional insulin therapy: a randomized controlled trial in medical and surgical critically ill patients. *Crit Care Med*. Dec; 2008 36(12):3190–3197. [PubMed: 18936702]
15. Van den Berghe G, Wilmer A, Hermans G, et al. Intensive insulin therapy in the medical ICU. *N Engl J Med*. Feb 2; 2006 354(5):449–461. [PubMed: 16452557]
16. Murad MH, Coburn JA, Coto-Yglesias F, et al. Glycemic control in non-critically ill hospitalized patients: a systematic review and meta-analysis. *The Journal of clinical endocrinology and metabolism*. Jan; 2012 97(1):49–58. [PubMed: 22090269]
17. Moghissi ES, Korytkowski MT, DiNardo M, et al. American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycemic control. *Diabetes Care*. 2009; 32(6):1119–1131. [PubMed: 19429873]
18. Agency for Healthcare Research and Quality National Quality Measures Clearinghouse. [November 12, 2013] Percent of cardiac surgery patients with controlled 6 A.M. postoperative blood glucose. 2012. <http://www.qualitymeasures.ahrq.gov/content.aspx?id=35532>.
19. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. The Diabetes Control and Complications Trial Research Group. *N Engl J Med*. 1993; 329(14):977–986. [PubMed: 8366922]
20. Turner R, Holman R, Cull C, et al. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet*. 1998; 352(9131):837–853. [PubMed: 9742976]
21. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med*. 2008; 358(24):2545–2559. [PubMed: 18539917]

22. Duckworth W, Abraira C, Moritz T, et al. Glucose control and vascular complications in veterans with type 2 diabetes. *N Engl J Med*. 2009; 360(2):129–139. [PubMed: 19092145]
23. Patel A, MacMahon S, Chalmers J, et al. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med*. 2008; 358(24):2560–2572. [PubMed: 18539916]
24. Association AD. Standards of medical care in diabetes—2014. *Diabetes Care*. 2014; 37(Supplement 1):S14–S80. [PubMed: 24357209]
25. National Committee for Quality Assurance. [November 12, 2013] HEDIS & Performance Measurement. 2013. <http://www.ncqa.org/HEDISQualityMeasurement.aspx>.
26. Hoerger TJ, Segel JE, Gregg EW, Saaddine JB. Is glycemic control improving in US adults? *Diabetes Care*. 2008; 31(1):81–86. [PubMed: 17934153]
27. Lipska KJ, Ross JS, Wang Y, et al. National trends in US hospital admissions for hyperglycemia and hypoglycemia among medicare beneficiaries, 1999 to 2011. *JAMA internal medicine*. 2014
28. Goldberg PA, Bozzo JE, Thomas PG, et al. “Glucometrics”—assessing the quality of inpatient glucose management. *Diabetes technology & therapeutics*. Oct; 2006 8(5):560–569. [PubMed: 17037970]
29. van Walraven C, Austin PC, Jennings A, Quan H, Forster AJ. A modification of the Elixhauser comorbidity measures into a point system for hospital death using administrative data. *Medical care*. Jun; 2009 47(6):626–633. [PubMed: 19433995]
30. Healthcare Cost and Utilization Project. [November 12, 2013] Clinical Classifications Software (CCS) for ICD- 9-CM. 2013. <http://www.hcup-us.ahrq.gov/toolssoftware/ccs/ccs.jsp>.
31. Krinsley JS, Meyfroidt G, van den Berghe G, Egi M, Bellomo R. The impact of premonitory diabetic status on the relationship between the three domains of glycemic control and mortality in critically ill patients. *Current opinion in clinical nutrition and metabolic care*. Mar; 2012 15(2):151–160. [PubMed: 22234163]
32. Van den Berghe G, Wilmer A, Milants I, et al. Intensive insulin therapy in mixed medical/surgical intensive care units: benefit versus harm. *Diabetes*. Nov; 2006 55(11):3151–3159. [PubMed: 17065355]
33. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group. *BMJ*. Sep 12; 1998 317(7160):703–713. [PubMed: 9732337]
34. Patel A, MacMahon S, Chalmers J, et al. Effects of a fixed combination of perindopril and indapamide on macrovascular and microvascular outcomes in patients with type 2 diabetes mellitus (the ADVANCE trial): a randomised controlled trial. *Lancet*. 2007; 370(9590):829–840. [PubMed: 17765963]
35. Collins R, Armitage J, Parish S, Sleight P, Peto R. MRC/BHF heart protection study of cholesterol-lowering with simvastatin in 5963 people with diabetes: a randomised placebo-controlled trial. *Lancet*. 2003; 361(9374):2005–2016. [PubMed: 12814710]
36. Colhoun HM, Betteridge DJ, Durrington PN, et al. Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): multicentre randomised placebo-controlled trial. *Lancet*. 2004; 364(9435):685–696. [PubMed: 15325833]
37. Cleeman J, Grundy S, Becker D, Clark L. Expert panel on detection, evaluation and treatment of high blood cholesterol in adults. Executive summary of the third report of the national cholesterol education program (NCEP) adult treatment panel (atp III). *JAMA*. 2001; 285(19):2486–2497. [PubMed: 11368702]
38. Gregg EW, Li Y, Wang J, et al. Changes in diabetes-related complications in the United States, 1990–2010. *New England Journal of Medicine*. 2014; 370(16):1514–1523. [PubMed: 24738668]
39. Berry C, Tardif JC, Bourassa MG. Coronary heart disease in patients with diabetes: part II: recent advances in coronary revascularization. *J Am Coll Cardiol*. 2007; 49(6):643–656. [PubMed: 17291929]
40. Inzucchi SE, Bergenstal RM, Buse JB, et al. Management of hyperglycemia in type 2 diabetes: a patient-centered approach position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care*. 2012; 35(6):1364–1379. [PubMed: 22517736]

41. Tseng C-L, Soroka O, Maney M, Aron DC, Pogach LM. Assessing potential glycemic overtreatment in persons at hypoglycemic risk. *JAMA Intern Med.* 2013; 174(2):259–268. [PubMed: 24322626]
42. Malmberg K, Norhammar A, Wedel H, Ryden L. Glycometabolic state at admission: important risk marker of mortality in conventionally treated patients with diabetes mellitus and acute myocardial infarction: long-term results from the Diabetes and Insulin-Glucose Infusion in Acute Myocardial Infarction (DIGAMI) study. *Circulation.* May 25; 1999 99(20):2626–2632. [PubMed: 10338454]

Figure 1. Quarterly in-hospital mortality for patients with and without diabetes from 2000 to 2010, stratified by ICU status and by service assignment.

Figure 2. Quarterly inpatient and outpatient glycemic control among ICU patients with diabetes from 2000 to 2010.

Table 1

Demographic characteristics of study sample

Characteristic	Overall, N=322,939 n (%)	Any ICU stay, N=54,646 n (%)	No ICU stay, N=268,293 n (%)	Medical service N=196,325 n (%)	Surgical service N=126,614 n (%)
Died during admission	7,587 (2.3)	5,439 (10.0)	2,147 (0.8)	5,705 (2.9)	1,883 (1.5)
Diabetes	76,758 (23.8)	14,364 (26.3)	62,394 (23.2)	55,453 (28.2)	21,305 (16.8)
Age, mean years (SD)	55.5 (20.0)	61.0 (17.0)	54.4 (21.7)	60.3 (18.9)	48.0 (23.8)
Age full range (interquartile range)	0-118 (42-73)	18-112 (49-75)	0-118 (40-72)	0-118 (47-76)	0-111 (32- 66)
Female	159,227 (49.3)	23,208 (42.5)	134,296 (50.1)	99,805 (50.8)	59,422 (46.9)
White race	226,586 (70.2)	41,982 (76.8)	184,604 (68.8)	132,749 (67.6)	93,838 (74.1)
Insurance					
Medicaid	54,590 (16.9)	7,222 (13.2)	47,378 (17.7)	35,229 (17.9)	19,361 (15.3)
Medicare	141,638 (43.9)	27,458 (50.2)	114,180 (42.6)	100,615 (51.2)	41,023 (32.4)
Commercial	113,013 (35.0)	18,248 (33.4)	94,765 (35.3)	53,510 (27.2)	59,503 (47.0)
Uninsured	13,521 (4.2)	1,688 (3.1)	11,833 (4.4)	6,878 (3.5)	6,643 (5.2)
Length of stay, mean days (SD)	5.4 (9.5)	11.8 (17.8)	4.2 (6.2)	5.46 (10.52)	5.42 (9.75)
Service					
Medicine	184,495 (57.1)	27,190 (49.8)	157,305 (58.6)	184,496 (94.0)	--
Surgery	126,614 (39.2)	25,602 (46.9)	101,012 (37.7)	--	126,614 (100%)
Neurology	11,829 (3.7)	1,853 (3.4)	9,976 (3.7)	11,829 (6.0)	--

ICU = intensive care unit

Table 2

Regression analysis of mortality trends

Independent variables	ICU Patients, N = 54,646 OR (95% CI)	Non-ICU Patients, N = 268,293 OR (95% CI)	Medical Patients, N = 196,325 OR (95% CI)	Surgical Patients, N = 126,614 OR (95% CI)
Model	1	2	3	4
Year	0.974 (0.963, 0.985)	0.925 (0.909, 0.940)	0.943 (0.933, 0.954)	0.995 (0.977, 1.103)
Diabetes	1.352 (1.562, 1.171)	0.958 (0.783, 1.173)	1.186 (1.037, 1.356)	1.213 (0.942, 1.563)
Diabetes*year	0.947 (0.927, 0.967)	0.977 (0.946, 1.008)	0.961 (0.942, 0.980)	0.955 (0.918, 0.994)
<i>c-statistic</i>	0.812	0.907	0.880	0.919

All models control for sex, age at time of admission, race, payer, length of stay in days, principal discharge diagnosis, and Elixhauser comorbidity variables. Models 1 and 2 additionally control for service assignment, whereas models 3 and 4 control for ICU status.

ICU = intensive care unit, OR = odds ratio, CI = confidence interval

Table 3

Regression analysis of mortality trends among intensive care unit patients with diabetes

Sample	Patients with inpatient glucose control measures (n=13,136)		Patients with inpatient and outpatient glucose control measures (n=4,989)		
Model	1	2	3	4	5
Independent variables	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Year	0.927 (0.907, 0.947)	0.926 (0.905, 0.947)	0.958 (0.919, 0.998)	0.956 (0.916, 0.997)	0.953 (0.914, 0.994)
% severe hyperglycemic days		1.016 (1.010, 1.021)		1.009 (0.998, 1.020)	1.010 (0.999, 1.021)
% hypoglycemic days		1.047 (1.040, 1.055)		1.051 (1.037, 1.065)	1.049 (1.036, 1.063)
% normoglycemic days		0.997 (0.994, 1.000)		0.994 (0.989, 0.999)	0.993 (0.988, 0.998)
SD of mean glucose		0.996 (0.992, 1.000)		0.993 (0.986, 1.000)	0.994 (0.987, 1.002)
Mean HbA1c					0.892 (0.828, 0.961)
c-statistic	0.806	0.825	0.825	0.838	0.841

All models control for sex, age at time of admission, race, payer, length of stay in days, principal discharge diagnosis, Elixhauser comorbidity variables, and service assignment. OR = odds ratio, CI = confidence interval, HbA1c = hemoglobin A1c